

A Manual of Poisonous Plants

Chiefly of Eastern North America, with Brief Notes
on Economic and Medicinal Plants,
and Numerous Illustrations

By
L. H. PAMMEL, Ph. D.
Professor of Botany, Iowa State College
Agriculture and Mechanic Arts



UNIVERSITY
OF NEBRASKA
LIBRARY

THE TORCH PRESS
CEDAR RAPIDS, IOWA

1911



Wild Barley on Summit Trail, H. Anderson Laboratory, La. Agr. Expt. Sta.

FOREWORD

During the last decade, there has been much interest manifested in regard to plants injurious to live stock. Numerous contributions have been made along this line, notably by Dr. Chesnut, formerly of the United States Department of Agriculture, Drs. True and Wilcox and their co-workers also of Washington, Dr. Schaffner of the Ohio State University, Dr. Jones of the Vermont Agricultural Experiment Station, Dr. Kennedy of the Nevada Experiment Station, Dr. Nelson of the Wyoming Station, Dr. Nelson of Washington, and Drs. Peters and Bessey of Nebraska. Other station botanists have also contributed to the same line of work. Much of the literature is scattered, however, hence an effort has been made to bring together in the following pages the results obtained.

Much information on this line of investigation has also been obtained from such works as Millspaugh's *Medicinal Plants of North America*, Dr. Johnson's *Manual of the Medical Botany of North America*, Lloyd's *Drugs and Medicines of North America*, Winslow's *Veterinary Materia Medica and Therapeutics*, Sayre's *Organic Materia Medica and Pharmacognosy*, Flückiger and Hanbury's *Pharmacographia*, Greenish's *Materia Medica*, Ellingwood's *Materia Medica, Therapeutics and Pharmacognosy*, Pereira's *Materia Medica*, Luerssen's and Czapek's publications and many others which give details in regard to the effects of poisonous plants.

Many persons may object to the great number of plants which are here regarded as poisonous or described as such in this work. I have placed the broadest interpretation on the subject and have, therefore, included all plants that are injurious although many of these are not known to produce poisons, some even being most useful economic plants and yet injurious to some people.

It has been thought best to arrange the manual so as to consider the plants in the same order as that given in Engler and Prantl's *Die Pflanzenfamilien*.

The Schizomycetes were contributed by my colleague, Dr. R. E. Buchanan, who has also favored me in many other ways.

The parts concerning the blue-green algae and algae, taking up the higher algae and their relation to the water slimes are given in their sequence under the Schizophyceae and Euphyceae.

The Eumycetes or true fungi are considered chiefly from the pathogenic standpoint; while other fungi are referred to and briefly considered under their respective groups. In regard to the higher fungi, such as the toadstools, such valuable information may be obtained from the works of Dr. Farlow, Prof. Peck and Prof. Atkinson.

The so-called Blastomycetic fungi have been arranged under the group of the imperfectly known forms. There are also brief characterizations of other groups of the cryptogams such as lichens, mosses, ferns and their allies.

The flowering plants or Spermatophyta are described more fully than the previous groups, and under the various orders and subdivisions we have also added notes on economic and medicinal plants. Those who desire fuller information in regard to other American species of the different orders described should consult the latest editions of Gray's or Britton's Manuals, in which the descriptions are full and complete. Of course, one cannot expect to add much to the excellent descriptions given in these treatises.

I have freely made use of published literature in the systematic portions as well as in the more technical matter pertaining to poisons, and I wish to acknowledge my assistance from these sources.

In order that the species named may be more readily recognized, a large number have been figured.

In addition to the descriptive part of the work there has been added a chapter on the active principles of plants, by my colleague, Prof. A. A. Bennett. I am also greatly indebted to Miss Harriette S. Kellogg who has carefully read the manuscript and assisted me in other ways besides preparing the bibliography. To Dr. Trelease of the Missouri Botanical Garden, I am indebted for the use of a number of books on the subject. Miss Charlotte M. King, Miss Ada Hayden, Mr. W. S. Dudgeon, and my daughter, Lois, have made drawings especially for this work. Other illustrations are taken from special works, due credit being given in each case.

I am indebted for the use of cuts to the following persons: Dr. C. F. Curtiss of the Iowa Agricultural Experiment Station, Dr. B. D. Halsted of the New Jersey Agricultural Experiment Station, Profs. S. B. Green and Washburn of the Minnesota Station and Prof. A. D. Selby of the Ohio Station, and to the United States Department of Agriculture. Some have been reproduced from Baillon's Dict., and from Bentham's Handbook of British Plants. I have endeavored in each case to give credit for the drawing or cut.

I am indebted to Drs. R. R. Dykstra and C. H. Stange, Profs. L. G. Michael, C. V. Gregory and A. A. Bennett for proof reading and to Dr. W. H. Stuhr for some matter in Part I.

The work does not pretend to be complete; we hope, however, that it may prove useful to the Veterinarian, Physician and Layman.

Ames, Iowa, June 1, 1909.

TABLE OF CONTENTS

PART I

CHAPTER I.	POISONS AND STATISTICS ON POISONS	1
	Ancient Use of Poisons, 2; The Rise of Chemistry and Poisons, 3; Ratsburg on Poisonous Plants, 4; Statistics of Poisoning, 5; Statistics on Poisoning and Actions of Poisons on Different Animals, 6.	
CHAPTER II.	BACTERIAL POISONS	9
	Impure Water the Source of Disease, 9; Bacterial Poisons, 10; Botulism, 11; Malignant or Pellagra, 11.	
CHAPTER III.	DERMATITIS	12
	Dermatomycoasis, 12; Dermatophytes of Domestic Animals, 13; Pityriasis, 17.	
CHAPTER IV.	FORAGE POISONING, ERGOTISM, AND ASPERGILLOSIS	20
	Forage Poisoning, 20; Poisoning from Silage, 24; Ergotism, 28; Aspergilliosis, 29.	
CHAPTER V.	POISONING FROM FUNGI	31
	Toadstools — <i>Amanita muscaria</i> , 31; <i>Amanita phalloides</i> , 31.	
CHAPTER VI.	POISONING FROM OTHER PLANTS. EQUISETARIA, LOCOISM, AND LUPINOSIS	37
	Equisetaria, 37; Locoinism, 37; Lupinosis, 40.	
CHAPTER VII.	DELPHINOSIS, LATHYRISMUS (LATHYRISM), ACONITISM, VERATRISM, UMBELLIFERAE, CONIUM, CICUTA	44
	Delphinosis, 44; Lathyrismus, 45; Aconitism, 46; Veratrum, 46; Poisoning from Umbelliferae, 49; Conium, 49; The Poison Hemlock, 51.	
CHAPTER VIII.	FISH AND ARROW POISONS, HYDROCYANIC POISONING—TOXALBUMINS—BLACK LOCUST, CASTOR OIL, AND JEGURITY	52
	Fish and Arrow Poisons, 52; Hydrocyanic Poisoning, 53; Poisoning from Toxalbumins, Black Locust, Ricinus, and Abrus, 55.	
CHAPTER IX.	POISONING FROM OPIUM, SOLANACEAE AND PLANTS THAT CONTAIN SAPONINS	59
	Opium, 59; Poisoning from Solanaceae; Jimson Weed; Hyoscyamine, 60; Poisoning from Plants that Contain Saponins; Bouncing Betty; Corncockle, 62.	
CHAPTER X.	POISONING FROM FLOWERS, POISONING FROM HONEY, MECHANICAL INJURIES	64
	Poisoning from Flowers, 64; Poisoning from Honey, 64; Mechanical Injuries; Wild Barley; Cheat; Needle Grass; Grass-wood; Crimson Clover, 66.	
CHAPTER XI.	CLASSIFICATION OF POISONS, SYMPTOMS, AND ANTIDOTES	72
	Poisons which Cause Coarse Anatomical Changes of the Organs, 73; Classification of Poisons and Poisonous Symptoms According to Bernard H. Smith — Narcotics, 73; Delirians, 73; Incubians, 74; Convulsives, 74; Depressants, 74; Asthenics, 75; Purgatives, 75; Abortives, 76; Irritants with Nervous Symptoms, 76; Simple Irritants, 76. Table of Symptoms Observed After the Administration of Poisons, Adapted from Castell After the Work of Kober, 77; Treatment for Poisoning, 79.	
CHAPTER XII.	THE PRODUCTION OF POISON IN PLANTS	82
	The Use and Actions of Poisons on Plants, 82; Distribution of Poisonous Substances in Plants, 82; Light, 83; Heat, 83; Seasons, 83; Climatic, 85; Culture, 87; Related Species and Toxic Substances, 88; Distribution of Chemical Substances, 88; Saponins, 89; Hydrocyanic Acid, 89; Cumarin, 90; Cytisin, 90.	

CHAPTER XIII. ALGAE IN WATER SUPPLIES	91
How Growth of Algae in Water May Be Prevented, 94; The Use of Alginates, 94.	
CHAPTER XIV. A CATALOGUE OF THE MORE IMPORTANT POISONOUS PLANTS OF THE UNITED STATES AND CANADA	96
Literature, 96; Euthallophyta, Schizophyta, Schizomycetes, Bacteria, 97; Euphyceae — Algae, 98; Eumycetes — Fungi; Phycomycetes, Black Molds and Downy Mildews, 98; Smuts, 98; Rusts, 99; Toadstools, 99; Ascomycetes, 100; Fungi Imperfecti, 100; Embryophyta, Pteridophyta, 100; Embryophyta siphonogama; Flowering Plants; Gymnospermae, Conifers and Allies, 101; Angiospermae, 102; Monocotyledoneae, 102; Grass Family, 102; Lily Family, 103; Dicotyledoneae, 105; Nettle Family, 106; Pink Family, 107; Crowfoot Family, 108; Poppy Family, 112; Rose Family, 115; Pulse Family, 117; Spurge Family, 121; Poison Ivy Family, 122; Carrot Family, 126; Heath Family, 127; Nightshade Family, 131; Composite Family, 137.	
CHAPTER XV. CHEMISTRY OF ALKALOIDS, GLUCOSIDES, ETC.	143
Composition and General Properties of Alkaloids, 143; Occurrence of the Alkaloids, 144; Classification of the Alkaloids, 145; On Alkaloids, 147; Glucosides, 148; Gluco-Alkaloids, 149; Saponins, 149; Pterotoxin, Clou-toxin and Toxicin, 149; Other Vegetable Poisons, 149; Amino, 149; Organic Acids, 150.	

TABLE OF CONTENTS

PART II

KEY FOR PLANT KINGDOM	153-158
MYXOTHALLOPHYTA	158-160
EUTHALLOPHYTA	160
SCHIZOPHYTA	160
SCHIZOMYCETES	161-184
SCHIZOPHYCEAE	184-188
FLAGELLATAE	188
EUPHYCEAE	188-194
PRIDINALES	188
BACILLARIALES	188
CONJUGATAE	189
CHLOROPHYCEAE	190
CHARALES	193
PHAEOPHYCEAE	194
DICTYOTALES	194
RHODOPHYCEAE	194
EUMYCETES	195-307
PHYCOMYCETES	195-209
ZYCOMYCETES	195-204
OOMYCETES	204-209
BASIDIOMYCETES	209
HEMIBASIDIUM	210-220
EURASIDIUM	220
ASCOMYCETES	247-281
FUNGI IMPERFECTI	281-306
LICHENS	307
EMBRYOPHYTA ZOOIDOGAMA	308-325
BRYOPHYTA	308-312
PYRIDIOPHYTA	312-325
FILICALES	313-322
EQUISETALES	322-325
EMBRYOPHYTA SIPHONOGAMA OF SPERMATOPHYTA	325-302
GYMNOSPERMAE	325-332
CONIFERAE	327
ANGIOSPERMAE	332
MONOCOTYLEDONNAE	332-395
PANDANALES	332
HELORHAE	332
GLUMIFLORAE	336-369

GRAMINEAE	336-367
CYPERACEAE	367-369
PRINCIPES	369-370
SPATHIFLORAE	370-372
FARINOSAE	372-374
LILIIFLORAE	374-390
SCITAMINEAE	390-392
MICROSPERMAE	392-395
DICOTYLEDONEAE	395-802
ARCHICHLAMYDEAE	395-664
VERTICILLATAE	395
PIPERALES	396
SALICALES	396-397
MYRICALES	397-399
BALANOPSIDALES	399
LEYNERIALES	400
JUGLANDALES	400-402
FAGALES	402-404
URTICALES	404-415
PROTEALES	415
SANTALALES	415
ARISTOLOCHIALES	416
POLYGONALES	417-423
CENTROSPERMAE	423-444
RANALES	444-479
RHORADALES	479-496
SARRACENIALES	497-500
ROSALES	500-574
GERANIALES	574-604
SAPINDALES	604-620
RHAMNALES	620-621
MALVALES	621-627
PARIENTALES	627-634
OPUNTIALES	634-637
MYRTIFLORAE	637-645
UMBELLALES	645-664
METACHLAMYDEAE	664-802
ERICALES	664-675
PRIMULALES	675-679
EBENALES	679-683
CONTORTAE	683-697
TURNIFLORAE	698-739
PLANTAGINALES	739-740
RUBIALES	740-748
CAMPANULATAE	748-802
CATALOGUE OF POISONOUS PLANTS	803
BIBLIOGRAPHY OF POISONOUS PLANTS	809-918
INDEX	919

LIST OF PLATES

PART I

Wild Barley or Squirrel Tail	Frontispiece
Anemoneella and Stemless Loco Weed	opposite page 37
Poison Ivy	" " 123
Mountain Laurel	" " 128
Jimson Weed	" " 133

PART II

Honey Colored Armillaria	" " 235
Amanita sp.	" " 236
Deadly Amanita	" " 239
A. Yucca. B. Greasewood and Tetradymia	" " 375
Death Camas	" " 378
Wild Indian Corn; Swamp Hellebore	" " 381
Pennsylvania Smartweed	" " 421
Wild Cherry	" " 515
Lupine in flower	" " 546
Great Basin Lupine	" " 548
Common Milkweed	" " 697
A. Rhododendron; Hydrangea. B. Mountain Laurel,	
Rhododendron	" " 667

ERRATA

On page 4, line 1, for "von Pragg," substitute "van Praag."

On page 26, line 40 should read, "the several investigators were not working," etc.

On page 29, line 14, for "tubercle," substitute "tuberculosis."

On page 89, line 22, for "Muscari," substitute "Muscaria."

On page 102, line 2, for "Swedish juniper," substitute "Savin."

On page 103, line 14, for "*G. gussoneum*," substitute "*H. gussoneum*."

On page 127, line 8, for "*Cornus paniculata* L'Her," substitute "*Cornus Amomum*, Mill, Kinnikinnik."

From page 189 to page 243, the heading "EUTHALLEPHYTA" should read "EUTHALLOPHYTA."

On page 201 omit *Mucor ramosus*, Lindt 1, and description following.

On page 243, lines 29, 33, for "Hand," substitute "Hard."

On page 344, line 13, for "Pamorosa," substitute "Palmarosa."

On page 395, line 31, for "oxogenous," substitute "exogenous."

On page 397, insert "MYRICALES" above Myricaceae, Sweet Gale Family.

On page 450, line 17, for "Dunstan and Anderson," substitute "Dunstan and Henry."

On page 522, Fig. 281 should be inverted.

On page 572, line 4, for "Convincin," substitute "Convicin."

On page 877, No. 186, for "Cooke, N. C.," substitute "Cooke, M. C."

On page 884, No. 329 should read "Gerard, A. W. . . . Wanika, a new," etc.

On page 901, No. 715, for "1901," substitute "191."

Typographical errors in scientific names occurring in the text are corrected in the index.

Part I

A General Treatise on Poisonous Plants

CHAPTER I

POISONS AND STATISTICS ON POISONS

A poison has been defined as "Any substance that, when taken into the system acts in a noxious manner by means not mechanical, tending to cause death or serious detriment to health."

Robert and other physicians define a poison as "A non-organized body, either organic or inorganic, which under certain conditions, affects temporarily or permanently one or more organs of the body, when in a state of health or in a healthy condition." Such poisons may develop in the body or may come from without. Some substances act injuriously in a mechanical way, that is, they may set up disturbances by irritating some parts of the body. Other substances, while poisonous to one who is ill, may be entirely harmless to persons or animals in a state of health.

Robert also defines poisons from a pharmacological standpoint as "All pharmacological agents which, in a given case, do not act beneficially but injuriously."

Toxicology is the science of poisons, the word being derived from the ancient word "tox," meaning bow, or arrow, probably from the ancient use of the arrow to kill.

In tracing the application of the word "tox," "arrow," to its later application, poison, Blyth says "Perchance the savage found that weapons soiled with the blood of former victims made wounds fatal; from this observation, the next step naturally would be that of experiment,—the arrow or spear would be steeped in all manner of offensive pastes, and smeared with the vegetable juices of those plants which were deemed noxious, and as the effects were mysterious they would be ascribed to the supernatural powers, and covered with a veil of superstition."

The different tribes of Indians in South America have from early days been skilful in preparing arrow poisons, the majority of which contain strychnin in some form. The following plants have, at various times, furnished poisons for arrow tips, not only in South America, but also in other countries: *Strychnos toxifera* (Strychnine), perhaps the most generally used of any; *Anhalis toxicaria*, an arrow poison of Java, Borneo, and North Africa; various *Leguminosae*, as *Erythrophloeum* in Angola, Sierra Leone, and Seychelles, a different species being used in each place, *Pithecolobium*, *Afelia*, and *Derris elliptica* of Borneo; of the *Menispermaceae*, two species of *Abuta* are used.

Perhaps in this connection, it would not be out of place to mention several fish-poisons, many of which are also legumes. Of this order are *Albizzia*, *Afelia*, *Bauhinia*, *Enterolobium*, *Leucosia*, *Milletia*, *Puccia*, *Acacia*, *Abrus precatorius*, *Clitoria*, *Mundelia*, *Derris*, *Lonchocarpus*, and *Tephrosia*. In preparing the last named, the leaves are crushed and mixed with quicklime before using. Among the *Menispermaceae* are the Indian Berry, Fish-berry or Levant Nut (*Anamirta paniculata*) of the East Indies which contains picrotoxin; and

Palmyra *arabica*, used also by the Malays to poison crocodiles. Of the *Rhamnaceae*, Kræmer mentions *Tapura*, *Gonima*, and *Zeryphus*. Among the *Tiliaceae*, Kræmer, also mentions species of *Croton* which are used as fish-poisons. The same authority adds species of *Barringtonia* of the order *Lecythidaceae* and *Laportea* *stimulans* of the *Urticaceae* to the above list.

ANCIENT USE OF POISONS

In his work on "Poisons their Effects and Detection," Blyth writes an excellent account of their history from which a few of the following more important data have been taken for the present work.

Their early history is involved in myth. Hecate was said to have been the discoverer of poisonous herbs and her knowledge passed in turn to Medea. The Egyptian kings, Menes and Attalus Philometer, not only had a knowledge of plants but the latter was also familiar with the uses of such plants as hyoscyamus, aconite, conium, and others of similar character. He experimented with poisons and compounded medicines. The Egyptians knew prussic acid, which was extracted from the peach and by means of which those who revealed religious secrets were put to death. The ancient Romans also must have been familiar with this poison, since a Roman knight once took poison and fell dead immediately at the feet of Samolus. The ancient Greeks knew about poisons and it was not considered a dishonorable thing to commit suicide.

Nicander of Colophon (204-138 B. C.) wrote two treatises on poisons, in one of which he described the effect of snake venom, in the other, henbane, aconite, conium, and lungi, were discussed. As antidotes for poisoning from any of these substances, he recommended such remedies as lukewarm oil, in order to excite vomiting.

Dioscorides (40-90 A. D.) divided poisons into (1) Animal poisons, as cantharides, poisonous snakes, the blood of an ox (probably putrid); (2) Poisons from plants as opium, hyoscyamus, conium, aconite (the latter coming from Akron in Heraclea), and colchicum; (3) Mineral poisons like arsenic and mercury (cinnabar).

Pliny mentions that the Gauls dipped their arrows in a preparation of veratrum.

Toffana of Naples sold under the name of Acquetta di Napoli a solution of arsenious acid, by which, it is said, 606 persons were poisoned, among them the popes, Pius III and Clement XIV.

Poisoning was much practiced in India for the purpose of revenge, robbery and suicide, every little quarrel being liable to end in assassination of one of the parties. Such poisons as arsenic, aconite, opium, and extracts derived from plants of the Solanum family, were also used in India to destroy cattle. It is said that gypsies used *Phycomyces nitens*, having knowledge of its properties from the same country. The spores of the fungus were administered in warm water and death, accompanied with all the symptoms of tuberculosis, followed in a few weeks. The Hebrews seem to have been familiar with certain poisons, as arsenic, aconite and, possibly, ergot.

The deaths of Socrates, Demosthenes, Hannibal, and Cleopatra, were due to the administration of poisons.

In the early part of the Christian era, there were many professional poisoners and their business flourished, kings, emperors, popes, and members of the nobility being among their victims. There were two great criminal schools

in Venice between the tenth and seventeenth centuries, the government secretly recognizing the operations of these criminals and paying a sum of money for the execution of prisoners of note. However, these efforts were not often successful. J. Baptista Porta, in the sixteenth century, wrote under the title of "Natura. Magic," a work devoted partly to cookery and partly to poisons and how to use them.

The early methods of detecting the presence of poisons were crude, the surroundings were always noted; the suspected poison was generally administered to an animal and, if it died, poison was sure to be diagnosed without further investigation since the early church forbade postmortem examination. Later, however, doctors were permitted to dissect and thus become familiar with pathological changes.

THE RISE OF CHEMISTRY AND POISONS

At the close of the eighteenth and the beginning of the nineteenth centuries, chemistry had advanced sufficiently to test for arsenic and the more important mineral poisons. Scheele discovered prussic acid, other chemists, as Berthollet, Lavoisier, and Stahl, added to the chemical knowledge of poisons. The father of modern toxicology, however, was Bonaventura Orfila, whose work was published in 1814. Derosne discovered the alkaloids of narcotin and morphin in 1818. Pelletier and Caventou discovered strychnin in 1818. Giesecke discovered conine in 1827, and Geiger and Hesse separated atropin and hyoscyamin in 1833.

The modern aspect of the subject began with treatises on poisons by such workers as Vogel¹ and Richard Mead,² and writings on the subject of chemistry through the works of Stahl, Scheele, Berthollet, Priestly and Lavoisier. Botanists, too, at this time began to be greatly interested in a study of poisonous plants. Thus we have the work of Bulliard³ and the work of Gmelin.⁴ The work of Bulliard discusses a large number of poisonous plants with excellent illustrations, and Gmelin treats quite fully of the then known poisonous plants of Europe.

The works of Gmelin, Bulliard and Plenck⁵ on Toxicology, and Buchner's Toxicology⁶ were frequently quoted by the older writers. The greatest of the older writers, however, was Orfila⁷ whose great work on toxicology became the recognized authority on toxicology. This work was first published in 1814, and passed through many editions. Orfila conducted actual experiments with different plants. This work of Orfila was also translated into different languages,⁸ Orfila was preceded by Podere.⁹ About that period other toxicologies were published in France and Germany, such as those of Sobernheim and Simon¹⁰

¹ The Usefulness of Natural Philosophy. 1654.

² Mechanical Theory of Poisons.

³ Historie des plantes veneneuses et suspectes de la France. Paris 1784. Folio X, 177 p., 72 tab. col.—Ed. II Paris 1798.

⁴ Abhandlung von den giftigen Gewächsen. Ulm. 1775. Allgemeine Geschichte der Pflanzengifte. Nürnberg 1777.

⁵ Toxicologia, seu doctrina de venenis et antidotis. Viennae, Graeffler 1785, 338 p.

⁶ Toxicologia. Nürnberg. 1827.

⁷ Traite des poisons, ou Toxicologie generale. Paris, 1813. Ed. III, ib. 1826. Orfila and Bonaventura. Traite des Poisons ou Toxicologie Generale. Paris, 1814. (Ed. 5) 1852.

⁸ Toxicologia. Seemann & Karle. 2 Vols. Berlin 1829-1831.

⁹ Nedec. leg. Ed. 2.

¹⁰ Handbuch der praktischen Toxicologie. Berlin, 1834. Toxicologie. Nürnberg. 1827, 2nd Ed.

Taylor,⁴ Hoffman,⁵ Th. Husemann & A. Husemann,⁶ van Praag⁷ and Opwyrdt Rebutau,⁸ Seign,⁹ Böhm and von Boeck,¹⁰ Dragendorff,¹¹ Falck,¹² and more recent works by Joshua Nunn,¹³ Smith,¹⁴ Hutyra and March,¹⁵ to say nothing of the recent contributions occurring in the American Veterinary Review, the Journal of the Chemical Society, American Journal of Pharmacy, besides treatises in many chemical and pharmaceutical journals.

The modern work of Blyth, though a somewhat exhaustive treatise on the subject of poisons, is not comprehensive so far as a large number of the poisonous plants are concerned. Many popular treatises on the subject of poisonous plants have appeared in nearly every European language both in ancient and modern times but perhaps no one has contributed more to the subject of poisons than Kobert, who has published several treatises, and one of his works "Practical Toxicology for Physicians and Students" was translated into English by Dr. Fr. Schubert. Such men as Dr. M. Greshoff of Haarlem, published a number of extended treatises on the subject "Poisons, especially Hydrocyanic Acids and Saponins, in Plants." His monograph on fish poisons and subsequent monographs two and three, really survey most of the poisonous plants of the world. Nor should we omit to mention the many treatises by Prof. Power of Wellcome laboratory and his students who have investigated a large number of poisonous plants or the work of Prof. Maiden of New South Wales, or of Cornevin of France.

RATZBURG ON POISONOUS PLANTS

Between 1834 and 1838 there appeared the first part of the work of Brandt and Ratzburg on Phanerogamous poisonous plants of Germany and in the year 1838 in the same work the poisonous Cryptogams by Phoebea. This like other works of the time contained numerous fine colored plates. This work pertaining to the flowering-plants, lists the following plants of Germany as poisonous:

Dandel (*Lolium temulentum*) *Juniperus Sabina*, Yew (*Taxus baccata*) *Arum maculatum* *Colchicum autumnale* *Fritillaria imperialis*, *Narcissus pseudonarcissus*, *Pulsatilla quadrifida*, *Veratrum album*, Aconite (*Aconitum Anthora*, *A. Lycoctonum* & *Commersonii*, *A. oligoneatum*, *A. variable*) *Anemone nemorosa*, *Galium palustre*, *Heliborus niger* II (*foetidus*, *H. viridis*, *Pulsatilla vulgaris*, Crowfoot (*Ranunculus acris*, *R. arvensis*, *R. alpestris*, *R. repens*, *R. bulbosus*, *R. flammula*, *R. Thora*), *Papaver somniferum*, *Euphorbia palustris*, *E. Cyathophorum*, *Rhus Toxicodendron*, *Acthusa Cynapium*, *Cicuta virosa*, *Conium maculatum*, *Coronilla varia* *Oenanthe fatulosa*, *Scum latifolium*, *Ledum palustre*, *Cyclamen europaeum*, *Daphne Mezereum*, *Nerium*, *Oleander*, *Cynanchum*, *Vincetoxicum*, *Atropa Belladonna*, *Datura Stramonium*, *Hyoscyamus niger*, *Mandragora verna*, *Scopolina atropoides*, *Solanum nigrum*, *Digitalis purpurea*, *Gratiola officinalis*.

⁴ Principles and Practice of Medical Jurisprudence. 3 vols. London, 1873.

⁵ Lehmann, Lehrbuch der gerichtlichen Medicin. 5th ed. Wien 1893-91.

⁶ Husemann & Husemann, Handbuch der Toxicologie. Berlin 1892.

⁷ Leerboek voor practische giftleer. In twee Theilen. Utrecht 1871.

⁸ Elements de Toxicologie et de Medecine Legale, appliquees a l'empoisonnement. Paris, 1873. 2nd ed. by H. Bouvier. Paris 1888.

⁹ Studi di Toxicologia Chimica. Bologna, 1871.

¹⁰ Böhm and von Boeck. Handbuch der Intoxicationen. (Bd. 15 of the German edition of Ziemssen's Cyclopaedia.)

¹¹ Die gerichtlich-chemische Ermittlung von Giften in Nahrungsmitteln, Luftgemischen, Speiseresten, Körpertheilen etc. St. Petersburg, 1906. 1st ed. Göttingen, 1888.

¹² Die giftigen Intoxicationen (Handbuch der spec. Pathologie u. Therapie vol. von R. Virchow, Bd. 2). Erlangen, 1894.

¹³ Veterinary Toxicology. Wm. R. Jenkins & Co., N. Y., 1901, 1907.

¹⁴ A Manual of Veterinary Hygiene. 5th ed. Wm. R. Jenkins & Co., N. Y. 1925 pages.

¹⁵ Specielle Pathologie und Therapie der Haustiere.

lis, and *Lactuca virosa*. Many poisonous fungi are enumerated in the second part.

It is interesting here to note their classifications of poisons. Sobernheim classifies the poisons into, A. vegetable poisons, B. nervous poisons, C. blood poisons. Orfila classifies poisons into four classes, (1) irritant, acrid, corrosive, (2) narcotic, (3) narcotic acrid, (4) septic poisons. In 1834 Brandt and Ratzeberg classified poisons as to their origin into (1) mineral, (2) plant, (3) animal. Brandt and Ratzeberg in their treatise on plant poisons make three divisions (1) narcotic (stupefying), (2) acrid (inflammatory), (3) narcotic (inflammatory). They use the classification of Buchner which is as follows - (1) narcotic, *a* HCN Prunus, *b*, volatile narcotic, *Lolium*, *c*, narcotic alkaloidal, Poppy, (2) acrid narcotic, *Cicuta*, *Conium*, *Ruta*, *Digitalis*; (3) irritant narcotic, *a*, Aconite, Oleander, *Rhus*, Smartweed, *b*, more volatile, hot acrid, *Dicra*, Pepper; (4) acrid, *a*, drastic resins, *Bryonia*, *Hypericum*, *Melia*, *b*, drastic coloring matter, *Abrus*, *Spartium*, Pokeweed, *c*, emetic alkaloids, *Iris*, *Colchicum*, *Narcissus*, *d*, unknown poisons, *Agaricus*, *Boletus*, *Phallus*, *Lycoperdon*. Fodere divided poisons into septic, narcotic, narcocracid, acrid, irritant and astringent.

STATISTICS OF POISONING.

The use of poisons for criminal purposes, although not nearly so extensive at the present time as during the middle ages, still plays an important part in criminal law. The following statistics afford some indication of the use of poisons for suicidal and homicidal purposes.

According to the last census of the United States, the number of persons reported as poisoned was as follows -

	By Active Poisons	By Gas ¹
1902.	1374	950
1903.	1551	1715
1904.	1632	2167
1905.	1269	1306
1906.	1734	1276

Intentional cases of poisoning in live stock are not nearly so frequent as are those in the human family, although there are many cases of the former on record. Poisoning of live stock is generally accidental, caused by consuming plants that are poisonous. Large losses occur annually in this way. In 1900 Prof. Chesnut and Dr. Wilcox investigated the conditions in Montana relative to this subject and published the following statistics resulting from their studies. They state that probably not more than one fourth of the actual cases occurring came under their observation.

¹ These numbers include deaths during, or as a result of conflagrations. The annual average was 1412 from active poisons and 1365 from gas. From 1900-1904, the number of deaths by poison averaged 4.5 per 100,000.

Poisoning cases among cattle, horses, and sheep in Montana observed during the season of 1900

	SHEEP		CATTLE		HORSES	
	Poisoned	Died	Poisoned	Died	Poisoned	Died
<i>Zygadenus venenosus</i>	3030	636				
<i>Zygadenus elegans</i>	40	15	6	2	4	3
Lupine	3600	1900			4	3
<i>Delphinium bicolor</i>			2	2		
<i>Delphinium glaucum</i>			100	56		
<i>Cicuta occidentalis</i>	105	80	36	30		
Loco weeds	3550	200	3		150	3
Total	9775	3331	147	90	154	6

Nearly every veterinarian has frequent calls to attend cases of poisoning from obscure causes. These can often be traced to plants that occur in the pasture or feed lot.

Accidental cases of poisoning from such wild plants as jimson weed and others are frequent in the United States, several cases occurring annually from cowbane in Iowa. Statistics in regard to such cases are, however, difficult to obtain.

H. W. Cattell, as senior coroner physician in Philadelphia, performed 799 postmortems, in 155 of which, death was due to poisoning. The poisons used were listed as follows: aconite, 1, ammonia, 1; arsenic, 5, carbolic acid, 10; chloroform, 1; creosote, 1, cyanide of potassium, 1, hydrocyanic acid, 2; illuminating gas, 12, lead, 1, oil of turpentine, 1, opium, 11, oxalic acid, 1, phosphorus, 1, silver nitrate, 1, stramonium, 1, strychnin, 2, sulphuric acid, 1.

In his work on poisons, Blyth states that the deaths from poisons in England and Wales during the ten years ending December, 1903, were 11,035. Deaths from laudanum were 1,505, cocaine, 12, atropia, 96 prussic acid and oil of almonds, 328, potassic cyanide, 207, strychnin and nux vomica, 244, aconite, 45; alcohol, 87, petroleum, 23, belladonna, 95; cocaine, 12.

STATUTES ON POISONING AND ACTION OF POISON ON DIFFERENT ANIMALS.

The statutes do not as a general rule define poisons, but in most of the codes the sale of certain poisonous substances is regulated by law. The following extract from the Iowa code illustrates this. Sale regulated of substances under Schedule A. Arsenic and its preparations, corrosive sublimate, white precipitate, red precipitate, biniodide of mercury, cyanide of potassium, hydrocyanic acid, strychnia and other poisonous vegetable alkaloids and their salts, essential oil of bitter almonds, opium and its preparations, except paregoric and other preparations of opium containing less than two grains to the ounce. Schedule B. Aconite, belladonna, colchicum, conium, nux vomica, henbane, safin, ergot, cotton root, cantharides, creosote, digitalis, and the pharmaceutical preparations, croton oil, chloroform, chloral hydrate, sulphate of zinc, mineral acids, carbolic acid, and oxalic acid.

Not all poisons act in the same way, some acting more quickly than others. Quality and quantity are prime factors in the results obtained. As an illustration of this fact, we may mention ricin which is obtained from the cas-

for oil bean, one gram of which, if properly diluted, is estimated as sufficient to cause the deaths of 1,500,000 guinea-pigs. The characteristics of the animal affected by the poison is also an important factor in the result. For instance, a fatal dose of strychnine in case of ruminants, when given by mouth is varying, when given hypodermically, it is a little larger than for horses; the minimum fatal dose for a horse being $1\frac{1}{2}$ to 3 grams when given hypodermically, and $\frac{3}{5}$ grams (or $\frac{1}{2}$ ounce of raw yucca) when given by mouth, but as much as 2 grams is permissible.

The snail is said to be capable of withstanding more strychnine than an adult man. The minimum dose for man is $\frac{1}{2}$ a gram, while $\frac{4}{7}$ grains constitute a lethal quantity.

Cardiac poisons produce no action upon insects. The rabbit can stand more morphine than a man. Robert says "*Amygdalin* does not affect dogs, but it kills rabbits. The hedgehog takes, with apparent enjoyment, a dose of *contheridex* that would kill several persons under excruciating pains. The bite of the most venomous snake does not harm him, he can even accommodate me in considerable quantity of *hydroxyonic acid*. Whereas the frog is extraordinarily susceptible to the *digitalis* poisons, they have no effect on the toad." "Poisons act more powerfully when absorbed from the subcutaneous connective tissue than when administered internally, with the following exceptions. The neutral *crotonalglycerol* which is found in large quantities in the fresh seeds of *Croton Tiglium*, but which is often lacking in commercial croton oil, is inactive when introduced under the skin. It possesses, however, terrific action when taken into the stomach. *Myronic acid* of mustard as an alkaline salt has no effect when it is injected under the skin. It has, on the other hand, a strong action when taken *per os* by herbivora, the same is true of *amygdalin*."

"In all three of the foregoing cases, the apparent exception to the rule is explained by the fact that the substance, in itself not poisonous, is split up in the intestinal tract, giving off, amongst others, a toxic substance. In the instance first mentioned, *crotonallic acid* is the poison thus freed, in the second, *etherical mustard oil*, and in the third, *hydroxyonic acid*. Some substances, such as *salts of manganese, iron, tungsten*, have no poisonous action when introduced into the intestinal tract, because under these conditions only very minute quantities are absorbed, others are rendered inert because they are excreted almost as quickly as they are taken up, curare being an example, and yet others, such as *snake poison, spider poison, quinine acid, sapotoxin, ergolic acid*, are converted into non-poisonous substances within the intestines."

Persons may become accustomed gradually to the use of poisons. Thus individuals who consume opium or its alkaloids may take large doses without apparent injury, although children are particularly susceptible. The former statement is equally true of those who daily use such poisons as hashish, nicotine, caffeine, cocaine, alcohol, or morphine. They must have the drug in order to keep up their condition. Many people exhibit idiosyncrasies with reference to food substances or drugs. Some people cannot inhale the odor of morphine, turpentine, or tobacco without becoming ill. Others are uncomfortably affected if the flowers of the common bird cherry or the haw are left in the room. Others become sick when in the presence of the flowers of the tuberose. Coming in contact with the castor-oil plant sometimes causes illness.

CHAPTER II

BACTERIAL POISONS

Impure Water In all ages great stress has been laid upon the value of the source a pure water supply. In ancient times, wherever there were of Disease, great centers of population, a large amount of labor as well as of money was employed to furnish water. Of this the Claudian aqueduct, built in Rome in the year 50 A. D., is an illustration. Prof W. P. Mason says "Not only was a generous daily per capita allowance sought for, but we note in the centuries gone by unmistakable evidences of a keen appreciation of the dangers lurking in a polluted supply; and upon this point many of the ignorant consumers of our day and generation would be benefited did they consult the wisdom of the past."

Of the value placed by the ancients upon the quality of water Prof Mason also says "In ancient times, the valleys of the Euphrates and Tigris, now almost a desert, were densely populated. Four thousand years ago the rulers of Assyria had converted those sterile plains and valleys into gardens of extreme productiveness by the construction of immense artificial lakes for the conservation of the flood-waters of the rivers, and as great distributing canals for irrigation. One of these canals, supplied by the Tigris, was over 400 miles long and from 200-400 feet broad, with sufficient depth for the navigation of the vessels of that time." "In India, tanks, reservoirs, and irrigating canals were constructed many centuries before the Christian era, and a great part of that country was kept in the highest state of cultivation. Some of the tanks or artificial lakes covered many square miles, and were often fifty feet in depth.

A great deal of interest has been manifested recently in all parts of the United States concerning water supplies. This has not been confined to the cities but the interest is manifested in the villages and rural districts as well. We are now demanding more than ever before, not only that a good wholesome supply of water be provided to the citizens of a city or village, but also that as good a supply be furnished the farmer. That such diseases as typhoid and cholera are water borne can not be doubted. Many others, as anthrax, hog cholera, and tuberculosis may also be conveyed by water. Animal parasites are also water borne. In addition to these, there are some poorly defined intestinal disorders that are caused by poor water.

That typhoid fever is quite as prevalent in the country as in the city admits no denial. A record of the cases of typhoid occurring during the fall and winter in any of our rural communities shows that the disease is as widely spread in the country as in the city.

A certain class of animal diseases is produced not by the invasion of micro-organisms, but is caused by the water supply being contaminated by the decomposing products of animals. The water may, for example, be highly charged with colon bacilli or other bacteria that produce poisonous products.

Several years ago Dr. Stalker traced an epidemic of horses, cattle, and pigs, all of which had been affected with similar symptoms, the animals uniformly dying after an illness of about two days. The disease was not contagious, the farm buildings were fairly comfortable and clean, and the trouble was, evidently, not due to the food consumed by the animals on the farm. Most of them, however, had been in the habit of drinking from a small creek which ran through the premises. The stream was supplied by a series of springs, and in ordinary seasons flowed for a portion of its course over a gravelly bed. This season the rainfall was light, and it so reduced the supply of water that it ceased to flow. Investigation made on these premises and on the adjoining farms indicated that dead animals were thrown down the steep bluffs into the bed of the stream. During the summer, chickens which had died from cholera and hogs dead from hog cholera had been dumped into the creek. In addition, the creek received the drainage from manure heaps. This was the kind of water that these animals had had to drink. Stock which did not have access to the creek but were watered from a well escaped the disease, while stock on other farms having access to the creek water suffered from the disease.

Dr. Lewis and Mr. Nicholson, in Bulletin 66 of the Oklahoma Agricultural Station, refer to certain troubles of live stock due to faecal contamination. In many cases the pond from which stock is watered is situated where plowed debris is carried into it by heavy rains, partially or completely filling it up, while the stock tramping down the banks soon complete the process. Stock standing in the pond also foul the water with excrement, and in hot weather, when the water is low, such a pond certainly can not afford a very satisfactory water supply. During the winter and spring months, when the rainfall is abundant, this condition is not so noticeable since the water is being continually changed by fresh water running in.

One of the dangers that follow allowing stock of all kinds to stand in a pond is that when the water is at a low stage, and foul, as it becomes in summer seasons, the cattle will not drink a quantity of the hot, foul, surface water, sufficient to prevent certain derangements of the digestive system such as impaction, "dry murrain," and other conditions that are usually ascribed to dry feed, but which are, in a large measure, brought about by insufficient water.

A type of injury resulting from the use of polluted water is illustrated in volume 19, page 74, of the "Journal of Comparative Pathology and Therapeutics." This record is in the form of evidence given in a case in which the plaintiff is the tenant of a farm on which is kept a dairy herd of from 30-35 cows. In 1903, there was no complaint but in 1904 the cows were put to grass in the middle of May and their condition became unsatisfactory at the end of July. Early in September, one of the cows aborted, six others lost their calves between that date and the 7th of October. On the 19th of November, another cow aborted; seventeen of the remaining ones carried their calves the full term and four were barren. The cows drank water from a small lake about one and a half acres in extent, which the town council of Maybole, who were the defendants in the action, used for the deposit of rubbish from the town. About the 18th of October, the cows were removed to another pasture with different water supply, and only one cow slipped her calf. (This occurred Nov. 19). The expert testimony was very conflicting. The plaintiff and expert

testimony held that septic poisoning resulted from the use of this water and thus caused abortion, Prof. Williams maintaining that water holding a large amount of vegetable matter is dangerous to pregnant cows, while the defendants held that this would not be a sufficient cause for the action. Judgment was rendered for the plaintiff.

Bacterial poisons are produced by two classes of bac-
Bacterial Poisons. *teria* the first includes such as are parasitic or pathogenic; the second, those which form poisonous products by the breaking down of dead animal or plant tissues. An illustration of the first class is seen in the products resulting from the tetanus bacillus and diphtheria bacillus which produce an extra-cellular toxin. Another type is the toxin known as endo-toxin. In the extra-cellular form, toxin exudes through the bacterial cell-wall and is found in the body; while in the endo-toxin form, the toxin remains wholly, or in part, in the cell during the life of the organism and is liberated only on the death of the bacteria.

There are on record numerous cases of poisoning as a result of eating certain foods of animal origin, and the same statement may be made in regard to foods of plant origin. Such foods as meats, fish, cheese, and milk, sometimes become injurious because of the products of bacterial growth which they contain. These products are classified as either ptomaines or toxins; a third class, the leucomains, result from the breaking down of tissues of the living animal body, being proteid bodies which have been broken down by enzymes, secreted by the cells of the body. These leucomains produce auto-intoxication. Ptomaines are soluble, basic substances formed by the action of bacteria on protein material. Dr Holland illustrates the action as follows: "The amino-acids, ornithin and lysin, constituents of pure protein, subjected to bacterial action, split off CO_2 , and change to putrescin and cadaverin." Some of these products, as methylamin, are harmless, while others are active poisons. The ptomaines are strongly basic, combining with acids to form salts. They are precipitated with chlorides of mercury and are of various kinds. Some are free from oxygen, while others contain that element, some, as typhotoxin, tetanin, pyocyanin, are unclassified, several are injurious in foods; some are produced in fresh oysters and muscles.

Ptomain poisoning. Symptoms. Gastro-enteritis is the most prominent symptom, with depression and nervous disturbances. In most cases, there are, also, marked thirst, salivation, nausea, and vomiting, diarrhoea, cramps in the legs, great prostration, feeble pulse, dilated pupils, delirium, paralysis, and collapse. The postmortem examination generally, but not always, shows inflammation of the stomach and bowels. Cholin, in large doses, nerval, diamin, amanitin, muscarin, all act as poisons; nerval is much more powerful than cholin, the symptoms being those accompanying obstruction of the bowels, together with nausea, pain, and depression, the diamins are all actively poisonous, dilated pupils, convulsions, diarrhoea, and paralysis being prominent symptoms. Muscarin, found in Fly Agaric and certain putrid products, is a much more powerful poison than cholin or nerval and produces vomiting, griping pains in the stomach and intestines, slow pulse, arrested action of the heart, contraction of the pupils and fatal collapse.

Toxins. These are poisonous bases produced by living bacteria or by saprophytic bacteria in the animal body and in higher plants.

Holland arranges the food toxins in two classes (1) The poisonous

products of specific bacteria, not putrefactive, growing in meat after slaughter, (2) Products of specific bacteria infecting the tissues of food-animals before slaughter. To these should be added (3) Toxins of higher plants (*phyto-toxins*), in which class would be placed *ricin*, a product of the castor-oil plant, *abrin* of the Jequirity plant, *crotin* of the croton, and *robin* from the black locust.

This belongs to the first group and is caused by the *Bacillus Botulinum*. *botulinus*, which contaminates ham, sausage, and fish, and is so poisonous that it frequently is the cause of death in the person using the affected meat or fish. Rabbits, guinea pigs, and cats are very sensitive to the poison and die when given the fluid culture, rabbits succumbing in 36-48 hours after injection of 0.0003-0.001 c c. *Proteus vulgaris*, growing in pork and beef sausage produces a similar poisoning.

Symptoms. Epigastric discomfort, belching nausea, vomiting, gripes, diarrhoea followed by constipation "nervous symptoms appear in a few days, these are dilated pupils, paralysis of the tongue and pharynx, loss of voice, death may follow delirium and coma, or, after some time, recovery may take place.

Poisoning from this cause manifests itself within a half hour occasionally, generally within twenty-four hours, although it may be delayed a week.

The toxins of the second class come from pathogenic bacteria and are due to their growth in living animals. The meat and milk from animals that have had septicæmia or pyæmia may cause such symptoms as headache, vomiting, profuse diarrhoea, gripes, chills and fever, the usual symptoms of these diseases.

In some of the European countries, and occasionally in the United States, there occurs a disease known as **Malignus or Pellagrus** or **Pellagra**. It is common in Northern Italy and Bessarabia and appears at times in France, Portugal, Spain, Roumania, and Wallachia.

Etiology. The disease results from the exclusive use of bread made from spoiled maize or Indian corn. It is caused by an unknown toxic substance, partly basic in its nature, called pellagrogen. When maize is kept in a moist place, this poison is likely to develop, probably through the presence of organisms, possibly bacteria or moulds.

Symptoms. The disease produces debility, reddening and swelling of the skin, an antipathy to food, profuse diarrhoea, delirium and raving mania, staxic walk, spasms and contractions, debility of muscles, and paraplegia. Suicide is not infrequent. In chronic cases, the central nervous system, the skin and digestive tract become the seat of this disease.

Postmortem. The intestines show multiple small ulcers and extensive catarrh. The posterior column and motor area of the lateral tract of the spinal cord show a diseased condition.

Treatment. Change of diet. Use wheat or rye bread or if maize is used it should be thoroughly dried.

A large percentage of cases prove fatal.

CHAPTER III

DERMATITIS

Skin diseases are produced by a variety of causes, some resulting from pathogenic organisms and others through the ingestion of food. On account of these differences, we have two classes sometimes given parasitic skin diseases and urticarial diseases. Under the second class are placed such eruptions as those produced by buckwheat and smartweed, known technically as *fagopyrismus* and *rhus venenosa* (dermatitis), and urticaria, the latter being produced by a large number of plants, especially the nettles. Or there may be internal causes due to innervation of vaso-motors. Of the parasitic skin diseases known under the general name of *dermatomycoses*, we have several types, the so-called *Tinea tonsurans* and the *Favus* organisms, the former occurring in cattle, dogs horses, sheep, swine, and poultry.




Fig. 1. Poison Ivy (*Rhus Toxicodendron*)
cause of dermatitis or Rhus poisoning (L. S.
Dept. Agr.)

This term is derived from two Greek words *Dermat-* meaning skin and *omycosis*, meaning fungus. The classification of the fungi concerned is not at all satisfactory, at present, however, they are generally included in the groups known as *Fungi Imperfecti*, the *Mucoraceae* and *Ascomycetes*.

The *Fungi Imperfecti* include a large group of fungi whose life history has not been worked out completely. The fungi of this class are form-genera, such as the *favus* fungus. The so-called *Achorion* and *Trichophyton* of various authors represent such form genera, the *Odium abicans* being another type, of these, some, perhaps, never produce any other kind of spore than the one commonly seen.

Many of these genera undoubtedly belong to the *Ascomycetes*, in which the spores are produced in little sacs called *asci*, the spores being known as *ascospores*. A kind of ringworm of the dog (*Eidamella spinosa*) belongs to

this group, also the forms that produce aspergillosis, one type of which occurs in the ear.

The *Mucoraceae* produce an unsegmented mycelium and septa only where the reproductive bodies are formed, the spores usually occur in sporangia, or occasionally small spores may be found in the mycelium, zygospores which result from fertilization also occur in some species. Several species produce surface lesions.

According to Neumann, the dermatophytes of domestic animals belong to six distinct genera of fungi as follows: *Trichophyton*, *Endamella*, *Microsporon*, *Achorion*, *Lophophyton*, and *Oospora*. These genera are not, however, all accepted by botanists.

Trichophyton was established by Malmsten in 1848 and is characterized by having a mycelium consisting of simple or dichotomously branched filaments and producing spores from 4-9 μ long. Sabouraud, however, divides the genus into several species depending on the position of the fungus with regard to the invaded hair. The *T. endothrix* lives inside the hair, the *T. ectothrix* develops outside of the hair, forming a sheath around it, and the *T. endo-ecto-thrix* develops both inside and outside the hair. This classification is scarcely tenable from either clinical or cultural characters. We have placed this genus with *Sporotrichum*.

The *Endamella spinosa* described by Matruchot and Dasseville in 1901, has a much branched mycelium. 15 μ in diameter, divided into short segments and splitting into somewhat squarish oval bodies, it is found on the dog.

The *Microsporon* discovered by Gruby in 1843, has a branched mycelium, the latter branches bearing conidia from 2-3 μ in diameter. This fungus has also been placed with *Sporotrichum*.

The *Lophophyton*, described by Matruchot and Dasseville in 1899, produces a mycelium with some tortuous filaments, others short curved, with thick curved walls, no spores produced, it occurs on fowls and is also referred to *Sporotrichum*.

The *Achorion* was described by Remak in 1833. The filaments of the mycelium are from 2-3 μ in diameter, flexuous or straight, variously branched; finally breaking up into spores. This fungus has been placed with the genus *Oospora*.

Oospora was described by Wallroth in 1833. Its mycelial threads are 2-3 μ in diameter, arranged in irregular chains.

The best expert account of dermatomycosis so far as it affects lower animals will be found in the treatise by Neumann¹ who includes also an excellent bibliography on the subject. Hutyrka and Marek have a German text which devotes considerable space to the subject. The work by Hyde and Montgomery treats the subject from a human standpoint. The work by Plaut is also an exhaustive treatise.

Etiology. The cause has been ascribed to various fungi which will be described later in the present work. The predisposing causes are uncleanness, weakness of animals (those that are worn out may offer a favorable medium for the attacks of the fungus). In the case of cows, Fleming observes that the disease is common in the winter when the stables are dirty, and disappears in the spring when the animals are turned out to

¹ "A Treatise on the Parasites and Parasitic Diseases of the Domesticated Animals" English Translation by George Fleming. Second Edition Revised and Edited by James MacQueen. 697, 365 p. 1907. New York.

pasture, the new condition being opposed to contagion. In the case of the cat, early age seems to be an essential condition. In the case of the rabbit a similar condition seems to hold true. In the dog, early age is not an especially predisposing influence but inoculation is successful in young dogs only according to Fiorani, so far as trichophyton is concerned. This latter statement does not hold true for favus.

Contagion. In ringworm of the horse, infection may occur from horse to horse. Megnin states that in one locality 200 horses became infected in this way, a saddle from an infected horse having carried the disease to other animals. In each case, the disease occurred on the side of the back. It has also been transmitted from an ox to a horse. The tinea tonsurans has been transmitted from the horse to calves and the tinea of the horse from horse to dogs, sheep, and pigs, and even to man. Neumann says: "The infection of man is exceptional when the frequency of tinea tonsurans in the horse is considered, as there is scarcely a regiment in which it is not a ways on some young horse." Grooming is the usual way in which the infection is carried and rubbing facilitates inoculation. The ease with which infection occurs in man depends on the character of the fungus, some forms adapting themselves to the conditions present more readily than others. In the case of the boy's nest the contagion may be direct. The virus may be preserved a long time in parts of stables where calves were affected with the tinea tonsurans. The infection spreads less readily to sheep and pigs but may be transmitted from hares to man. In the case of the dog, it is transmitted from dog to dog, from rats and mice to the dog and, occasionally, from dogs to man. In cats, favus is largely transmitted from mice and it is certain that this form can be transmitted to man. In general, it may be said that the transmission of favus from the rat or mouse is frequently brought about through the domestic cat. The tinea of the fowl is transmitted by contact with a diseased fowl. The favus fungus of the fowl cannot be inoculated on the rat or dog but when inoculated in man, it produces lesions similar to favus. Man may be inoculated very easily by handling a fowl on which large erythematous patches occur. Similar patches have occurred in man when inoculation from a fowl was very probable.

Symptoms. Two forms of skin dermatomycosis in the horse have been recognized: (1) called microsporesis, and (2) trichophytosis. The more important symptoms of the first as given by Neumann are: "It appears in patches which are more especially seated on the upper part of the body—on the shoulders, back, loins, croup, sides, and flanks. These patches may, however, be met with on any part of the body, though they are rare on the lower parts of the legs. What are first noticed are the circular patches, the diameter of which is generally about that of a shilling, they are distinguished from the healthy skin by the dullness and erectness of the hairs covering them. Some time before the circular patches appear a very small tuft of hairs—probably from half a dozen to a dozen—may be seen slightly, but markedly, raised in the form of a fine pencil, and feeling as if they had a somewhat hard base, or were matted together at the bottom, when the finger is passed over them. These tufts may be several in number and are usually best seen in hindquarters at the very commencement of the disease, or in the vicinity of the patches, of which they are the initial symptom. The hairs fall off in a

few days, and this is often the first symptom that attracts attention. The epidermis of the patch falls off at the same time as the hairs, it appears to be softened, and the surface of the skin has then a dark-grey tint and is slightly moist, which might be attributed to the rupture of vesicles, though their presence has never been demonstrated. It cannot, therefore, be said, as Ralliet remarks, that the disease presents itself in the form of *herpes* as what is so called in human pathology includes a phase marked by the appearance of vesicles. The humidity of the patch is ephemeral. Its surface generally soon dries, and is covered with epidermic scales of varying thickness, which are agglutinated into flat crusts that are shed and renewed incessantly. These crusts have—more frequently than in the ox—a shining appearance and a grey or yellowish color like flax. At the same time, the lesion progresses by peripheral extension until it attains the diameter of a five-shilling piece or more, and on each zone invaded, successive symptoms are observed. Pruritus is nearly absent in Microsporosis, and is scarcely even shown to exist, except by movements indicating satisfaction on the part of the animal when the patches are gently scratched.

"2. *Trichophytosis* (*Sporotrichum*) A. — *Trichophyton flavum* — The lesions consist of large patches, at least 8 to 10 cm. broad, of a more or less regular form, greyish, and quite smooth. The hairs, raised and matted at their base by a greyish-yellow crust, fall away very rapidly with the crust. The naked surface is not prominent, and shows no trace of suppurative folliculitis (Bodin).

"B. *Trichophyton equinum* occurs usually in numerous patches, some isolated, scattered over the croup and shoulders, and attaining at the most 3 cm. in diameter. At first they can be detected only by touch, but later they become visible by the flattening of the hair. The least traction or slight friction carries off a scaly crust which brings away the diseased hairs. The skin then appears smooth, moist, pinkish, or light grey. Very soon the patch becomes dry, scaly, or powdery, and at its base a slaty grey. The lesions spread by the falling out of the marginal hairs (Matreuchot and Dassonville).

"C. *Trichophyton verrucosum*, var. *equi*, occurs in numerous patches which average 5-6 cm. in diameter, and are localized on the shoulders, neck, withers, flanks, in fact, everywhere where the harness touches. By their confluence, these patches may produce large, irregular lesions. At first the hairs are raised, not broken, and are matted together at the base by a soft grey crust. This crust falls off in less than a week, carrying away the hairs, and leaving an absolutely bald, grey surface covered more or less with dry, greyish scales, and without follicular suppuration. In young horses, however, the shedding of the crust leaves a vesicle or pustule, then the surface of the patch is slightly raised, red, and indurated (Bodin).

"D. *Trichophyton verrucosum*, var. *anis* — The lesions are the same as in the preceding *Trichophytosis* of the Horse, but are generally confined to the neck, head, and ears (Bodin).

"E. *Trichophyton mentagrophytes* — This *Trichophytosis* occurs usually on the nostrils or head. It forms patches up to 5 or 6 cm. in diameter, which may be mistaken for pustules of horse pox undergoing regression. Over the whole of these patches the hairs are matted together at their base by a soft, brownish crust of unequal thickness. Slight traction on the hair brings away the crust, exposing a bare, slightly raised surface, which is red, inflamed, and pitted with

small depressions, grey at their base. These result from the opening of the pustules, of which some may be found at the margin of the patch. The hairs are not broken, but shed, and the condition is, in fact a suppurating folliculitis."

The symptoms observed in the bovine, as described by Neumann, are as follows. "The commencement is manifested by a slightly salient ring, on the surface of which the hairs are erect. An active proliferation of the epidermis causes the rapid formation of scales more or less adherent to each other, and crusts of 2 mm. to 7 mm. thick—hence; the name *dartre croutense* was given to the affection by the older (French) veterinarians. According to Gerlach the crusts are thicker on dark skins, on which they have a greyish-white, fibrous appearance, resembling the amaranthus (*porrigo asbestinea*) on white skins, which are usually finer, the crust is thinner and a little yellow in color."

Diagnosis. The diagnosis based on clinical symptoms should be verified by microscopical examinations. It is best to take material from the younger and deeper parts of the crust which may be moistened with water, or a better examination can be made if it is boiled with a 40 per cent solution of potash after which the particles can be dissected and the fungus threads and spores made out. The different forms cannot readily be distinguished except by cultural methods. The organism grows readily on nutrient media that are neutral or with slightly alkaline reaction. Sabouraud recommended the following

Pure glycerine, glucose, lactose, or maltose	4	grammes
Granulated potash	1	gramme
Distilled water.	100	grammes
Gelose	1.50	grammes

Solid media like potato, agar, and peptonized bouillon are favorable media while liquid media are less favorable. Growth may occur at 15°C, the optimum being 30°C.

Prognosis. The duration of the disease depends on circumstances, it gradually diminishes and may disappear without medical aid. The average length of time of the disease is 40-50 days. Cleanliness has much to do with its disappearance. It lasts longer in thick coated animals than in those with thin coats.

Treatment. Cleanliness and sanitary surroundings, disinfection of stables, careful and regular grooming (all articles used in this process having been thoroughly disinfected, especially when they have previously been used on a diseased animal), avoiding any substance that causes irritation. The following preparations have been used with success. Mercury 1/500, carbolyzed glycerine, alcohol, an ointment composed of 1 part of carbolic acid, hard and soft soaps, each 20 parts. Fournie and La Caive recommend pure carbolic acid, tincture of iodine, and chloral hydrate in equal parts. The applications should be made once or twice a day or every other day depending upon the irritant properties of the preparations used.

For bovines, the remedies named above will prove efficacious, for the dog, application of an ointment prepared from 1/5 per cent of nitrate of silver, is satisfactory.

Dr. Stuhr has contributed the following account of favus in animals. "Favus is a contagious, vegetable parasitic disease of the skin, characterized by the formation of more or less circular, cup-shaped crusts, varying in size from very small up to that of a dime. It has been observed in almost every species of domestic animals. The disease is quite rare among horses and cattle although dogs and cats are frequent sufferers. The domestic fowl and pigeon are also susceptible.

Of the laboratory animals, mice, rats, rabbits, and guinea pigs harbor the




Fig. 2. Favus and Herpes Fungus. A. Spores germinating grown in gelatine. B. Hypha breaking into segments. C. Formation of spores in chains (a) formation of buds (b) chlamydospores. D. Herpes, threads of the mycelium and formation of apores. E. *Odium favus* spore with germ tube. After Crawitz.

disease. Young age, thin skin, and debility predispose. Favus is communicable from lower animals to man and vice versa. Man frequently contracts the disease from cats, the latter becoming infected from eating mice and rats.

Etiology. Favus is caused by a vegetable parasite, (*Oospora porriginis*) which invades the cutaneous structures, especially the epidermal portion. The *Achorion Schoenleinii* was first discovered by Schoenlein in 1839,

although Remak was the first to demonstrate its pathogenic character by direct inoculation. It consists of mycelium and spores, existing in such profusion that it is readily detected. Skin abrasions are an important accessory cause.

Symptoms. The disease is characterized by dry scabs, brownish or yellowish, gray or silver white on the surface, and white or sulphur yellow in their deep layer. These scabs have a circular form, with a diameter not to exceed that of a dime, and a thickness varying up to one fifth of an inch. They occasion atrophy of the hair and a slight depression of the skin. These scabs are usually found on the forehead, cheeks, ears, face of women, external side of the hind legs and in the neighborhood of the claws. At first the scabs are perforated by hairs which soon fall out. Later the skin exfoliates under the scab and leaves a pit. In the horse the scabs may become confluent and form bands as wide as the finger. Itching is observed in the dog. In most cases the progress of the disease is quite rapid although the prognosis is favorable unless the disease has become too far advanced.

Lesions. According to Robinson the parasite first obtains a lodgment in the funnel shaped depression in the epidermis, through which the hair shaft emerges upon the surface. It grows luxuriantly in the upper part of the hair sac and innervates itself on all sides between the superficial layers of the epidermis. When it reaches a short distance on all sides of the hair to one it breaks up the lower layers and appears on the surface producing the characteristic cup-shaped bodies. It also invades the hair shaft itself penetrates between the cellular layers of the root sheath and by its mechanical pressure upon the papillae interferes with the nutrition of the hair and causes it to fall out. If the pressure is sufficient to cause atrophy of the papilla, a new growth does not occur.

In the skin the parasite usually confines itself to the upper corneous cells and does not extend to the living tissues. In cases where the surface is covered by irregular confluent masses of the parasites, the entire upper layer of the epidermis will be found infiltrated with the achorion.

The corium itself is usually in a state of chronic inflammation, and suppuration, which may be quite abundant, often occurs under crusts. Even in the absence of pus, the pressure of the parasite causes atrophy of the skin, and at last pit-like depressions or more extensive reddened scars are left. The disease ends with the destruction of the glandular structures of the skin.

Treatment. This is purely local except when debility complicates the disease, in which instance tonics should be administered to build up the system. As for the local treatment, its aim is to destroy the parasite and relieve the cutaneous irritation. The dry scabs may be softened and removed by thoroughly washing with soft soap and water. The removal of the hairs, by extraction, from the affected part has been attended with good success since, in so doing, many of the parasites are disposed of. A liniment composed of liquid tar and green soap two parts and alcohol one part will prove beneficial since it is antiparasiticide, disinfectant, dessicating, emollient and cleansing.

Mercuric chlorid in one to two per cent aqueous solution, tincture of iodine, sulphur iodid ointment, red iodid of mercury ointment 1-8, sulfur ointment, etc. are all useful applications. It is advisable to clip the hair from unaffected parts adjacent to the diseased foci, so that any spread of the malady may be immediately detected. Whatever the treatment, it is a good plan to wash daily with soft

soap and water before applying it. Cleanliness is extremely essential." Hyde and Montgomery state that the parasitocides are corrosive sublimate in the strength of 1-4 grains (0.066-0.266) to the ounce; formalin (1-4 per cent), sodium hyposulfite in saturated solution, spirit of green soap.

CHAPTER IV

FORAGE POISONING, ENZOOTIC, AND ASPERGILLOSIS

We have several excellent illustrations of how other **Forage Poisoning** external known parasitic organisms may produce disease.

Catarrhal stomatitis, for instance, may be produced by the ingestion of fodder which has become infected with any one of several fungi belonging to distinct orders. Among these are the rust of clover, bacteria, mildew of grass, and the rape-destroying fungus, *Polydesmus exiosus*, even the common grass rust and other rusts upon grasses as well as the bunts and smuts are known to produce this form of disease. Among higher plants, such products as the pungent spices of pepper and of the roots of horseradish and radish are treated at length in such pathologies as the Friedburger and Frohner Veterinary Pathology.

Serious diseases of the stomach are caused not only by pathogenic germs but also by the ingestion of various foods. Many foods, such as unclean, or damaged fodder, poor water, musty hay, mouldy corn, decomposing potatoes are responsible for gastro-intestinal catarrh. Many fodders, also, contain irritant substances. There are several forms of gastro-enteritis. Among forms of the third class (including those caused by ingestion of lower organisms such as fungi or poisonous substances) we may mention botulism, fish poisoning, injuries produced by mould fungi, smuts, rusts, and, finally, the so-called toxic gastro-enteritis produced by numerous poisons. These have sometimes been classed as irritant poisons and narcotic irritant poisons. The vegetable poisons under this head are numerous and have been treated under the different plants. Some pathologists, however, mention especially lupinosis of sheep and equisetosis.

The terms applied to this disease are Cryptogamic Poisoning, Forage Poisoning, Enzootic Cerebritis, Epizootic Cerebro-Spinal Meningitis, Leuco-Encephalitis, etc.

Characterization. So-called forage poisoning among horses and mules is a non-communicable disease, which undoubtedly belongs to a group of cryptogamic poisonings. Horses seem to be slightly more susceptible than mules, although it usually terminates fatally in both species.

The disease is characterized by symptoms which are referable to a disturbance in the central nervous system, and by lesions which, if present, are also found there. The course of the disease may be very acute, or it may be greatly lengthened, depending upon the suddenness of the onset. The mortality is very high and but few well developed cases ever recover. Suckling foals do not contract the disease.

History. This disease has prevailed quite generally throughout the Eastern and Central parts of the United States for many years, but until recently has not attracted any considerable attention. During the past few years, however, it has occurred with unusual frequency in the Central West, and, because of the extensive losses directly attributable to it, has

become of great economic importance. In various parts of Iowa, for instance, individual stock-owners have lost several thousand dollars from its ravages. In the different localities the disease has been known by various names, such as "grass staggers," "choking distemper," and "putrid sore throat," and because it apparently presents some of the distinguishing characters of a specific infectious disease, has been frequently recognized as "infectious cerebro-spinal meningitis."

A noteworthy fact however is, that thus far no evidence has been discovered which would indicate that the disease is transmissible from animal to animal, or that it is even inoculable.

On the other hand, an outstanding feature in every outbreak is, that the affected animals have had access to unwholesome food, either while at pasture or in the stable.




Fig. 3. Common *Aspergillus* on mouldy corn. 1. General appearance, showing long conidiophore and sterigmata on end. 2. Perithecial sac with one of its asci and ascospores. 3. Contents from an unripe perithecial sac. 4. A small part of the mycelium with conidiophore c and spore bearing sterigmata, young sporangium or. 2, 3, 4 after DeBary.

Geographical distribution. The disease has been reported from nearly every part of the United States. It never becomes epizootic, but is usually confined to isolated localities.

While forage poisoning is not necessarily peculiar to low, poorly drained districts, it is at least most frequently observed in those places where conditions are most favorable for the development of cryptogamic growth.

Etiology. The disease seems especially likely to appear when horses or mules are fed on grain or fodder which has become overgrown with

moulds, or when at pasture, they have had access to grass which, for various reasons, has become fermented or mouldy. Various micro-organisms have been found to be associated with the disease, but as yet none have been proven to possess any etiological significance. Cultural and histological studies have all proved negative. Dr. Moore has in one instance succeeded in obtaining a pure culture of the colon bacillus from the brain.

Symptoms. Depending upon the severity of the attack, the disease may manifest itself in any one of three forms, namely, acute, subacute and abortive. It is possible to observe all of these forms in a single outbreak, as the suddenness of the onset is apparently regulated by the amount of the poison laden food which the animal has ingested.

The acute type is characterized by the abruptness of its appearance, and the grave general disturbances which immediately manifest themselves. There is sometimes violent trembling and twitching of the muscles over the entire body, but most commonly the acute form is ushered in by stupor.

There is manifested a weak, staggering gait and the pharynx is either partially or completely paralyzed. The tongue may also be partially paralyzed and protrude from the mouth, and saliva falls in strings from the lips. The pupils are dilated and the conjunctiva is, as a rule, highly congested. The pulse is variable and may be very rapid and hard, or scarcely perceptible, the respiration is hurried and jerky. The temperature may be slightly elevated, but is most frequently subnormal. Intestines and bladder are paralyzed.

In this form there may be slight muscular rigidity affecting the muscles of the back, neck and jaws, although in many cases this symptom never manifests itself. There is no rigidity of the ocular muscles. The animal soon becomes so weak that he is no longer able to support himself and falls. Delirium may manifest itself, in which the patient may perform a series of movements as if trotting, or become so violent as to do himself serious injury, but most often coma and complete paralysis supervene and death results in from four hours to two days from the commencement of the attack.

The subacute form is much the same as the preceding, except that it develops more slowly and the symptoms are not so violent. It is first noticed by a slowness in mastication and a difficulty in swallowing. A further indication of approaching paralysis is seen in the frequent knuckling and the loss of control over the tail. The temperature is subnormal and the pulse and respiration are but slightly altered. The bowels and bladder are inactive and it is seldom that voidance of urine and faeces occurs voluntarily. There is but slight rigidity of the muscles if indeed there is any, and no evidence of pain is apparent. These symptoms may last two or three days, when gradual improvement takes place, or the paralysis becomes more complete, the general weakness more marked, paroxysms of delirium develop, with inability to stand, breathing becomes more labored, coma comes on and death results apparently without a struggle. This form lasts from six days to two weeks.

In the abortive form there are no well marked constitutional symptoms. The appetite may be somewhat lessened, the ability to swallow slightly impaired, and the animal's movements a little uncertain, but no very noticeable symptoms appear to attract the attention. Improvement usually takes place on the third or fourth day, and recovery is the usual result.

Lesions. As a rule, post-mortem examination reveals no naked eye changes in the tissues of animals dead of forage poisoning. There

may be congestion of the brain and cord with extensive effusion into the ventricles and subarachnoid spaces. Few small hemorrhages and parenchymatous degenerations within the various organs have been mentioned. MacCallum and Buckley have found in the brains of horses dying of this disease, areas of softening in the frontal region on each side, anterior to the motor region of the cortex." This lesion was practically confined to the white matter immediately under the cortex. In the affected areas there was "complete destruction of the brain substance, in which the anatomical structures are disintegrated and largely replaced by a colloid-like material. The neighboring blood vessels were acutely inflamed, with cellular infiltration of leucocytes and red corpuscles into the perivascular spaces and tissues. In a later outbreak these writers failed to find the brain lesion, but did observe the vascular changes above described.

McCarthy and Ravenel, in a study of fifteen animals found certain lesions in the upper gastro intestinal tract and in the central nervous system. These were

(1) In the intervertebral and Gasserian ganglia, where a pericapsular, small round cell accumulation was present. The cells were all of the same type, the nucleus and protoplasm being about the size of a red corpuscle. There was no evidence that these cells were the result of proliferation of the original layer of capsular cells.

(2) Cortical lesions. — These consisted of congestion of the cerebellar and cerebral cortex. There were also capillary hemorrhages. The meninges were normal.

(3) Changes in the choroid plexus. — In three cases the choroid plexus was changed into a triangular tumor-like mass, of a yellowish red color and of a firm consistency. The increase in size was found to be due to a proliferation of the elastic tissue surrounding the vessels.

(4) Changes in the nerves. — There was a distinct degeneration of the nerves supplying the larynx and neck. This was present in the nerve up to the ganglion, but was not found in the posterior roots. Other slight changes were detected.

Moore failed to find any gross lesions in the nervous system and other organs in the cases examined by him. In one case the brain, spinal cord and organs were studied histologically with negative results.

Differential diagnosis. A very important point in the recognition of forage poisoning is the history which has been referred to previously. It must be distinguished from inflammations of the brain and meninges, and from rabies.

Treatment. In the acute cases this is seldom successful, although quick-acting stimulants to arouse the patient may be tried. In the subacute cases a purge should always be given to rid the intestines of the poison. Strychnin in large doses, to overcome the extreme depression of the nerve centres, and atropin to support a failing circulation may be administered hypodermically at frequent intervals with benefit.

In the very mild cases, all that is necessary is to empty the bowels with a purge.

It is of the utmost importance, in all cases, with the return of the appetite, to supply only such food and water concerning the wholesomeness of which there can be no question.

Prevention. Since it seems to be quite generally accepted that this disease is brought about by the ingestion of mould-contaminated food

the prophylaxis is apparent. Whenever the disease makes its appearance either in a stable or a pasture, the animals should be immediately removed from further exposure by changing the food supply. The food should come, preferably, from a clean, new source and the water should not be contaminated by surface drainage. It is also well to thoroughly disinfect the mangers and feed-boxes, and render innocuous the soiled litter.

There is no known means of artificial protection and the disease will recur if the animals are again allowed access to spoiled food. (Stuhr)

Poisoning from Silage *History.* During the winter of 1908-1909, several cases of poisoning from spoiled silage were reported to Dr. Stange of the Iowa State College. Other cases have no doubt been encountered. In every instance, as in the case reported by Dr. Beaumont, below, moulds occurred in the silage. Dr. Beaumont says: "I am sending you under separate cover by mail a specimen of corn silage upon which you will notice a growing some form of mould which in my opinion is accountable for a very peculiar disease existing among a herd of young horses and mules belonging to a farmer living here."

Dr. R. E. Buchanan found these moulds occurring in spoiled silage to be a species of *Monascus*. Other moulds, *Mucor*, *Penicillium glaucum*, and *Verticillium* were also present, but there was a preponderance of *Monascus*.

Symptoms. "The first animal, a three year old filly, was taken sick about April 1st, showing symptoms as follows: Gaunt, depressed, stiffness of gait. When lying was unable to rise but when assisted to rise would stand and show inclination to eat but was unable to masticate and swallow food. Temp. 103.5 F. Pulse 86, Respiration 36. friction sounds distinctly heard at each heart beat. A whistling sound was emitted during expiration and there was also a suppressed painful cough. Animal died in about five days.

"A two-year-old male and one two-year-old filly were attacked with disease. The male is improving and will recover but the two-year-old filly shows exactly the same symptoms as Case No. 1, aside from being especially stiff and lame in one fore shoulder, and I think will die within two days."

Treatment. The treatment as followed by Dr. Beaumont is described in detail in his paper before the Missouri Valley Veterinary Association, June 16-17, 1909. Briefly, the method was as follows:

Tincture *Strophanthus* in two-dram doses, every two hours (given as a cardiac stimulant, the heart action being very weak). 1 quart of raw linseed oil given in two doses, six hours apart (as general laxative). Potassi Nitras in half to one ounce doses, dissolved in water and given as a drench, every three hours (alternative diuretic, and respiratory stimulant). After the first twenty-four hours the Tr. *Strophanthus* was discontinued and he began giving Iron Quinine and Strychnin tonic in one-ounce doses three times daily. This was continued with the Potassi Nitras until the animal showed marked improvement when both remedies were discontinued and he prescribed Fowler's Solution (Liquor Potassi Arsenici) in half-ounce doses three times daily during convalescing stage of the disease which lasted about ten days or two weeks.

Dr. C. H. Stange has contributed the following on forage poisoning and especially with reference to silage:

"Numerous cases have been reported of an affection of the central nervous system, the symptoms being in general quite similar but different and varying

causes are assigned. Dr. Francis reports that in the fall of '03, spring of '04 four to five thousand horses and mules died with a nervous disorder characterized by structural changes in the brain which cause incoordination, delirium, coma and usually death. He concludes that the disease is not caused by moulds but is the result of animals having free access to a labor diet when kept in idleness. He was unable to find the germ described by Wilson and Brimball.

"Professor Harrison of the Ontario Agricultural College reported several cases and as a result of his investigations he concluded that the disease was due to a coccus isolated from the meningeal fluids. Pearson studied an outbreak in seven horses, five of which died. The outbreak occurred soon after opening a new silo, the ensilage from which was mouldy. The symptoms observed were very similar to those observed by Professor Harrison and he emphasized the paralysis of the pharynx and great muscular weakness. He concluded as a result of feeding experiments that the so-called cerebro-spinal meningitis was a forage poisoning. Dr. Dow of Connecticut describes two cases which were attributed to watering from a tub containing a mouldy silage. Dr. Ferguson of Texas describes three cases of forage poisoning due to moulded corn. There was vertigo, coma, low temperature, pulse in later stages rapid and irregular. In 1901 Dr. Hickman investigated an outbreak among horses in North Carolina in which a large number of horses died. In 1906 another outbreak occurred at the same place (Hyde Co.) in which about forty horses and mules died in about three weeks. The cause in these cases seemed to be moulds on vegetation. On the whole the country is low and swampy. The pathological changes of Epizootic Leuco-Encephalitis were described by McCullum and Buckley in 1902. Muller of Germany reported an outbreak among horses, cattle and sheep due to mouldy straw (*Berliner-Tierärztliche Wochenschrift*). Drs. McCarroll and McMullen describe an outbreak of cryptogamic poisoning in horses due to feeding mouldy beet tops. Dr. Lockhart describes several cases in Canada. The prominent symptom seemed to be the inability to swallow.

"Two outbreaks have come under our observation during the past year. The first consisted of eleven head of horses, two horses were being fed for market, the others were fed in the same manner during the nights and turned out during the day. The first animal affected was one being fed for market. It ate part of its feed in the morning but in a few hours showed symptoms of ptyalism, depression and paresis of the hind quarters. By noon the animal was down, unable to rise and struggling some and died that night. The next animal to show symptoms was its mate. The symptoms shown in this case were similar to acute cases of the so-called cerebro-spinal meningitis, coming on with trembling and weakness causing the animal to stagger. An early symptom in all cases coming under our observation is the ptyalism due to inability to swallow (*Dysphagia*). As a result the saliva collects in the mouth and hangs from it in strings. Muscles of different regions of the body are liable to contract. The breathing is rapid and in some cases may be of the Cheyne-Stokes variety. The temperature in this case was sub-normal. In some of the more chronic cases and when the animal has been down for some time with considerable struggling the temperature was somewhat elevated. The pulse was variable, being about normal in some cases and rapid and almost imperceptible in others. The animal became quite violent at times and finally died living but a few hours longer than the first animal. The other seven animals showed

a more chronic course, showing inability to swallow, slow, weak pulse, difficult, noisy respiration, weakness and paralysis, spasm of muscles of head, neck and back, death taking place in from two to six days. The other two animals showed a mild type of the disease as slight loss of control, some exophthalmia, loss of appetite and thirst and loss of condition. These animals were placed on potassium iodid and nux vomica and recovered.

"This outbreak was attributed to mouldy silage, which was being fed to the horses, but in order to be more certain 150 pounds of silage were shipped to the college and fed, first to one horse which died in two days from an acute form of the disease. Another horse was fed but would not eat the silage so well, consequently did not die quite so soon, living for several days. In both cases the symptoms resembled those seen in the original outbreak. Post mortem revealed no changes except a few petechia along the small intestine, a few infarcts in the kidney and slight softening of the brain. This however was not very marked, probably due to the fact that they were acute cases. Microscopic examination revealed the presence of mould in the mucosa and submucosa of the intestine, also mycelial threads growing between and around the renal tubules.

"The other outbreak consisted of four horses, three of which died of an acute form of the disease, the symptoms being similar to those already described. The fourth being of a more chronic nature was placed on potassium iodid and nux vomica and recovered. In this outbreak the hay was found to contain a fine mould and was cut from an old pond which had been plowed up and seeded. The water had overflowed this, however, and stood for some time. The symptoms and post mortems were similar to those described in the first outbreak, with the exception that no histological examinations were made.

"A form of cerebro spinal meningitis is quite common in Germany. It has also been described in Australia, Great Britain and Russia. It may be that these outbreaks are due to other causes than those already described. Sigamgroizky and Schlegel found a form of coccus in the sub-arachnoid fluid, but it was necessary to make sub-dural injections of cultures of this organism to cause meningo encephalitis. Johnne found a diplococcus in the cerebro-spinal fluid of affected horses.

"Ostertag found a diplococcus similar to the one found by Johnne in the cerebro-spinal fluid in the so-called Borna's disease. They were pathogenic for horses and sub dural injections produced symptoms and death similar to cases of Borna's disease. Hutyrá and Marek call attention to the fact that bacteriological investigations have not been followed by the same result but possibly the several investigators were working with the same organism. Nevertheless it remains to be shown whether all cases of cerebro-spinal meningitis are due to the same cause and resemble Borna's disease. On the other hand it is possible that epizootic cerebro-spinal meningitis of domestic animals has no specific cause.

"It is apparent that mouldy food and water has caused several outbreaks in this country. Natural infection in European outbreaks is also supposed to take place through infected food and drinking water. The disease is not transmitted from one animal to another. Mohler calls attention to the very interesting work of Schlegel and the Berliner Tierärztliche Wochenschrift who



ERGOT ON VARIOUS GRASSES

FIG. 4-1. Manna Grass (*Glyceria serotina*). 2. Blue Grass (*Poa*). 3. Spikelet of Bottle Grass (*Aspella Hystrix*). 4. Reed Canary Grass (*Phalaris Arundinacea*). 5. Wild Rye (*Elymus robustus*). 6. *Koeleria cristata*. 7. Wheat Grass (*Agropyron Smithii*). 8. Red Top (*Agrostis alba*). 9. Blue Joint (*Calamagrostis canadensis*). 10. Timothy (*Phleum pratense*).

associates with the affection an organism which he termed *Streptococcus melanogenes*. Mohler states, however, that whether the disease is of microbial origin or an intoxication has not yet been definitely established."

Ergotism is a disease of bovines caused by the ingestion of *Ergotium* considerable quantities of food contaminated by ergot. Equines are apparently less susceptible than bovines, although the horses have been known to suffer severely from the disease. Ergotism in man is not an uncommon occurrence, and in nearly every instance it has resulted from eating bread made of ergotized grain. The disease makes its appearance among cattle chiefly in the winter and spring seasons and has at times been the cause of serious losses throughout the central and western states.

Ergot is the sclerotium of a parasitic fungus, *Claviceps purpurea*, which infects many species of native and cultivated grasses, and appears on some of our grains, especially rye. The sclerotium represents a stage in the life history of the fungus, which is intermediate between that of the mycelium or spawn, and that of the spore-bearing thallus. It flourishes particularly well on rich soil and in warm, damp seasons. The chemistry of ergot is not exactly known, although Robert succeeded in separating three bodies, namely, ergotinic acid, cornutin, and sphacelinic acid.

Ergotinic acid is a protoplasmic poison, and when injected intravenously produces inflammation of serous and mucous membranes, disintegration of red blood cells, and wide-spread ecchymoses, cornutin excites the central nervous system and causes general convulsions, and sphacelinic acid induces gangrene.

Symptoms. Ergotism manifests itself among animals chiefly in the chronic form, since, as a rule, the poison is acquired in small amounts and accumulation takes place slowly. Two distinct types of the disease are recognized, namely: spasmodic and gangrenous. Symptoms referable to the digestive tract, such as nausea, vomiting, colic, diarrhoea or constipation appear in both forms. Pregnant animals very frequently abort.

In the spasmodic type of the disease, symptoms due to over stimulation of the central nervous system, appear. These are tonic contraction of the flexor tendons of the limbs, anaesthesia of the extremities, muscular trembling, general tetanic spasm, convulsions and delirium. Death usually occurs from secondary causes.

Gangrenous ergotism is attributed to prolonged constriction of the arterioles, and more directly perhaps to degenerative changes in the vessel walls, and the consequent formation of hyaline thrombi.

It is characterized by coldness and anaesthesia of the extremities, followed ultimately by dry gangrene of these parts. The effects of this dry gangrene are often very serious and amount to sloughing of the feet, tips of the ears, tip of the tail, shedding of the hair, teeth, etc. Death takes place from exhaustion.

Lesions. With the exception of the gangrene which may vary greatly in severity, there are no lesions of especial significance. Degenerative changes in the sensory area of the cord and in the vessel walls have been observed in animals slowly poisoned with ergot.

Treatment. The first essential in the treatment of ergotism is to remove the cause. In well established cases treatment does not as a rule prove satisfactory.

Tannic acid is the chemical antidote, and should be given to neutralize the unabsorbed portion of the poison. Chloral is the physiological antidote. In

addition to giving the antidote, the treatment is entirely symptomatic. (Stuhr).



FIG. 5. A. *Aspergillus fumigatus* showing conidiophore on right with sterigmata and spores attached on left. B. *A. niger* showing conidiophore, sterigmata, and spores attached in chains. After Siebenmann.

Pneumonomycosis is a not uncommon disease of domestic animals caused chiefly by the mould, *Aspergillus fumigatus*, although the *Aspergillus niger* is also pathogenic for birds. This disease is most frequent in birds, both domestic and wild, occasionally observed in horses and cattle, and rarely in man. Respiratory diseases and lowered vitality predispose. In all species the disease is characterized by purulent local inflammations in the lungs or other tissues, and a purulent and necrotic pseudomembrane upon the bronchial, tracheal, and other mucous membranes upon which it grows. The appearance of the pulmonary lesions sometimes resembles tubercle, sometimes actinomycosis.

Pneumonomycosis has been experimentally produced in birds (pigeons and geese) by compelling them to inhale aspergillus spores for a few minutes, after which they usually die of pneumonia in a few days. Rabbits have also been successfully inoculated by intravenous injection of spores.

Etiology In mammals the *Aspergillus fumigatus* and in birds the *Aspergillus fumigatus*, *niger* and *florescens* seem to be pathogenic species.

Infection takes place most commonly by inhalation of the spores which often are suspended in the air, or by taking them in with the food. Intestinal infection has not been observed. The spores are widely distributed in nature and exist in vegetable matter and grain abundantly. They possess remarkable vitality and exhibit considerable resistance to destructive agencies. The pathogenic power of the mould does not depend upon any product which it elaborates but upon the reactions which result from its penetration into the tissues. Peck observed the disease in seven subjects, in a stable where horses were fed on mouldy hacked hay.

Symptoms. The disease is of slow development in the larger animals and may not be observed until well advanced. In general the symptoms are of a pneumonic nature and in addition there is progressive emaciation. A case in a

Jersey cow, described by Pearson and Ravenel presented the following symptoms: the animal had been in poor condition for six months before it was examined. It was weak and depressed, did not eat, breathed with difficulty and, at times, coughed violently. Percussion of the chest gave sounds clearer and louder than normal and auscultation revealed the lung and bronchial sounds much intensified. Six days later these symptoms became more pronounced, the respiration and pulse very rapid. The animal grew rapidly weaker and died ten days after first being seen. The symptoms in birds are much the same as those in mammals except, that the disease runs a more rapid course. Emaciation advances rapidly and febrile diarrhoea may set in and continue until death in from a week to two months. At times emaciation is the only symptom. Fowls emit a glairy discharge from the nostrils which may contain the spores. In the prevention of the disease in fowls therefore, it is necessary to isolate or destroy the sick fowls together with the carcasses and fumigate the poultry houses. The roosts may be whitewashed.

Lesions. The lesions take the form of a military suppurative process, the foci varying in size from very small up to that of a pea. These may exist in large numbers and be scattered throughout the entire lung. Sometimes they become confluent and produce large areas of disease. The process starts in the bronchial mucous membrane, and later involves the bronchioles and alveoli. A very important feature is the intense amount of emphysema which is apparent on external examination of the lung. The lobules are often widely separated and can be readily seen in outline when a portion of the tissue is examined by transmitted light. In these emphysematous interlobular spaces, and in the air passages are seen whitish, mouldy looking patches. They are composed of denuded epithelium, inflammatory exudate, fruit hyphae and spores.

The lesions spread by penetration of the mycelium causing a destruction of tissue. Spores are not found within the tissues. In rare cases there is diffuse pneumonia characterized by hepatization and interstitial infiltration. On this latter account the disease has been described as being similar to contagious pleuro pneumonia of cattle. There may be pulmonary gangrene from secondary invasion of putrefactive organisms acting upon the devitalized tissue. An interesting feature is that this disease may interfere with the tuberculin test. This was shown in the case, above referred to, in which the test was used without success, and lesions of tuberculosis found in the lung on postmortem examination.

Treatment. This must of necessity be unsatisfactory since it is quite impossible to destroy the moulds which have penetrated the lungs. (Stuhr)

CHAPTER V

POISONING FROM FUNGI

That fungi of various kind are injurious, was known to the ancients. Prof. Ford¹ says, "The most interesting cases of mushroom or, as commonly described, toadstool poisoning and one of the first authentic cases on record, occurred in the family of the Greek poet, Euripedes, who lost in one day, wife, daughter, and two sons, who in the poet's absence partook of the deadly species. Among the great ones whose lives were sacrificed to the same ignorance may be mentioned Pope Clement VII, the Emperor Jovian, the Emperor Charles VI Berronial of Naples and the widow of Tsar Alexis. The death of the Emperor Claudius is also assigned to this cause, but the reason and manner of the accident are not certain.

In addition to poisoning from toadstools, it has long been known that Ergot (*Claviceps purpurea*) is injurious to man and lower animals. In recent years Ergotism has not been so serious as formerly.

Other fungi also may be responsible for the death of animals by poisoning. The Fly Agaric (*Amanita muscaria*), a beautiful species, is common in many parts of the United States. I have described it in detail in another part of this work. In this connection I shall quote freely from the detailed and excellent account of poisoning as given by Prof. V. K. Chesnut, and the excellent report given of *A. phalloides* by Prof. Ford, who has written the most recent account of poisoning from this fungus.

The symptoms and treatment are thus described by Mr. V. K. Chesnut:

"The symptoms of poisoning from the fly amanita, as deduced from a number of cases, are varied. In some instances they begin only after several hours, but usually in from one-half to one or two hours. Vomiting and diarrhoea almost always occur, with a pronounced flow of saliva, suppression of the urine, and various cerebral phenomena beginning with giddiness, loss of confidence in one's ability to make ordinary movements, and derangement of vision. This is succeeded by stupor, cold sweats, and a very marked weakening of the heart's action. In case of rapid recovery the stupor is short and usually marked with mild delirium. In fatal cases the stupor continues from one to three days and death at last ensues from the gradual weakening and final stoppage of the heart's action.

"The treatment for poisoning by *Amanita muscaria* consists primarily in removing the unabsorbed portion of the amanita from the alimentary canal and in counteracting the effect of muscarin on the heart. The action of this organ should be fortified at once by the subcutaneous injection, by a physician, of atropin in doses of from one one-hundredth to one-fiftieth of a gram. As a stimulant emetic, mustard is particularly valuable. If this is not effective apomorphin should be administered by a physician. In case of profound stupor however, even this may not produce the desired action. Tannin is of little

¹ Science N. S. 30: 97, 98.




Fig. 6. Deadly Amanita (*Amanita phalloides*) U. S. Dept. Agri.

or no value in rendering the muscarin insoluble in the stomach. If vomiting has not taken place, recently burned charcoal, or two grains of a one per cent alkaline solution of permanganate of potash may then be administered, in order, in the cases of the former substance, to absorb the poison, or, in case of the latter, to decompose it. This should be followed by oils and oleaginous purgatives, and the intestines should be cleaned and washed with an enema of warm water and turpentine.

"Experiments on animals poisoned by the fly amanita and with pure muscarin show very clearly that when the heart has nearly ceased to beat it may be stimulated to strong action almost instantly by the use of atropin. Its use as thus demonstrated has been the means of saving many lives. We have in this alkaloid an almost perfect physiological antidote for muscarin, and therefore in such cases of poisoning its use should be pushed as heroically as the symptoms will warrant. The presence of *phalloin* in *Amanita muscaria* is possible, and its symptoms should be looked for in the red color of the blood serum discharged from the intestines. Its treatment, which is difficult, is discussed under *Amanita phalloides*.

"It is well known that in some parts of Europe the fly amanita, after the removal of the poison by treatment with vinegar, is a common article of food. It was interesting to discover not long since that among some of our own people a similar practice prevails. Though most of the colored women of the markets look upon the species with horror, one of them recited in detail how she was in the habit of cooking it. She prepared the stem by scraping, the cap by removing the gills and peeling the upper surface. Thus dressed the mushrooms were first boiled in salt and water, and afterwards steeped in vinegar. They were then washed in clear water, cooked in gravy like ordinary mushrooms and served with beefsteak. This is an exceedingly interesting operation from the fact that although its author was wholly ignorant of the chemistry of mushroom poisons, she had nevertheless been employing a process for the removal of these poisons which was scientifically correct. The gills, according to various pharmacological researches, are the chief seat of the poisonous principles in this plant and their removal at once takes away a large part of the poison. The salt and water would remove *phalloin* or any other toxalbumin the mushroom contained, and although the presence of *phalloin* or any of this class of poisons has not been demonstrated in *Amanita muscaria*, there is a strong suspicion that it may occur in slight amount. The vinegar, secondly, removes the alkaloid poison, muscarin, and the mushroom after the two treatments is free from poisons. This process is cited, not to recommend its wider use but as a matter of general interest. The writer's recommenda-

tion is that a mushroom containing such a deadly poison should not be used for food in any form, particularly at a season when excellent non-poisonous species may be had in abundance.

"It is surprising that cases of poisoning are not more frequent. At Tacoma Park, D. C., on November 9, of last year, a lady who has a thorough knowledge of edible and poisonous mushrooms met a family, consisting of a man, woman, and two children, who had just completed the gathering of a basketful of the fly amanita and the death cup, described below, which they were taking home to eat. In reply to questions the woman stated that they had often eaten this kind purchased dry at an Italian store, but that they had never gathered fresh ones before. Of course they had mistaken the species, or possibly the dried ones were fly amanitas from which the poison had been removed by treatment with vinegar. After considerable persuasion the people consented to throw the lot away.




Fig. 7. Fly Amanita (*Amanita muscaria*). C. B. Dept. Ag 1

"It is impossible to say what amount of the fly amanita would prove fatal, but in this connection it is of interest to note the custom reported by Krasheninnikoff, a Russian who travelled in Siberia and Kamchatka from 1733 to 1743, namely that the natives of the latter country, particularly the Koraks, used the fly amanita as an intoxicant, three or four specimens constituting a moderate dose for one habituated to its use, but ten being required for a thorough drunk.

The same observations, with varied details, have been made by others, particularly by Langsdorff, who traveled around the world with the Russian navigator Krusenstern from 1803 to 1806, and in more recent times by Kennan in his first Siberian journey of 1865-67.

"The plant may be taken fresh, but its taste is so disagreeable that only with great difficulty can a sufficient amount be eaten to produce the intoxicating effect. The Koraks have two principal methods of taking it. First, by swallowing pieces of the dried caps without chewing them, second, by boiling the dry caps in water and then drinking the liquor thus produced mixed with the juice of berries or herbs to disguise the taste. The intensity of the poisonous character of the fly amanita undoubtedly varies at different ages, with different individuals, and with different methods of preparation. The amount of the poison that can be taken into the system with impunity varies, too, with the person who takes it. The fact that a Korak, who has long used the plant as an intoxicant, can eat ten specimens and merely become drunk, does not prove that a similar number would not be fatal to an American who had never eaten it before.

"Very diverse statements concerning the properties of this fungus have been recorded. While some have attributed to it edible qualities, others have asserted that it is a most active poison and has caused numerous accidents by being confused with the Orange amanita. It is said to have caused death even when eaten in small quantities, and again it is said to have been eaten in abundance without any evil results. According to Quelet, it acts as a cathartic if eaten in small quantity but causes death if eaten freely. One of my own correspondents assures me that he has eaten of the yellow variety *Var formosa*, without evil results, and that he regards it as very good. But there is no disputing the fact that the species possesses intoxicating and poisonous properties. It has long had the reputation of possessing properties fatal to flies that sip its juice. This suggests the names muscaria, Fly amanita, Fly agaric and Fly killer by which it is known. I have myself seen the cap of a single specimen surrounded by a circle of lifeless flies that had sipped the viscid juice from its moist surface and fallen victims to its virulent properties before leaving the place of their fatal repast.

"Some have attempted an explanation of the contradictory statements concerning this plant by supposing that its poisonous properties are not always developed, that in some localities or under favorable circumstances it is harmless. This explanation violates our sense of the constancy of Nature, and is not at all satisfactory. In the case of my own correspondent, the caps were peeled before cooking. May it not be that much of the noxious quality resides in the epidermis and the viscid substance upon it, and that by discarding this the dish is rendered less dangerous? In some cases it is said that those who eat it freely and without harm boil it a long time in water and throw away the water. In this way, doubtless, much of the poison is abstracted. Long soaking in salt and water, also in vinegar, have been recommended as a means of rendering suspected or noxious species harmless, and may have been practiced in some of the cases in which this fungus has been eaten with impunity. Whatever may be the explanation of the contradictory statements, the only safe way is to consider this species as deleterious and avoid its use under all circumstances. There is no need of taking any risks, with suspected species, -

since there are so many good ones against which no charge of evil has ever been established."

A second very poisonous species is the White or Deadly Amanita (*Amanita phalloides*), common also in some parts of the United States. This species is described in another part of this work. This and allied species are eaten ignorantly by persons who do not know the nature of the powerful poison found in the plant. Prof. Ford says, "A small amount of the fresh material is sufficient to cause profound illness with fatal outcome, so potent is the poison contained in its meshes, and the raw plant seems usually more toxic than the cooked specimens."

"Two or three 'deadly amanitas' suffice to bring on disastrous results, and Plowright reports the death of a child of twelve from eating a third of the pileus of a small raw plant. The extreme toxicity of this species illustrates the dangerous consequences which the admixture of two or three specimens to a dish of edible mushrooms entails."

"Following the consumption of the fungi there is a period of six to fifteen hours during which no symptoms of poisoning are shown by the victims. This corresponds to the period of incubation of other intoxications or infections. The first sign of trouble is sudden pain of the greatest intensity located in the abdomen, accompanied by vomiting, thirst and choleraic diarrhoea with mucous and bloody stools. The latter symptom is by no means constant. The pain continues in paroxysms often so severe as to cause the peculiar Hypocratic facies, "la face vultueuse" of the French, and though sometimes ameliorated in character, it usually recurs with greater severity. The patients rapidly lose strength and flesh, their complexion assuming a peculiar yellow tone. After three to four days in children and six to eight in adults the victims sink into a profound coma from which they cannot be roused and death soon ends the fearful and useless tragedy. Convulsions rarely if ever, occur and when present indicate, I am inclined to believe, a mixed intoxication, specimens of *Amanita muscaria* being eaten with *phalloides*. The majority of individuals poisoned by the "deadly amanita" die, the mortality varying from 60 to 100 per cent. in various accidents, but recovery is not impossible when small amounts of the fungus are eaten, especially if the stomach be very promptly emptied either naturally or artificially."

Kobert isolated from the fungus, a substance which he called *phalun*, and which had the property of dissolving the red blood corpuscles. Such substances are called hemolysins. Prof. Ford says, "Very minute traces of this substance brought in contact with the red blood cells of man or with those of animals, produced within a short space of time, fifteen minutes to one or two hours, a complete solution of these corpuscles—a laking of the blood. So powerful was the hemolytic action that even in a dilution of 1-125,000 it was still operative upon the red cells of ox blood."

In a recently published statement by Prof. Ford it appears that¹ the fungus always contains another poison which differs from hemolysin in being resistant to heat and digestion, "the blood-laking substance *phallin*, being destroyed by heating to 70° C., and by the action of the digestive ferment. This substance he called *Amanita-toxin*, and the blood-laking substance *Amanita-hemolysin*. Abel and Ford² have shown that the so-called *phallin*, regarded by Kobert as


¹ Science N. S. 30:101

² Jour. Biol. Chem. 2:273, 1907.

a toxalbumin is a glucoside. Prof. Ford has obtained an anti-poison or an anti-hemolysin with a high grade of immunity. According to Schlesinger and Ford¹ the *Amanita*-toxin in a purified state is one of the most powerful of organic poisons—four tenths of one milligram killing a guinea pig within twenty-four hours. Ford believes that the hemolysin plays no part in human intoxication, but that the toxin is the active principal which resists the action of the gastric juice and boiling. He finds that the *Amanita rubescens* considered an edible species by some, contains an hemolysin as powerful as the Deadly *Amanita*. He found a toxin and an hemolysin in *Amanita virosa*. The latter substance in a dilution of 1-200 killed a guinea pig. The *A. sprete* produced intoxication and according to Ford must be classified with the "deadly poisonous" as the *A. verna*.

The *A. strabuliformis*, *A. chlorinorma*, *A. radicata*, and *A. porphyria*, do not contain hemolysins but small quantities of a toxin probably identical with *amanita-toxin*. The *Amanita solitaria*, regarded as edible, causes the blood corpuscles to adhere in clumps much as agglutination occurs with typhoid bacilli when brought in contact with the blood of a typhoid patient.

¹ Jour. Biol. Chem. 2279, 1909.



CHAPTER VI

POISONING FROM OTHER PLANTS. EQUISETOSIS, LOCOISM, AND LUPINOSIS

Equisetosis. It has been recently proven by direct experimentation that the common horsetail (*Equisetum arvense*) when ingested in sufficient amount, is capable of producing fatal poisoning among horses. This discovery is of great importance since the plant has a wide distribution, and at times is the cause of extensive losses. The common horsetail thrives best in moist sandy soils or in low, damp meadows, which are not frequently cultivated, and often constitutes a large part of wild hay. The dried plant alone seems to be poisonous. Young horses seem to be the most susceptible. Sheep are supposed to be slightly susceptible although cattle eat the hay in which the plant occurs in large proportion, with impunity. The toxic principle of the plant has not been determined.

Symptoms. The effects of poisoning from eating horsetail appear at times varying from two to five weeks, depending upon the age of the animal, and the amount of contaminated hay ingested.

The first symptoms are usually unthriftiness, general bodily weakness and emaciation. The animal seems to have a depraved appetite, preferring the plant to wholesome feed. As the disease progresses the muscular weakness becomes more pronounced, the animal loses muscular control and exhibits in-coordinate movements. During this stage the pulse and temperature are depressed, extremities are cold and the visible mucosae are pale. Appetite usually remains good until the end and consciousness is apparently retained. Finally the animal falls, manifests nervous excitement, paroxysms of convulsions appear and death results from exhaustion. In the final stage the pulse becomes accelerated and the temperature elevated.

Hypostatic pneumonia is a frequent complication.

Treatment. The first step in the treatment is the removal of the cause. A cathartic should be administered to rid the bowels of the irritant and nerve and heart stimulants given to combat the symptoms of depression. In case the patient is unable to stand, it would be advisable to give some support. When the animal is down it becomes necessary to guard against the development of hypostatic pneumonia.

Where cases are not too far advanced and appropriate treatment is instituted, recovery is the usual result. (Stuhr)

Stock poisoning by the loco weed is a frequent and serious condition with which the stock-owners of the western half of the United States have to contend. Montana and Colorado, especially, sustain heavy annual losses. Similar diseases occur in other parts of the world. In Australia other plants of the order *Leguminosae* like *Gastrolobium* produce similar symptoms. Maiden¹ states that the "Nenta Lessertia disease of S. Africa is identical with a disease of the Pea-eating animals of Australia and

¹ *Minicell*. Pub. Dept. Agr. N. S. Wales, 477-11.

with the Loco disease of the United States. Many forage plants of excellent repute such as white clover, alfalfa, lotus and other plants, may produce tympanites




Fig. 8. Common Horsetail (*Equisetum arvense*), the plant causing Equisetosis. 1. Fertile stem terminating in cone. 2. Sterile stem. a. rhizome tubers. 3. Sporophyll with sporangia. 4. Sporangia opened to discharge spores. 5, 6, 7. Spores with spiral elaters. After Wossidlo.

Symptoms—difficult breathing the poison enters the circulation and stops the action of the lungs and heart when the animals stagger and die

In the advanced stages the animals become frantic, hence the name "loco" or crazy. Horses and sheep are the most susceptible, although cattle are also affected.

Of the various species of loco weed, the stemless loco *Oxytropis Lamberti* Pursh, and the woolly loco weed (*Astragalus mollissimus* Torr), are the most injurious. These weeds grow luxuriantly on sandy ranges and appear early in the spring when other vegetation is scarce, and since they retain their fresh



Fig. 9. Loco Weed (*Astragalus mollissimus*), U. S. Dept. Agr.

green color during the entire summer they prove especially attractive to stock.

Recently the poisoning has been attributed by Dr Crawford and others of the U. S. Dept. of Agrl., to mineral salts in the plant.

The period of greatest danger is chiefly during the month of May.

Symptoms. The symptoms, which are referable to the nervous system, are attributable to the narcotic effect of the plant. They appear slowly and are apparently divisible into two stages. The first stage is characterized by the following symptoms. Stupor, defective vision, unnatural movements and apparent hallucinations. When excited the animals become frenzied. The coat becomes shaggy the teeth grow long and become loose, and a depraved appetite which is very marked, is developed. The animals prefer the loco weed to wholesome food, and will dig up the roots and eat them to satisfy their craving.

In the second stage there is emaciation, exhaustion, feeble movements and finally death from starvation. The course of the disease is quite variable

and may last from a few months to one or two years. Sheep manifest symptoms very similar to those above described.

Treatment. When the disease has reached the advanced stage, treatment is of no avail as recovery does not occur. If, however, the afflicted animals are taken early in the course of the disease and placed on pasture where loco weeds do not exist, and are given good nourishing food, there is hope of recovery.

Prevention, which of course is the most desirable, is not always practicable. Animals do not as a rule become addicted to the loco habit when they have plenty of wholesome food and salt.

There are no demonstrable lesions other than emaciation. (Stuhr).

The recent investigations of Marsh and Crawford lay considerable stress on the presence of barium in the plants and the Bureau of Animal Industry, Washington, D. C., recommends the following treatment: for cattle, strychnin in doses of three-twentieths to four-twentieths of a grain daily, administered hypodermically, for horses, Fowler's solution of arsenic in half-ounce doses daily in the drinking water or in the grain. This treatment should be continued for at least a month. To correct the constipation which is almost universal in locoed animals, magnesium sulphate (Epsom salt) may be administered as a drench in two-ounce doses. Epsom salt may also serve to some extent as an antidote to the poison produced by the weeds. Beneficial results have also been obtained by giving horses daily a drench containing two ounces of Epsom salt with ten drops of dilute sulphuric acid, and by giving cattle tri-weekly three or four ounces of Epsom salt with a proportional increase in the quantity of dilute sulphuric acid.

As the foregoing treatments are in the experimental stage, the Bureau of Animal Industry of Washington, D. C., would be glad to receive reports from their use.

The value of keeping stock away from these poisonous plants is indicated in some investigations that have been carried on by the Bureau of Forestry and the Bureau of Plant Industry. In many cases the ranges are becoming practically useless on account of these poisonous plants and if used the losses are so heavy as to materially reduce the profits of the business. In the Manti Forest Reserves in Utah for instance, it was found that the death of sheep was due to their browsing upon the chokecherry. Certain portions of the old trail were abandoned, and along other portions the chokecherry bushes were cut out. The method of handling the sheep was also changed. Instead of large bands which could be moved but slowly, smaller bands were trailed, and so far as possible they were allowed to fill up on healthy forage before entering the dangerous area. The trail was also improved wherever practicable and by this means it was possible to get the sheep through in much better shape and with little or no loss. The Department has also, in some instances, adopted the plan of flagging the area in which these injurious plants occur.

This is a disease of sheep and horses especially, caused by **Lupinosis**, eating the seeds and straw of the lupine. Cattle and goats are also susceptible and the dog has been poisoned experimentally. There are many species of the lupine growing in various parts of the United States, although the yellow lupine (*Lupinus luteus*) is the most toxic. The nature of the toxic agent found in the lupines has not as yet been determined. Arnold and Schneidemühl succeeded in isolating a chemical poison and



Fig. 16. *Lupinus* (*Lupinus leucophyllus*) common lupinus U. S. Dept. Agr.

gave to it the name lupinotoxin. They described its physical properties but failed to determine its chemical composition. Attempts to associate a fungus with the plant have failed. Lupinosis is characterized by jaundice, acute yellow atrophy of the liver, and parenchymatous inflammation of other internal organs.

Symptoms in Sheep. The disease appears in either the acute or chronic form, depending upon the amount of poison ingested. These two forms have been experimentally reproduced by giving carefully regulated amounts of lupinotoxin.

In the acute form the disease appears suddenly. There is loss of appetite, fever, hurried and difficult breathing, rapid pulse, stupor, vertigo, and not infrequently swelling of the lips, ears or face. The initial temperature may be as high as 104° to 106° Fahr., but is intermittent and gradually falls just before death. The pulse may reach 130 per minute and the respirations 100. A bloody froth may issue from the nostrils. Icterus which may be detected in the conjunctiva and the urine, usually appears on the second or third day. In certain cases this latter symptom fails to manifest itself and therefore is not constant. There is grinding of the teeth and sometimes trismus. The animal apparently prefers the recumbent position, extends the head on the ground and seems entirely oblivious to all surroundings. At first there is constipation the faeces being hard and scanty and covered with yellow mucus. Later diarrhoea may set in and the excreta be tinged with blood giving them a dark brown color. Emaciation develops rapidly. In case of recovery the symptoms gradually abate and improvement takes place slowly. Cachexia is a common sequel. In the chronic form the symptoms are not so violent. Jaundice may be entirely absent and emaciation and anemia may be the chief signs. Inflammatory tumefaction of the lips, eyelids, and ears with the formation of ulcers and scabs is described by various writers.

Course. Death may supervene within twenty-four to forty-eight hours, although frequently the disease lasts four or five days. The immediate cause of death is rapid emaciation and extreme weakness. Horses contract the disease from eating oats contaminated with the seeds or from eating the straw of the plant. The symptoms which they manifest are essentially the same as those above described. Horses seldom die from the effects of lupines.

Lesions. The cadavers are emaciated and decompose rapidly. The muscles are of a grayish yellow color, the fibers having become fatty and having lost their striations. The subcutaneous tissue of the abdomen and the omentum and mesentery are yellowish. The most important lesion in both the acute and chronic forms is found in the liver. The alterations in this organ are those of acute hepatitis. The liver cells have become swollen and granular on account of the parenchymatous change, or they may be more or less completely degenerated into fat. The gland is soft and friable and may be somewhat swollen. The interlobular connective tissue is greatly increased in amount due to inflammatory hyperplasia. In the course of a few days the liver undergoes acute yellow atrophy as a result of the absorption of the degenerated cells and the contraction of the hyperplastic stroma. In the chronic form the changes are those of chronic interstitial hepatitis. The icterus is of hepatic origin and due to catarrh of the bile ducts. The gall bladder is distended with bile and its lining membrane is congested and swollen. The kidneys and bladder may show changes, more or less marked, due to inflammation. The blad-

der is, as a rule, empty. In the digestive tract we observe frequently yellowish discoloration of the mucosa, hemorrhages in the small intestine with catarrhal lesions of the entire canal. The heart is pale and friable and the blood which it contains is dark and thick. Capillary hemorrhages are quite generally observed throughout all of the tissues.

Treatment. This is chiefly preventive since there is no specific antidote. Attempts should be made at once to prevent further absorption of poison by administering some acid as acetic or hydrochloric, well diluted with water. Alkalis should be strictly avoided as the poisonous principle is very soluble in alkaline solutions. It is advisable to evacuate the bowels by giving a purgative, preferably oil. Potassium permanganate is recommended by some as an antidote. Further than this the treatment is entirely symptomatic. (Stuhr).



Fig. 1. A. Larkspur (*Delphinium bicorne*). B. *D. carolinianum*. Larkspur poisoning (*Delphinosis*) is caused by various species of *Delphinium*. A. U. S. Dept. Agr. B. Ada Hayden.

CHAPTER VII

DELPHINOMIS, LATHYRISMUS (LATHYRISM), ACONITISM, VERATRISM, UMBELLIFERAE, CANTH, CICUTA.

The purple larkspur (*Delphinium Montanum* D C), and *Delphinocula*, other species which are found in the northwestern part of this country and especially in Montana are plants very dangerous to stock. Drs. Cheever and Wilcox have proven the toxicity of the above species by direct experiment and have called attention to the serious losses which they occasion annually.

The latter experimenter has fatally poisoned a yearling lamb within two hours by administering, *per os*, the extract made from less than an ounce of the dried leaves. The weed appears early in spring, in advance of the forage plants and it is during this period that the greatest harm results. The poison is found both in the leaves and the roots although the latter are not frequently eaten on account of their woody fibrous nature. The poisonous principle has not been isolated. Cattle and sheep are most susceptible although horses frequently suffer.

Symptoms. The first indication of poisoning is a general stiffness and a straggling gait, especially in the posterior limbs. Walking appears to be difficult, and is evidently painful. At this stage the pulse and respiration are much depressed, and the temperature is lowered. The appetite is retained in most cases until the appearance of the final stage of the poisoning. This is manifested by irregular muscular twitching of all of the muscles which finally becomes frequent and violent, and by incoordinate movement. There are attempts at vomiting and the animal froths at the mouth. Finally all of the muscles of the body contract spasmodically, the animal falls and dies in violent spasms. The pulse and respiration become very weak and rapid just before death. There is no aberration of the special senses. The course of the poisoning is quite rapid and death usually takes place within a few hours.

Lesions. The direct cause of death from larkspur poisoning is probably failure of respiration due to paralysis of the centre and the alterations therefore are those of asphyxia. The lungs are congested and dark-colored and the right heart, veins and capillaries are distended with dark colored blood. Other organs and tissues are normal with the exception of the general venous and capillary congestion.

Treatment. Potassium permanganate is the chemical antidote and should be given as early as possible in doses of fifteen to twenty grains to horses, thirty to fifty grains to cattle, and five to ten grains to sheep, dissolved in a copious amount of water. To combat the extreme depression of the circulation and respiration, atropin sulphate has proven very efficacious. Even after the final convulsions have begun this drug has been of good service. It should be hypodermically administered in doses of three-fourths to one grain to the larger animals and one-twentieth to one-fifteenth of a grain to sheep. In the late stages sheep should be given as much as one-sixth to one-

fourth of a grain. In the convulsive stage when there is impending failure of respiration, inhalations of ammonia may be resorted to with good results.

A feature of the treatment quite as important as the medication is the care of the animals. Complete rest and freedom from excitement are very essential since exercise or fright is likely to induce fatal spasms.

Finally the danger from poisoning can be largely obviated by preventing access to the young plant in the early spring.

In ancient times, this disease was quite common, affecting **Lathyrismus**, both man and the horse. It is very frequent today in Spain, Italy, Russia, and India.

Etiology. It is caused by eating bread made of flour derived from some species of Lathyrus or Vetch (*L. Cicera*, *L. Clymenum*, or *L. sativus*), or, in case of horses, by the consumption of Fodder Pea. The seeds of these species contain a toxic substance formed without the action of bacteria. Man, the horse and the pig are subject to this disease.

Symptoms. In domestic animals, the symptoms are debility of the rear and lower extremities, producing motor paralysis. In lower animals, the normal functions of the larynx become impaired because of paralysis.

Treatment. Change of fodder, providing good food, application of massage and electric treatment.



Fig. 12. Aconite (*Aconitum napellus*). Aconitism is caused by this and other species of Aconitum. Charlotte M. King.




Fig. 13. Green Hellibore (*Veratrum viride*). Common Eastward. (U. S. Dept. Agr.).

Prognosis. Fatal cases are not frequent.

Post-mortem. The ganglion cells of the anterior horns are atrophied, also the recurrent nerves and the muscles of the larynx.

(Adapted from Friedberger and Fruhner)

Various species of aconite (*Aconitum*) are known to be **Aconitum**, poisonous. One species, the *Aconitum napellus*, has long been used in medicine. The most common species in North America is the Columbia Aconite (*A. columbianum*), which is found in the Rocky Mountains and on the Pacific coast. Several other species occur in North America, one extending into northeastern Iowa, the European Aconite (*A. napellus*) is frequently cultivated in gardens. Cases of poisoning are largely due to the administration of over doses in medicine. In the Rocky Mountains accidental poisoning among live stock is confined to animals that graze at higher altitudes. All parts of the plant are poisonous, one tenth of a grain of the drug is a poisonous dose for some animals. The smallest fatal dose recorded in man is a teaspoonful of the tincture of aconite, which is equivalent to about XXX gr of the crude drug. The minimum lethal quantity is 1-16 of a gr for man.

Symptoms. The effects of the poisoning are a tingling sensation on the end of the tongue, which shortly gives rise to a burning sensation followed by pronounced constriction in the throat. It reduces the pulse and frequency of the cardiac pulsations. The action of the heart is lessened and the pulse is weak, irregular and intermittent, at first slow and then rapid, tingling and prickling over the entire body is characteristic, vision is confused, there are abdominal pains, vomiting and diarrhoea. Death is caused by the stoppage of respiration, but is preceded by numerous twitchings, in the case of the horse the animal falls and is unable to rise. The symptoms are different in cats and rabbits.

Treatment. No specific antidote is known, but physicians use atropin or digitalis and nuxie of amyl. The stomach should, however be evacuated at once; cardiac and respirative stimulants are given subcutaneously.

The Hellebores belonging to the genus *Veratrum* are common **Veratrum** in the mountain regions of the west and one species is abundant in the east. They are found in swampy places, in wet meadows and along brooks. The most frequent cases of poisoning come through the administration of the drug; although in the Rocky Mountains considerable quantities of the plant are consumed by sheep where grazing is close. All parts of the plant are poisonous although the root is more poisonous than the seeds and leaves which contain several alkaloids.

Symptoms. *Veratrum* is a powerful irritant and when inhaled in minute quantities produces sneezing. When injected under the skin it causes restlessness, when consumed in large quantities it produces salivation, frequent vomiting with purging pain, and collapse, the temperature falls. *Veratrum* is a drug poisonous to the motor and sensory nerves. Death is caused by paralysis of the heart, 1/16 of a grain has produced alarming symptoms in man and 1 gr injected subcutaneously produces poisonous symptoms in a horse.

Treatment. The stomach should be emptied immediately, then give stimulants, such as brandy and strong coffee.



Fig. 34 Larkspur (*Delphinium glaucum*) equals *Delphinium* (C. S. Dent, Agr.)



Fig. 5. *Urtica dioica* L. (Nettle) - a densely glandular plant. (U. S. Dept. Agr.).

The Cowbane (*Cicuta maculata*) and related species belong to the Carrot family. The Cowbane is common in low grounds in the Northern States and in the Rocky Mountains; another species (*C. rugosa*) occurs on the Pacific Coast, and a third species (*C. bulbifera*) is a common bog plant in the Northern States.

The Cowbane is frequently mis-called Wild Parsnip. The latter, however, has a conical root and is the feral form of our cultivated species, and, although it may be somewhat injurious, as indicated elsewhere, is not poisonous like the Cowbane (*Cicuta maculata*).

Cowbane is a smooth, marsh perennial plant from 2.5 feet high, with fleshy, fascicled roots and a pungent odor, leaves are pinnately compound with coarsely-serrate leaflets, flowers are white and small, fruit broadly ovate to oval and small. The European species has long been recognized as poisonous. Many cases of poisoning of man and lower animals by this plant are on record.

Cases of poisoning of children by Cowbane are not of infrequent occurrence, several being reported each year in the daily press.

The following item appeared in the Des Moines Register and Leader of May 23, 1909, and is but one of many that have come under the observation of the writer during the last twenty years.

Boone May 22.—"Virgil Hyatt, a high school boy, was poisoned last night while walking to the Ledger, a summer resort near Boone. He fell to the ground unconscious, and a companion carried him to a nearby farm house, and summoned medical aid and his mother from Boone. The trip was made in an automobile at record breaking speed. The boy was brought to a hospital here (Boone) but died just as he was being carried into the institution.

It seems that the young man, while walking across a plowed field with a companion picked up some of the weed whose roots had been exposed and ate freely of them. In an hour he complained of illness, and fell into the creek from which his companion rescued him. However, as reported above, all attempts to save his life were unavailing and he died from convulsions in a few hours.

As a sequel to the above, the daily papers of June 3, of the same year report a second death in the same locality from the same cause. In this case a young man who was sent to secure some of the weed for examination, became poisoned in some way, possibly by the juice of the plant coming in contact with some abrasion of the skin. The lad died after a short illness, having shown symptoms similar to those of the previous case.

Symptoms. The first symptoms are pain in the bowels, urging to ineffectual attempts to evacuation, burning in the stomach, nausea, vomiting, tetanic convulsions which may be severe resembling those produced by strychnin, or there may be coma without convulsions.

Dr. Hazeltine says that, in one case, he found the patient "showing convulsive agitations consisting of tremors, violent contractions and distortions with imperfect relaxations of the whole muscular system, astonishing mobility of the eyeballs and eyelids, with wide dilated pupils, frothing at the mouth and nose mixed with blood, and, occasionally, genuine, violent epilepsy."

The convulsive agitations were so violent that the pulse could not be examined with sufficient accuracy to determine its character. There is a profuse sweat. In fatal cases, the respiration is stertorous, the pulse small, and the face cyanotic. Not many cases among animals have been recorded. In a case

reported to the writer by an Iowa farmer, a cow which had eaten freely of the roots, fell into a spasm when brought into the barnyard. The animal, however, rose, walked one hundred feet and fell again, got up again, walked about thirty rods, fell and died in about thirty minutes. In a second case, a yearling owned by the same man had been in good healthy condition but began to display the same symptoms and died in twenty minutes.

Dr E. S. McCord, on September 31, of the same year, gave an old horse six drachms, hypodermically, of a strong decoction of the root. In fifteen minutes the animal showed uneasiness, pulse was full and fast, in a short time the animal laid down, and the pulse decreased, the horse was in great pain and kept moving the extremities, the pulse was weak but the patient finally recovered. The botanist of the Oregon Experiment Station found that the root has less of the toxic substance in the summer than in the winter and spring, which may account for the failure in this last case to produce death. In frogs, frequency of breathing is increased, tetanic convulsions follow, gradually paresis of the extremities, and lastly full paralysis and death. Cicutoxin,



FIG. 16a. Poison Hemlock (*Cicuta maculata*), native to Europe; naturalized in the U. S. (U. S. Dept. Agri.).

the characteristic poison of *Cicuta*, acts especially upon the medulla oblongata, the brain and spinal cord are merely secondary seats of its action.

Treatment. The stomach should be effectually evacuated by the use of the stomach pump or by a strong emetic. External and internal stimulants such as whisky should be applied; anaesthetics and narcotics used to control the spasms, hypodermic injection of morphia aids in recovery. It usually hap-

pena, however, that the veterinarian or physician is called too late to accomplish much.

The Poison Hemlock (*Conium maculatum*) is indigenous to Europe and has long been known as a poisonous plant. It is a fetid smelling herb from 2-3 feet high, with a spotted stem, compound leaves, and small, white flowers in umbels. The plant is not uncommon in waste places in the East and in the Rocky Mountains, especially in Utah. It has long been used as a poison.

Symptoms. In lower animals, there is observed a dilatation of the pupil, followed by weakness of the limbs, passing into paralysis, labored respiration, frequency of breathing diminished, heart action irregular, death preceded by convulsions. In man, there are weakness in the lower extremities, staggering gait, in two hours paralysis of upper and lower extremities and slight convulsions, death occurs in a few hours usually caused by cessation of respiration.

Treatment. The stomach should be evacuated by means of a pump or tube; or a hypodermic injection of 4-5 drops of a solution of apomorphin given, or emetics of sulphate of zinc or mustard administered. The temperature of the body should be kept up by hot applications.

Stimulants may be given, and, if necessary, artificial respiration applied. As a drink, strong tea, tannin, or any harmless vegetable decoction containing tannin may be administered.

CHAPTER VIII

FISH AND ARROW POISONS, HYDROCYANIC POISONING—TOXALBUMINS—BLACK LOCUST,
CASTOR OIL, AND JEQUILITY

Fish and arrow poisons have played an important part with the aborigines of all countries and they are still used to a considerable extent by primitive people. Thus Merrill¹ mentions the use of the *Antaria toxicaria* in the Philippines and other plants used in the same way which are being worked up by Dr. R. F. Bacon.

Radlkofer² some years ago published a long list of plants which are used to poison fish, and added a history of the earlier literature. He lists some 134 species which have been used in various parts of the world for this purpose and these plants belong to the following orders and genera. The species are listed under the poisonous species in another part of this work.

Dileniaceae (*Tetracera*) Menispermaceae (*Anamirta*, *Abuta*, *Pachygone*), Cruciferae (*Lepidium*), Labiatae (*Cleome*), Rutaceae (*Pangium*, *Hydnocarpus*), Ternstroemiaceae (*Lycopodium*), Tiliaceae (*Gratia*), Meliaceae (*Visnago*), Chaietaceae (*Chaietia*, *Tapura*), Rhamnaceae (*Gouania*), Sapindaceae (*Serjania*, *Poultonia*, *Lupinus*, *Dodonaea*, *Harporhiza*, *Mayonia*), Hippocastanaceae (*Pavia*), Leguminosae (*Tephrosia*, *Milletia*, *Orobanchia*, *Abrus*, *Centrosema*, *Crotalaria*, *Camptosema*, *Phaseolus*, *Lonchocarpus*, *Derris*, *Piscidia*, *Rottledia*, *Cassia*, *Bauhinia*, *Leucaena*, *Aibizma*), Myrtaceae (*Barringtonia*, *Gustavia*), Compositae (*Clibadium*, *Ichthyothere*), Campanulaceae (*Tupa*), Ericaceae (*Rhododendron*), Primulaceae (*Cyclamen*), Myrsinaceae (*Aegiceras*, *Jacquinia*), Sapotaceae (*Bastia*), Ebenaceae (*Phospyrus*), Apocynaceae (*Melodinus*, *Thaetelia*, *Cerbera*, *Aspidosperma*), Loganiaceae (*Buddleia*, *Strychnos*), Solanaceae (*Hyoscyamus*, *Nicotiana*), Scrophulariaceae (*Verbesum*, *Digitalis*), Bignoniaceae (*Bignonia*, *Tecoma*, *Jacaranda*), Labiatae (*Freemostachys*), Chenopodiaceae (*Chenopodium*), Polygonaceae (*Polygonum*), Aristolochiaceae (*Aristolochia*), Piperaceae (*Piper*), Thymelaeaceae (*Daphne*, *Winkstroemia*), Euphorbiaceae (*Euphorbia*, *Phyllanthus*, *Securinega*, *Piranhea*, *Croton*, *Joomana*, *Manihot*, *Jatropha*, *Excoecaria*, *Hura*), Comletae (*Tamus*), Liliaceae (*Veratrum*).

Ernst³ lists only sixty species that are used as fish poison. There must, however, be considerably more as indicated by Radlkofer.

W. M. I. Brost Pauwels⁴ in his contribution on the Surinamic fish poisoning⁵ contributes an interesting article on the subject.

Pauwels who made an investigation of Nekoe (*Lonchocarpus toxicus*) states that it is a powerfully toxic substance. He found that Nekoe will poison fish in proportion of 1,500,000, and that a second substance il-

¹ Philip Journ. of Sci. 2:111. Sect. C.

² Nov. Math. Phys. Class. k. k. Acad. d. Wiss. München, 1886, 179.

³ Memoire Bot. et Ficharbancaire o van la Pense per mediu de Plante venenose.

⁴ It is a tot de Kerk van der Surinamische Afschrijvingen. M. Greshoff has likewise published a number of works on fish poisoning plants.

⁵ Here and Swatters found in the *Piscidia* *Erythema piscidia* $C_{10}H_{16}O_6$ and Greshoff found in *Pachyrhizus angustatus* *pachyrhizus* $C_{10}H_{16}O_6$ (OC_2H_5).

Nekoesid will poison fish in proportion of 1-10,000,000. The poison will take effect in one hour. The water poisoned with the substance will cause the fish to make an effort to get away from the poison, they are in a horizontal position, breathe heavily, come to the surface of the water and try to jump out and finally breathing becomes increasingly difficult and at last they turn on their backs and die.

Under poisoning from cherry sorghum and a few other Hydrocyanic plants, an account has been given of poisoning from hydro-Poisoning. cyanic acid. It may be convenient to bring together some of the plants from which the very poisonous substance, prussic acid, has been obtained.

Maurus Greshoff of the Colonial Museum in Holland has taken the pains in a paper on Cyanogenesis to give the distribution of Prussic acid in the vegetable kingdom, the Hydrocyanic acid being found in a great many different plants. The following list gives the orders in which this substance occurs.

Ranunculaceae (*Aquilegia vulgaris*, *Thalictrum aquilegifolium*), Berberidaceae (*Nandina domestica*), Cruciferae (*Lepidium sativum*), Bixaceae (*Gynocardia odorata*, *Hydnocarpus venenata*, *Hygelaria africana*, *Pongium edule*, *Ryparosa castia*, *Taraktogenos Blumeri*, *Trichadenia seylanica*), Sterculiaceae (*Sterculia*), Tiliaceae (*Echinocarpus*), Linaceae (*Linum catharticum*), Rutaceae (*Citrus medica*), Dipsacales (*Chalcidha cymosa*), Olacaceae (*Ximenia americana*), Celastraceae (*Knerimia seylanica*), Rhamnaceae (*Rhamnus Frangula*), Sapindaceae (*Lupinus Schlenkeri*, *trifolium*), Anacardiaceae (*Corynocarpus laetigula*), Leguminosae Papilionaceae (*Lotus arabicus*, *Indigofera galegoides*, *Phaseolus lunatus*, *Luna sativa*, *Dalman*, *Lobelia*), Rosaceae (*Amelanchier vulgaris*, *Chamaecrista*, *Cotoneaster integerrima*, *Crataegus Oxyacantha*, *Eriobotrya japonica*, *Nuttallia cerasifera*, *Ostrya*, *Phytol*, *Pyrus*, *Prunus Amygdalus*, *Pygeum africanum*, *Spiraea Aruncus*), Saxifragaceae (*Ribes aureum*), Combretaceae (*Combretum constrictum*), Myrtaceae (*Myrtidium montanum*), Melastomaceae (*Memecylon*), Samydicaceae (*Homalium*), Passifloraceae (*Passiflora quadrangularis*, *Tecoma*), Caprifoliaceae (*Sambucus nigra*), Rubiaceae (*Plectranthus discolor*), Compositae (*Chardonia xeranthemoides*, *Xeranthemum annuum*), Sapotaceae (*Isomandra*, *Lacuna bonplandii*, *Payson latifolia*), Asclepiadaceae (*Gynemna latifolia*), Convolvulaceae (*Ipomoea disticta*), Bignoniaceae (*Diospyros nocturna*), Euphorbiaceae (*Euphorbia avata*, *Latexospermum*, *Tupia*, *Hevea brasiliensis*, *Jatropha angustifolia*, *Mimosa cathartica*, *Ricinus communis*), Urticaceae (*Sporus virgata*), Araceae (*Arum maculatum*, *Colocasia gigantea*, *Cyrtosperma latifolia*, *Lasia aculeata*), Gramineae (*Glyceria aquatica*, *Panicum*, *Sorghum vulgare*, *Stipa hystrix*), Fungi (*Hygrospora aquosissima*, *Mucorinus oroides*, *Phallotia radicata*, *Rustia foetida*). He makes the following statement with regard to the presence of this substance in plants.

"Many plant physiologists in Europe, with more experience with *Prunus* or *amygdalin* than with the tropical *Pongium*, incline to the view that hydrocyanic acid in these plants has nothing to do with either the building-up or the breaking-down of proteids, but that this substance is made by the plant from sugar and nitrate by a special process, and serves no other purpose than to defend the plants against the attacks of animals. It is above all the incompleteness of our physiological knowledge which makes decision between these theories difficult.

"In the study of this question it is important to remember the possible diversity of origin of this body, and every cyanogenetic plant will be required to be examined on the lines laid down by Treub."

The wide distribution of glucosides that yield hydrocyanic acid is evident from the list above. Dunstan and Henry¹ discovered three glucosides, *dhurria* $C_{17}H_{27}O_7N$ in the common sorghum, *lotus* $C_{22}H_{31}O_8N$ in a species of lotus of Egypt and *pharro-luathin* $C_{19}H_{27}O_8N$ in wild beans of *Phaseolus lunatus*, the common lima bean. Brumich² attributed death from the feeding of immature sorghum to *dhurria*. Power and Lees³ isolated from the seeds of *Gynocardia odorata* a glucoside to which they gave the name *gynocardin* $C_{13}H_{19}O_9N$. All of the above glucosides yield on hydrolysis, hydrocyanic acid. The most important and best known of all the glucosides that yield hydrocyanic acid is *amygdalin*.

Creshoff discovered an amygdalin-like glucoside in two tropical trees, *Pygeum parviflorum* and *P. latifolium*. The same author found glucosides in a member of the milkweed family *Asclepiadaceae*. The *Pongium edule* of the tropics contains a large quantity of a glucoside capable of being converted into hydrocyanic acid and a large amount can be prepared from a single plant. The *Hydnocarpus rubrians* also contains a large quantity of a glucoside which yields hydrocyanic acid. It is used to destroy fish. The common linseed cake contains a glucoside which yields prussic acid. Francis found prussic acid in the sweet cassava root 0.68 per cent, and in the bitter cassava 0.275 per cent. It is well known that fresh bitter cassava root is bitter poison. The above facts are brought together by Blyth in his work on poisons and may be consulted for more of the details.

The statistics on poisoning seem to indicate that it occupies third place among poisons in the order of frequency in Great Britain. In that country there are about forty deaths annually from this poison according to Blyth. It is responsible for the loss of a great many cattle in sections of the country where the wild cherries are abundant and also from sorghum poisoning. It is frequently used for criminal poisoning, at one time more frequently than now. It is nearly always taken by the mouth into the stomach, but occasionally the vapors produce death. It is generally used by Entomologists to kill insects.

Blyth gives the symptoms of poisoning as follows: Cold blooded animals require a larger relative dose than warm blooded animals except the birds which are slightly less sensitive but the action is essentially the same. Hydrocyanic acid acts in two ways:

- 1 It profoundly interferes in the ordinary metabolic changes in animals.
- 2 It causes a paralysis of the nerve centers.

Normal blood decomposes with great ease hydrogen peroxide into oxygen and water. If it is normal venous blood and a little hydrogen is added it becomes bright red, but if a trace of prussic acid be present it is a dark brown color.

The blood corpuscles lose their power of conveying oxygen to all parts of the system and asphyxiation results. The main symptoms in animals are as follows:

The main differences between the symptoms induced in cold-blooded and warm-blooded animals, by a fatal dose of hydric cyanide, are as follows:

The respiration in frogs is at first somewhat dyspnoic, then much slowed, and at length it ceases. The heart, at first slowed, later contracts irregularly, and at length gradually

¹ Proc. Roy. Soc. lxviii and lxix (See Blyth "Poisons" p. 204).

² *Ib.*, lxxvii. (See Blyth "Poisons" p. 204)

³ Journ. Chem. Soc. lxxix. (See Blyth "Poisons" p. 204)

stops, but it may continue to beat for several minutes after the respiration has ceased. But all these progressive symptoms are without convulsion. Among warm-blooded animals, on the contrary, convulsions are constant, and the sequence of the symptoms dyspnoea, slowing of the pulse, giddiness, falling down, then convulsions with expulsion of the urine and faeces.

When the dose is short of a fatal one, the symptoms are as follows. Evident giddiness and distress, the tongue is protruded, the breath is taken in short, hurried gasps, there is no action, and convulsions rapidly set in, preceded, it may be, by a cry. The convulsions pass into paralysis and insensibility. After remaining in this state some time, the animal again wakes up, as it were, very often howls, and is again convulsed, finally, it sinks into a deep sleep, and wakes up well.

Dr K Winslow in his work on Veterinary Materia Medica and Therapeutica, gives the Toxicology of Hydrocyanic Acid as follows. "Three stages may be distinguished in fatal poisoning. First, a very short period elapses before the symptoms appear. There are giddiness, difficult breathing, and slow pulse in this stage. Second, the pupils dilate, vomiting may occur, and the animal utters loud cries. Spasmodic defecation, micturition and erections may be present, with convulsions and unconsciousness. Third, the last stage is characterized by collapse, spasms, general paralysis and death. The subacute form of poisoning may ensue and prove fatal, or, owing to the volatile character of the drug, complete recovery may take place within one-half or three-quarters of an hour. Occasionally dogs continue to be paralyzed for several days and get well. The minimum fatal dose recorded in man is 9/10 of a grain of pure acid, or about 50 drops of the medicinal solution. Four to five drachms of the diluted acid frequently, but not invariably, causes subacute poisoning and death in horses, within an hour. One or two drachms of the pharmacopoeial preparation usually kills dogs within ten minutes."

Poisoning from Toxalbumins Black Locust Ricinus and Abrus

In recent years much work has been done with a class of poisons, known as toxalbumins. These are of especial interest because many of the bacteria produce such poisons. Some of the fungi responsible for "forage poisoning" produce, it is thought, toxalbumins. In recent years a number of cases of horse poisoning from Black Locust bark have been reported. The poisoning from castor oil bean (*Ricinus*) and from *Abrus* are also of this class.

Castor Oil Seed and Abrus. One of the best known of the toxalbumins is that occurring in the castor oil seed, known as ricin. This albuminous substance is very poisonous, more so than strychnin and prussic acid. Ricin coagulates the blood. Blyth in his work on poisons states

If castor-oil seeds are eaten, a portion of the poison is destroyed by the digestive processes, a part is not thus destroyed, but is absorbed, and produces in the blood vessels its coagulating property. Where this takes place, ulcers naturally form, because isolated small areas are deprived of their blood supply. These areas thus becoming dead, may be digested by the gastric or intestinal fluids, and thus, weeks after, death may be produced. The symptoms noted are nausea, vomiting, colic, diarrhoea, tenesmus, thirst, hot skin, frequent pulse, sweats, headache, and on and death is convulsions or from exhaustion. Animals may be made immune by feeding them carefully with small doses, gradually increased.

The post-mortem appearances are ulceration in the stomach and intestines. In animals the appearances of hæmorrhagic gastro-enteritis with diffuse nephritis, hæmorrhages in the mesentery, and so forth have been found.

A toxalbumin also occurs in the Jequrity seed (*Abrus precatorius*) which causes similar effects and symptoms. That the poisons are not the same have been shown by experiments with animals. It is known that animals may become immune by repeated doses of Jequrity against *abrus* and the principle of castor oil does not produce immunity against *abrus*, nor does *abrus* confer immunity against the ricin of the castor oil bean. The *abrus* when applied to the con-



Fig. 16b Castor oil plant (*Ricinus communis*). The seeds furnish the castor oil of commerce, and also contain an acrid poison. (Schubert, U. S. Dept. Agr.)

before. This was a sapling of about four inches in diameter and had probably made a very rapid growth and the bark, from this reason was tender and easily peeled. They had done a good job of peeling but as they had their bits in they were not able to swallow much. The poison obtained from this bark is, in my opinion, the cause of the trouble.

Dr. Waldron says that the symptoms otherwise are about the same as those occurring in cases of belladonna poisoning and are about as follows:

Extreme lassitude, which includes almost imperceptible pulse and which, when found, is weak and prolonged, respiration less than normal by one-third and sonorous, temperature normal, no pain, no appetite, mucous membrane congested, of a blue rusty, or yellow color. Mucous membrane of the mouth some swollen, caused by the congestion of the capillaries, slight pyrexia, and above all, the dilation of the pupil of the eye, in fact I should judge, we have nearly the same symptoms we get in belladonna poison.

He also says that he had not known before that locust bark was poisonous and although he had searched for literature upon the subject had found but one reference, that being in the U. S. Pharmacopoeia, which records a case reported in Jan., 1887, when 33 children were said to have been poisoned by chewing locust bark. In mild cases there were "flushed faces, dryness of the throat and mouth, and dilation of the pupils. In severe cases, were added epigastric pain, extremely intermittent heartbeats, and stupor." It is evident from the fact that there is not much literature on the subject that such poisoning does not occur often.

Dr. H. S. Murphy has kindly contributed the following case on locust bark poisoning:

Anamnesis: Grey mare twelve years old, pregnant ten months, has been at light work

junction causes coagulation in the vessels and a secondary inflammation. The disease is known as Jequirity ophthalmia. More details in regard to the poisons of these plants are given under the plants of the families in which they occur.

Black Locust Poisoning. The Black Locust which is commonly planted as an ornamental tree has in a number of instances caused death.

Dr. Waldron in the American Veterinary Review, writes thus of locust bark poisoning, referring especially to the beating of the heart of a horse that had been poisoned by the locust bark. This beating shook the horse and could be heard outside the stable.

The sound was caused by the action of the diaphragm. It was greater when the ribs were at their fullest expansion and could be heard most distinctly at a distance of ten feet. I tried to locate or rather find out what produced the sound, but in that I am as ignorant as I was then. . . In questioning where the team was hitched at the mill, it was found that the driver had tied them to a young locust tree that had been cut down a few years

continuously, is driven to town afternoon of March 11, and tied to a locust tree which is 6-8 inches in diameter; owner noticed that animal had pecked bark from tree also noticed animal pant on road home. Anorexia and stupidity were all owner noticed until 8 p. m. when a faint noise was heard this gradually grew worse until I arrived at 7 a. m. April 1.

Symptoms: 1. Animal stands back from manger with legs well apart resting nose and mouth on floor, most of the time, a part of the time head is held 18-24 inches from floor.

2. A continual "thud" is heard which is not synchronous with respiration but is synchronous with a vibration of thorax.

3. Animal will not obey commands and when pushed over in stall loses balance and nearly falls.

4. Fetus is seen kicking in mare's left hypo-chondriac region.

5. Temperature 99 C Perb.

6. Pulse 54 full, bounding.

7 Respiration, 14, very feeble.



Fig. 14c. Black Locust (*Robinia Pseud-acacia*) Bark, leaves and flowers poisonous. (After Paget).

8. Mucous membranes dry, very yellow a dirty yellow (in distinction from bright yellow of icterus) swollen, only moderately sensitive (probably due to general depression)

9. Intense dilation of pupil.

10. No evacuation of feces; auscultation reveals gurgles of bowels, ropy urine is passed.

11. Internal cause of thud was determined as due to heart beat in the following manner

1. Pulse over maxillary artery is synchronous with "thud" and thoracic vibration. 2. Owner counts aloud while auscultation of heart is conducted and heart beat is synchronous with thud. 3. Auscultation of heart and radial pulse are synchronous, also heart sounds are muffled but very loud. (Palpitation).

Treatment: (Symptomatic) $\frac{1}{4}$ gr. of Digitalin (P. D.) every fifteen minutes subcutan-

ously until four doses are given, also $\frac{1}{2}$ potassi iodidum is given per os at once and the following prescription left.

Rx Strychnini sulphatis gr. IV

Fid. ext. digitalis $\frac{1}{2}$ l.

Potassi Iodidum II

Aqua $\frac{1}{2}$ XVI

Sig. A table spoon full every 2 hours.

Result: After the second hypodermic was given a slight improvement noticed (lessening in force and frequency of thud, which was quite marked after fourth hypodermic. Owner reports thud gone at end of 24 hours but stupor still present though not so marked.

Recovery uneventful and at end of gestation parturition proceeds normally, and foal born.

Toxicity of bark proven in the following manner. Two cc of 10% tincture (bark grated) killed half grown kittens which showed above symptoms in an aggravated form.

One ounce necessary to kill a 25 pound dog.

This tincture dropped in eye acted much as atropin but no toxic symptoms developed. Simply a dilation of pupil. Grated root inlard produced slight symptoms in two dogs. P. m. on cats revealed a generalized dirty so. m. but only a very slight yellow on cartilages. Swollen liver and a few petechia on serous membranes. Blood quite dark.

The cadaver, resembled somewhat that of one dead of septicaemia.

CHAPTER IX

POISONING FROM OPIUM, SOLANACEAE AND PLANTS THAT CONTAIN SAPONINE.

The use of opium by Chinese and other races is as Fluckiger and Hanbury say "in the words of Pereira, the most important and valuable used in medicine of the whole *Materia Medica*, and we may add, the source by its judicious employment of more happiness by mankind." Blyth in his work on Poisons, states that in England and Wales 1905 deaths were attributed to the use of opium or its active constituents between the years 1898-1903. Of these 882 were accidental or because of negligence, 621 were suicidal. In France opium and morphin poisons are said to cause about 1 per cent of the cases of poisoning. Various patent medicines contain opium or some of its products and in the past have been the cause of frequent cases of poisoning. The use of the drug in patent medicines for children in the United States in the form of soothing syrups was once more common than now. The use of opium for infants is a common practice in India, according to Blyth who quotes from Dr. Chevers.¹

In general the opium and morphin poisoning are as follows. The beats of the heart are at first accelerated and then diminished. Large doses introduced into the circulation diminish the pulsations without acceleration and may even cause heart paralysis. "The arterial blood pressure, at first increased is afterwards diminished. If morphin is in sufficient quantity thrown into the circulation, then tetanus at once occurs. Depression and stimulation depend on dosage. The common form occurring in 99 per cent of the cases, excitement, narcosis, and coma, bowels nearly always constipated. (2) A very sudden form in which death occurs rapidly, the person sinks into a deep sleep almost immediately. (2) An abnormal form in which there is no coma but convulsions.

Blyth in referring to opium eating says:

The consumption of opium is a very ancient practice among Eastern nations, and the picture, drawn by novelist and traveler, of poor, dried-up, yellow mortals addicted to this vice, with their faculties torpid, their skin hanging in wrinkles on their wasted bodies, the conjunctives tinged with bile, the bowels so inactive that there is scarcely an excretion in the course of a week, the mental faculties verging on idiosyncrasy and imbecility, is only true of a percentage of those who are addicted to the habit.

In the case of opium poisoning the stomach tube should be used to empty the stomach, and wash with warm water, then coffee may be given. Permanganate of potash is a perfect antidote and should be given when at hand.

The alkaloid codein also found in opium produces sleep but its effects are different. Large enough doses produce death and cause epileptiform convulsions. Thebain found in opium produces symptoms that resemble those produced by strychnin, namely tetanic spasms. Apomorphin found in opium is an active emetic. Papaverin causes paralysis of respiration in guinea pigs.

¹ Jurisprudence 122 (3rd ed.).




FIG. 164. Garden Poppy (*Papaver rhoeas*). A poisonous plant producing a large number of alkaloids like morphine, codeine, etc. (Strasburger, Vohl, Schenck and Schimper).

Poisoning from Solanaceae

A number of plants of the Solanaceae are known to be poisonous; among them the common thorn-apple or Jimson weed (*Datura Stramonium*), the atropa (*Atropa Belladonna*) and hyoscyamin (*Hyoscyamus niger*) besides such suspected plants as the common black nightshade (*Solanum nigrum*), horse nettle (*Solanum carolinense*), bittersweet (*Solanum dulcamara*) and scopolia. The cases of poisoning from atropin are more frequent, perhaps, than statistics seem to indicate. The English death statistics for ten years, ending 1903, according to Blyth show 95 per cent of the deaths from atropin, 35 per cent were suicidal. Most of the accidental cases arise from mistakes made by the pharmacist or physician. Criminal poisoning is carried on to a less extent in Europe and America than in India. Blyth states that of the 120 cases recorded in works on Indian toxicology no less than 63 per cent were criminals, 19 per cent suicidal, and 18 per cent accidental. The most important alkaloids found are atropin, hyoscyamin, scopolamin and solanin. Solanin is poisonous and is regarded as a nitrogenised glucoside. In man the symptoms of atropin poisoning are Dilating of the pupils, dryness of the mouth and throat; the

mucous membrane is reddened, inability to swallow, deranged vision, breath at first a little slow and then rapid, the nervous system is affected in a marked degree the lower extremities are often partly paralyzed. There is want of coördination. "The person reels like a drunken man." In adults this takes on a hilarious pleasing form. The symptoms of poisoning from hyoscyamin are similar to those of atropin. The absence of delirium and excitement, however, makes it decidedly different. The symptoms in animals for both of these substances do not differ essentially from those given above



Fig. 165. Nightshade (*Solanum nigrum*)
From Darlington's Weeds and Useful Plants.



Fig. 161. Jimson Weed (*Datura Stramonium*)
a, leaf and flowers b, fruiting capsule.
(U. S. Dept. Agri.)

Atropin may be absorbed by the skin, and enough may be absorbed, if it is broken, to cause death. Blyth quotes Ploss¹ to the effect that atropin sulfate applied as an ointment to the abraded skin was fatal. Atropin has also been absorbed from the bowel as recorded by Blyth.

A elyater containing the active principles of 5.2 grms. (80 grains) of belladonna root was administered to a woman 27 years of age, and caused death. Allowing the root to have been carefully dried, and to contain .21 per cent of alkaloid, it would seem that so little as 10.9 mgrms. (.16 grains) may even prove fatal, if left in contact with the intestinal mucous membrane. Belladonna berries and stramonium leaves and seeds are eaten occasionally by children. A remarkable series of poisoning by belladonna berries occurred in London during the autumn of 1846.

¹ Zeitschr. f. Chir. 1863. Blyth, Poisons, Their Effects and Detection.

Poisoning from Plants that contain Saponin.

In recent years our knowledge of the Saponins has been greatly extended; many of these studies have been made by Kobert or his students in the laboratory at Dorpat. The term saponin has been applied to a class of substances of a glucosidal nature which are poisons and when dissolved in water form a solution which froths much like soap-suds. These substances are not all the same chemically, but have the general formula $C^xH^{2x}O^{10}$. Blyth gives the following list with their formulae

Saponin 1.

Saponin seegeni	}	$C_{12}H_{24}O_{10}$
Quillaja saponin		
Sapindus saponin		
Gypsophila saponin		
Agrostemma saponin		

Saponin 2

Asamin	}	$C_{18}H_{36}O_{10}$
Digitonin saponin		

Saponin 3.

Quilagic acid	}	$C_{18}H_{36}O_{10}$
Polygalic acid		
Herniaria saponin		
Cyclamin	}	$C_{20}H_{40}O_{10}$
Sarsaparilla saponin		

Sarsa saponin. $C_{22}H_{44}O_{10}$

Parillin $C_{24}H_{48}O_{10}$

Melanthin $C_{26}H_{52}O_{10}$

The suggestion is made that possibly ducanarin $C_{22}H_{44}O_{10}$ and syringin $C_{18}H_{36}O_{10}$ may belong to the same series.

One of the oldest of the known saponins was isolated from the Bouncing Betty, *Saponaria officinalis*, and later from the corn cockle *Agrostemma Githago* and many other plants. This saponin is a white amorphous powder, very soluble in water, is neutral and reacts without odor, it causes sneezing when applied to the mucous membrane of the nose; tastes at first sweetish, then becomes sharp and acid. The saponin when rubbed on the skin exerts no action because not absorbed, when injected subcutaneously into frogs it becomes quickly absorbed and acts upon the nerves and muscles. In warm blooded animals there is little or no absorption because of an aseptic abscess which forms. Intravenous injections in small amounts in the laboratory of Kobert proved fatal for cats and dogs. It acts injuriously on the striated muscle and heart muscle. The sensor and motor nerve fibers are also affected in a serious way. On the digestive tract it causes inflammation and peristalsis. The saponin substances dissolve the blood corpuscles of all animals and thus penetrate the corpuscles. It is thought that the hemolytic action of these substances is due to the liquefaction of the cell membrane.

Ransom found that the saponin may become bound to the corpuscles and the serum. That this action depends on the cholesterol, saponin so bound will not act on the red corpuscles. The saponin cholesterol mixture exerts no action on dog's blood.¹ Dr R. F. Bacon and H. T. Marshall who made a study

¹ The toxic action of saponin. Phil Jour Sci. 1 1937 Dr R. F. Bacon and H. T. Marshall.

of the saponin found in *Entada scandens* proved that it was highly toxic for rabbits and guinea pigs.

When diluted with normal salt solution to a concentrated form 1:200 and injected into the peritoneal cavity 2-3 mgrms. of saponin to 100 grms. of animal was uniformly fatal, while a quarter of this amount killed in several cases. Where smaller quantities were used and animals living for a longer period of time localized peritonitis was found.

"The saponin is powerfully haemolytic" 0.005 cubic cm. of $\frac{1}{2}$ of 1% solution (0.025 mgrms.) completely dissolved one cubic cm. of a 5% suspension of rabbit corpuscles which were freed from serum and the half of this amount produced haemolysis of the serum of the free corpuscles of the guinea pig. "Saponin, however, loses its haemolytic power after the addition of serum." Immunity could not be produced in rabbits or guinea pigs recently treated with intraperitoneal doses of saponin or saponin serum mixture.

Blyth has studied the general action of saponin on kittens. He states that when 13 to 22 mm. ($\frac{1}{5}$ to $\frac{1}{2}$ gr.) is injected underneath the skin of a kitten immediately symptoms of local pain occur, in 5 to 10 minutes the respiration is quicker and the animal falls into a lethargic condition with signs of muscular weakness, just before death breathing became rapid with all the signs of asphyxia. The appearances after death were fullness in the right side of the heart and congestion of the intestinal canal. In man the taking of saponin causes an increase of mucus secretion and nausea.

Saponin or saponin-like substances occur in the following families:

Phytolaccaceae, (*Phytolacca abyssinica*), *Caryophyllaceae*, (*Gypsophila struthium*, *Agrostemma Githago* *Lychnis Saponaria officinalis*, *S. rubra*, *Hemeris*); *Berberidaceae*, (*Coniophyllum thalictroides*) *Leguminosae*, (*Entada scandens*, *Gymnocladus dioica* *Gleditsia*, *Enterolobium*) *Oleaceae* (*Chaenactis virginica*, *Syringa vulgaris*), *Amaryllidaceae*, (*Asape*) *Liliaceae* (*Yucca glauca*? *Chlorophatum pomeridianum*), *Rosaceae*, (*Quillaja Saponaria*), *Sapindaceae*, (*Sapindus trifolatus*, *S. saponaria*, *Pometia pinnata*, *Magonia*), *Hippocastanaceae*, (*Aescula Hippocastanum*, *A. Pavia*), *Theaceae*, (*Thea Sasanqua*, *T. asamica*), *Sapotaceae*, (*Omphalocarpum procerum*), *Polygalaceae* (*Polygala Senega*).

In all more than 200 species of plants contain saponin.

See Chapter XIV, and Pl. II, for a list of others.

CHAPTER X

POISONING FROM FLOWERS, POISONING FROM HONEY, MECHANICAL INJURIES

Poisoning from Flowers. The odors from a large number of flowers are more or less injurious, especially to some individuals. The flowers of the common wild black cherry (*Prunus serotina*) when past their prime give off a cyanogenic odor that is quite objectionable to many people, causing headache. The carion-like odors, like those produced by *Stapeia*, the carion flower (similar herbaceous) and *Aristolochia* are sickening to many people, causing headache and a depressing feeling. Many flowers that are ordinarily sweet scented like the tuberose (*Polyanthes tuberosa*) often give people the headache if the flowers are abundant in the room they are sitting in. The flowers of Wistaria are injurious to some people. The flowers of *Magnolia grandiflora* are "overnutering" according to some authorities.

Poisoning from Honey. The honey obtained from the flowers of some plants is said to be poisonous, for instance the honey collected by insects from the oleander was long ago recorded as being injurious, and it is said that the honey collected from the mountain laurel (*Kalmia latifolia*) appears to be poisonous under some conditions according to Chesnut. According to Chesnut the honey collected from the flowers of the snow-on-the-mountain, (*Euphorbia marginata*) is bitter and disagreeable, but does not appear to be a serious poison. Several cases of poisoning from the nectar of the flowers of *Datura metel* and *D. Wrightii* are on record and the flowers of the Brazilian magnolia of the family *Sapindaceae* produces poisonous honey, also the flowers of *Rhododendron* are said to contain andromedotoxin. Toxic honey has also been gathered from the flowers of Black Locust (*Robinia pseudo-acacia*) and Lily of the Valley (*Convallaria majalis*).

Prof. Lyman F. Kelder who has made a somewhat extended investigation with poisonous honey¹ has given an excellent bibliography with reference to the earlier literature on the subject. It has been known for centuries that the honey collected from *Eriacaceae* acts as a narcotic irritant, producing giddiness, vomiting, and purging. Poisonous honey was described by Xenophon. He gives a fairly accurate description of how the soldiers of his army acted that ate honey that was poisoned. He states that they lost their senses, vomited and were afflicted with purging, and those who had eaten but little were intoxicated, but when they had eaten much they were like mad men. Strabo and Pliny spoke of poisonous honey the latter writer, an early naturalist noted for his accurate observations, records poisonous honey which he called "aegolethron" (goat's death) which bees collected at Hieraclea. He gives a description of the honey which is said to have had a peculiar smell and produced sneezing. It is generally supposed that this honey came from a species of *Rhododendron*, the *R. pontica*. This and allied species are the chief source of poisonous honey in Asia and Asia Minor.

¹ Poisonous Honey. Proc Amer Pharm Assoc 1896 167-173.

but it may be said in this connection that honey collected from the Heather in Scotland is not poisonous.

Barton, an early American botanist, reported poisonous honey in New Jersey as early as 1794. Subsequently Coleman reported a large number of cases in 1852, and Gammer in *Gleanings in Bee Culture* and several writers in the *American Bee Journal* reported poisonous honey. Other writers like Cheesnut and Crawford have reported on the occurrence of poisonous honey in the United States and Kebler reports no less than eight cases for New Jersey in 1896 and believes that it is much more common than the records seem to indicate. Kebler was fortunate enough to investigate some of the poisonous honey following a case of poisoning in New Jersey. He examined a part of the comb of the dark honey which had a light brown color and a nauseating odor, pungent taste, caused a burning sensation in the back of the mouth similar to that of acetic. Persons who partook a small amount of this honey began to cough immediately. He also made a chemical analysis of the now suspicious honey digesting with alcohol then evaporated, the residue was again treated to alcohol and evaporated and administered to two cats. One received a small dose and the other a larger one. The results from the cats were so interesting that I quote from Prof. Kebler¹

The small dose produced partial relaxation, relaxation of the voluntary muscles and general depression. The large dose in a short time produced restlessness, vomiting, purging, prostration and almost complete loss of the voluntary muscles, showing that the honey contained a prompt and potent poison. The animal could scarcely be induced to move, and when motion was attempted, first the forelimbs would fail, and then the back limbs would give way. First one portion of the body would sway in one direction, then the other portion in another, reminding one of a highly intoxicated person. Had the entire dose been retained, death undoubtedly would have followed. As it was, the cat had regained her normal condition only at the end of twenty-four hours.

Along with this we may append the symptoms as recorded by the physician who attended the persons who were poisoned.

Mr. and Mrs. Chambers took but a small quantity, yet each noticed a peculiar pungent, burning taste in the comb as soon as it had passed their lips. In fifteen or twenty minutes afterwards, Mrs. C. was taken with nausea, abdominal pain, and vomiting, soon followed by loss of consciousness, coldness of extremities, feebly acting heart, and complete collapse. While ministering to her, Mr. Chambers, who had also experienced the initiatory symptoms of pain and nausea, suddenly exclaimed, "I cannot see," and soon sank to a state of syncope to the floor. In each case the symptoms were similar. Retching, vomiting, purging, acute gastric and abdominal pain, and continued cramps for some hours, with surface redness, and deadly pallor, and the general symptoms of collapse.

Kebler was, however, unable to definitely locate the andromedotoxin found by Plugge.² This author recorded it for a large number of plants as follows: *Andromeda japonica*, *A. polifolia*, *A. Coteshana*, *A. calyculata*, *Asimina latifolia*, *Menotropa uniflora*, *Pieris formosa*, *P. ovalifolia*, *Rhododendron grande*, *R. barbatum*, and *R. fulgens*. It has been recorded for additional plants by Greshoff who mentions the following plants which produce poisonous honey, *Nerium oleander*, *Cytisus Laburnum*, *Pieris acutifolia*, *Callitriche procera*, *Daphne*, *Ponthea*, *Buxus laevis*, *Clerodendron serratum*, *C. Rhomantarium*, *Sapindus emarginatus*. (It is said that thousands of bees are killed by this honey.) *Centaurea scabiosa*, *Carduus marianus*, *Scutellaria succisa*. A South African species of *Euphorbia* also produces a poisonous honey which was not noted by Greshoff.

¹ Proc. Amer. Pharmac. Assoc., 1896:170.

² Arch. & Pharm., 29 - 2 1891. Am. Jour. Pharm. 63 603. Shresh, J. C. Notes on Trebizonde Honey. Pharm. Jour. and Trans., 16 397, 404, 1867-68.



Fig. 16g. The flowers of Black Locust (*Robinia pseudo-acacia*) produces a toxic nectar. (Ada Hayden).



Fig. 16h. The wilted and old flowers of Wild Cherry (*Prunus*) produce a cyanogenic poisonous odor. (Ada Hayden)

We have a long list of plants that act injuriously **Mechanical Injuries** in a mechanical way. Among the best known of these are Wild Barley or Squirrel-tail Grass (*Hordeum jubatum*) and the related species, which, by mechanical means, injure sheep, horses, and cattle. The awned heads, when eaten with hay or grass, break up into sections, the awns working their way into the mucous membrane, insinuating themselves around the teeth, thus causing inflammation and deep ulcerating sores, with the formation of pus. The teeth may consequently become loosened and fall out.

A Cheat or Brome Grass (*Bromus tectorum*) which is common in Utah, parts of Colorado, and westward, produces similar injuries.

The Needle Grass, common in the dry gravelly hills and sandy plains of Northern Mississippi Valley has a fruit with a sharp pointed callus, with hairs above the pointed callus projecting upwards. The sharp-pointed callus of the "seed" enters the skin, especially of sheep, where it produces an irritation which is sometimes followed by death. This has sometimes been eaten with forage, thus entering the intestinal tract, perhaps perforating it, causing death, especially when they have pierced the walls of the intestines. Several allied species as Black Oat (*Stipa arenastra*) produce similar injuries. The Western Needle Grass (*Stipa comata*), however, is less injurious than our Common Needle Grass.



Fig. 161. Wild Barley (*Hordeum jubatum*). b. Wild Barley (*H. nodosum*) cause of mechanical injuries to animals.

Similar and allied species in other parts of the world are known to be injurious in the same way. Other plants are injurious by means of their sharp-pointed fruit. Among those of the last named type are members of the Geranium family notably the Stork's bill (*Erodium cicutarium*), common in the west, which frequently gets into the wool of sheep and may produce local irritation. Mechanical injuries are also produced by the Sand Bur (*Cenchrus tribuloides*) whose spiny involucre may work into the flesh of animals and induce an irritation followed by inflammation, and the formation of pus. Hogs and cattle often receive mechanical injuries from the Cocklebur (*Xanthum canadense*) and allied species.

A recent number of Horticulture (Jan. 1, 1910) notes that rose thorns have caused the poisoning of the hands of clerks who handle the roses.

Prof. J. Dávy states that in South Africa the burrs of a clover *Trifolium terrestris* var. *hispidissimus* are said to injure young lambs.

The Burdock (*Lappa major*) sometimes produces no little irritation and the same may be said of the Spanish Needle (*Bidens frondosa*). The sharp, stiff




Fig. 17 A. Fruit of the Cocklebur (*Xanthium condensae*), causes mechanical injuries to dogs when eaten by them. B. Sandbar (*Cenchrus tribuloides*). C. Burdock (*Arctium major*). D. Boat Jack (*Bidens frondosa*). E. One of the Borages (*Cynoglossum*). F. Carrot (*Daucus Carota*).

branches of Greasewood (*Sarcobatus vermiculatus*) easily penetrate the skin and sometimes induce pus infection. The bristles and spines of the rose and stiff bents of the Russian Thistle may be the cause of injury to animals and men. The small, barbed trichomes of the calyx of Crimson Clover (*Trifolium incarnatum*), according to Prof. Coville, produce phytobezoars similar to those commonly produced by hair. Dr Trelease has described similar bezoars from the barbed trichomes and spines of cacti. Millet and barley awns are known to produce similar balls in horses as well as in cattle. Corn stalks, when eaten with an insufficient amount of water, produce impaction. This has, however, been attributed to other causes also. Bloat is known to follow the use of such forage as white and red clover, alfalfa, pigweed and many other plants consumed under the same conditions.




Fig. 17a. Fresh water cord grass (*Spartina cynosuroides*) with sharp leaves often injurious. 17b. Sand Bur (*Cenchrus tribuloides*) U. S. Dept. Agrl.




Fig. 18. Needle Grass (*Stipa comata*). U. S. Dept. Agrl. Fig. 18a. Needle Grass (*Stipa spartea*, inflicting mechanical injuries. U. S. Dept. Agrl.




Fig. 18d. Urticating hairs and cutting leaves. a. Urticating hair of nettle. b. Bristles of hollyhock. c. barbed margin of a leaf of hedge. d. barbed margin of a leaf of grass.

CHAPTER XI

CLASSIFICATION OF POISONS, SYMPTOMS AND ANTIDOTES

Blyth classifies poisons as follows:

A. POISONS CAUSING DEATH IMMEDIATELY OR IN A FEW MINUTES. Prussic acid, cyanides, oxalic acid and occasionally, strychnia.

B. IRRITANT POISONS. Symptoms mainly pain, vomiting, and purging. Savin, ergot, digitalis, colchicum, yew, laburnum, and putrid substances.

C. IRRITANT AND NARCOTIC POISONS. Symptoms of an irritant nature, with more or less cerebral indications. Oxalic acid or oxalates.

D. POISONS MORE ESPECIALLY AFFECTING THE NERVOUS SYSTEM

1. Narcotics. Chief symptoms insensibility which may be preceded by more or less cerebral excitement. Opium.

2. Delirants. Delirium, for the most part, a prominent symptom: Belladonna, hyoscyamus, stramonium, and other Solanaceae, poisonous fungi, Indian hemp (*Cannabis*), darnel (*Lolium temulentum*), camphor and *Oenanthe crocata*.

3. Convulsives. Almost every poison has been known to produce convulsive effects, but the only true convulsive poisons are the alkaloids of the strychnin class.

4. Complex nervous phenomena. Aconite, digitalis, poison hemlock, Calabar bean, tobacco, *Lobelia inflata*, and curare.

Kobert's classification is as follows:-

I. POISONS WHICH CAUSE COARSE ANATOMICAL CHANGES OF THE ORGANS.

A. Those which especially irritate the part to which they are applied

1. *Acids*

2. *Causive alkalis.*

3. *Causeic salts*, especially those of the heavy metals.

4. Locally irritating organic substances which neither can be classified as corrosive acids nor alkalis, nor as corrosive salts, such are *cantharadin*, *phrymin*, and others in the animal kingdom, *croton oil* and *asinin* in the vegetable kingdom, locally irritating colors, such as the *anilin dyes*.

5. Gases and vapors which cause local irritation when breathed, such as *ammonia*, *chlorin*, *iodin*, *bromin*, and *sulphur dioxide*.

B. Those which have but little effect locally, but change anatomically other parts of the body, such as lead, phosphorus, and others.

II. BLOOD POISONS.

1. Blood poisons interfering with the circulation in a purely physical manner, such as *peroxid of hydrogen*, *ricin*, *abrin*.

2. Poisons which have the property of dissolving the red corpuscles, such as the *saponins*

3. Poisons which, with or without primary solution of the red blood corpuscles, produce in the blood methaemoglobin, such as *potassic chlorate*, *hydroxine*, *nitrobenzene*, *anilin*, *picric acid*, *carbon disulphid*.

4. Poisons having a peculiar action on the coloring matter of the blood, or on its decomposition products, such as *hydric cyanid*, and the *cyanides* and *carbon monoxid*.

III. POISONS WHICH KILL WITHOUT THE PRODUCTION OF COARSE ANATOMICAL CHANGE.

1. Poisons affecting the cerebro-spinal system; such as *chloroform*, *ether*, *nitrous oxid*, *alcohol*, *chloral*, *cocain*, *atropin*, *morphin*, *nicotin*, *conium*, *aconitin*, *strychnin*, *curarin*, and others.

2. Heart poisons; such as *digitalis*, *helleborin*, *muscarin*.

IV. POISONOUS PRODUCT OF TISSUE CHANGE.

1. Poisonous albumin.

2. Poisons developed in food.

3. Auto-poisoning, e. g. *uræmia*, *glycosuria*, *oxaluria*.

4. The more important products of tissue change; such as, *fatty acids*, *oxy-acids*, *amido-fatty acids*, *amines*, *diamines*, and *ptomaines*.

CLASSIFICATION OF POISONS AND POISONOUS SYMPTOMS, ACCORDING TO BERNHARD H. SMITH

POISONS ACTING ON THE BRAIN

I

NARCOTICS

Symptoms. Giddiness, dimness of sight, contracted pupils; headache, noises in the ears; confusion of ideas, and drowsiness, passing into insensibility.

Treatment. (Immediate).

EMETICS (especially mustard, a tablespoonful in half a tumbler of warm water).

STOMACH TUBE. (Wash out at half-hourly intervals with Potash Permanganate solution B. P. 1 per cent solution diluted three times with warm water). Dash cold water on face and chest. Ammonia or Amyl Nitrite to nostrils.

(Later:)

TANNIN (Hot, strong coffee, tea, or infusions of nut-galls or oak-bark).

Strychnin solution B. P. (1 per cent solution) two minims subcutaneously.

Atropin, half grain subcutaneously, repeated if necessary.

Faradic current. Oxygen.

Artificial respiration if necessary.

DEMLLCENTS. (Milk, eggs, oil, etc., swallowed and injected)

Note—All vegetable poisons act more quickly on the empty stomach, our treatment therefore is directed towards (1) *diluting* and *evacuating* the poison by means of washing out with warm water containing powdered charcoal; (2) *neutralising* the poison by means of an antidote such as Tannin.

Plant producing poisons that act on the brain as narcotics—Poppy (*Papaver somniferum*)

II

DELIRIANTS

Symptoms. Spectral illusions; delirium; dilated pupils; thirst, and dryness of the mouth, inco-ordination. Occasionally, though rarely, there are paralysis and tetanoid spasms.

Treatment. (Immediate:)

EMETIC and STOMACH TUBE.

Pilocarpin Nitrate, one-third of a grain subcutaneously, repeated if necessary.

Ammonia or Amyl Nitrite to nostrils.

TANNIN

(Later)

Stimulants, Castor Oil.

DEMULCENTS (swallowed and injected).

Note—Muscarin poisoning give Belladonna Tincture B. P fifteen minims

Plants furnishing poisons that act as delirians—Thorn Apple (*Datura Stramonium*), Black Nightshade (*Solanum nigrum*), Hemp (*Cannabis sativa*), Darnel (*Lolium temulentum*), Several Fungi as Fly Agaric (*Amanita muscaria*)

III

INEBRIANTS

Symptoms. Excitement of cerebral functions, and of the circulation; loss of power of co-ordination, and of muscular movements, with double vision; leading to profound sleep, and deep coma.

Treatment. (Immediate:)

EMETIC and STOMACH TUBE.

TANNIN.

(Later:)

Epsom Salts.

DEMULCENTS (swallowed and injected).

Plants furnishing poisons that act on the brain as inebriants: Wormwood (*Artemisia Absinthium*), Jamaica Dogwood (*Piscidia Erythrina*).

POISONS ACTING ON THE SPINAL CORD

CONVULSIVES

Symptoms. Clonic (intermittant) spasms, extending from above downwards. Opisthotonos very violent; but trismus (lock-jaw) rare. Swallowing spasmodic. Death, usually, in less than three hours, or rapid recovery.

Treatment. (Immediate:)

EMETIC and STOMACH TUBE. TANNIN.

Chloral Hydrate five grains subcutaneously (a weak solution as it is an irritant), repeated if necessary. Chloroform inhalation. Artificial respiration.

Potassium Bromide, one drachm in water every half-hour. Morphia (?).

(Later)

DEMULCENTS (swallowed and injected).

Castor Oil. Chloroform inhalation if convulsions return.

Plants furnishing poisons that act upon the spinal cord as convulsives: Nux-vomica Tree (*Strychnos Nux-vomica*), St. Ignatius's Bean (*Strychnos Ignatii*)

POISONS ACTING ON THE HEART

I

DEPRESSANTS

Symptoms. Vertigo, vomiting; abdominal pain; confused vision; convulsions, occasional delirium; paralysis; syncope; sometimes asphyxia.

Treatment. (Immediate.)

EMETIC and STOMACH TUBE. TANNIN.

STIMULANTS. Strychnin solution B. P., two minims subcutaneously.

Atropin, half a grain subcutaneously, repeated if necessary.

(Later.)

Stimulants. Hot fomentations.

Artificial respiration if necessary. Castor Oil.

DEMULCENTS (swallowed and injected).

Plants furnishing poisons that act on the heart as depressants: Tobacco (*Nicotiana Tabacum*), Hemlock (*Conium maculatum*), Indian Tobacco (*Lobelia inflata*).

II

ASTHENICS

Symptoms. Numbness, and tingling in the mouth, abdominal pain, vertigo; vomiting, purging, tremor, occasional delirium; paralysis; dyspnoea, ending in syncope.

Treatment. (Immediate:)

EMETIC and STOMACH TUBE. TANNIN.

STIMULANTS. Cold affusion.

Faradic current. Atropin, half a grain subcutaneously.

(Later.)

DEMULCENTS (swallowed and injected).

Continued recumbent position.

Artificial respiration if necessary. Castor Oil.

Note—In Aconite poisoning inject Digitalis Tincture B. P., twenty minims subcutaneously (12½ per cent strength).

Plants furnishing poisons that act on the heart as asthenics: Lima Bean (*Phaseolus lunatus*), Taproot (*Jatropha manihot*), Aconite (*Aconitum Napellus*), Cohosh (*Cimicifuga racemosa*), Oleander (*Nerium Oleander*), Fox-glove (*Digitalis purpurea*), White Hellebore (*Veratrum album*), Green Hellebore (*Veratrum viride*).

VEGETABLE IRRITANTS

I

PURGATIVES

Symptoms. Abdominal pain; vomiting, and purging, cramps; stranguary and tenesmus, followed by collapse, and sometimes accompanied by drowsiness, and slight nervous symptoms.

Treatment. (Immediate.)

EMETIC and STOMACH TUBE. TANNIN.

(Later:)

Opium to relieve pain.

Stimulants to counteract collapse.

DEMULCENTS (swallowed and injected).

Plants producing poisons that act as purgatives. Castor Bean (*Ricinus communis*), Green Hellebore (*Helleborus viridis*), May Apple (*Podophyllum peltatum*), Marsh Marigold (*Caltha palustris*).

II

ABORTIVES

Symptoms. Nausea; vomiting; stupor, polyuria, sometimes tenesmus.

Abortion may or may not occur; coma.

Treatment. (Immediate:)

EMETIC and STOMACH TUBE. TANNIN.

Ammonia or Amyl Nitrite to nostrils.

(Later.)

Opium to relieve pain.

Stimulants to counteract collapse.

DEMULCENTS (swallowed and injected).

Plants producing poisons that act as abortives Ergot (*Claviceps purpurea*) Herb of Grace (*Ruta graveolens*), Cotton root (*Gossypium herbaceum*), Pulsatilla (*Anemone patens*, and its variety).

III

IRRITANTS WITH NERVOUS SYMPTOMS

Symptoms. Abdominal pain; vomiting and purging; dilated pupils; headache; tetanic spasms, occasional convulsions, sometimes rapid coma.

Treatment. (Immediate:)

EMETIC and STOMACH TUBE. TANNIN.

Opium to relieve pain.

Stimulants to counteract collapse.

Bleeding if necessary.

(Later.)

Castor Oil.

DEMULCENTS (swallowed and injected).

Plants furnishing poisons that act as irritants, causing also nervous symptoms Indian Pink (*Spigelia marilandica*), Cut-leaved Water Parsnip (*Berula erecta*), Fool's Parsley (*Aethusa Cynapium*).

IV

SIMPLE IRRITANTS

Symptoms. Burning pain in the throat and stomach; thirst; nausea, vomiting; tenesmus; purging; dysuria; dyspnoea and cough occasionally Death through shock; convulsions; exhaustion; or starvation due to injury to throat or stomach.

Some few (i. e. the Nettles) cause smarting pain on the merest contact with the secretions of the plant; quickly followed by erythema and urticarial rash, which slowly subsides.

Treatment. (Immediate:)

EMETIC and STOMACH TUBE. TANNIN.

Opium to relieve pain.

Stimulants to counteract collapse.

Chloral and Bromides if convulsions.

(Later.)

Castor Oil.

DEMULCENTS (swallowed and injected).

Plants producing poisons that cause the above symptoms Arrow Root

(*Arum maculatum*), Wood Anemone (*Anemone nemorosa*), Cursed Crowfoot (*Ranunculus sceleratus*), Buttercup (*Ranunculus acris*), Bouncing Betty (*Saponaria officinalis*), Kinnikinnik (*Arctostaphylos Uva-ursi*), Sundew (*Drosera rotundifolia*), Poison Ivy (*Rhus Toxicodendron*), Nettle (*Urtica dioica* U. *gracilis*), Wood Nettle (*Laportea canadensis*), Bull Nettle (*Jatropha stimu-losa*)

V

SIMPLE IRRITANTS WHEN TAKEN IN LARGE QUANTITIES

Symptoms. Burning pain in throat and stomach. Vomiting, purging; difficulty in swallowing. Recovery usual.

Treatment. (Immediate.)

EMETIC and STOMACH TUBE. TANNIN.

Opium to relieve pain.

Stimulants to counteract collapse.

(Later.)

Castor Oil.

DEMULCENTS (swallowed and injected)

Plants affording poisons that act as simple irritants when taken in large quantities only. White Mustard (*Brassica alba*), Black Mustard (*B. nigra*), Black Pepper (*Piper nigrum*), Common Ginger (*Zingiber officinale*), Cayenne Pepper (*Capicum annuum*).

TABLE OF SYMPTOMS OBSERVED AFTER THE ADMINISTRATION OF POISONS, ADAPTED FROM CATTELL AFTER THE WORK OF KOBERT.

ACUTE SYMPTOMS PRESENT	WE SHOULD THINK OF
1. Death within a few seconds or minutes.	Hydrocyanic acid; potassium cyanid, carbonic acid, carbolic acid.
2. Deep coma.	Alcohol, morphin, opium.
3. Collapse.	Nicotin; colchicin.
4. Feverish rise of temperature.	Cocain, enzymes.
5. Mania, furious delirium; psychic excitement.	Chronic alcoholism, atropin, camphor, physostigmin; veratrin.
6. Mental disturbance of the most diverse kind.	Alcoholism, morphinism, cocainism; pellagra, ergotism.
7. Violent at times, tetanic convulsions.	Strychnin, toxin of tetanus, cytisin, cornutinin; picrotoxin; cicutoxin, active principle of digitalis; cocain; santonin; aconitin, gelsemin; flicic acid.
8. General paralysis, for the most part ascending.	Comin, curarin, colchicin.
9. Dilation of the pupil.	Atropin, hyoscyamin; scopolamin; cocain, ephedrin, aconitin.
10. Contraction of the pupil.	Muscarin, pilocarpin; nicotin, codein; opium, physostigmin.
11. Anæsthesia.	Quinin; extract of male fern; belladonna; uræmic poisoning.
12. Diplopia and ptosis.	Sausage (botulism) and fish poisoning.
13. Conjunctivitis.	Irritating vapors, ethereal oil of mustard;

- 14 Moist skin.
Opium, morphin; aconitin; pilocarpin; nicotin, physostigmin, lobelin.
- 15 Skin conspicuously dry even in a warmed bed. Mouth and throat parched.
Atropin, belladonna, stramonium; hyoscyamus, hyoscyamin, scopolamin, sausage and fish poisoning.
- 16 Urticaria or scarlatiniform erythema.
Atropin; hyoscyamin, antipyrin, quinin, balsam of copaiha, cubebe, morphin, bandling of nettles; buckwheat; smartweed.
- 17 Eczematous eruptions of skin.
Croton oil, curcas oil, cardol, Rhus Toxicodendron, powdered cinchona bark, carbolic acid; tar.
- 18 Diffuse dermatitis with perspiration of the hands.
Anilin colors; aurantia; butter yellow.
- 19 Acne pustules.
Powdered ipecac.
- 20 Blisters on the skin or the mouth, containing clear serum.
Ranunculus acris; R. sceleratus, etc.
- 21 Gangrenous ergotism, carboliism.
Ergot.
- 22 Cyanosis.
Antifebrin; exalgin; anilin.
- 23 Yellowish-brown discoloration of the conjunctiva, in combination with that of the skin.
Helvellic acid, lupinotoxin (strychnin).
- 24 Discoloration primarily of the tongue and mucous membrane of the mouth.
Carbolic acid.
- 25 Salivation.
Pilocarpin; muscarin; arecolin; nicotin; cornutin, physostigmin, cyllium saponin.
- 26 Metallic cough and aphonia.
Atropin, hyoscyamin, scopolamin; sausage poisoning.
- 27 Oedema of the glottis.
All caustic poisons.
- 28 Oedema of the lungs.
Muscarin; morphin, pilocarpin, nitric acid vapors.
- 29 Increased dullness of the liver.
Agaricus bulbosus; poley oil, alcohol.
- 30 Diarrhoea with vomiting.
Digitalin, pilocarpin; nicotin, muscarin; colchicin, corrosive poisons, colocynthin, emetin, cephalin; croton oil.
- 31 Vomiting without diarrhoea.
Apomorphin; lobelin; cyttisin.
- 32 Diarrhoea without vomiting.
Jalap, podophyllotoxin, croton oil, calomel.
- 33 Pulse continuously and markedly becoming slower.
Opium, morphin, muscarin, arecolin, physostigmin, all narcotics.
- 34 Pulse first slower, then irregular, finally accelerated.
Digitalin, hellebore, adonis; coronilla; cheiranthin, nerwin; scilla, strophanthus; convallaria, pilocarpin, nicotin; scopolamin.
- 35 Pulse greatly accelerated.
Belladonna, hyoscyamin; atropin.
- 36 Abortion.
Sabina, thuya, rue; mentha, pulegium, ergot, cotton root.

- 37 6-12 hour period of good health between the poisoning and the appearance of the symptoms. Most of the poisonous fungi, but especially *Amanita phalloides*.

TREATMENT FOR POISONING. (CHIEFLY AFTER KOBERT).

KIND OF POISON

TREATMENT

- Aconite.** Use stomach pump at once, give emetics of sulphate of zinc, or a hypodermic injection of apomorphin, patient should recline, when stomach has been evacuated, give atropin (4 drops U. S. P. solution) hypodermically or by the mouth, or 20 drops of tincture of belladonna, if there is a tendency to heart-syncope, give tincture of digitalis, in $\frac{1}{2}$ drachm doses by mouth, or hypodermically in doses from 10 drops upwards, apply mustard plasters to pericardium, and vomiting by plenty of water if necessary apply artificial respiration.
- Alcoholism, acute.** Wash out the stomach with a siphon tube; cause vomiting by emetics, cold and hot effusions alternated; strychnin hypodermically.
- Aspergillous.** Treatment unsatisfactory; moulds that have entered lungs cannot be destroyed.
- Atropin.** Wash out the stomach with a solution of tannic acid or cause evacuation with an emetic; hypodermic injection of strychnin to stimulate respiration, administer tea or whiskey.
- Carbolic acid.** A liberal dose of whiskey or alcohol as a diluent, use soft stomach tube to wash out the stomach with sodium sulfate; sodium sulfate, raw eggs, milk and saccharate of lime are antidotes.
- Cicuta virosa. C. maculata**
cowbane.
Cicutoxin. Tannin and narcotics according to symptoms, especially chloral hydrate.
- Cocain.** Chloroform, chloral hydrate, artificial respiration, stimulants like whiskey.
- Colchicin.** Evacuate the stomach, then give vegetable astringents, iodin 1 gr and potassium iodid 10 gr dissolved in water, digitalis and amyl nitrite are given for syncope; give stimulants like oxygen and whiskey for cyanosis.
- Coniin.** Mucilaginous potions, morphin, warm compresses on the abdomen.
- Conium.** Wash out the stomach after giving tannic acid or some other astringent, strong coffee and whiskey, strychnin hypodermically; apply artificial respiration if necessary.
- Cytisin.** Tannin, stimulants, especially camphor.
- Delphinosis.** When spasms occur, give narcotics, artificial respiration in case of paralysis of the organs of respiration.
- Dermatitis. Rhus.** Potassium permanganate is the antidote in extreme depression of circulation and respiration, atropin sulfate administered hypodermically.
- Antidote is acetate of lead.

Dermatomycosis.

Digitalis.

Equisetosis.

Ergotism.

Fagopyrism.

Forage Poisoning.

Formic Acid. (In stinging net-
tles)

Gelsemin.

Helleborus niger.

Hydrocyanic Acid.

Lathyrism.

Locoism.

Lupinosis.

Maydism.

Morphin.

Muscarin.

Nicotin.

Phallin.

Pterotoxin.

Pilocarpin.

Cleanliness; sanitary surroundings; disinfection; preparation of bichloride of mercury.

No antidote; treat symptomatically.

Administer cathartic, also nerve and heart stimulants to combat symptoms of depression; change of food.

Administer purgatives and stimulants, especially camphor; tannic acid is chemical antidote and will neutralize unabsorbed portions of poison; chloral is physiological antidote; further treatment symptomatic.

Change of food.

Change of food.

Cooling compresses externally, chalk, soda, or magnesia internally.

Wash out the stomach thoroughly; give stimulants and hot applications to the epigastrium and extremities, digitalis to strengthen the heart action and atropin to increase respiration.

Use stimulants, especially camphor.

Wash stomach with 0.5 per cent potassium permanganate solution or with hydrogen peroxide; the latter may be given hypodermically in small doses but with great caution, artificial respiration.

Treat paralyzed parts electrically; massage.

In advanced stage of disease, treatment of no avail; in early stage, removal to an uninfected pasture with plenty of good nourishing food besides may benefit.

Chiefly preventive; no specific antidote; to prevent further absorption of poison, administer an acid; also give purgative.

Change of diet, transfer to hospital.

Wash out stomach with siphon tube using water containing potassium permanganate in the proportion of 20 gr. of the permanganate to 1 tumbler full of water; or use in the same manner an infusion of tea or tannic acid; emetics like mustard, using one or two teaspoons of each; apomorphin 5-10 minims of a 2 percent solution; hypodermic doses of strychnin 1-20 gr. or sulphate of atropin 1-60 gr.

Atropin used hypodermically, stimulants like strychnin may be given, wash out stomach.

If free vomiting has not occurred, wash out stomach with warm water or tea, give stimulants like whiskey or use hypodermic injection of strychnin nitrate 1-25 gr.

No known antidote; undigested material should be removed from stomach and same remedies as those suggested in muscarin poisoning may be administered.

Chloroform, chloral hydrate, artificial respiration.

Evacuate the stomach and wash out with so-

Physostigmin.

Ranunculus.

Santonin.

Strychnin.

Toxina.

Veratrin.

lution of tannin; atropin is antagonistic, give 1-60 gr. hypodermically, use whiskey and ammonia as stimulants.

Scopolamin hypodermically; artificial respiration.

According to symptoms, tannin.

Chloroform, chloral hydrate, artificial respiration.

Control spasms by inhalation of chloroform; use stomach tube with warm water containing potassium permanganate, 4 gr in 1 l fl oz. water.

Botulism produced in sausage poisoning may be treated by removing the poison, by giving free potations of warm water and salt, irrigation of intestines with enemata; vomiting and purging can be relieved by hypodermic injections of morphin, inject normal salt solutions.

Wash out stomach thoroughly with the siphon tube, or give emetics, give tannic acid or vegetable astringents which will precipitate the alkaloids, atropin or strychnin will counteract the cardiac depression¹.

¹ In the above table use has been made of the English translation of Robert's Practical Toxicology by Friedburg, as well as the original.

CHAPTER XII

THE PRODUCTION OF POISON IN PLANTS

THE USE AND ACTION OF POISONS ON PLANTS

An extraordinarily large number of poisonous substances belonging to the alkaloids, glucosides, saponins, and toxins, occur in the vegetable kingdom. In addition there is a larger number of substances not strongly poisonous which are curative in their nature. These substances so widespread in the vegetable kingdom are the products of metabolism and probably in some cases, waste products, although according to Weevers, may act as reserve food substances. Treub¹ states that the hydrocyanic acid in the *Pinguicula vulgaris* is of importance in the metabolism of the plant. It occurs not only in certain parts of the fibro-vascular bundle but in the cells from the leaf and certain specialized cells of the epidermis, occurring both in a free and in an unstable combination. His conclusion is that the hydrocyanic acid is the first recognizable product of nitrogen assimilation. It is certainly true that in some cases the poisonous products formed in the plant do not undergo any further change. Undoubtedly the substances serve as a protection to the plant and it may be interesting to note that large quantities of these substances may be excreted and occur in an insoluble form in the cell sap and do no injury to the plant. *Digitatin*, *morphin*, *atropin*, *escerin*, *muscarin*, and *veratrin*, seem to exert little or no poisonous action on most plants; while *strychnin* may act as a strong poison.² There are also other alkaloids that when applied to the plant are poisonous to the plants from which they have been obtained. *Morphin* is said to poison the poppy and the motile spores are speedily killed by the same substance according to Strasburger. Too little, however, is known about this subject to make any extended remarks. It is interesting, however, to observe in this connection that as in the case of man and other animals, plants can be gradually accustomed to doses which would probably prove fatal in many cases. The Blue Mould (*Peridermium glaucum*) and some species of *Aspergillus* can accommodate themselves to strong solutions of copper and formalin. There are some reasons for believing that the protoplasm of different plants is not a uniform substance but varies, and that one substance may be toxic to the plant while harmless to another and even act as a stimulant.

DISTRIBUTION OF POISONOUS SUBSTANCES IN PLANTS

The seed may contain a toxic substance and upon the germination the poisonous material may occur not only in its juvenile stage but at maturity. In some cases the seed and the juvenile form may be non-poisonous, but as the plant becomes older the poisonous substance is elaborated as in the *latex* of some plants that contain narcotic principles. In some cases the seed is poisonous and the young plants apparently do not contain a toxic material, the

¹ Ann. du Jardin Bot. de Buitenzorg 13. Phym. Review 14 372.

² Pfeffer, Physiology of Plants. English Translation. 2 360, Schwarz, Wirkungen von Alkaloiden auf Pflanzen, Erlanger Dissertation. 1897.

poisonous substance apparently being broken up to serve as a nourishing material for the growth of the plants.

Cornevin* in his work upon poisonous plants says "In one group "the poisonous substance does not exist in the plants themselves, but, in some parts or tissue elements are present which are not really poisonous in themselves but become so when the parts or tissues come in contact with one another. An example is seen in the glucoside *amygdalin* which in contact with *emulsin* forms *hydrocyanic acid*."

The activity of vegetable poisons may depend upon the age of the plant producing them. No definite rule can be established in regard to the age at which a plant produces its poisonous substance. Sometimes a younger plant is more actively poisonous than when older, sometimes, also, the poison is stored in certain tissues.

Poisonous principles are found in various parts of the plant, such as the root, stem, flowers, fruit, leaves, bark, tubers, seed and bulbs.

In many aerial parts of plants the poisonous substances are more ephemeral than they are in organs of the plant that serve as store houses of food.

Cornevin says

It sometimes happens that the subterranean part only is poisonous as in *Atracylis gym-nifera*. This is true, also, of the common European Violet. On the other hand when subterranean organs of certain plants come in contact with the light a poisonous substance may be produced. This is true of the tuber of potato which when green is poisonous. Poisonous substances are elaborated both in evergreen and deciduous leaves; no rule of comparison has, however been established.

Certain variations of the plant are dependent upon its environment, certain conditions of which play an important part especially in the elaboration of poisonous substances.

These conditions are light, heat, season, climate, soil, culture and fertility

LIGHT It is a well known fact that light seriously interferes with the growth of parasitic fungi and bacteria so that pathogenic species may become quite harmless when placed in direct sunlight. The poison *atractylin* is formed only in darkness, on the other hand *solanin* is formed only in the light, as in the case of the green potato.

HEAT Cornevin says:

The action of heat upon plants should be considered with that of light, of the seasonal variation, and of climate, and not as an isolated cause. When it is prolonged it induces desiccation of the plant and as a consequence evaporation follows and destruction of the poison when it is volatile. This result is produced in some *Ranunculaceae*, *Chenopodiaceae* etc.

Moist heat, that is boiling, conduces to the same result in some poisons. Three-sided mercury (*Mercurialis annua*) becomes inoffensive when it has been submitted to the action of heat, because of the volatilization of its toxic substance *mercurous*.

SEASONS. Seasons cause considerable variation in the poisonous material produced in plants, the amount of poison contained often varying with the advance of the season. In *aconite* the poison, at first contained in the leafy organs becomes concentrated little by little in the seed.

The time of the year may have an important bearing upon the amount of poisonous material found in the plant. According to Prof. Hedrick the Cowbane or Musquash root (*Cicuta rugosa*) of the west is much more toxic in the fall, winter, and spring, than in the summer, and this is partially confirmed by an experiment conducted with our common Cowbane (*Cicuta maculata*). It is also well known that the mature bulbs of *Colchicum* contains a much larger amount of the toxic substances than the growing bulbs and that for medicinal

* Des plantes vénéneuses et des empoisonnements qu'elles déterminent, 524. Paris, 1893

purposes the bulbs are usually collected between the decay of the foliage and the production of the flowers.

Opium is obtained from the capsules of the opium plant a few days after the petals have fallen, the seed containing comparatively little of the narcotic substances. It is well-known also that the poisonous principles of the Larkspurs are much more active in the spring than in the summer. Dr. Albert C. Crawford¹ says referring to the *Delphinium consolida*, with which he conducted some experiments and was able to kill several guinea pigs with toxic material obtained from the plant collected on April 26th and May 16th but failed to get positive results of material collected in June.

There is no question as to the fact that *Delphinium* when injected subcutaneously will kill, and these experiments also establish the fact that the plant loses much of its toxicity as it approaches the flowering stage. It has been noted that *Delphinium consolida* is also less active when mature.

Just after flowering, the purple larkspur turns yellow and ceases to be attractive, so that there is less danger of poisoning, although Chomut and Wilson report death in cattle from eating *Delphinium glaucum* in September. The great danger early in the season seems to arise from the fact that the *Delphinium* appears early in the spring, and the ground may again be covered with snow, so that it is the only green plant in sight, and therefore when in an especially poisonous stage it is eaten by cattle.

Botanical and other writers have frequently called attention to the fact that the greatest amount of poisoning in the west occurs in early spring. Of course, this may be because there is less green food and live stock may consume more of this plant than at other seasons. However, there seems scarcely any reason to doubt that the plant does contain a larger amount of the acrid toxic substances in the spring than in the summer, as proved by the experiments of Dr. Crawford.

The same author who investigated the Mountain Laurel² calls attention to the well-known fact that most of the cases of poisoning from Mountain Laurel occur in the winter. Undoubtedly the animals will eat more of the tough and leathery leaves in the winter because there is very little green for them; but may they not also have a larger amount of the toxic material? The plant is evidently also poisonous in the summer, as indicated by numerous reports of the experiments by Dr. Crawford, who conducted an experiment with material collected in the summer, in May and June, death occurring in a sheep weighing 49 lbs., that had received 90 grams of powdered dried laurel leaves.

The late lamented Dr. Greshoff has called attention to the peculiar distribution of hydrocyanic acid in plants. In referring to a species of *Hydrangea* of the *Saxifragaceae* he states that he sometimes found considerable quantities of hydrocyanic acid in some of the well-known ornamental plants like the *H. hortensis* and sometimes he did not find it. He surmises that the cyanogenic principle disappears from the leaves in the autumn and that the young leaves have much more of the HCN than the older ones. In the case of the Plane Tree (*Platanus*) he found considerable of the same acid in the young leaves but as the leaves grow older the HCN content falls off to small traces. He states further that in the ordinary plane tree of the London Streets there is so much HCN that the amount from each London Plane Tree leaf would be enough to kill a London sparrow.

¹ Bull. Bur. Plant Ind., U. S. Dept. of Agric. 111 Part II, 7

² Danmann, C. Gesundheitspflege. 1886 1072.

³ Bull. U. S. Dept. of Ag., Bur. Plant Industry. 121:21.

Miss Alice Henkel in a paper on American Root Drugs¹ notes the importance of collecting medicinal plants and drugs at the right time. The roots of the American Hellebore (*Veratrum viride*) should be collected in the autumn after the leaves are dead. Generally speaking the drugs contain more of the medicinal virtues after the period of cessation of growth of the plant. Dr C Mueller² calls attention to the fact that *Colchicum* is much better if the rhizome is collected at time of flowering. On this point all authorities do not agree. Cornevin who has made a study from month to month of the migration of poison in plants such as the *Laburnum* found the following conditions. On May 20th, 2 grammes of the dried leaves of *Cytisus Laburnum*, administered to some carnivorous animal, were sufficient to induce vomiting. On June 10 (at which time pods were forming) 4 grammes were required to produce the same result. July 28 (when the pods were fully formed) 12 grammes were required. September 28 (pods were beginning to dry) 20 grammes were required.

The experiments show that at all periods the leaves were poisonous but as the poison became concentrated in the pod the leaves became less toxic.

Similar experiments resulted in demonstrating that while the poison concentrates in the pod it also loses toxicity as the season advances, enormous doses taken from dried seed in October failing to produce death, while a 2-gramme dose from a June pod proved fatal.

CLIMATIC. Latitude has an influence upon the formation of poisons. There are more poisonous plants in tropical regions than in colder regions. Certain plants which are poisonous in temperate regions lose their poisonous properties when taken into colder regions. Examples of this are aconite and cherry laurel.

It is difficult to make a comparison between the total number of poisonous plants in tropical and temperate regions, but it is probable that the warmer regions will show the greater number of poisonous species.

It might be well, however, to note in this connection that the poisonous properties of *Rhododendron Chrysanthemum* and *R. colymbicæ* as well as of *Aconitum* are developed in colder regions while such toxic plants as the calabar bean (*Physostigma venenosum*), *Strychnos nux-torica*, the Upas tree (*An-haris toxicaria*) and numerous others are inhabitants of warm climates.

It is well known that the sorghum grown in dry climates produces a greater amount of a glucoside which is capable of being converted into hydrocyanic acid than in more moist regions. Some writers even assert that in dry seasons the drought depauperates the plant and that in the nodes there are considerable quantities of potassium nitrate³.

Again Dr Ludwig Bernegau notes that the amount of alkaloid in Cola Nut varies considerably, it depending upon the source of the nut.

With reference to the presence of alkaloids in different kinds of opium, it is known that the Smyrna opium is of superior quality for medicinal purposes. It is said to be superior to the opium obtained from India, which is in part attributed to climatic conditions, and probably in part to the method of collection. According to Blyth, the amount of morphin varies as follows:

Crude morphin (containing about 7-10 of pure morphin). Highest, 12.30 lowest, 6.76, mean, 9.92 per cent, which equals 12.4 per cent of the dried drug.

¹ Bull. U. S. Dept. Agri., Bur. Plant Industry, 167. Farmers Bull. 168.

² Pharm. Review, 14:113. Dic. Natur. 44:552.

³ An English writer, Frost, attributed the death of animals to this substance rather than to the hydrocyanic acid. Pharm. Review, 15:266.

The nicotine of tobacco depends largely on climatic and soil conditions. The Havana, Porto Rico, Sumatra, Connecticut Seed Leaf and Wisconsin have quantities all peculiarly their own.

Blyth records the following percentages of nicotine in various tobaccos as given by Cox¹

Variety examined	Nicotin per cent
1. Syrian leaves (a),	612
2. Syrian leaves (b),	1 093
3. Gold Flake (Virginia),	2 501
10. "Navy-Cut" (Light colored),	3 640
15. "Best Shag" (b),	5 000
17. Algerian tobacco (a),	8 813

According to Professor Garner² the nicotine contents vary as follows

Nicotin soluble in petroleum ether in domestic filler tobacco 2.20 percent.
Imported Cuban Santa Clara tobacco 1.33 percent.

That climate plays an important part in connection with chemical products is illustrated in the essential oils. The most important of these such as bergamot, cassia, cinnamon, jasmine, fennel, lavender, orange, rosemary, attar of roses, and many of the others are imported into the United States. These are produced in localities that are more favorable for their production than in many parts of this country. However, some of the essential oils like those from peppermint, sweet birch, and sassafras are produced extensively in the United States. The time of collecting and many other factors as well as the matter of labor enter into the production of these oils. The variation in composition has been indicated by Dr. Edward Kremers and his students in various publications.³

It is well known that the locality for attar of roses which supplies most of the commerce of the world is a small district of country on the southern side of the Balkan Mountains. The best localities according to those best informed on the subject are those occupying southern or southeastern slopes. The flowers attain perfection in April or May and are gathered before sunrise. It is concluded from some experiments carried on in some of the northern countries in Europe, that a cool northern climate is not conducive to the production of highly odorous oils.⁴

It is interesting to note that the oil of orange flowers comes chiefly from the southern part of France, and that the flowers of sweet orange afford about one-half the amount of oil that those of the bitter orange do. Lavender oil, made from *Lavandula vera*, is very variable in quantity, depending upon its source, although not grown to any great extent in England. Flückiger and Hanbury state that the Mitcham oil grown in the Surrey, is of a superior and high quality. The above is used here to illustrate the fact that the chemical products of plants vary depending upon the different climatic conditions.

¹ Pharm. Journ., Jan. 20, 1894. Blyth, Poisons, Their Effects and Detection. Ed. 2, 274.

² Bull. U. S. Dept. Agr., Bur. Plant Ind. 141, Pl. I 12.

³ Kremers and Schreiner. The Quantitative Estimation of Carvone in Volatile Oils. Pharm. Review 14 76.

Edward Kremers and Florence M. Caze. Notes on Two Oils Containing Pulegone. Pharm. Review 16 413.

Kremers and Sievers. Oils from Mistle. Pharm. Review. 25 215.

The Volatile Oils, by E. Gildemeister & Fr. Hoffman. English Translation, E. Kremers.

⁴ Flückiger & Hanbury, 262.

Dr Rodney H True in an article in the *Cyclopedia of American Agriculture*,¹ says

The sources of our crude drugs and condiments are very widely separated depending in large part on climatic conditions. Common drug plants belonging to the temperate zone, such as digitalis, burdock and caraway are in very large part produced in northern and central Europe frequently in more or less localized regions. Caraway comes chiefly from Holland in small quantities from Norway east Prussia and southern Germany. Fennel is cultivated in Saxony, Galicia, Macedonia and Italy. Digitalis leaves and berries reach the market of northern Germany, Austria, Belgium, Holland and England. Peppermint oil is produced chiefly in Japan and the United States. Other plants demanding tropical conditions are obtained from regions in which their culture has been undertaken. Cassia bark from which quinine is obtained, came formerly from the slopes of the Andes. Cultivation of this plant in India, Java, and other parts of the Orient has succeeded in so far as to cause the practical disappearance of the wild barks of South America from the market. Ipecacuanha, likewise a native of northern South America, is apparently repeating this history. Black and white pepper are chiefly produced in southeastern Asia, coming on the market through Singapore and Penang. Cloves are in large part supplied by Zanzibar where the crop constitutes one of the royal monopolies. Some products are derived from well more localized regions, as buchu leaves from the vicinity of Cape Town, South Africa, and olives from South Africa, the island of Socotra in the Red Sea, and the Barbados islands. Some are cultivated, as may be seen in numerous cases cited above, and some are wild products. Camphur until recently has been derived from an essentially wild tree growing in Japan, China and Formosa. The great depletion of the natural forests has led the Japanese government to make extensive plantings. Several African sorts of the red peppers of the market are collected by natives from the wild plants and brought long distances to market.

CULTURE. Cultivation often affects the amount of poison contained in a plant. A wild vetch of Europe (*Lathyrus*) is extremely bitter, but through cultivation the poisonous material has been largely eliminated and the vetch has become a useful cultivated forage plant. It is stated also that the aconite (*Aconitum Napellus*) when cultivated loses some of its toxic properties and that this loss of toxic action occurs in a few generations. Prof. S. M. Tracy informs me that the cultivated forms of the Cassava are not injurious to stock but the wild form, as is well known, contains toxic materials. The wild forms of the lima bean (*Phaseolus lunatus*) contains much more HCN than the cultivated forms.

Dr Rodney H True in speaking of the physiological action of the betel nut states that its physiological action² depends on several factors. The green nuts produce temporary dizziness. The poisonous variety according to Indian authorities is one that is reverted to its former wild condition, while the common or ordinary betel nut which has been cultivated for hundreds of years is a mild narcotic stimulant bringing about a feeling of general comfort, good humor and exhilaration.

It is also well known that when certain toxic bacteria as *Streptococcus pyogenes* are cultivated they lose some of their pathogenic properties.

We must not, however, conclude that because a plant is cultivated it loses its poisonous properties, because there are certain cultivated ornamental plants that are as poisonous in their cultivated form as when grown wild. For example the tobacco which has been cultivated for several hundred years contains as much nicotine in its cultivated as in its primitive form.

SOIL. It is believed also that the soil plays an important part in connection with the amount of poison produced in the plant. There can be no question that the soil plays an influence upon the quality of the fruit as well as upon

¹ *Cyclopedia of American Agriculture*, 2:458.

² *Pharm. Review*, 14:130.

the quality of the grain. Why should it not also influence the poisonous constituents of the plant? Cornevin mentions that soil affects the color of *Digitalis* it being much paler when grown in calcareous than when grown in granitic soil.

RELATED SPECIES AND TOXIC SUBSTANCES. In some cases the same toxic substance is widely distributed in different families of plants. On the other hand closely related plants frequently have entirely different amounts of toxic substances. The bitter and sweet almond illustrate this in a very marked degree, the bitter almond having considerable amounts of the glucoside amygdalin, while the sweet almond is entirely harmless.

Frank Rabak¹ has made an investigation of the amount of the chemical substances found in the kernels of the fresh apricot and plum, in which it appears that the amount of hydrocyanic acid by per cent contained in these plants is as follows:-

	Per Cent
Peach	2.20
Apricot	2.40
Apricot	2.05
Prune	1.75
Bitter Almond	4.80

According to Dohme and Engelhardt² the thin green or young bark of *Prunus serotina* is richer in hydrocyanic acid, than the dark brown or older bark.

It may be interesting to note that certain species of Cacti are used for stock food,³ and others contain powerful drugs. For instance, the Mescal Bean, which is derived from several species of *Anhalonium* which contain pelletin, is a narcotic of considerable potency.⁴

According to Dr. Peinemann⁵ the alkaloidal-content of *Datura* varies as follows, the seeds, 0.541 percent.; root, 0.315; leaves, 0.41 percent. of atropin. The *Datura alba* is richer in alkaloids than the native species, *D. Stramonium*.

Dunstan and Henry⁶ are authorities for the statement that two forms of Mandrake, the American (*Podophyllum peltatum*) and the Indian (*P. emodi*), produce the same principle podophyllotoxin $C_{22}H_{32}O_8$, a strong purgative and also the so-called podophyllin which consists of a resin. The podophyllotoxin occurs in the Indian plant from 9-12% and in the American plant from 4-5%.

DISTRIBUTION OF CHEMICAL SUBSTANCES The same chemical substance is frequently found in plants that are not closely related. For instance, Edward Kremers and many other pharmaceutical chemists have found the essential oils in widely separated families, and Greshoff states that a leaf of Five-finger (*Potentilla daurica* of China) which is closely related to our shrubby Cinquefoil (*P. fruticosa*) produces a powerful odor of roses. Geraniol occurs, for instance, in the young foliage of willow and many other plants.

The substance berberin is found in a large number of plants like the Barberry, the Mandrake, Twin Leaf (*Jeffersonia diphylla*), Meadow Rue (*Thalictrum flavum*), *Toddalia aculeata*, *Hydrastis canadensis*, *Argemone mexicana*, etc.

¹ Bull. U. S. Dept. Agr., Bur. Plant Industry 133.

² Pharm. Review. 14 13.

³ Griffiths and R. F. Harc, The Yucca as Food for Man. Bull. U. S. Dept. Agr., Bur. Plant Ind. 116.

⁴ Pharm. Review. 14 151.

⁵ Pharm. Review. 14 213.

⁶ Proc. Chem. Soc. 189 42-44.

Greshoff¹ has indicated that a species of *Thymus*, the *Thymus Serpyllus* can be readily distinguished from the variety *album* phytochemically, the common form of this species (*Thymus Serpyllus*) produces thymol in its leaves, whereas the other has the odor of lemons.

SAPONINS. We may note here the very wide distribution of saponins in different plants. Greshoff in several papers² gives a long list of the orders in which saponin occurs,³ many of the others are mentioned in the list of species at the end of this manual.

The substance saponin occurs in many different plants of which the following is a partial list:

Amaryllidaceae *Agave*, *Fourcroya*, Araceae *Arum* Araliaceae *Aralia*
 Begoniaceae *Begonia*, Berberidaceae *Berberis*, Caulophyllum, *Leontice*
 Bromeliaceae *Bromelia*, Caryophyllaceae *Acanthophyllum*, *Arrenaria*, *Dianthus*,
Gypsophila, *Lychnis*, *Melandryum*, *Polycarpus*, *Saponaria*, *Silene* Chenopod-
 iaceae: *Chenopodium*, Commelinaceae *Tradescantia*, Compositae *Arnica*,
Grindelia *Mutina*, Convolvulaceae *Ipomoea*, Cuscutaceae *Echinocystis*,
Luffa *Trichomanthes*, Dioscoreaceae *Dioscorea*, Euphorbiaceae *Phyllanthus*?
 Ficoidae: *Trianthema*, *Isiles* *Polypodium* Gramineae *Panicum*, *Lolium*,
 Hippocastanaceae *Asclepias*, Illecebraceae *Hernandia*, Leguminosae *Acacia*
Albizia, *Cassia* *pinus*, *Eutoda*, *Euterolobium*, *Gleditsia*, *Gymnocladus*, *Milletia*,
Mimosa, *Pithecolobium*, *Prosopis*, *Calliandra*, *Tetrapleura*, *Xylo* & c. Lili-
 aceae *Chamaecrista*, *Chlorogalum*, *Heliconia*, *Medeola*, *Muscari* *Paris*, *Poly-*
gonatum, *Smilax*, *Trillium*, *Yucca*, & c. Loganiaceae *Buddleia* Magnoli-
 aceae *Illicium*, Meliaceae *Trichilia*, *Walsura*, Melianthaceae *Berseria*,
 Menispermaceae *Stephanandra*? *Cocculus*? *Coccythium*? Myrtaceae *Barringtonia*,
Caraya, Oleaceae *Chionanthus* *Syringa* Passifloraceae *Carica*? *Medeola*,
 Phytolaccaceae *Piscaria* Pittosporaceae *Pittosporum* Polemoniaceae *Cam-*
panula, Polygalaceae *Moumou*, *Polygala* Primulaceae *Anagallis* *Androsace*,
Cyclamen, *Primula*, *Soldanella*, *Trientalia*, & c. Ranunculaceae *Ficaria* *Nigella*,
 Rhamnaceae *Gonoma*, *Zizyphus* Rosaceae *Quillaja*, *Spiraea*, Rubiaceae
Boussacantha, *Cephalanthus*, *Mitchella*, *Mazenda*, *Randia* Rutaceae *Zan-*
thoxylum, Sapindaceae *Blighia*, *Dodonaea*, *Gonophyllum*, *Korlventeria*, *Ma-*
gonia, *Nephelium*, *Paulinia*, *Poncirus*, *Pometia*, *Sapindus* *Serjania*. More
 than 100 species of this family contain saponin. Sapotaceae *Achras* *Bassia*
Lucuma, *Omphalocarpum*, Saxifragaceae *Hydrangea*, Scrophulariaceae
Digitalis, *Leptandra*, *Limosella*, Simarubaceae *Balanites*, Solanaceae *Ac-*
nastus, *Lycopersicum*, *Scopolia*, *Solanum*, Ternstroemiaceae *Comelia*, *Cary-*
ocarp, Tiliaceae *Greina*, Urticaceae *Ficus*, Zygophyllaceae *Gnaparum*, *Tri-*
bulus.

HYDROCYANIC ACID. A list of the plants containing a glucoside cap-
 able of being converted into HCN has been given in another connection, but we
 may note in this connection plants belonging to many different orders like the
 Compositae, Euphorbiaceae or Spurge Family, Gramineae or Grass Family,
 Ranunculaceae or Buttercup Family, Fern, *Fucus*, Droseraceae or Sundew Fam-
 ily, Saxifragaceae or Saxifrage Family. A common Rocky Mountain shrub, the
Jamena, of this family, is said to contain considerable quantities of HCN, and
 the Mountain Mahogany (*Cercocarpus parvifolius*) of the Rose Family (Rosa-

¹ Bull. Min. Information, Royal Bot. Garden Kew 1909 413.

² Kew Bot. Garden 1919 397 Phytochemical Information at Kew. Mededelingen uit

Land Plantentuin. 29 1909.

case), contains HCN, and Greshoff says must henceforth be counted among the poisonous plants. Of the Grass Family (*Gramineae*) quite a number of species such as the *Stipe lessingiana*, are now known to contain HCN. Some of them like our *Stipe robusta*, have been known to be toxic for a long time, but such genera as the Quaking Grass, Briza (*Catabrosa*), Wild eye (*Elymus*), Blanna Grass (*Glyceria*), Salt Grass *Holcus lanatus* all contain HCN.

CUMARIN. The substance cumarin also occurs in many different plants which we may list as follows:

Tournefortia (*Magnoliaceae*), *Phoenix* (*Palmae*), *Dipteryx*, *Toluifera*, *Melilotus* (*Leguminosae*), *Prunus* (*Rosaceae*); *Cerastophyllum* (*Saxifragaceae*), *Asperula*, *Razoumarantha*, *Borreria*, *Diadia*, *Galium*, *Mitracarpum*, *Spermatoca* (*Rubiaceae*), *Ageratum*, *Chrysanthemum*, *Eupatorium*, *Humea*, *Teihia*, *Liatris* (*Compositae*); *Alyssa* (*Apocynaceae*), *Hemidecimus* (*Asclepiadaceae*); *Rhinocanthus* (*Acanthaceae*), *Acerus*, *Angraecum*, *Orchis* (*Orchidaceae*), *Hierochloa*, *Anthoxanthum* (*Gramineae*); *Adiantum*, *Cheilanthes*, *Polypodium* (*Filices*); *Lactorius*, *Russula* (*Fungi*).

CYTISIN. Greshoff reports cytisin as occurring in the following plants of the Pulse family: *Cytisus Adamsi*, *C. alpinus*, *C. Alschingeri*, *C. biflorus*, *C. hirsutus*, *C. Laburnum*, seed 1.8 per cent. *C. nigricans*, *C. polytrichus*, *C. proliferus*, *C. Weldenii*, *C. Atkinsonii*, *C. condensis*, *C. formosissimus*, *C. monosperma*, *C. Ruthenicus*, *C. scoparius*, *Ulex europaeus*, seed 1 per cent, *U. Jussieu*, *Genista ephedroides*; *G. monosperma*, seed 1.9 per cent, *G. florida*, *G. germanica*; *G. spicata*, *G. tinctoria*, *G. romanensis*, *Sophora secundiflora*, seed 3.5 per cent; *S. tamentum*, *S. fluorescent*, *S. sericea*; *Baptisia perfoliata*, *B. tinctoria*, *B. alba*, *B. australis*, *B. leucantha*, *B. vesicolar*, *Euchresta Horsfieldii*, *Anagyris foetida*, Rauwerda reported it in 28 species of *Lotus*, *L. snaytolens*, *Thermopsis Caroliniana*, and *Th. montana*.

In some instances apparently the same alkaloidal substance was first reported in some plants, but later investigations have shown that these substances are not identical. For instance Schlotterbeck who investigated the alkaloids found in the poppy family, especially *Argemone Mexicana*, reported that this plant does not contain morphin but protopis and berberis.

CHAPTER XIII

ALGAE IN WATER SUPPLIES

Algae have long been known to be injurious to water supplies and numerous papers in the United States and Europe have discussed some of the problems arising therefrom. Algae are generally small, microscopic plants found in water and belong to the great class known as Thallophyta. These frequently cause the pollution of water supplies and are therefore of first importance in this connection. Some algae contain a green coloring matter called chlorophyll and hence they are able to make their own food out of water and carbon dioxide. Many of the algae, however, are not green, some being blue, others red, and still others brown. The Sea Mosses are algae also, and occasionally attain great size and length. Some algae are supposed to be injurious to stock, Mr. George Francis in Australia having attributed a disease of horses, sheep, dogs and pigs to some blue-green algae. The organism causing the trouble in this case was referred to *Nodularia spumigena*, which floats on the water and, being wafted to the lee shores, forms scums from two to six inches thick. He says that, when animals drink the water, it acts as a rapid poison and causes death. It first produces stupor and unconsciousness, the animal falling and remaining quiet, as if asleep, unless touched, when convulsions come on, with head and neck drawn back by rigid spasms which subside before death. This, in sheep, takes place in from 1-6, or 8 hours, in horses, in 8-24 hours, in dogs, in 4-5 hours; and in pigs, in 3-4 hours. A post mortem was made on a sheep that had 30 ounces of fresh scum administered by mouth, death occurring in 15 hours. The post mortem, 6 hours later, showed that the stomach had none of the green scum left, all having been absorbed, the abdominal cavity contained 2 pints of yellowish material, the heart was flaccid with effusion of serum around it; lungs, liver, kidneys, and brain normal, dura mater enlarged, blood of ventricles and arteries black. It was thought that poisoning was due to the decomposition of the algae which smelled like putrid urine, previous to this it had an odor like butyric acid. Account was not taken of the products of decomposition by bacteria. The trouble may have been caused by the poisonous products of bacteria rather than these algae. This seems not have occurred to Dr. Francis.

Many of the algae, especially the following, *Anabaena*, *Clothrocystis* and bacteria-like *Bruggston*, produce what is known as the working of the lakes and pig pen odors. Dr. Farlow has described these from Massachusetts and Dr. Trelease has described them for the lakes about Madison, Wisconsin. The greenish-yellow scum occurs on the surface of the lakes, especially noticeable during hot weather of the summer. Dr. Trelease says

When a lot of it is present, it appears as fine granules suspended in the water, scarcely visible to the naked eye except as they reflect the light, when they call to mind the dancing notes in a beam of sunlight. Under the influence of a gentle, but continuous breeze, these particles are collected into scummy masses and are driven ashore, so that they accumulate along the margins of the lake, forming a slimy scum, which quickly putrefies, giving off a very disagreeable odor. During this change, its color changes to a decided




Fig. 19 Algae found in water supplies. *a*, *Prorococcus*; *b*, *c*, *Coscinodiscus*, one of the desmids; *d*, *Pediastrum*; *e*, *Oscillatoria*; *f*, *Acetabularia*; *g*, *Closterium*, chlamydomonadophore; *h*, *Botrydium*; *i*, *Volvox*; *j*, *Volvox*; *k*, *Volvox*; *l*, *Volvox*; *m*, *Volvox*; *n*, *Volvox*; *o*, *Volvox*; *p*, *Volvox*; *q*, *Volvox*; *r*, *Volvox*.

blue green, which stains the pebbles, sticks, etc., over which it is scattered. The appearance of this scum is sometimes spoken of as the working of the lakes, from the resemblance to the collection of scum on cider, etc., when fermenting, or, as an idiom expresses it, "working." The odor given off by this putrefaction can be detected from some distance.

In speaking of the water bloom, Prof Dwight C. Marsh says

This phenomenon is especially marked in Lake Winnebago in some summers. It is due, of course, to the enormous growth of the plants of the plankton, that growth being particularly fostered by the hot weather of midsummer. The plants especially concerned in forming the bloom are *Clathrocystis*, *Anabaena*, *Aphanizomenon*, *Oscillatoria*, *Lyngbya*, and *Gleotrichia*. The times of occurrence of these plants have already been noticed in the discussion of the individual constituents of the plankton. At the middle of August, in some summers, on a still day, the surface of Lake Winnebago is apparently a solid, opaque green. Some of this material decomposes, and as the currents slowly move along the surface material, it shows a wavy streaked appearance like the surface of polished malachite. Its intrinsic beauty, however, does not attract the average person, for he looks upon it as "scum," and he thinks of it simply as an evidence of filth. This material is thrown upon the shores by the waves until the rocks everywhere are completely covered with it, and it may in its decay become very offensive. Following its maximum period of the "bloom" *Cladophora* appears and covers the bottom rocks with a thick mat of green. This great growth of "bloom" nature attracts the attention of the non-scientific observer and many absurd explanations of its appearance are given. The most common one in Oshkosh is that it is a mass of seeds coming from the marshy shore of the Fox and Wolf above Oshkosh. Doubtless the *Anabaena* and *Gleotrichia* have given rise to the supposition that the bloom is a mass of seeds. The decomposition of *Gleotrichia* produces a blood red coloring matter which is sometimes very noticeable on the shores of Lake Winnebago, and has led people to question as to whether the lake is not affected by one of the plagues of Egypt.

When the water is still the plants of the bloom are in greatest abundance close to the surface, and are distributed very uniformly over the lake. Presently, in the latter part of July and in August, there may be seen floating about yellowish green masses of a more or less spherical outline, perhaps as much as three inches or more in diameter. These masses, which are composed of aggregations of *Aphanizomenon* mixed with scattered fronds of *Gleotrichia* and *Anabaena*, have very little coherence and easily disintegrate by falling in places almost at a touch. As is evident from the discussion of the occurrence of the usual constituents of the plankton, the bloom is not a permanent feature of the deep lakes,—in fact, in some years the growth of these algae is hardly noticed by the ordinary observer—and of the shallow lakes few seem to produce as large an amount as is seen in Lake Winnebago. Of the lakes under observation, Shumano and Pelican were the only ones that could be compared at all with Lake Winnebago.

Odors produced by the decomposition of organic matter in water are easily detected. Sometimes they are grassy and sometimes musky or musty. Sometimes the odor is decidedly fishy, which is especially true of some of the higher plants like *Myriophyllum* or *Potamogeton*. Mr. Whipple* says that the cucumber taste found in the Boston water supplies was due to a fresh water sponge. He made a series of experiments with oil of peppermint and other oils and he found that the oil of peppermint could be recognized when the dilution was 1 to 50,000,000; oil of cloves, 1 to 1,000,000, and liver oil, 1 in 8,000,000. Kerosene oil could not be detected when diluted to 1 in 1,000,000. According to Whipple, aromatic odors are produced by certain diatoms, grassy odors are produced by certain blue-green algae like *Anabaena* and *Rivularia*, the sweet grassy odor is produced by *Clathrocystis*; *Volvox* produces a fishy odor, while *Pandorina* and *Eudorina* produce a faintly fishy odor. Certain protozoa produce fishy and oily odors.

Perhaps the most extensive investigation of microscopical organisms in water are those reported by the Massachusetts State Board of Health. Special methods for determining the amount of impurities found in water are given in these reports. Prof. Parker has shown that when the odors are pronounced the organisms are always present in considerable numbers. He says

The majority of organisms which have been recorded are of microscopic size. At first sight it might appear that these organisms would be insignificant in proportion to their mass, and that more attention should have been devoted to the larger plants and animals. Very few cases have been observed, however, in which very serious trouble in water supplies can be referred directly to the growth of large plants or animals whereas, in many instances, waters with very objectionable qualities contain nothing to which these qualities can be attributed except microscopic organisms. Further, although the organisms here dealt with are of such small dimensions, the immense numbers in which they occur more than counterbalance their small size and I am therefore, of the opinion that, of those organisms which cause objectionable qualities in water, the microscopic ones are the more important.

HOW GROWTH OF ALGAE IN WATER MAY BE PREVENTED

The growth of algae may be prevented by covering the reservoir, since light is essential for their growth, they will not develop in a closed reservoir, and we hear of little complaint under such conditions. Large reservoirs for public water supplies are not generally covered. Often reservoirs are much troubled by algal contamination, in some cases it becomes necessary to remove these by a laborious method. The removal of organic matter by keeping the source of the water supply in as pure condition as possible will no doubt do something toward keeping algae out, but nearly all water contains sufficient organic matter for the growth of algae, especially water coming from water sheds.

THE USE OF ALGICIDES

Within recent years much work has been done in preventing the growth of algae and bacteria by copper sulphate solutions. Moore and Kellerman in a recent paper arrive at the following conclusions:

The diarrhoeic odors and tastes so often present in drinking water are due almost exclusively to algae, although the economic importance of studying these plants has not been recognized until recent years. These algal forms are widely distributed, and reservoirs in many states have been rendered unfit for use by their presence. It has been found that copper sulphate in a dilution as great as to be colorless, tasteless, and harmless to man is sufficiently toxic to the algae to destroy or prevent their appearance. At ordinary temperatures one part of copper sulphate to 100,000 parts of water destroys typhoid and cholera germs in from three to four hours. The ease with which the sulphate can then be eliminated from the water seems to offer a practical method of sterilizing large bodies of water when this becomes necessary. The cost of material for exterminating algae will not exceed fifty to sixty cents per million gallons and will usually be less. The destruction of pathogenic bacteria requires an expenditure of from \$5.00 to \$8.00 per million gallons, not including the cost of labor.

It has been found that *Spirgyra* will die in water containing one part of copper to one billion parts of water. Some of the algae like *Anabaena* are destroyed in dilutions from one to five million, although Moore and Kellerman found that one species of *Spirgyra* requires a greater strength of solution. There is a wide-spread opinion that metallic copper and copper salts are injurious, which is certainly true when the same are taken in larger quantities. According to Tschiesch, 15 to 20 of copper sulphate causes vomiting and diarrhoea. In the paper cited, Moore and Kellerman write as follows:

It is evident that there is still a considerable difference of opinion among eminent authorities as to the exact amount of copper which may be injurious, but as a very conservative limit we may accept 0.02 gram as the amount that may with safety be absorbed daily. According to Merck's Index, the National Dispensatory and the United States Dispensatory, the dose of copper sulphate for tonic and astringent purposes is one-fourth grain, or 0.016 gram as an emetic, a dose of five grains, or 0.33 gram. Thus it is seen that even if the maximum concentration of copper sulphate necessary to destroy algae in reservoirs were maintained indefinitely, the absorption from daily use would be very far below an amount that could produce the least injurious effect. Taking a solution of one

to one million, which in all cases would be sufficient to prevent the growth of a polluting algal form, it would be necessary to drink something over twenty quarts of water a day before an amount which is universally recognized as harmless would be introduced into the system, while more than fifty quarts would have to be consumed before there would be danger of producing an unpleasant or undesirable effect. As will be seen from the preceding tables the use of copper sulphate at this maximum strength of one to one million would need to be resorted to only in extreme cases, and for a very short length of time. For the reservoir once entirely free from the organisms, a very much weaker solution would be sufficient should any further application be necessary.

The Report of the Massachusetts State Board of Health for 1905 stated as follows:

The objectionable tastes and odors of pond and reservoir waters, which are often attributed to decaying fish and other causes, are, in practically all cases which have been brought to the attention of the Board, caused by the presence of organisms of some kind of kinds which attach themselves to the sides and bottom of the reservoir but in the majority of cases of those kinds which live in suspension in the water. Early in 1903 Dr. George T. Moore formerly Assistant of the United States Department of Agriculture, brought to the attention of the Board the results of investigations which he had made, indicating that the microscopic organisms which are the chief cause of objectionable tastes and odors in the waters of ponds and reservoirs could be destroyed by applying sulphate of copper or blue vitrol to the water in very small quantities and information was also submitted tending to show that bacteria were also destroyed in water brought in contact with metallic copper.

The results of some of the experiments and investigations so far as obtained tend to support the conclusions which had been reached when the matter was first brought to the attention of the Board as to the practicability of the copper treatment for the removal of growths of organisms and bacteria, but the results of other experiments conflict with some of these conclusions. Further study and experiment are necessary before the probable results of the use of copper in preventing objectionable conditions resulting from growths of organisms, or the probable effect of the use of this substance in public water supplies upon the public health can be conclusively determined.

Dr. Moore states that

Spirogyra is often the cause of considerable trouble in a mechanical way and on account of its method of forming resting spores is usually able to withstand the most unfavorable conditions to which it may be subjected in a pond or reservoir. In at least one instance this alga has been the cause of the loss of thousands of dollars by the damage it produced in smothering out the young water-cress plants in the artificial beds constructed for the winter propagation of this vegetable. When the cress is cut for market it necessarily leaves the plants in a weakened condition, and if the *Spirogyra* gets a start it will form a thick heavy mat over the water, which is sufficient to prevent the growth, if not entirely to kill the cress plants.

CHAPTER XIV

A CATALOGUE OF THE MORE IMPORTANT POISONOUS PLANTS OF THE UNITED STATES AND CANADA

For the purpose of facilitating a study of the poisonous plants of this country a partial catalogue is given here. It is based in part on a catalogue of the poisonous plants of Iowa,* issued by the writer and Eusebe D. Fogel (Buchanan). To further facilitate the study of the poisonous plants the following papers on the subject should be mentioned:

Chestnut,¹ Chestnut & Wilcox,² Wilcox,³ Nelson,⁴ O'Garra,⁵ Kennedy,⁶ Peters, Slade & Avery,⁷ Glover,⁸ Crawford,⁹ Heald & Peters,¹⁰ White,¹¹ Johnston,¹² Beasey,¹³ Rusby,¹⁴ Gutenberg,¹⁵ Pammel,¹⁶ Millspaugh,¹⁷ Coville,¹⁸ Peck,¹⁹ J. U. & C. G. Lloyd,²⁰ Stebler & Schroeter,²¹ Schaffner,²² R. Schimpfky,²³ Winslow,²⁴ Farlow,²⁵ Maiden,²⁶ Atkinson,²⁷ Greshoff,²⁸ Wilton,²⁹ Cornevin,³⁰ Halsted,³¹ Robert,³² Crawford,³³ and Marsh.³⁴

*Proceedings of the Iowa Academy of Science 14:147-173. Contributions, Department of Botany, Iowa State College 37:147-176.

1 Principal Poisonous Plants of the United States, Div. Bot. U. S. Dept. Agr. 20:40, (34).
2 Thirty Poisonous Plants of the United States, U. S. Dept. Agr. Farmers Bull. 26:32 f., 4. Rep. Bu. An. Ind. 15:347-420, 169.

3 The Stock Poisoning plants in Montana, Bull. Div. Bot. U. S. Dept. Agr. 26:150, pl. 36.
4 Rep. Bu. An. Ind. 37:91-125, p. 32.

5 Feeding Wild Plants to Sheep, Rep. Bu. An. Ind. 15:421-426.

6 Rep. Neb. Agr. Expt. Sta. 16:14-16, f. 13.

7 Bu. Nev. Agr. Expt. Sta. 1:37, p. 26.

8 Poisoning of Cattle by Common Sorghum and Kaffir Corn, Neb. Agr. Expt. Sta. 77:16.

9 Bu. Cal. Agr. Expt. Sta. 1:1-24, pl. 4.

10 The Poisonous Action of Johnson Grass, Bull. Bur. Pl. Ind. 30, part 4.

11 Ergot and Ergotism, Proc. Bull. Neb. Agr. Expt. Sta. 74:1-11.

12 Dermatitis Venenosa, An Account of the Action of External Irritants upon the Skin, 216:1887.

13 A Manual of the Medical Botany of N. Amer. p. 292, pl. 9, f. 140.

14 Bull. Dept. Bot. 1924. Proc. Soc. Penn. Agr. Soc. 23:33-41.

15 Poisonous Plants of the Vicinity of New York City, 19.

16 Poisonous Plants which Grow in and around Y. C., 21:23.

17 Bull. In. Agr. Expt. Sta. 70:423-444. Poisoning from Combane, Bull. In. Agr. Expt. Sta. 70:213-223, f. 2.

18 Medicinal Plants, 1:99, pl. 99. 2:100-100, pl. 100.

19 L. in. Mus. Herbaria, Cir. Div. Bot. 11:24.

20 Edible and Poisonous Fungi of New York Ann. Rep. State Bot. of State of N. Y. 1894, 103:248, pl. 43. Rep. State N. Y. Mus. Nat. Hist. 48.

21 Drugs and Medicines of N. Amer. Ranunculaceae f. 104, p. 24, f. 185.

22 Beiträge zur Kenntnis der Giften und Weiden der Schweiz. Landw. Jahrb. d. Schweiz. 5:141-155, pl. 20, f. 4.

23 Poisonous and other Injurious Plants of Ohio, Ohio Naturalist 4:16, 32, 48.

24 Deutschlands Wichtigste Giftgewächse im Wirt und Feld nebst einer Abhandlung über Pflanzengifte, pl. 27.

25 Veterinary Materia Medica and Therapeutics, 775.

26 Some Edible and Poisonous Fungi, Bull. U. S. Dept. Agr. Div. of Veg. Path. and Physiology 13:433-476.

27 Plants Reported to be Poisonous to Stock in Australia. N. S. W. Dept. of Agr. Mus. Pub. 477.

28 Mushrooms Edible Poisons, etc. 322. Ithaca, N. Y. (2nd Ed.).

29 Phytotoxins & investigations at Kew No. 10, 1909. Monog. de plantes veneneuses et sapientibus, etc. Batavia, 1909.

30 On the Poisoning of Live Stock, Bull. Ariz. Agr. Expt. Sta. 59:301-167.

31 Des plantes vénéreuses et des empoisonnements qu'elles determinant, 624. Paris, 1893.

32 The Poisonous Plants of New Jersey, Bull. N. J. Agr. Expt. Sta. 135:28.

33 Lehrbuch der Toxikologie, Steigk, 1902.

34 Practical Toxicology for Physicians and Students, Trans. and ed. by L. H. Friedberg.

201. N. Y. 1892.

35 Laboratory work on loco-weed investigation, Bull. U. S. Dept. Agr. Bur. Pl. Ind.

121:39-48.

36 The Loco-weed Disease, Proc. Bull. U. S. Dept. Agr. 386:16.

Many of the plants in this list are certainly not strongly toxic but they frequently produce injurious symptoms in man and animals. The list is by no means complete, it will, however, serve the purpose of calling attention to some of the injurious plants in North America.

It may be noted in this connection that experiments recorded by no means agree as to the effect of certain poisonous plants upon animals. Notice the very discordant results obtained with reference to the poisoning from loco weeds. We may note also the opposite results obtained by experiments reported by Dr S. V. Nelson¹ with reference to feeding *Delphinium Menziesii* to sheep. This writer concluded that when fed to sheep early in May it is not poisonous to them. In the same report² Dr E. V. Wilcox reports a very different experience. This writer reports an arrested heart action and respiration and paralysis of the spinal cord. The animals were fed early in the season, about the same time of the year that the unsuccessful experiments were reported by Dr Nelson. Either these writers were dealing with different plants or the plant is more toxic in some localities than in others. Future experiments only can determine.

EUTHALLOPHYTA

SCHIZOPHYTA

SCHIZOMYCETES, BACTERIA.

Bacteria produce disease in two ways—*First*, as parasites when they derive their nourishment from the living animal; in this case they may cause embolism as in the case of *Bacillus anthracis*, or they may produce within the body products, toxins that are poisonous, as the diphtheria bacillus does. The tetanus bacillus though parasitic produces powerful poisons that when injected even in minute doses cause a fatal termination, producing all the symptoms found in animals having the disease. *Second*, many saprophytic bacteria produce poisonous substances, especially such as occur in putrid flesh, fish, and other decaying substances. The list of such organism is a long one and cannot be given in this connection.

SCHIZOPHYCEAE.

Oscillatoriaceae. Blue-green algae.

Oscillatoria.

Several species in the U. S., probably somewhat injurious.

Nostocaceae.

Nostoc caeruleum Lyngb.

N. muscorum, Ag.

N. commune Vauch.

Dr J. C. Arthur some years ago thought that one of the common blue-green algae, a species of *nostoc*, found in lakes in Northern Iowa and Southern Minnesota, was poisonous. Its poisonous nature was not conclusively demonstrated. These algae have been suspected in other parts of the world. Mr George Francis calls attention to the *Nodularia* occurring in a fresh water lake in Australia. Thirty ounces of a scum fed to sheep produced death. It is also poisonous to horse, dogs and pigs. Many of the algae of this group

¹ Rept. Bur. An. Ind. 1898:421.
² *Id.* 437

produce very disagreeable odors when decomposition occurs, and this plant is no exception to the rule. A few of these algae may be mentioned.

Anabaena flos-aquae Breb.

Anabaena stagnalis, Kg. Both of these are found floating on the water. In their decomposition they produce pig-pen odors.

EUPHYCEAE—Algae

CHLOROPHYCEAE.

Volvocaceae.

Pandorina. Common in stagnant pools, especially in barnyards. The water is repulsive. Cattle will not drink it unless driven to do so. May be injurious. *Volvox* may be placed in the same category.

EUMYCETES---Fungi

PHYCOMYCETES. BLACK MOULDS AND DOWNY MILDEWS.

Mucoraceae.

The species of this family are common; among them are. *Mucor mucedo* L., found on horse manure.

Mucor corymbifer F. Cohn, and *Mucor rhizopodiformis* F. Cohn. Both species are pathogenic.

Mucor stolonifer Ehrh.

Common in the United States. Not pathogenic. Mouldy vegetables, etc.

Mucor racemosus A. Fres.

Is widely distributed in North America. Mouldy vegetables, etc.

BASIDIOMYCETES.

Ustilaginaceae.

Ustilago Zeae (Beckm.) Ung. Corn smut is supposed to be poisonous to cattle, but the evidence is not very conclusive.

Ustilago avenae (Pera.) Jena. The common loose smut of oats is supposed to be injurious in large quantities, the same may be said of other smuts occurring upon cereals. Among these are barley smut *Ustilago hordei*, *U. nuda*, *U. tritici*.

Ustilago neglecta Niesl.

Pigeon Grass smut is thought by many farmers in Iowa to cause poisoning, especially abortion. Prof. Power, formerly of the University of Wisconsin, found present in this smut a small quantity of ergotin.

Ustilago utriculosa Tul. Smartweed Smut.

This common smut is said to produce irritation and frequently is the cause of trouble at husking time. This fungus is widely distributed in North America.

Tilletiaceae.

Tilletia foetens (R. & C.) Trel.

Stinking smut of wheat is not common in this state, but when it occurs in flour it causes bad odors. The spores give to the flour a dark color and make it unsalable.

Melampsoraceae.

Coleosporium solidaginis (Schw.) Thum.

Parasitic on golden rod and some other plants of the order. Common in

the state. A number of horses in Black River Falls, Wisconsin, a few years ago, became diseased, it is thought, by means of this rust. It may produce a form of Mycotic stomatitis.

Pucciniaceae

Puccinia graminis Pers.

Wheat and oat rust, especially the uredo stage, produces inflammation of the mucus membrane of the mouth and nose. The dust coming from the straw when the grain is threshed often causes serious disturbances. Other rusts might be mentioned in this connection, like *Puccinia coronata*, Cda., the uredo-spores of which have an effect similar to that of the common grass rust. The above rusts are widely distributed in North America. *Uromyces trifolii*, Clover Rust, is widely distributed on red clover and, according to several authorities, is responsible for mycotic stomatitis.

Polyporaceae

Boletus felleus Bull.

It has a bitter taste and is poisonous.

Agaricaceae

Lepiota morgani Pk.

This fungus is very common in the fall in meadows, pastures, and lawns. The cap is from 5 to 11 inches across. The gills or radiating plates beneath the cap are brown when mature. The lower part of the stipe is somewhat enlarged but no cup occurs. This when eaten by some people is known to produce poisoning.

Amanita muscaria L. Fly agaric.

Used to poison flies in Asia. Poisonous to persons.

Amanita phalloides Fr. Amanita.

This species is very poisonous and no one should eat so-called mushrooms unless he is familiar with them.

Amanita spreta Pk.

This species is said to be poisonous in Eastern North America.

Amanita rubescens Fr.

This species is said to be edible, but authorities recommend that it should be used with great care. The same is true of *A. solitaria*. Both species are found in E. North America.

Amanita verna Bull.

This species is closely related to *A. phalloides*, but it is pure white in color, it is deadly poisonous. *A. virosa* Fr belongs to the same category. *A. frostiana* Peck is also poisonous.

Clitocybe illudens Schw. Jack-my-Lantern.

This species is common in Eastern North America and is phosphorescent. While it is not dangerously poisonous, it is not edible.

Phallaceae

Ithyphallus impudicus (L.) Fries

The common stinkhorn is probably poisonous. Its disagreeable odor, however, would seem to render it distasteful to animals.

Lycoperdaceae

Lycoperdon Bovista L.

The giant puffball is edible in fresh condition, when the flesh is white, but

in the mature form is considered poisonous; the same may be said of other puffballs.

ASCOMYCETES

Aspergillaceae.

Aspergillus glaucus Link.

This fungus is supposed to produce staggers. Frequently found in mouldy hay and gives rise to digestive disorders. Mycotic stomatitis.

Aspergillus niger van Tieghem.

This mould also occurs in mouldy hay and other mouldy substances and like the preceding species is injurious.

Aspergillus fumigatus Fr

Commonly found on decaying substances, especially mouldy hay. It is pathogenic. Found in the ear of man as a parasite.

Aspergillus flavus Link.

Also pathogenic.

Hypocreaceae.

Claviceps purpurea Fr

Found on many different grasses, especially wild rye, cultivated rye timothy and quack grass. Produces a disease known as ergotism. The fungus is very poisonous, causes dry gangrene and abortion.

FUNGI IMPERFECTI

Many of the imperfect fungi undoubtedly produce Mycotic stomatitis.

Oidium albicans Robin.

Commonly found in the mouth of sucking animals of different kinds, especially calves. Blastomycosis produced by *Oidium* or *Saccharomyces*.

Fusarium equinum Norg. Umatilla horse mange.

Dr Mohler and others have demonstrated that the fungus is parasitic and produces this mange.

Fusarium roseum is commonly found on corn and other mouldy grain. It is probably concerned in forage poisoning. Dr Burrill and Prof. Barrett report several forms of *Fusarium* on corn in Illinois.

Diplodia zeae Lev.

This disease is widely distributed on the sheaths and blades of corn in the United States. It is reported as common in Illinois by Burrill and Barrett and in Nebraska by Heald and Wilcox. It is also common in the state of Iowa. Dr Erwin F Smith suggests some connection between pellagra and this mould.

Sporotrichum Furfur Rob. Favus.

Sporotrichum tonsurans. Barber's Itch.

Polythrincium trifolii Kunze. *Helminthosporium gramineum* and other fungi may be responsible for mycotic stomatitis. The rape fungus, *Polydesmus estiosus* Kuhn has long been associated with this disease.

EMBRYOPHYTA

PTERIDOPHYTA

Polypodiaceae. Ferns.

Pteris aquilina L.

The common brake is found in eastern states and across the continent,

Oregon and Washington. It produces a large, strong root-stock and is said to be poisonous to cattle and horses.

Adiantum pedatum L. Maiden Hair Fern.

Said to be poisonous. Widely distributed in Northern States.

Aspidium marginale L.

The root-stock is used as a remedy for the expulsion of tapeworm and undoubtedly sometimes produces poisoning. E. N. Am., Canada to Mo.

Cystopteris fragilis (L.) Bernh. Bladder fern.

This and many other ferns, according to Greshoff, contain HCN

Osmundaceae. Royal Ferns.

Osunda claytoniana L.

This fern is widely distributed in the eastern states. The odor is not pleasant and the plant is undoubtedly more or less poisonous.

Equisetaceae. Horsetails.

Equisetum arvense L.

The common horse-tail is supposed to be injurious to horses, as reported in Vermont by Dr. Richman and Professor Jones, and reports of poisoning in other parts of the country are recorded. In Europe it has long been suspected of being poisonous.

Equisetum robustum A. Br.

It is common in the state and, like the preceding, is considered poisonous.

A suspected case of poisoning recently was reported to the writer from Iowa.

EMBRYOPHYTA SIPHONOGAMA

(Flowering Plants)

GYMNOSPERMAE Conifers and Allies

Taxaceae. Yews.

Taxus canadensis Marsh. American yew.

Found in northern U. S., especially on calcareous sandstone rocks or in sandy woods. The European species has long been regarded as poisonous to stock. Cases of poisoning have been reported in this country as well. It contains toxic substances, one of which is known as *taxin*.

Taxus brevifolia Nutt. Yew. Common on Pacific Coast.

Pinaceae. Pines and Junipers.

Juniperus communis L. Common juniper.

Poisonous, especially the oil obtained from the juniper berries. Rusby and others refer to the poisonous nature of this plant, and Schaffner records that goats are poisoned by eating the leaves. Common only in lime and sandstone rocks in Northern U. S. and Rocky Mountains.

Juniperus virginiana L. Red Cedar.

Common in Northern and Eastern N. Am. to Texas. The leaves contain the same principle as that found in the juniper, and according to Schaffner, are poisonous to goats. The oil produces abortion and poisoning has also resulted from its use. The oil of Cedar has well known antiseptic properties. It contains an aromatic body, cedren, oil of cedar from which cedren-camphor has been obtained.

Juniperus scopulorum Sarg.

Probably poisonous, like the preceding.

Juniperus sabina L. Swedish juniper.

Poisonous, like the preceding species. Cultivated.

Juniperus horizontalis Moench. American Savin.

Poisonous, Canada to Minn., in swamps.

Juniperus occidentalis Hook and var. Red cedars

Common from the western slope of the Rocky Mountains westward

Sequoia sempervirens Endl.

Redwood leaves are said to be poisonous.

ANGIOSPERMAE

Typhaceae.

Typha latifolia L. Common Cat-tail.

Said to be poisonous to cattle.

MONOCOTYLEDONEAE.

Alismaceae.

Alisma Plantago-aquatica L. Water plantain

Recorded as being poisonous. Swamps.

Sagittaria latifolia Willd. Large arrow head

The milky juice is somewhat bitter. The plant is edible, when cooked

The root stocks of several species of the genus are eaten by the Indians and in China. Swamps.

Gramineae.

Zea Mays Gärtn. Numerous cases of poisoning have been recorded from the consuming of corn. This, however, is probably due to the production of a poison from the growth of mould or bacteria. The feeding of large amounts of dry fodder produces impaction of the stomach.

Andropogon sorghum Brot. Sorghum.

Second growth sorghum has frequently been reported as poisonous to live stock; this is due to the formation of HCN in the wilted leaves. *Calamagrostis*, *Milium* and some other grasses contain the same substance.

Setaria italica, Kunth. Millet.

Both the German Millet and the Hungarian Grass are poisonous to horses, acting especially on the kidneys. The poisoning is probably due to a glucoside.

Stipa spartea Trin. Needle Grass, or Porcupine Grass.

The sharp pointed callus often inflicts serious injuries; the fruits work their way under the cuticle into the flesh of the animals, and in some instances they have even penetrated the intestines.

Stipa comata Trin. Western needle grass.

Somewhat injurious, like the preceding. West of Missouri River

Stipa robusta Vasey. Sleepy grass.

This grass has been suspected of producing stupor in horses. Rocky Mountains. Some species produce HCN.

Avena sativa L. Common oats.

The chaff of this grass sometimes produces balls in the stomachs of horses, known as phytobezoars.

Avena fatua L. Wild oats.

Common in a few counties in northern Iowa. Common in N West., Cali-

fornia, Rocky Mts. Sometimes causes mechanical injuries on account of the pointed callus of the fruit.

Lolium temulentum L. Darnel.

The grain of this grass is injurious when ground in with flour. It produces stupor and symptoms resembling drunkenness. The poisoning is due to the fungus found in the seed. Principle lolin, a glucoside.

Hordeum vulgare L.

The chaff and awns of barley are often injurious, especially when coming in contact with the mucous membrane, not only in man, but in lower animals.

Hordeum jubatum L. Squirrel-tail. Wild Barley.

Common throughout the west. This grass produces mechanical injuries in animals that feed on hay containing it, the awns working their way in between the teeth and maxillae, where they cause inflammation and the formation of pus. *G. pussonianum*, *H. cespitosum*, *H. secalinum*, etc., produce similar injuries.

Sitamon Elymondes Raf. Squirrel tail.

Common Rocky Mts. to Pacific Coast. Troublesome like Wild Barley.

Agropyron repens Beauv. Quack grass.

Widely distributed in northern Iowa. Produces a slight irritation of the mucus membrane. Contains tannin.

Araceae.

Arisaema triphyllum (L.) Schott. Jack-in-the-pulpit.

It is widely distributed in the north. The corm is known to be very acrid and poisonous, but when boiled or roasted the poisonous substance is expelled.

Arisaema Dracontium (L.) Schott. Dragon's head.

Widely distributed, especially in eastern and central Iowa and northern states. The corm is somewhat acrid and is used to destroy insects, it is said to be a good vermifuge. The action of the plant in fresh condition is somewhat similar to that of ammonia.

Symplocarpus foetidus (L.) Raf. Skunk Cabbage.

Local only in a very few places. Said to be poisonous, causes vomiting, and temporary blindness. The juice is acrid and the plant has a very disagreeable odor.

Liliaceae

Zygadenus elegans Ph. Swamp Camas.

Common in northern Iowa, Minnesota and westward. In the western states it is regarded as poisonous to cattle and sheep, occasionally causing death. It is not as poisonous as some other species of the genus.

Zygadenus venenosus Wats. Death Camas. Common in the Rocky Mountains and westward. The bulb is especially poisonous.

Melanthium virginicum L. Bunch-flower.

Common on low grounds in eastern Iowa and South and East. The root stocks are regarded as poisonous, but reports have come to us of the poisonous effect on horses, of the leaves and stems, when occurring in hay.

Veratrum woodii Robb. False Hellebore.

Southeastern Iowa to Mo. Poisonous like the eastern white Hellebore and the western California Hellebore. Probably contains jervin, cevadina and cevadin.

Veratrum viride Ait. Common swamp hellebore.

• Eastern North America; swampy places. Seeds are poisonous, also herbage under some conditions.

Veratrum californicum Durand. California hellebore.

Rocky Mountains to Pacific Coast.

Amanthum Muscaetoxicum (Walt.) Gray. Fly poison. Crow poison.

Common along the Atlantic Coast and Long Island southward.

Erythronium purpurascens S. Wat. Dog-tooth Violet.

California. Contains saponin. Said to be poisonous.

Leucocrinum montanum Nutt. *Leucocrinum*.

Northwestern United States. Thought to be poisonous to sheep.

Nothoscordum bivalve (L.) Britton. Crow poison.

Southern United States.

Aletris farinosa L. Cohic root.

Said to be poisonous. Sandy soil. E. N. Am.

Medeola virginiana L. Indian cucumber root.

From New Brunswick to Minn., and Florida. Said to be poisonous.

Chamaelirium luteum (L.) Gray. Blazing Star.

Said to be poisonous. Mass. to Ark. and Fla. low grounds.

Chlorogalum pomeridianum Kunth. Soap plant.

Pacific Coast. Contains saponin.

Allium canadense L. Wild onion.

Widely distributed, common in low pastures, N. U. S. Milk is flavored where cattle feed on the plant. The *A. unifolium* of California is poisonous.

Allium tricoccum Ait. Wild Leek.

Eastern and northern States. Taints milk like preceding.

Lilium superbum L. Turk's-cap lily.

According to Schaffner, this species produces dermatitis. The bulbs produce mental exhaustion and headache.

Asparagus officinalis L. Asparagus.

According to Dr. White, in his *Dermatitis Venenata*, persons who constantly work with asparagus may have the skin somewhat blistered.

Convallaria majalis L. Lily-of-the-valley.

All parts of this plant are very poisonous to man and domestic animals. Contains two glucosides, convallamarin $C_{44}H_{44}O_{14}$, and convallarin $C_{24}H_{24}O_{11}$.

Trillium grandiflorum, Salisb. Large flowered Trillium.

This is used as an emetic and contains a principle which has been called trillin, found in a few other species of the genus.

Trillium erectum L. Erect Wake-robin.

The root stock of this species is somewhat poisonous.

Smilax rotundifolia L. Round-leaved Greenbrier.

Widely distributed in the northern states. Dr. Schaffner reports a case of poisoning from eating the young leaves of these plants. The spines are injurious in a mechanical way; they cause inflammation and pus formation.

Haemodoraceae.

Lacianthes tinctoria (Walt.) Ell. Red root.

Mass. to Florida. It is said that white hogs are subject to poisoning from this plant.

Amaryllidaceae.

Narcissus poeticus L. Narcissus.

This plant and other species regarded as poisonous in Europe

Agave americana L. Century plant.

Used as a fish poison in some countries. Cultivated.

Zephyranthes atamasco (L.) Herb. Atamasco lily

From Pa. to Florida. Said to produce the disease "staggers."

Belamcanda chinensis (L.) D C Blackberry lily.

Rhizome acrid, said to be deleterious, E. N. Am. Naturalized

Dioscoreaceae.

Several species of *Dioscorea* or yams are poisonous.

Iridaceae.

Iris versicolor L. Large Blue-flag.

The underground root stocks of this plant are known to be very poisonous.

Dr. Rusby thinks there is some danger "that it might be eaten in mistake for *Calamus*, which is commonly known as Sweet-flag. If so it would prove seriously if not fatally poisonous, as its well-known emetic-cathartic properties, even when toned by drying and keeping, are powerful, and in a fresh state would be decidedly violent." It contains the resinous body iridin

Orchidaceae.

Cypripedium spectabile Swz. Showy Moccasin flower

Causes dermatitis resembling that produced by Poison Ivy. A great many persons are more or less susceptible to this form of dermatitis. Swamps in Northern States.

Cypripedium pubescens Willd. Yellow Lady-slipper

Poisonous like the preceding. This species is more widely distributed

Cypripedium candidum Muhl. Small white Lady-slipper

Less poisonous than the preceding. At one time common in the marshes in the north, but rapidly disappearing.

DICOTYLEDONEAE.

Salicaceae

Populus balsamifera L. Balsam Poplar

Said by Dr. Halsted to be poisonous.

Juglandaceae.

Juglans nigra L. Black walnut.

It is commonly believed that the bitter principle *Juglanton* found in the fruit and leaves of this black walnut are more or less poisonous at least it is thought that different species may poison the soil and prevent the growth of other plants, this is, however, not well substantiated. Probably contains nuclei, an acrid body, causing an irritation which resembles scarlatina

Fagaceae. (Beech Family)

In Europe beech nuts are known to produce some form of enteritis and the burr of the chestnut, *Castanea dentata*, produce mechanical injuries. Species of oak (*Quercus*) particularly the acorns of *Quercus rubra* are injurious to cattle and the nuts of European oaks produce death in hogs and cattle. This may possibly be due to the tannin and glucoside contained in the seed. The *Q. Chrysolepis* of California is poisonous.

Urticaceae.

Maclura pomifera (Raf.) Schneider Osage orange.

Cultivated especially in the southern part of Iowa to Nebraska and Southward. The leaves and fruit are more or less poisonous. The thorns upon the plant produce serious injuries giving rise to inflammation.

Humulus Lupulus L. Common Hop.

Hop pickers often have an inflammation of the hands. The plant is a sedative and contains cholin, lupulic acid and oil of humulus.

Cannabis sativa L. Hemp.

Naturalized in many parts of the North. The narcotic effect of the resin of the plant is well known and in India an intoxicating drink is made from the juice of the leaves. Contains the substances cannabin, and oxycannabin.

Urtica dioica L. Stinging nettle.

The urticating properties of our common nettle are known to all who have had any experience in collecting the plants; there is at first a reddening, followed by a swelling, intense burning, and a small amount of itching. It is said to contain formic acid. The Western Nettle *U. holosericea* of Utah has similar properties.

Urtica gracilis L. Slender nettle.

Injurious like the preceding but more widely distributed.

Laportea canadensis Gand. Wood nettle.

This plant is even more widely distributed than the preceding nettles. It is found in deep woods. It produces an irritation of the skin like that caused by common nettle.

Loranthaceae.

Poisoning has been reported from the European mistletoe *Viscum album* and there are a few references in America regarding poisoning by the false mistletoe *Phoradendron flourensii*. Possibly the *Arceuthobium* may also be injurious.

Aristolochiaceae

Arristolochia Clematidis L.

Atlantic States from New York to Md. Some of the European species of *Aristolochia* are poisonous like the *A. Clematidis* which is naturalized along the Atlantic Coast.

Polygonaceae.

Rumex acetosella L. Sheep sorrel.

The plant is widely distributed in the U. S. and is becoming more common. Said to be poisonous to horses and sheep. Contains oxalic acid.

Rumex crispus L. Sour dock.

The plant is an astringent and is looked on with suspicion as are some of the other species of the genus.

Fagopyrum esculentum Moench. Buckwheat.

A dermatitis produced by the eating of buckwheat cakes is well known to most people and occasionally where screenings of this material are fed in quantities to hogs a similar rash is produced. Buckwheat straw is also considered poisonous. The plant contains the glucoside indican.

Fagopyrum tataricum Gaertn.

Poisonous like the preceding.

Polygonum acre H B K. Smartweed.

The acrid properties of many of the species of Polygonaceae are well known. This species is widely distributed in the north. Contains probably polygonic acid

Polygonum hydropiper L. Smartweed.

Poisonous like preceding.

Chenopodiaceae

Chenopodium anthelminticum L. Worm-seed.

Occasionally reported in the state. Cases of poisoning from the oil of the seeds have been reported in medical literature. Contains the volatile oil of worm-seed. This is a narcotic-acrid poison.

Chenopodium ambrosioides L. Mexican Tea

This species is occasionally reported with properties like the preceding.

Kochia scoparia (L.) Schrad. Summer cypress.

This plant contains saponin and, according to Greshoff, the extracts from the seeds of another species froth in a solution up to 1 to 700. These plants must be regarded as slightly toxic in their effect. We may add here that saponin has also been found in *Eurotia ceratoides*. *Eurotia lanata* is used as a forage plant in the West; it is commonly called winter fat.

Chenopodium mexicanum Moq. Mexican Lamb's Quarters.

This plant contains saponin and is known to be poisonous.

Amaranthaceae

Amaranthus retroflexus L. Green pigweed.

Common everywhere in eastern north America, also in the great basin. O'Garra reports bloat from it in Nebraska.

Amaranthus spinosus L. Thorny pigweed.

The species are injurious.

Greshoff states that the leaves of *A. hypochondriacus* give an extract which froths strongly and contains saponin. The saponin is only slightly toxic.

Atriplex Nuttallii S. Wats. Salt bush.

The leaves of the salt bush contain saponin and Greshoff also found the same substance in *A. halimifolia*, *A. hortensis* and *A. laciniata*. He states that the haemolysis of the seeds of some of the species is moderately great. In China a skin disease known as Atriplicismus is caused by a species of *Atriplex*.

Beta vulgaris. Mangolds and Sugar Beets.

The feeding of mangolds and sugar beets to sheep causes renal calculi.

Sarcobatus vermiculatus Hooker. Grease wood.

Frequently produces mechanical injuries. According to Chestnut one man lost over 1000 sheep, probably due to bloat, caused by this plant.

Phytolaccaceae

Phytolacca decandra L. Pokeweed.

The roots and seeds contain a very poisonous substance. The young shoots are eaten as greens, probably the poisonous principle is dissipated on boiling the plant. Found from southern Iowa Eastward and Southward. Contains phytolaccin.

Caryophyllaceae

Stellaria media L. Chick-weed.

This has been reported as poisonous, although the seeds are eaten by birds.

Agrostemma githago L. Corn cockle or cockle.

Generally found in wheat fields. Screenings are often sold as stock food.

and several cases of poisoning from food that contained screenings of cockle have been reported. When cockle is in flour, it is poisonous. Several cases of poisoning from flour containing cockle are on record. Cockle is said to be especially poisonous to poultry. Contains the substances saponin, sapogenin and the alkaloid agrostemmin.

Silene antirrhina L. Sleepy catchfly.

Very widely distributed in the north. Said to be poisonous.

Silene noctiflora L.

Widely distributed in the north. Said to be poisonous.

Widely distributed in northern states. Clover fields. Native to Europe, probably also poisonous.

Saponaria officinalis L. Bouncing Betty.

This plant is said to be somewhat poisonous. Naturalized in the East Ry Mts. and Pacific Coast.

Vaccaria vulgaris Host. Cow cockle.

Common only in grain fields, seeds said to be poisonous, like corn cockle.

Nymphaeaceae

Nelumbo lutea Pers. The American Nelumbo.

The root stock is used by the Indians for food. According to Schaffner it is said to be used to destroy cockroaches. Roasting dispels the poisonous principle.

Ranunculaceae

Hydrastis canadensis L. Orange Root. Golden Seal.

In northeastern Iowa and eastward. Contains the alkaloids hydrastin, berberine, and xanthopuccine. Hydrastis causes severe ulceration and catarrhs, inflammation.

Caltha palustris L. Marsh marigold.

The leaves of the marsh marigold are eaten, but the poisonous principle is dissipated on boiling. Plant found on low grounds, especially in northern states. The related species with whitish flower *R. leptosepala* of the Ry Mts. must be regarded with suspicion.

Actaea alba Mill. White baneberry.

More or less poisonous, but generally not eaten by live stock. Found in woods more or less widely distributed in northern states.

Actaea rubra Willd. Red baneberry.

Widely distributed in the state, but never abundant. Berries poisonous.

Delphinium consolida L. Field Larkspur.

Naturalized from Europe. Poisonous and fatal to cattle, frequently cultivated as an ornamental plant. It contains several poisonous alkaloids. The alkaloids, delphinin, delphusin, delphinoidin and staphisagrin occur in *D. Staphisagria* and may be looked for in some of our native larkspurs.

Delphinium carolinianum Walt. Carolina Larkspur.

Native to prairies, especially gravelly knolls. Reported as fatal to cattle.

Delphinium exaltatum Aut. Tall Larkspur.

Frequently cultivated, native to Europe.

Delphinium tricornis Michx.

Produces fasciated tuberos roots. Common in southern Iowa. Very poisonous to cattle.

Delphinium hesperium Gray. Larkspur.

California. Said to be poisonous to cattle. Also the *D. nudicaule* which has narcotic properties.

Delphinium menziesii DC. Western purple larkspur.

Common in Montana and Westward. According to Drs. Wilcox and Chestnut it is poisonous, although Dr. Nelson fed 24½ pounds of fresh leaves to sheep without injury. Montana and west.

Delphinium geyeri Greene. Wyoming Larkspur

Common in Wyoming, Colorado, and Nebraska. It is known as the poison weed.

Delphinium recurvatum Greene. Larkspur

Common in Southern California. Said to be fatal.

Delphinium scopulorum Gray. Tall Mountain Larkspur

Rocky Mountains to Canada. The Canadian Department of Agriculture states that it is poisonous to cattle.

Delphinium trollisfolium Gray

Common along the Pacific Coast, especially California, Washington, and Oregon. It is known as cow poison in Humboldt county, California. Chestnut says perhaps it is not equally poisonous throughout all stages of its growth.

Aconitum uncinatum L. Wild Monk's-hood.

Native to a limited area in northeastern Iowa, Wisconsin and Eastward. Contains aconitin. Root, flowers and leaves are poisonous.

Aconitum napellus L. European aconite.

Commonly cultivated in gardens. This plant is powerfully toxic; it contains several important alkaloids, as aconitin, pseudoaconitin, and aconin. Bodily heat is reduced by aconite. Winslow states that the smallest fatal dose to a man is a teaspoonful of aconite, equivalent to about gr XXX of the crude drug. The minimum lethal quantity of aconitin is 1-16 of gr for man. Large doses produce death by paralyzing the heart.

Aconitum columbianum Nutt

Common in the Rocky Mountains and westward. In swamps near springs and banks at higher altitudes in Colorado and Utah.

Aconitum noveboracense Gray. Aconite.

Northern States.

Aconitum reclinatum Gray. Trailing Wolf's-hane

Alleghany Mountains.

Helleborus viridis L. Green Hellebore

This plant is sometimes an escape from cultivation from Long Island to West Virginia. All parts of the plant are poisonous and have long been so recognized in Europe.

Anemone nemorosa L. Wood Anemone.

The common wind flower is said to be a local irritant. Common in woods of N. States.

Anemone patens, var. *Wolfgangiana*, (Bess.) Koch. Crocus, Sand flower. Pasque flower. Wind flower.

This plant is local in central and southern Iowa, but in northern Iowa it is common on gravelly knolls, also in Ill., Wis., British America, Rocky Mountains, Nebr., Col., New Mexico. It is a well known irritant containing the bitter substance anemonin.

Clematis virginiana L. Virgin's bower

This plant is widely distributed in the state along with other species of the

same genus. The herbage is said to be acrid and caustic. The juice of some species of the genus causes blisters, or even ulcers. The fresh leaves of the *C. erecta* are used as a vesicant in Europe, especially by beggars, hence sometimes called beggar's weed.

Clematis Fremonti Wat. *Clematis*.

Common from Missouri to Kansas.

Clematis Pitcheri Torr. & Gray.

From Southern Indiana to Texas.

Clematis ligusticifolia Nutt.

Very abundant in the Rocky Mountains west to the Pacific Coast along streams at a lower altitude. Greshoff reports saponin in a large number of species, notably *C. Pitcheri*, *C. erecta*, and others, also hydrocyanic acid in *C. Fremonti*.



Fig. 49a. Common Crowfoot (*Ranunculus acris*). An acrid poison. (U. S. Dept. Agr.).

Ranunculus acris L. Tall buttercup.

Poisonous, causes inflammation when it comes in contact with mucous membrane. Rarely found in Iowa, but sometimes naturalized, common eastward.

Ranunculus septentrionalis Poir. Creeping Buttercup.

Widely distributed in low grounds. Acrid like the preceding species.

Ranunculus fascicularis Muhl. Tufted buttercup.

Found in eastern Iowa, as far west as the Iowa River, common north and east. Probably poisonous like the preceding species.

Ranunculus bulbosus L. Bulbous Crowfoot.

Common eastward, naturalized from Europe.

Ranunculus abortivus L. Crowfoot.

Common weed in many parts of the north, the leaves are quite acrid and have a sharp, peppery taste.

Ranunculus accleratus L. Cursed crowfoot.

A very poisonous species, especially to cattle, since it grows in marshes along with other herbage and is often eaten with other forage plants. Contains anemonol and anemonie acid.



Fig. 19b. Common Poppy (*Papaver somniferum*). Narcotic. (Louis Pammel)

Anonaceae. Custard Apple Family

Asimina triloba Donal. Pawpaw

It occurs in eastern Iowa as far north as Clinton and Dubuque. Common Southward. The pawpaw is commonly eaten, but a case of poisoning is reported.

Trollius laxus Salisb. Spreading globe flower

From western Connecticut to Michigan and the Rocky Mountains. Several species contain saponin and *T. europaeus* is considered poisonous in Europe. It contains saponin.

Thalictrum revolutum DC. Meadow Rue.

Common in low moist meadows in Western North America. All of the species of meadow rue are more or less acrid. The *T. polycarpum* of California is poisonous to man according to Chesnut

Magnoliaceae. Magnolia Family

Magnolia grandiflora L. Magnolia.

The flowers of this plant are said to be injurious

Lilium floridanum Ellis. Amise tree.

Native to Florida. The leaves are poisonous to stock.

Berberidaceae. Barberry Family

Berberis repens Lindl. Trailing Mahonia.

Cultivated. Poisonous. According to Schaffner, the berries are injurious to birds. Probably contain the alkaloids berberin and oxycanthin which occur in the common barberry.

Berberis Aquifolium Pursh. Oregon grape

This plant also causes poisoning. Cultivated. In the Cascade Mts., Oregon, Washington.

Podophyllum peltatum L. May Apple or Mandrake.

This plant is widely distributed east of the Missouri divide. The roots and leaves are drastic and are known to be poisonous. It is said also that when the leaves are eaten by cows, they produce injurious milk. The roots, according to Dr White, are irritating to the eye, nose, mouth and skin, and contain picropodophyllin, podophyllotoxin. The podophyllin is a resinous mixture.

Canthophyllum thalictroides Mich. Blue Cohosh.

Widely distributed from Iowa eastward and northward. Said to be poisonous.

Menispermaceae.

Menispermum canadense L. Canadian Moonseed

This plant is widely distributed in woods in Iowa and the north. Contains menispermicin, and menispermicin. A case is reported of the death of three boys from eating the berries in mistake for grapes. The *Cocculus indicus* is a well-known remedy for the destruction of pediculi and is known to be poisonous. It contains picrotoxin, cocculin and an alkaloid menispermicin.

Calycanthaceae. Allspice Family

Calycanthus floridus L.

Cultivated in Southern Iowa and in the southern states. The aromatic properties of the flowers resemble those of strawberries. This makes it a very desirable cultivated ornamental plant. It contains an active principle calycanthin. Chesnut records it as poisonous.

The other species of *Calycanthus* may also be regarded as poisonous.

Lauraceae. Laurel Family.

Sassafras officinale Nees. Sassafras.

According to Schaffner the berries of this plant are reported to be poisonous. This species is native to southeastern Iowa, southward and eastward.

Umbellularia Californica (Arnot) Nutt. California Laurel.

California. Leaves smart. Used to drive fleas away.

Papaveraceae. Poppy Family.

Papaver somniferum L. Opium Poppy.

Opium is obtained from the common garden poppy. This species is widely cultivated in the state, and is frequently spontaneous. The more common alkaloids found in the poppy plant are morphin, codein and narcotin. The seeds of poppy are sometimes used to spread on top of cookies and bread.

Papaver rhoeas L. Corn Poppy.

This species is occasionally cultivated and sometimes spontaneous, the milky juice contains some poisonous alkaloids, as papaverin and codein.

Papaver dubium L. Long Smooth-fruited Poppy

Cultivated, native to Europe, naturalized in Eastern North America.

Papaver orientale L.

This handsome, showy plant is frequently cultivated for ornamental purposes and contains a large amount of milky juice. This plant must be regarded as suspicious.

Argemone mexicana L. Mexican Prickly Poppy

The Mexican prickly poppy, with pale yellow or yellowish petals, is cultivated in gardens. The latex not only contains narcotic principles, but the prickly pods sometimes cause mechanical injury and set up inflammation. The latex contains morphin, common in Texas to Kansas.

Argemone intermedia Sweet.

This prickly poppy of the plains contains a narcotic substance in its latex. The prickly leaves of the plant cause serious mechanical injury and inflammation. Common from Nebraska to Utah, Kansas and west.

Sanguinaria canadensis (Mill) L. Bloodroot

This widely distributed plant of Eastern North America is well known to nearly everyone. The root is poisonous and contains the principle sanguinarine, found in the red latex. It is sharply irritating, especially to mucous surfaces. Contains the alkaloid sanguinarin.

Chelidonium majus L.

Celandine is occasionally cultivated and spontaneous eastward. The orange yellow juice or latex is an irritant. The fresh juice produces inflammation and blisters, and it poisons the skin if handled so as to crush the leaves or stem. Contains chelerythrin $C_{21}H_{11}NO_4$, with a burning taste, and glaucin.

Stylophorum diphyllum (Michx.) Nutt. Celandine Poppy

Common from Wisconsin to Tennessee and Pennsylvania. According to Schlotterbeck this plant contains the following substances which are regarded as injurious: protopin, stylopin, sanguinarine and others.

Eschscholtzia Douglassii (Hock & Arn.) Walp.

Narcotic used to stupefy fish. California.

Fumariaceae.

Fumaria officinalis L. Fumitory

It has long been regarded as poisonous in Europe. Naturalized in waste places eastward.

Corydalis aurea Willd. Golden Corydalis

A related exotic species which is said to be very poisonous. The above species and others of the family have been considered poisonous.

Cruciferae. Mustard Family

Lepidium sativum L. Garden Cress

Native to Europe, but occasionally escaped from cultivation. The Garden Cress is used as greens. It is a local irritant.

Lepidium apetalum Willd. Small Pepper-grass.

Widely distributed in this state, and like the other pepper-grasses, may produce synapsm.

Radicula armoracia (L.) Robinson. Horse-radish.

This is poisonous, like the preceding species of this order that have been

mentioned. Dr. Rusby refers to its irritating properties when taken in excess especially because of its action upon the urinary organs. One case referred to by Dr. Johnson was extreme and serious.

Sisymbrium officinale Scop. Hedge Mustard.

This plant is widely distributed in this state, as a weed, and produces more trouble than mustard.

Sisymbrium altissimum L. Tumbling Mustard.

This weed is common in grain fields in the north and may thus find its way into wheat screenings which should be fed with caution. It probably produces sinapism, as do the other plants of the order. It is also found in Iowa.

Brassica arvensis (L.) Ktze. Charlock.

This weed produces sinapism. After the application of the powdered material, there is a sense of burning. The volatile oil of mustard is a powerful irritant, and caustic, and should be used with caution.

Thlaspi arvense (Tourn.) L. Penny Cress.

This weed is said to be injurious to animals in Canada and is more or less avoided by them in pasture. It is widely distributed in Northern United States, but is more abundant in Canada.

Capparidaceae. Caper Family

Cleome serrulata Pursh. Stinking Clover

This plant is more or less pungent and acrid; it is widely distributed from Missouri River westward across the continent, especially along railways, etc. The flowers are showy and purple.

Cleome lutea Hook. Western Cleome.

This plant is a western species with yellow flowers and has similar properties to the one named above.

Polanisia graveolens Raf

Fetid annual, with glandular hairs and common in sandy soils in the eastern States. The *P. trachysperma* T. & G. is similar to the preceding. This plant is found from Iowa to Kansas.

Sarracenaceae.

Sarracenia purpurea L. Side-Saddle Flower

Probably not native to Iowa, although it is found distributed with *Drosera* in sphagnum bogs. It contains the substance sarracenin. The root produces diuresis, gastric excitation, and an increased, irregular action of the heart. It produces papular eruptions changing to vesicular with depression as in smallpox. The plant was formerly used medicinally by the Indians.

Droseraceae. Sundew Family

Drosera rotundifolia L. Sundew.

Sundew is said to be poisonous to cattle. In bogs in the northern states. Greshoff reported hydrocyanic acid in this species and in *D. intermedia*. Several other species in Europe were reported as harmful and toxic to cattle.

Crassulaceae. Orpine Family.

Sedum acre L. Stone Crop.

Produces inflammation when applied to the skin of many persons. The juice is acrid and biting.

Other species of this genus are known to be acrid. Several species with yellow flowers are common in rocky soils in the Rocky Mountains.

Saxifragaceae.

According to Greshoff several species of the genus *Saxifraga* of Europe contain hydrocyanic acid. The order also contains the currant and gooseberry which are well known food plants.

Hydrangea arborescens L. Wild Hydrangea.

Common in the Southern States and woods from Southern Iowa eastward to New York. According to Greshoff this species contains saponin; hydrocyanic acid occurs in the well known ornamental plant *H. Hortensis*.

Jamesia americana T. & G. *Jamesia*.

This plant is widely distributed from New Mexico to Montana. The leaves of this Rocky Mountain shrub contain hydrocyanic acid.

Philadelphus grandiflorus Willd. and *P. coronarius* L.

Mock Orange or Syringa. Native in the mountains from Virginia to Florida. The well known cultivated syringa contains saponin, according to Greshoff the Rocky Mountain *P. microphyllus* also contains saponin.

Dentzia. Several species of *Dentzia*, as *D. staminea*, contain saponin.

Chrysosplenium oppositifolium L.

This plant contains some saponin and the *C. tetrandum* Fries is known to be poisonous to sheep.

Rosaceae. Rose Family

Fragaria vesca L. European Strawberry.

Found on sand-stone and limestone rocks. In some people it produces irritation of the stomach.

Fragaria virginiana Mill. Wild Virginia Strawberry

Similar to the preceding.

Fragaria chiloensis Duchesne. Cultivated Strawberry

Like the preceding. There are people who can not eat strawberries or pick them without being irritated.

Rosa arkansana Porter Arkansas Rose

Widely distributed in the west, especially in prairie regions. Not poisonous, but the bristles and prickles often enter the skin and produce serious inflammation. Other species produce mechanical injuries.

Rosa rubiginosa L. Sweetbriar

A frequent escape in pastures. The recurved spines and prickles are injurious like those of the preceding species.

Pyrus Aucuparia Meyer. Mountain Ash.

The berries are poisonous to man, but not to birds. However, they are readily disseminated by birds.

Pyrus communis L. Pear.

Dr Schaffner states that horses are reported to have been killed by eating rotten pears.

Nuttallia cerasiformis T. & G. Oso Berry.

It is found in moist places and on north slopes in western U. S., and contains amygdalin. It is poisonous.

Gillenusa stipulata (Muhl) Treil. Indian Physic

From New York to Kansas. This species and *G. trifoliata*, known as Bowman's Root, are said to contain a poisonous glucoside

Filipendula ulmaria (L.) Maxim Queen of the meadow

It is cultivated and occasionally an escape, said to be poisonous
Sanguisorba canadensis L. Burnet.

From Labrador to N. Mich. and Alleghany Mts. Said to be poisonous.



Fig. 19c. Coffee Bean (*Gymnocladus dioica* (L.) Kach.
 The sweet juice in pod is poisonous. (Ada Hayden)

Pyrus malus L. Apple.

The seeds are poisonous and contain the glucoside which is changed into hydrocyanic acid.

Crataegus mollis L. Haw.

Cases of poisoning are reported by persons eating the fruit of *Crataegus mollis*, the injurious effects being probably due more to the inedible seeds than to the fruit.

Prunus americana Marsh. Wild Plum.

The shoots and seeds contain the principle amygdalin, which is converted into hydrocyanic acid. The fruit, it should be said, is entirely harmless.

Prunus pumila L. Sand Cherry.

This cherry is not widely distributed in the state, but is found along the Missouri river and in sandy soil eastward. The fruit is slightly acid and somewhat astringent, but is not poisonous, except the wilted leaves and the seeds. The astringent qualities in our wild fruit are undesirable. The same is true of *P. Besseyi* of Nebraska, the Dakotas and the Rocky Mountains.

Prunus pennsylvanica L. Wild Red Cherry

Common especially in the eastern part of Iowa and northern states. The leaves are poisonous, as well as the seeds. The fruit is edible.

Prunus virginiana L. Choke Cherry.

The leaves and seeds are poisonous. The fruit is so astringent it often produces very unpleasant conditions when eaten in any considerable quantity. Choke Cherry is widely distributed in the north. The leaves in the wilted condition contain hydrocyanic acid.

Prunus demissa (Nutt) Walp. Choke Cherry

Common in the Rocky Mountains. Cases of poisoning from this species have frequently been reported.

Prunus serotina Ehrh. Wild Black Cherry.

Most poisonous species in the north. The half wilted leaves are much more poisonous than the fresh leaves, and the seeds are very poisonous, all producing hydrocyanic acid.

Prunus caroliniana (Mill) Ait. Laurel Cherry

Laurel cherry is distributed in Southeastern United States. It is frequently cultivated for hedges. The leaves contain prussic acid.

Cercocarpus parvifolius Nutt. Mountain Mahogany

It is poisonous and the same may probably also be true of *C. ledifolius*, both being common in the Rocky Mountains, the former contains hydrocyanic acid.

Amygdalus persica L. Peach.

The leaves and seeds are poisonous. They contain amygdalin from which hydrocyanic acid is derived.

Leguminosae, Pulse Family

Cassia chamaecrista L. Partridge Pea

Widely distributed in the north especially on sandy, gravelly soils. A case was reported where a great many sheep had the scour, the ailment being attributed to this plant. Many species of the genus are known to be laxative.

Cassia marylandica L. Wild or American Senna.

Found in the southeastern part of the state, and probably acts like the preceding. This plant contains saponin.

Gymnocladus dioica (L.) Koch. Kentucky Coffee-tree

This is widely distributed in the Miss. Valley, especially along the river courses in Eastern Iowa, although growing as far north as Sioux City in the northwestern part of the state, and along the Mississippi into Minnesota. It is abundant in Missouri and Illinois. The fruit contains a sweetish but disagreeable pulp which, as well as the leaves, is poisonous. In the South the leaves are used as fly poison.

Baptisia leucantha T & G. Large White Wild Indigo

It is widely distributed in prairie regions but is generally avoided by stock. Two eastern species are regarded as poisonous, being emetics. It is probable that this species must be regarded as suspicious.

Baptisia bracteata (Muhl) Ell.

This plant contains the same active substances as the preceding species, including haptitoxin. It is common in Eastern North America. The *B. australis*, or the blue indigo of the Southern States, is also regarded as poisonous.

Thermopsis rhombifolia (Nutt.) Richards.

Common in the Rocky Mountains and in the plains region. The seeds are said to be poisonous and the plant is very bitter. Several other Rocky Mountain species and one Eastern species must be put in the suspected list. These plants produce handsome yellow flowers.

Sophora secundiflora (Cav.) DC. Coral Bean.

Common in Central and Southern Texas. It contains a powerful poisonous substance. Reports of its poisoning occur in Mexico and Texas.

Sophora sericea Nutt. Silky Sophora.

Common on the plains of Colorado also from the Dakotas to Mexico. Said to be poisonous.

Crotalaria sagittalis L. Rattle-box.

This is found in the western part of the state in the Missouri River Bottoms and produces a disease known as the Missouri Bottom disease, called crotalism, by Dr. Stalker. The seeds contain an unnamed alkaloid found by Dr. Power. The plant is not only poisonous in the meadow, but also in hay.

Cytisus scoparius (L.) Link. Scotch Broom.

Found along the Pacific and Atlantic oceans. It is common in Washington and Oregon, also in Massachusetts and Virginia. The flowers are yellow. The plant contains cytisin which also occurs in several other genera of this family like *Ulex*, *Thermopsis* and *Baptum*.

Lupinus albus L. Lupine.

The European lupine is occasionally cultivated and in Europe it produces a disease known as lupinosis. One native species, *L. perennus*, is also regarded as poisonous. Contains lupinin, lupanidin, lupinin.

Lupinus leucophyllus Dougl. Western Lupine.

It is said to be poisonous in Montana, although this is disputed by some. Other suspected species of the blue lupines are *L. argenteus* and *L. argophyllus*.

Trifolium incarnatum L. Italian or Crimson Clover.

According to Coville, it produces "hair" balls. This plant is cultivated as a cover crop in the south and east.

Trifolium repens (L.) White clover.

This plant and several other species such as *T. pratense*, at times, produce bloat.

Melilotus alba Desv. White Sweet Clover.

Widely distributed in the U. S. as a weed. The honey bees collect considerable quantities of honey from the Sweet Clover blossoms. It has been looked upon with suspicion. Dr. Schaffner states that the seeds impart a foul odor to flour.

Melilotus officinalis Willd. Yellow Sweet Clover.

This is also widely distributed in the state and is objectionable like the preceding.

Medicago sativa L. Alfalfa.

Alfalfa may cause bloat.

Psoralea tenuiflora Pursh. Slender Psoralea.

Common from Illinois to Texas. Perennial herb sprinkled with little glandular dots. The Silvery Psoralea, *P. argophylla* was reported from Iowa as poisonous. The tuberous roots of *P. esculenta* Pursh. were eaten by the pioneers and Indians.

Tephrosia virginiana Pers. Goat's Rue.

It grows only in sandy soil from Wisconsin to Iowa, east and south. It was used by Indians as a fish poison. Several other species in South America and Mexico have been used in a similar way. One is called *T. toxicaria*, and is a well known fish poison. Other N. Amer. species probably also poisonous.

Sesbania vesicaria Muhl.

An annual vine of the Southern States, Carolinas and Westward. The seeds are said to be poisonous.

Robinia neo-mexicana A. Gray.

Common in New Mexico and frequently cultivated as a hedge plant in Southern Colorado. The leaves somewhat resemble those of the black locust.

Robinia pseudacacia L. Locust-tree. Black locust.

This plant is frequently cultivated in the north and west, and in numerous places is an escape from cultivation. Native to N. Y. and Alleghany Mts. The roots, leaves and bark are very poisonous to man. Contains robinin.

Robinia viscosa Vent. Clammy locust.

Cultivated as an ornamental plant, the roots are somewhat poisonous.

Astragalus molissimus Torr.

♣ A loco-weed from Neb. to New Mexico and Wyoming

Astragalus hornii A. Gray.

Poisonous in Arizona and adjacent regions.

Oxytropis lamberti Pursh. Stemless Loco Weed.

Found in the western part of Iowa, along the Missouri River and its tributaries, very abundantly. One of the conspicuous loco or crazy weeds of the west.

Coronilla varia L. Coronilla.

An escape from cultivation in the West. It has long been regarded as a poisonous plant in Europe.




Fig. 19d. Chick Pea (*Cicer arietinum*) said to be poisonous.

Phaseolus lunatus Linn. Lima bean.

Investigations carried on in Europe seem to indicate that the lima bean leaves in the wilted condition contain hydrocyanic acid. According to Guignard practically all varieties, whether wild or cultivated, were found to contain a principle which when acted upon by an enzyme yields hydrocyanic acid. Prolonged boiling, however, extracts the greater part of it, but it is not destroyed, consequently this water should not be used, as it contains the substance which is converted into hydrocyanic acid.

Phaseolus multiflorus Willd. Scarlet Runner

Commonly cultivated in gardens. The root is poisonous.

Vicia sativa L. Common Vetch.

A frequent weed in grain fields. The seeds of this are said to be injurious to pigs. It is not injurious to cows. Contains vicin.

Prosopis juliflora DC. Mesquit Tree.

The seeds of this tree contain a small amount of saponin, probably it is not very strongly poisonous as it is used as food for cattle in Texas.

Cicer arisetinum L. Chick Pea. Idaho Pea.

Commonly cultivated in Southwestern U. S. Said to be poisonous in Europe.

Geraniaceae. Geranium Family

Erodium cicutarium (L.) L'Her and *E. moschatum* (L.) L'Her. Stork-bill.

Both of these plants are widely distributed, especially on the Pacific Coast. The former is becoming abundant in Utah. The pointed callus of the seed sometimes inflicts mechanical injuries.

Oxalidaceae. Wood Sorrel Family

Oxalis violacea L. Wood Sorrel.

Dr. Schaffner reports a case of a boy who was poisoned from eating a considerable quantity of the leaves. The leaves are frequently eaten as a salad.

Tropaeolaceae. Nasturtium Family

Tropaeolum majus L. Nasturtium.

This plant is commonly cultivated and has more or less acrid properties.

Linaceae. Flax Family

Linum usitatissimum L. Flax.

Said to produce death to cattle, probably due to the formation of HCN in the wilted leaves. People working with the fiber of the plant often have a form of dermatitis. Flax-seed, when fed in considerable quantities to live stock, especially hogs, produces death. *Linum catharticum* contains a bitter principle *linin*, and *linamarin*.

Linum rigidum Pursh. Large-flowered Yellow Flax.

This plant is reported as poisonous to sheep in some parts of the country. Found westward.

Zygophyllaceae.

Tribulus terrestris (L.) Caltrop.

Caltrop is found chiefly from Nebraska to Kansas and occasionally eastward. *T. maximum*, sometimes called soap-brush, is found in the west. The prickly fruit is more or less injurious, it presumably contains saponin. This plant should be looked upon with suspicion as one exotic species is regarded as poisonous.

Rutaceae. Rue Family

Ruta graveolens L.

This plant is more or less acrid and produces blisters. Several species of the family are regarded as poisonous.

Zanthoxylum scandens and *Z. alatum* are used as fish poisons in India. It is not known whether our native prickly ashes are poisonous or not, but they may be looked upon with suspicion.

Simarubaceae

Ailanthus glandulosa Desf. Tree-of-Heaven.

This plant is occasionally spontaneous in the United States, southeastern Iowa to Texas and eastward, where it is cultivated as an ornamental plant. It is supposed to produce poisoning when people come in contact with it. The odor of the flowers is very disagreeable. It is said also, according to Dr. Rusby that water coming in contact with the leaves is poisonous.

Polygalaceae.

Polygala Senega L. Seneca Snakeroot.

The roots of this plant are used in medicine as an emetic. The plant is common in eastern North America. Probably other species of the genus *Polygala* are injurious.

Meliaceae.

Melia azedarach L. Chinese Umbrella Tree.

This plant is commonly planted as an ornamental tree in the South and on the Pacific Coast. It has long been regarded as poisonous in Europe. Prof. Fawcett writes that it is so regarded in Florida and Cheanut reports that hogs have been poisoned by eating the seeds.

Euphorbiaceae.

Croton capitatus Michx. Hogwort, Croton.

Native to Southeastern Iowa and south. Many species of the genus contain very active poisonous principles. While this plant has not been suspected, the related species, *C. texensis*, is known to be poisonous.
Croton texensis (Klotzsch) Muell. Texas Croton.

This plant is common from Western Nebraska to Colorado and South. It is poisonous. The *S. setigerus* of California is used as a fish poison.

Ricinus communis L. Castor-Oil Plant.

It is widely cultivated as an ornamental plant. The seeds contain a deadly poisonous substance *ricin*.

Euphorbia corollata L. Flowering Spurge.

Widely distributed in the Mississippi Valley, upon sandy or gravelly soil. Produces inflammation of the skin.

Euphorbia Preshii Guss. Spurge.

Widely distributed in meadows and fields. It has been sent to the writer as supposedly poisonous to live stock.

Euphorbia marginata Pursh. Snow-on-the-Mountain.

The honey coming from the plant is poisonous. The milky juice produces dermatitis. At one time the plant was used for branding cattle.

Euphorbia lathyris L. Caper or Myrtle Spurge.

Native to Europe, occasionally cultivated. The seeds of this plant are poisonous. *Euphorbia rennifera* contains *euphorbon*.

Euphorbia cyparissias L. Cypress Spurge.

Frequently escaped from cultivation, especially near cemeteries. Poisonous to the skin, producing dermatitis.

Probably we should add to this list other species like *E. maculata* which has several times been suspected of being poisonous in the state of Iowa. *E. obtusifolia*, common East of the Rockies, and *E. heterophylla*, known in the East as painted leaf, and common from Central Iowa eastward to Florida and

westward to Texas. *E. maculata* is common everywhere east of the Rockies.

The *E. Cyparissias* was recently sent to me from Logan Iowa, where it is said to have killed lambs.



Fig. 19c Flowering Spurge (*Euphorbia corollata*) (Leah Pamme) Fig. 19f Three-Headed Mercury (*Acalypha virginica*). Both plants poisonous.

Acalypha virginica L. Three-Headed Mercury

Common in fields and open places from Canada and Minnesota to the Gulf. The leaves of this plant turn purple in the Autumn.

Jatropha stimulosus Michx. Spurge or Bull Nettle.

This plant has stinging bristles and is common in sandy soil from Texas to Missouri, Virginia and Florida.

Tragia pepetacifolia Cav. Nettle Spurge.

With stinging hairs, common in the south. Other species in the Rocky Mountains and south also have stinging hairs.

Buxaceae.

Buxus sempervirens L. Box.

Cultivated in the eastern states for hedges. It is a well known poisonous plant.

Anacardiaceae.

Rhus toxicodendron L. Poison Ivy.

The leaves and stems are poisonous to many people. The form of dermatitis produced, and the seriousness of the case varies according to the susceptibility of the individual. The plant is widely distributed in the state.

Rhus vernix L. Sumac.



FIG. 1. *ST. 11.*

This plant occurs in swamps in the Northern States and is the most poisonous species.

Rhus Rydbergii Small. Poison Ivy.

This plant is common from Montana and British Columbia to Colorado and Nebraska. It is poisonous to touch.

Rhus diversiloba Torr & Gr. California Poison Ivy

Common on the Pacific Coast. It is poisonous to the touch.

Celastraceae. Bitter Sweet Family

Euonymus atropurpureus Jacq. Burning Bush, Wahoo.

The European species has long been regarded as poisonous.

Celastrus scandens L. Climbing Bittersweet

The aril is red and has a somewhat disagreeable, sweetish taste. The leaves are said to be poisonous to horses. Bittersweet is quite widely distributed in Eastern North America.

Aquifoliaceae. Holly Family

The berries of the European *Ilex Aquifolium* commonly cultivated in the East are said to be poisonous. This plant is called Poison Berry. The *Ilex Cassine* Yaupon or Black Vomut may be poisonous. This occurs from Virginia to Texas.

Hippocastanaceae. Buckeye Family.

Aesculus Hippocastanum L. Horse-Chestnut

The seed of this species has long been recognized as poisonous in Europe. Frequently cultivated in the U. S. It contains aesculin, and argyrescin.

Aesculus glabra Willd. Ohio Buckeye.

Found in Southern Iowa, along river courses as far north as Fort Dodge and common South and East. Cases of poisoning have been reported, especially where cattle eat the young shoots and seeds of the plant.

Aesculus Pavia L. Red Buckeye

Common from Virginia to Missouri.

Aesculus octandra Marsh. Buckeye.

Common Southward.

Aesculus californica (Spach.) Nutt. California Buckeye.

The leaves and fruit of this tree are regarded as poisonous to stock. The California Buckeye especially is said to cause abortion.

Sapindaceae.

Many members of this family are poisonous, among them several species of *Paullonia* sometimes cultivated in North America, also a species of the genus *Sapindus*. One species occurs in the Southern States, *S. marginatus*, known as Soap-tree or Soap-berry.

Balsaminaceae.

Impatiens fulva Nutt. Spotted Touch-me-not

Widely distributed in the Northern States, especially on low grounds. The leaves are acrid and the plant is suspected of being poisonous to live stock.

The cultivated *I. Noli-tangere* is alterative and is regarded with suspicion. The exotic *I. Royali* is considered poisonous.

Rhamnaceae.

Rhamnus cathartica L. Buckthorn.

Frequently cultivated and used as a hedge plant. The ripe fruit is said to be poisonous. It contains the glucosides rhamnin, rhamnetin, and rhamn-cathartin.

Rhamnus lanceolata Pursh. Buckthorn.

Native to southern Iowa, east and south. Has the same properties as the preceding species.

Rhamnus Frangula L.

This plant is considered as poisonous. The *R. Frangula* of Europe is commonly cultivated. The tropical *Colletia* is poisonous.

Karwinskia Humboldtiana Zucc. Cajotillo.

Southwestern U. S. Dr. Mitchell of the 3rd U. S. Cavalry says that it is poisonous to goats. The berries have long been regarded as very poisonous.

Ceanothus velutinus Dougl. Ceanothus.

This species with shiny leaves and small white flowers occurs abundantly in the Rocky Mountains from New Mexico to British America, to Oregon and Washington. According to Greshoff, it contains saponin in considerable quantities. The same authority states that several other species examined by him contained the same substance, among them were *C. aculeus* Desf., and *C. thyrsiflorus* Eschw. He also found the same substance but in less quantity in the seeds of the New Jersey tea *C. americanus* L. and *C. ovatus* Desf., two shrubby plants of Eastern North America, the latter being also common on the east slope of the Rockies.

Vitaceae. Grape Family

Pedern quinquefolia (L.) Greene. Virginia Creeper

Common from New England to the Rocky Mountains, especially east of the Missouri River. This plant is regarded as poisonous by some. The leaves and fruit abound in raphides. In this connection it may be of interest to state that the fruit of *Cissus nictitans* of the old world produces poisonous fruits, and that the *C. pruriens* with a pleasant, acid, sweet taste produces a painful, burning sensation later. The same is true of the southern Mustang Grape (*V. rotundifolia*).

Malvaceae.

Abutilon Theophrasti (Medic) Gaertn. Velvet-Leaf. Indian Mallow. Butter Print.

Widely naturalized in the state. The plant gives off a very disagreeable odor, and is suspected of being poisonous.

Gossypium herbaceum L. Common Cotton.

The root of cotton is well-known as an abortive. Feeding excessive amounts of cotton seed meal frequently produces death in animals, particularly in hogs.

Dr. Crawford states that this is due to meta and pyrophosphoric acid in cotton seed meal.

Sida urens L. The Stinging Sida.

Occurs in tropical America and produces mechanical injuries because of the hairs with which it is covered. This is also true of the *S. paniculata* L. Probably other species of *Sida* of the southern states may produce similar mechanical injuries.

Hypericaceae

Hypericum maculatum Walt. Spotted St. John's-Wort

All the species are suspected of being poisonous to horses. Vesicant.

Hypericum perforatum L. Common St. John's-Wort

Naturalized in Eastern U. S. A well known vesicant

Hypericum Ascyron L. Great St. John's Wort

Most widely distributed species in the state, in woods or borders of woods

Violaceae

Viola odorata L. Sweet Violet.

Commonly cultivated in greenhouse; Said to be somewhat poisonous

Underground parts of the plant are emetic

Viola cucullata Ait. Common Blue Violet.

The most widely distributed species in the eastern states, the roots, like the preceding, are emetic

Loganiaceae

Mentzelia ornata Torr & Gray *Mentzelia*.

The backwardly-barbed trichomes sometimes produce mechanical injuries.

Found in Northwestern Iowa on the Big Sioux near Sioux City, and west and southwest

Datiacaceae

Datura glomerata (Presl) Benth & Hook.

California fish poison

Cactaceae

Opuntia Rafinesquii Engelm. Cactus.

Found in sandy soil in the state. The barbed trichomes cause mechanical injuries.

Other spiny species of *Opuntia* *Mammillaria* and *Cereus*, largely developed in southwestern United States, produce similar mechanical injuries

Anhalonium Lewinii Hemi. Mescal Bean.

This cactus contains a narcotic substance which has long been used by the Indians of Mexico and adjacent regions to produce narcosis

Cereus grandiflorus Mill. Night blooming *Cereus*

It has been regarded with suspicion where the plant is cultivated and native. Frequently cultivated in the United States

Thymelaeaceae *Mezereum* Family

Daphne Mezereum L. Spurge Laurel.

This is a well-known poisonous plant of Europe and is occasionally cultivated. It contains the glucosides daphnin and daphnetin

Other cultivated species in North America are known to be poisonous and several of the related genera native to Asia and Australia are poisonous

Dicra palustris L. Leather-Wood, Moose-Wood.

Found in the North to Iowa. The bark is acrid, the berries narcotic and poisonous.

Elaeagnaceae. *Oleaster* Family

Hippophae rhamnoides Lam.

A native of Europe but cultivated in the United States and said to be narcotic.

Araliaceae. *Ginseng* Family

Palsia horrida (Smith) B. & H.

From Isle Royal, Lake Superior region, to the Rocky Mountains and Alaska, and California. It causes mechanical injuries.

Umbelliferae. Parsley Family

Conium maculatum L. Poison Hemlock.

Introduced here and there in Iowa and eastward, common in Utah. The plant contains the alkaloids conium, conydrin, methyconium, and a bitter principle cicutoxin. A very poisonous plant both to man and lower animals.

Petroselinum hortense Hoffm. Parsley.

Some people are suspicious of parsley. Dr. Schaffner states that the seeds are injurious to birds. He reports a case of poisoning of several parrots from eating the leaves of this plant. Cultivated.

Apium graveolens L. Celery.

Several cases are known where persons who have handled celery have had a form of dermatitis. Some persons cannot eat celery because a rash forms.

Cicuta maculata L. Water Hemlock. Cowbane.

The roots of this plant are very poisonous. The plant is widely distributed in the north, especially in low grounds. The European cowbane, *C. virens*, contains the alkaloid conium, a substance which probably also occurs in our plant. The poisonous principle is cicutoxin.

Cicuta bulbifera L. Bulb-bearing Hemlock.

The roots of this, and the whole plant are supposed to be very poisonous. In swamps and northern states.

Cicuta vagans Greene. Cowbane.

Occurs in Washington, Oregon and California. Occasionally said to poison cattle drinking water in which they have trampled roots of this plant, thus expressing the extract. Very poisonous.

Cicuta Bolanderi A. Gray.

Found in marshy regions in California.

Cicuta occidentalis Dougl. Cowbane.

Found in the Rocky Mountains and considered poisonous.

Sium cicutaefolium Gmelin. Water Parsnip.

Common in many parts of the north in low grounds. Said to be poisonous.

Aethusa Cynapium L. Fool's Parsley.

A poisonous herb native to Europe, with a disagreeable odor. Possibly occurs in a few places in the state. Contains the alkaloid cynapin and another conium-like alkaloid.

Angelica atropurpurea L. Purple-stemmed Angelica.

Found in low grounds in North and Northeastern Iowa. Possibly poisonous. Cattle do not relish it.

Oxypolis rigidior (L.) Coult. & Rose. Cowbane.

Aquatic herb with white flowers, leaves simple pinnate with 3-9 linear-lanceolate leaflets.

Pastinaca sativa L. Parsnip.

Persons are often poisoned by handling the plant, which causes inflammation and vesication. Mr. F. C. Stewart, in a letter to the writer, states that in one case, the eyes became swollen, vesication occurred from poisoning caused by the flowers.

Heracleum lanatum Michx. Cow Parsnip.

Supposedly poisonous, although the leaves of the fresh plant are eaten by the Indians. This species is widely distributed in Iowa, especially in rich woods. Contains the bitter principle heracelin.

Daucus carota L. Carrot.

The carrot, like the parsnip, causes vesication. Dr Schaffner says that persons handling the plant are often poisoned, especially when the plant is wet with dew.

Cornaceae. Dogwood Family

Cornus paniculata L'Her Dogwood.

Widely distributed in the north. Some regard it as poisonous.

Cornus paniculata L'Her

The fruits of this species are considered by some people to be poisonous, but there are no authentic cases.

Garrya Fremontii Torr Dogwood.

Contains an alkaloid. It may be of interest here to state that the *Marlea piliensis* of Australia and the Pacific Islands is poisonous. It would not be strange, therefore, to find that some of the other members of this family are poisonous.

Ericaceae, Heath Family

Epigaea repens L. Trailing Arbutus.

North and East, Alleghany Mountains. Supposed to be poisonous. Contains the glucoside ericolin.

Andromeda polifolia L. Wild Rosemary.

Arctic America, Adirondack Mountains and in Europe and Asia. *A. glaucophylla* Lank. is the Bog Rosemary, with whitish leaves found in bogs and wet shores from Labrador to Minnesota, and the *A. floribunda* Pursh, is found in moist hillsides in the Alleghany Mountains from Virginia to Georgia. These plants contain andromedotoxin.

Rhododendron maximum L. Large Rhododendron.

Small shrub or tree from Nova Scotia to Ohio and the Alleghany Mountains. *R. catawbiense* Michx., in the high Alleghanies from Virginia to Georgia known as the Mountain Rosebay, and the California Azalea (*R. occidentale*) are known to be poisonous, the latter especially to sheep.

Rhododendron californicum Hooker California Rhododendron.

From San Francisco to British Columbia. Cases of poisoning have been reported from the Pacific Coast.

Kalmia angustifolia L. Lambkill. Laurel.

Common on hillsides, pastures, and bogs, from Labrador southwest through the Alleghanies. Known to be very poisonous. The leaves contain andromedotoxin, frequent cases of poisoning on record.

Kalmia latifolia L. Laurel

From New Brunswick to Ontario to Pennsylvania and south to the Alleghanies. A well-known poisonous shrub. Many cases of poisoning of sheep and cattle are on record from this species, probably the most poisonous of all the members of this genus. *K. latifolia* Wang., found in bogs, with rose-purple flowers and pale colored leaves, is poisonous.

Ledum glandulosum Nutt. Labrador Tea.

Common in bogs of the Northern Rockies, the Cascade Mountains to California. It is said to be poisonous. *L. groenlandicum* Oeder., with white flowers and rusty wool underneath, occurs in bogs from New England to Wisconsin and Minnesota. Said to be poisonous.

Leucothoe Catesbaei (Walt.) Gray.

With white flowers in axillary or spiked racemes, shrub. From Virginia to Georgia in the mountains; an allied species *L. racemosa* (L.) Gray occurs in moist thickets from Massachusetts to Louisiana, near the coast. Both species are said to be fatal to stock, the latter especially to calves.

Lyonia mariana (L.) D. Don. Stagger-bush.

From the Atlantic Coast to Tennessee and Arkansas. Said to produce intoxication, hence the name.

Primulaceae. Primrose Family

Anagallis arvensis L. Poor Man's or Shepherd's Weather Glass. Pimpernel. Possibly growing in the state. Known to be poisonous. Contains glucoside cyclamin.

Primula obconica Hance.

Poisonous to the touch, very much like poison ivy. This plant is commonly cultivated in greenhouses.

Primula Parryi A. Gray. Parry's Primrose

Common in higher altitudes in the Rocky Mountains especially near brooks or springs. The root has the odor of musk, and is said to be poisonous.

Cyclamen Europaeum L. Cyclamen.

This beautiful cultivated plant has long been regarded as poisonous in Europe. The *C. persicum* Mill. is also regarded as poisonous.

Plumbaginaceae. Plumbago Family

Lamornum carolinianum (Walt.) Britton.

It is not known whether this plant is poisonous or not but a related plant, *Statice pectinata* Ait. of the Cape Verde Islands is poisonous.

Sapotaceae. Sapodilla Family

Achras Sapeia L.

A tropical fruit cultivated in Florida. It is said to be injurious and according to Greshoff contains saponin. *Lycuma* forms a well-known genus of West Indian fruit trees. In *L. Bonplandii*, H. B. K., Altamirano demonstrated amygdalin as early as 1876, and in another species a cyanogenic glucoside was suspected, but the experimental proof was wanting. Greshoff reports hydrocyanic acid in the *L. mammosa*.

Ebenaceae. Ebony Family

Diospyrus virginiana L. Persimmon.

Common in the Southern States as far north as southern Iowa. It is used as an anthelmintic, but it is not positively known whether any part of this plant is poisonous. Several exotic species, however, blister the skin and one species in Madagascar is said to be very poisonous.

Oleaceae. Olive Family

Ligustrum vulgare L. Privet.

The privet is frequently cultivated in the U. S. The leaves and fruit of the plant are said to be poisonous. Prof. Chesnut says that accidents have been occasioned in children, both by fruit and the leaves. It contains the bitter principle syringoperin.

Forsythia suspensa Vahl. Forsythia.

Commonly cultivated in northern states. The root is slightly poisonous. The *F. viridissima* Lindl. is very bitter. Greshoff has found saponin in the seeds of the former species.



Fig. 2. *Phlox paniculata* 'Majestic'



Fig. 19g. Oleander (*Nerium Oleander*). Cultivated poisonous. Fig. 19h. False Jessamine (*Gelsemium sempervirens*). Contains several poisonous alkaloids. (Cheatnut, U. S. Dept. Agr.).

Chionanthus virginica L. Fringe Tree.

Common in the Adirondack Mountains, New Jersey to Florida and Missouri. This is said to be a narcotic.

Loganiaceae. Logania Family

Gelsemium sempervirens (L.) Ait. Yellow Jessamine

This climbing shrub with yellow fragrant flowers is common in the South. It contains several potent alkaloids, among them gelsemin. The symptoms of poisoning from consuming this plant are muscular weakness, especially in the fore legs, followed by convulsive movements of the head, fore legs and hind legs.

Spigelia marilandica L. Indian Pink or Pink Root

Common from Ohio to Missouri, Florida and Texas. This plant is used in medicine and is known to be toxic. Strychnin belongs to this same family.

Gentianaceae. Gentian Family

Menyanthes trifoliata L. Buckbean.

The plant has bitter properties and is rauseous. Contains menyanthin.

Apocynaceae. Dogbane Family

Apocynum androsaemifolium L. Spreading Dogbane

Widely distributed in Iowa, probably poisonous.

Apocynum cannabinum L. Indian Hemp.

Like the preceding. Contains apocynin, which is poisonous.

Nerium Oleander L. Common Oleander

Cultivated. The leaves are deadly poisonous to stock. Contains conessine, and nerun, which has the properties of digitalin. *Neranthus* bears a resemblance to digitalin.

Dr. Wilson of the Arizona Experiment Station has recently demonstrated the very poisonous nature of this plant in Arizona.

Asclepiadaceae. Milkweed Family

Asclepias tuberosa L. Pincus-Root.

Widely distributed in Iowa, especially on gravelly knolls and prairies. The leaves are more or less poisonous to stock. However, honey bees collect considerable honey from this plant.

Asclepias incarnata L. Swamp Milkweed.

Poisonous probably like the preceding. The root is emetic and cathartic.

Asclepias syriaca L. Milkweed.

Poisonous. Contains the glucoside asclepioside, an amorphous bitter substance.

Asclepias speciosa Torr. Showy Milkweed.

Poisonous. This species is found in Western and Northwestern Iowa to Utah.

Asclepias campestris, DCne. Milkweed.

Commonly cultivated in gardens southward. It is said to be poisonous.

Asclepias eriocarpa Benth. Milkweed.

Common in California and adjacent regions. According to Chesnut sheepmen in California very much fear this weed. It has broad mullein-like leaves.

Asclepias mexicana Cav. Narrow Leaved Milkweed.

Native to California, Oregon and Nevada. According to Chesnut sheep and calves are not infrequently poisoned by eating this plant and cows have been poisoned by eating hay contaminated with it.

Convolvulaceae. Convolvulus Family

Ipomoea pandurata Meyer. Wild Potato Vine. Man of the Earth.

The large root is poisonous. Contains the glucoside ipomoein.

Convolvulus sepium L. Hedge Bindweed.

The plant produces a somewhat disagreeable odor. Dr. Schaffner states that it is supposedly poisonous to swine. Jalap contains several glucosides which also probably occur in our Morning Glory. One is convolvulin.

Cuscuta epithymum Murr. Clover and Alfalfa Dodder.

Dr. J. Q. Taylor of Lisbon, Ohio, in writing to Prof. A. D. Selby stated that dodder produced a bowel trouble in horses, and Dr. Jenkins of New Haven writes that clover hay containing a great deal of dodder produced scours. He added that the hay had moulded badly, although the injury could not be definitely traced to dodder. Some exotic dodders are poisonous. The dodder must, therefore, be looked upon with suspicion.

Polemoniaceae. Polemonium Family

Gilia aggregata Spreng. Cypress plant.

Common in the Ry. Mts. to Neb. This species according to Greshoff contains a considerable amount of saponin and is very poisonous. He lists several other species which contain this substance.

Hydrophyllaceae. Waterleaf Family.

Phacelia circinata Jacq. Rough Phacelia.

This plant is common in the Rocky Mountains. It produces rough bristles and causes considerable irritation and inflammation, other species act in a similar way.

Boraginaceae. Borage Family

Heliotropium Europaeum L. Wild Heliotrope

This plant contains a poisonous alkaloid and as well as the *H. indicum* L. has long been regarded as poisonous.

Cynoglossum officinale L. Hound's Tongue

Poison acts much like Curare.

Cynoglossum virginicum L. Wild Comfrey

Supposed to be poisonous.

Lappula officinalis Lehm. Stickweed.

The fruit of this plant gets into the wool of sheep and sometimes produces mechanical injuries.

Echium vulgare L. Viper's Bugloss.

Occasionally spontaneous, probably in the Eastern part of the state. Contains a poisonous alkaloid.

Verbenaceae. Verbena Family

Callicarpa americana L. Mexican Mulberry

Common in the southern states. Several Asiatic species of the genus used as fish poisons. No report of poisoning from the American species is recorded.

Labiatae. Mint Family

Scutellaria galericulata L. Marsh Skullcap.

Common in wet places in the North. Suspected of producing poisoning.

Stachys arvensis L. Corn or Field Woundwort

Waste places, especially along the Atlantic Coast. Suspected.

Galeopsis tetrahit L. Dead Nettle

Common hemp nettle with purplish flowers and stiff calyx bristles. Common Eastward and in the Northwest in British Columbia and Washington. Causes irritation.

Nepeta hederacea (L.) Trevisan. Ground Ivy

Widely naturalized in the state. It is said to be poisonous to horses.

Hedeoma pulegioides Pers. Pennyroyal.

Common, especially in clay soils in Eastern Iowa. The oil is known to cause poisoning.

Leonurus Cardiaca L. Common Motherwort

Known to produce mechanical injuries and dermatitis. Widely distributed in the U. S., naturalized in Europe.

Solanaceae. Nightshade Family

Nicandra Physalodes (L.) Pers. Apple of Peru.

Cultivated here and there in Iowa. Said to be poisonous. Used as a fly poison in parts of the United States.

Solanum nigrum L. Black Nightshade.

The leaves and other parts of the plant are reputed to be poisonous to calves, sheep, goats and swine, and the green berries are known to be poisonous to man. The fruit of a form of this species is cultivated as an esculent.



Fig. 191. Black Nightshade (*Solanum nigrum*). Contains a poisonous alkaloid. (Chevrolt, U. S. Dept. Agr.).

The writer has not only eaten berries of this, but has seen others eat berries of this and the common Black Nightshade without injurious results. Contains the alkaloid solanin, with a hot, bitter taste.

Solanum triflorum Nutt. Nightshade.

Common from Nebraska to Rocky Mountains to Alberta, south to Texas.

Poisonous.

Solanum tuberosum L. Potato.

At certain times the tubers of the potato are poisonous, especially when green. The writer knows of an instance where the eating of potatoes acted as a poison. The substances produced in the young shoots of the potatoes are solanin and solanadin.

Solanum carolinense L. Horse-nettle.

Southern United States North to Minnesota and East to New York.

Fruit with disagreeable odor. Said to be poisonous. Narcotic.

Solanum Dulcamara L. Bittersweet.

The berries are poisonous, as are also the leaves. Cattle are known to have been poisoned by it. The bitter substance contained in it is known as dulcamarin.

Lycopersicon esculentum Mill. Tomato.



FIG. 2. *W. coccinea* (H.)

The green parts of the plant contain saponin and solamin.

Physalis heterophylla Nees. Ground Cherry.

From New Brunswick southward and westward. Suspected plant as is *P. virginiana* Mill.

Cestrum caubiflorum Jacq. Cestrum.

West Indies. The *C. nocturnum* L., frequently cultivated in greenhouses. Several species are poisonous.

Atropa Belladonna L. Belladonna, Deadly Nightshade

Occasionally cultivated and possibly also an escape. Very poisonous.

Nicotiana Tabacum L. Tobacco.

Cultivated. Narcotic and poisonous, and produces the alkaloid nicotine a very poisonous substance

Nicotiana glauca Link & Otto. Flowering Tobacco

Poisonous like the preceding.

Nicotiana glauca R. Grah. Smooth Tobacco

An escape from cultivation in California. According to Dr. G. Burr Davy poisonous.

Hyoscyamus niger L. Black Henbane.

Probably occasionally found in Iowa, Utah, Mont., Atlantic States. Known to be poisonous to stock and also to hogs. Universally recognized as a poisonous plant in Europe and this country. Probably one of the most deadly poisonous plants in the United States. Seeds are poisonous to chickens. Contains the alkaloid hyoscyamin.

Datura Stramonium L. Jimson weed.

Naturalized in various parts of the U. S. All parts of the plant are narcotic and poisonous, especially the seed. Several cases of poisoning in children are reported in Iowa. The plant produces a very disagreeable odor, and the hay containing the plant is poisonous to cattle. It contains the alkaloid, atropin, and hyoscyamin.

Datura Tatula L. Purple Jimson weed.

Poisonous like the preceding, and the following species. Naturalized in U. S.

Datura Wrightii, DC. Wright's Datura.

Frequently cultivated as an ornamental plant and known to be poisonous. The nectar from the flowers which is produced in great abundance is known to produce poisoning in children in this country

Datura aiba Nees. Thorn Apple.

Cultivated. Plant said to have been much used in India for criminal purposes. Very toxic

Capsicum annuum L. Red or Cayenne Peppers.

Well known remedy used as a stimulating plaster externally if the pepper is applied long enough it produces vesicles. Red pepper is often injurious when taken in too large doses internally. The active poison is capsaicin with a strong odor and burning taste.

Capsicum frutescens L. Shrubby Pepper

Southwestern United States, said to be poisonous. The fruit has a sharp pungent taste.

Lythrum halimifolium Mill. Matrimony Vine.

Cultivated and commonly naturalized. Somewhat spiny thorns. Said to be poisonous.

Scrophulariaceae. Figwort Family

Verbascum Thapsus L. Moth Mullein.

Naturalized on the Atlantic coast and common in Utah and the west. Said to be poisonous.

Linaria vulgaris Hill. Butter-and-eggs. Toad Flax

A weed, especially northward. Suspected of being poisonous

Scrophularia marilandica Gray. Simpson Honey Plant.

Widely distributed in E. U. S., pastures and woods. Not eaten by stock. According to Millspaugh, the physiological effect of this plant is bleeding of the gums, colic, and sleepiness. Contains a crystalline bitter substance, scrophularin.

Digitalis purpurea L. Purple Foxglove.

This plant is widely cultivated in the U. S., and naturalized on the Pacific Coast.

Is poisonous to man and live stock, especially horses. It contains the glucosides digitalin, which dilates the pupil, digitoxin, and digitonin.

Gerardia tenuifolia Vahl. Slender Gerardia.

Said to be poisonous to sheep and calves. Probably other species are likewise poisonous, like *G. grandiflora* and *G. purpurea*.

Pedicularis lanceolata Mx. Lousewort.

Widely distributed in low grounds and swamps. Said to be poisonous

Pedicularis canadensis L. Lousewort.

Widely distributed in the state in gravelly soils and on knolls. Said to be poisonous. Sheep, however, eat large quantities of the *P. groenlandica* without apparent injuries.

Pedicularis groenlandica Retz. Mountain Lousewort.

Common at higher altitudes in swamps. Suspected of being poisonous.

Melampyrum lineare Lam. Cow Wheat.

Common in open woods eastern states to Tennessee. The European species, *M. nivaticum*, is regarded as poisonous in Europe. Our American plant has not, however, been reported.

Gratiola sp. Hedge Hyssop.

The European *G. officinalis* is said to be poisonous to cattle. Several species common in eastern North America.

Veronica virginica L. Culver's Root.

Common in swamps in eastern North America especially northward. Contains arponin.

Chelone glabra L. Balmony

In swamps northern United States. Contains an alkaloid. Suspicious.

Bignoniaceae. Bignonia Family

Catalpa speciosa Warder. Hardy Catalpa.

Widely cultivated in U. S. Odor coming from the fragrant flowers is poisonous and Dr. White in his *Dermatitis Venenata* states that the flowers are irritating to many persons. Dr. Millspaugh, on the other hand, states that it is said to be dangerous to inhale the odor of the flowers for a long time, which, however, is probably not generally true. The allied Caroba contains the bitter principle carobin.

Catalpa bignonioides Walt. Common Catalpa.

Occasionally cultivated in the north, but scarcely hardy. Poisonous like the preceding.

Bignonia capreolata L. Cross Vine

A tall climbing vine with large orange colored flowers, from Virginia to Louisiana. Said to be poisonous.

Tecoma radicans (L.) Juss. Trumpet Creeper

Climbing vine, large tubular orange and scarlet flowers. New Jersey to Iowa, Texas to Florida. Said to be poisonous, several exotic species are poisonous.

Rubiaceae. Madder Family

Cephalanthus occidentalis L. Buttonbush.

This plant commonly occurs in low grounds, especially along river courses. The leaves contain a poisonous principle, cephalanthin, a very bitter glucoside.

Caprifoliaceae. Honeysuckle Family

Triostema perfoliatum L. Feverwort, Horse-gentian.

Widely distributed in woods. Some species of the genus were used by the Indians as a cure for fevers and early practitioners in this country used the root as an emetic. The physiological action of the plant is to produce vomiting.

Sambucus canadensis L. Elderberry

Dr. Rusby states that the plant is poisonous. The elderberry is widely distributed in the U. S. and the flowers of this are commonly used to prepare a tea. Contains conin.

Sambucus racemosa L. Red-berryed Elder

Northern states and Rocky Mountains. Northeastern Iowa. Suspected.

Symphoricarpos orbiculatus Moench. Coral berry

Common in the south, occurring on sterile or rocky soils and on borders of woods. Reported as poisonous.

Symphoricarpos mollis Nutt. Coral Berry.

This plant is said to contain saponin according to Greshoff. The *S. racemosa* Michx. Snowberry. Across the continent northward. Greshoff found saponin in leaves but not the fruit. Cases of poisoning have been reported in the old world.

Lonicera involucrata (Richards) Banks. Rocky Mountain Honeysuckle

Common in the Rocky Mountains. It is said to contain saponin in considerable amounts. Several other species contain saponin and may be regarded as poisonous.

Diervilla lonicera Mill. Bush Honeysuckle

Common in woods, north. Not reported as poisonous, but the Japanese *D. japonica* DC is regarded as poisonous. It contains saponin.

Cucurbitaceae. Gourd Family

Some of the exotic genera of this family, like the *Bryonia dioica* known as the poison berries, and the *Trichosanthes palmata*, are poisonous.

Sicyos angulatus L. Bur Cucumber

Eastern North America. The barbed prickles are irritating and injurious to many persons according to Dr. Halsted.

Boeria sp.

One or more species of the genus regarded as poisonous in California according to Chesnut.

Campanulaceae Bluebell Family

Lobelia cardinalis L. Cardinal-flower

This is listed as one of the poisonous plants by Dr. Schaffner. Cardinal flower is very abundant in the swamps along river courses in Eastern Iowa, and N. Miss. Valley.

Lobelia siphilitica L. Blue Lobelia.

Also listed as a poisonous plant by Dr. Schaffner.

Lobelia spicata Lam. Pale Spiked Lobelia

Everywhere on prairies of the northern states. Probably poisonous like the preceding.



Fig. 19. Greater Ragweed (*Ambrosia trifida*). Said to cause hay fever. (Hewey U. S. Dept. Agr.)

Lobelia inflata L. Indian Tobacco.

Widely distributed, occurring in woods. The leaves of this plant were used by the Indians. The plant has long been used in medicine. *L. inflata* is very poisonous and is used for its action upon the pneumogastric nerve, the toxic doses produce exhaustion and dilation of the pupils. Death is usually preceded by insensibility and convulsions. Contains the acid lobelin.

Lobelia Kalmii L. Kalm's Lobelia.

Found in swamps from N. D. to N. J. west to northern Iowa, Minnesota, and Man. Probably poisonous like the preceding.

Compositae. Composite Family.

Lactuca Scariola L. Prickly Lettuce

Common across the continent. also the var *integra* Said to be poisonous

Cichorium Intybus L. Chicory

It has become widely naturalized in the north and west. When fed in large quantities to dairy cattle it imparts a bitter flavor to the milk and butter. It contains the bitter glucoside chicoricin.

Iva xanthifolia Nutt. Marsh Elder. Half-breed Weed.

Western Wisconsin, Red River Valley, south and west to the Rocky Mountains. Said to cause hay fever.

Ambrosia artemisiifolia L. Common Ragweed.

The pollen of this plant is suspected of causing hay-fever.



Fig. 192. Spiny Clothier (*Xanthium spinosum*). Injurious in a mechanical way. Fig. 191. Prickly Lettuce (*Lactuca Scariola*). Said to be poisonous. (Genth)

Ambrosia trifida L. Great Ragweed.

The pollen of this species is said to produce an irritating action upon the mucous membrane.

Xanthium spinosum L. Clothier

Maine to Texas. It causes mechanical injuries.

Xanthium canadense Mill. Cocklebur

Young seedlings of this plant are poisonous to horses. Several cases of poisoning to hogs have been reported in this state.

Xanthium strumarium L. Cocklebur

Poisonous like the preceding. This species is not common in the state. Contains the glucoside xanthostrumarin.

Eupatorium perfoliatum L. Boneset.

Commonly found in low grounds and marshes. It is an emetic when given in large doses.

Eupatorium urticaefolium Reichard. White Snake root

Widely distributed in woods in this state. It is said to produce a disease known as milk fever. No reports of this kind of poisoning have come to us in Iowa. The *E. cannabinum* contains the alkaloid eupatoria and the glucoside eupatorin.

Trifolium odoratissimum (Walt.) Cass.

It is said to be injurious and has the odor of sweet clover

Grindeia squarrosa (Pursh) Dunal. Tar Weed.

From Wisconsin, Iowa, westward and northward. Is said to be injurious. The *G. lanceolata* Walt., occurs from Tennessee to Texas. It has large heads and resinous viscid leaves.

Xanthoxylum texana D. C.

It is said to be poisonous and contains saponin. Southward.

Erigeron canadensis L. Horse Weed.

It is a widely distributed troublesome weed in the north. The physiological action of the drug obtained from this plant is to produce smarting of the eyes, soreness of the throat, and prostration.

Erigeron ramosus (Walt.) B. S. P. White Weed

Common in meadows westward. The *E. annuus* of the same distribution is found also in meadows. Both are regarded with suspicion.

Baccharis halimifolia L. Salt Groundsel.

The European *B. cordifolia* is said to be poisonous. It occurs in salt marshes along the Atlantic sea coast.

Solidago canadensis L. Golden-rod.

It is widely distributed in the U. S., and is one of the most common of our golden-rods. The golden-rods are generally regarded as harmless plants, but in a few cases they are suspected of being poisonous. A disease of horses near Black River Falls, Wisconsin, was attributed to a golden rod. Chestnut thinks the disease due to a rust on the plant. As a general thing stock does not relish the golden-rod.

Madia glomerata Hooker. Tarweed.

A glandular viscid, heavily scented herb common from Saskatchewan to Colorado, Utah, Oregon and Washington. Probably poisonous or at any rate it is avoided by cattle.

Hemizonia macrademia DC. Tarweed.

Common on the Pacific Coast from San Francisco southward. A strongly and unpleasantly scented herb avoided by stock. Many of the species of the genus occur in California and most of them are strongly scented.

Aster Parryi Gray. Woody Aster

Common in the Rocky Mountains, Utah and Wyoming. Thousands of sheep in Wyoming where this plant occurs suffer with the disease which has been attributed to "grub in the head." Healthy sheep often die within a very short time after the first symptoms appear. This plant is affected with a fungus, *Puccinia xylophagae*, and Prof. Aven Nelson suggests in his account of this disease, "The chances are rather better that the suggested poisonous qualities are due to the fungus. Some other parasitic fungi have been proven poisonous

and we may well in the absence of evidence suspect this one." The aster is confined to Gumbo clay soil intermixed with gravel and soil that contains more or less alkali and other salts.

Dr. O. L. Frier is investigating the disease.




Fig. 19m. Fetid Marigold (*Dysodia chrysanthemoides*). Pungent odor said to be injurious. (Charlotte M. King.) Fig. 19n. Dog Fennel (*Anthemis cotula*). Contains a pungent principle.

Rudbeckia lacinata L. Cone-flower

In moist grounds throughout the north. Dr. Schaffner says it is supposed to be poisonous to sheep.

Bidens frondosa L. Black Beggar-ticks.

Common in the north. The downwardly barbed awns are irritating.

Coreopsis discoidea T & G. Small Beggar-ticks.

Very common in the east. It is a local irritant.

Helenium autumnale L. Sneezeweed.

It is common in low grounds throughout Iowa and is used by the Indians to produce sneezing. The whole plant and flowers are poisonous to cattle and sheep.

Helenium tenuifolium Nutt. Sneezeweed.

Common in the south and said to be fatal to horses and mules. It imparts its bitter flavor to milk.

Helenium Hoopesii A. Gray Mountain Sneezeweed

Common in the Rocky, the Uintah and Wasatch Mountains. Said to be injurious to sheep.

Achillea millefolium L. Yarrow

It is used as a forage plant, but causes an irritating sensation of the membranes and much pain in the gastric and abdominal regions. It contains the glucoside achillein, an amorphous bitter substance, and the alkaloid moschatin.

Anthemis Cotula L. Mayweed.

Has a very disagreeable odor and causes blistering of the skin. The plant is carefully avoided by stock.

Anthemis arvensis L. Corn chamomile.

Occasionally escaped from cultivation. Seeds of this and other species contain HCN.

Dysodia chrysanthemoides Lag. Fetid Marigold.

Common in the west, Dak., Ia., Neb. to Mo., Tex. The leaf bracts and other parts of the plant are provided with large pellucid glands which produce the characteristic odor of the plant.

Tanacetum vulgare L. Common Tansy

Introduced into many parts of the north. Many serious and a few fatal cases of poisoning are recorded by the use of tansy oil. The symptoms of poisoning are varied, convulsions, violent spasms, dilation of the pupils, frequent and feeble pulse. Eleven drachms of the oil in a girl produced death in three and one-half hours. The effect on animals is salivation, vomiting, dilation of the pupils, muscular twitchings, followed by chronic spasms, death appears to be caused by paralysis of the heart and lungs.

Artemisia biennis Willd. Biennial Wormwood.

Probably poisonous.

Artemisia tridentata Nutt.

Sage brush used as forage by sheep.

Artemisia Absinthium L. Common Wormwood.

Occasionally cultivated. The volatile oil of the plant is a violent, narcotic poison, and contains the glucoside absinthin, the alkaloid abrotin, the bitter principle santonin.

Other species probably also poisonous.

Arnica cordifolia Hook. Arnica.

This species with yellow flowers is common in the Rocky Mountains. The bruised leaves give off the odor of arnica. The European *A. montana* is suspected.

Senecio Jacobaea L. Staggerwort.

Occasionally found in the east and causes the Picton disease. The Squaw Weed (*S. aureus*) is common in the north. The *S. plattensis*, common in western Iowa to Montana and east to Ontario, has been associated by Dr. Day with the Missouri Bottom disease. The species are numerous in the Rocky Mountains and may be responsible for some diseases. It is to be noted that the *S. guadalupensis* of Mexico is fatal to stock. The exotic *S. tolucaeus* contains an alkaloid with tetanus like action.



Fig. 19c. Absinthium (*Artemisia absinthium*). Plant, leaves, and flowering branch. (Faguet).

Arctium Lappa L. Burdock

Produces itching. Contains the alkaloid lappin. Common weed in the U. S.

Cirsium lanceolatum (L.) Hill. Bull Thistle.

Commonly naturalized in the north from the Atlantic to the Pacific. It acts injuriously in a mechanical way. Other species like *C. Nelsoni*, Canada Thistle (*C. arvense*), *C. scariosum*, etc., all act in a similar way. Some species contain HCN.

Centaurea solstitialis L. Knapp Weed.

Common in alfalfa meadows westward. Acts injuriously in a mechanical way. Some species, according to Greshoff, contains HCN.



Fig. 19p. Sneezeweed (*Helenium tenuifolium*)
(*Cirsium lanceolatum*) Mechanical injuries.



Poisonous. Fig. 19q. Bull Thistle.

CHAPTER XV

CHEMISTRY OF ALKALOIDS.

PROF. A. A. BENNETT

The term alkaloid is a relic of an early method of nomenclature, namely, the method of naming substances without reference to their fundamental properties. For example, the name, oil of vitriol, does not describe sulfuric acid except that it has a deceptive resemblance to an oil, and was originally produced from green vitriol. The word alkaloid literally signifies a substance resembling an alkali. They do form salt-like compounds with acids but here the likeness as to specific properties ends.

It is only about one hundred years since the facts as to alkaloids began to accumulate. The first separation of these compounds was made in 1803 by Derosne but their basic character was not noted until three years afterwards by Serturner while studying opium. Before this time many plant extracts were known to contain some very active compounds usually called principles, but their isolation and the determination of their composition and properties date from the first quarter of the nineteenth century.

The first alkaloid that was prepared and reported according to the usual method of procedure of the chemists, namely, obtaining the pure compound and then determining its chief properties, was morphin. Although this was done about 1806 it was not until some eleven years afterwards that the report attracted the attention of chemists, sufficiently to start the investigation of other substances for the presence of similar compounds with the result that new compounds of this class have been separated and described each year since this date. The property that especially characterised these compounds was their basicity, i. e., they formed salt-like compounds with acids, although they were but weakly alkaline to the usual indicators of alkalinity.

COMPOSITION AND GENERAL PROPERTIES OF ALKALOIDS.

The alkaloids all contain nitrogen, carbon and hydrogen and all but two of them contain oxygen. A large proportion of the alkaloids are non-volatile, solid, crystalline compounds, while a few are volatile liquids but contain no oxygen. They are generally insoluble in water but are generally soluble in alcohol and possess varying degrees of solubility in ether, chloroform, amyl alcohol and carbon disulfide. These latter facts of solubility are often made use of to separate the alkaloids from each other, and from other substances. The salt-like compounds, on the other hand, generally possess a measurable degree of water solubility but not in the other solvents mentioned. They differ from the true alkalis, like potassium and sodium hydroxide, in the fact that the molecules of acids and alkaloids union with each other without forming other products. In this property, they resemble ammonia, HN_3 , when it forms such salts as ammonium chloride, NH_4Cl or NH_4Cl . However, this property does not characteristically belong to the so-called true alkaloids, since the amines, purin bases,

and ptomaines react in a similar manner. The volatile alkaloids, represented by nicotine and cocaine, possess disagreeable odors suggesting in part that of ammonia. The solid alkaloids, represented by strychnine, morphine, are odorless, but have a bitter taste often very characteristically bitter.

Physiologically, the alkaloid is quite generally a very active intoxicant, acting directly on the nervous tissues and producing results all the way from the atonic to the strongly tonic effects. The extreme effects are seen in the muscular excitation of strychnine and the depressing action of morphine, or, in the heart stimulation of atropine, and the depressing effects of cocaine.

OCCURRENCE OF THE ALKALOIDS.

The alkaloids are peculiarly a plant product and probably may be regarded as a protective agency to preserve a given species of plants. They are deposited in various parts of the plant but commonly in the seeds. The seeds containing, as they do, the vital parts of annual plants, (and also perennials) are protected from destruction by micro-organisms and by animals using them for food, by the intoxicating property, just referred to. A similar case is that of the glucoside amygdalin, as found in the seeds of several of the Rosaceae. The enzyme present in the seed, under proper conditions of temperature and moisture, decomposes it into glucose benzaldehyde and hydrogen cyanid or prussic acid, this latter compound acting as the intoxicant.

The alkaloids are not widely distributed in the plant world, although they are found in several orders or families of plants. Three families are especially characterized by the presence of alkaloids, namely, the poppy family or Papaveraceae, night shade family, Solanaceae, and the Rubiaceae.

The basic property of the alkaloids suggests the probability of their occurrence in combination with acidic compounds more or less characteristic of the plants in which they are found. In some cases the so-called alkaloid appears to be similar to glucoside, i. e. it can be hydrolyzed. For example, cocaine can be hydrolyzed into ecgonine, $C_8H_{11}NO_6$, benzoic acid, $C_6H_5CO_2H$ and methyl alcohol. Others are real glucosides like digitalin and solanin. It is undoubtedly true that the latter two should not be classed with true alkaloids, but with the glucosides, and are like caffeine theobromine in this sense, i. e. that they are substances that have been classed with the alkaloids on superficial grounds, such as bitter taste, but really have no close chemical relation with them. Caffeine and theobromine are now known to belong with the purine compounds. The purine compounds are basic, and hence their classification with the alkaloids.

The acids with which the alkaloids are often united are somewhat common in plants, or in a few cases they are characteristically found in combination with the given alkaloid, e. g., meconic acid in combination with morphine in opium, or aconitic acid united with aconitine. Other acids form compounds with the alkaloids in the various plants in which they are found. Among these acids are tannic, citric, malic and quinic. The combinations are easily broken up by strong bases, like potassium or sodium hydroxid, and in this way they may be separated from the acids. Since the true alkaloids are generally soluble in water while alkali salts of the acids are not soluble, the alkaloids may be separated by treatment with potassium or sodium hydroxid and filtering out the insoluble alkaloid. They are then further purified by the formation of the soluble salt and re-precipitating the alkaloid by an alkali. The alkaloid is then dissolved by the appropriate solvent and crystallized. Again it is interesting

to note that an alkaloid rarely exists alone in a given plant, but is accompanied by several others. For example, aconitin, as extracted from the roots of the aconite plant, *Aconitum napellus*, contains nine alkaloids, in the extract from poppies, called opium, upwards of seventeen alkaloids have been separated and studied, in the so-called chinchona or Peruvian bark extracts, some thirty-three distinct alkaloids have been isolated; strychnin is accompanied by brucin in *Strychnos ignatii*, or Saint Ignatius bean.

CLASSIFICATION OF THE ALKALOIDS.

Owing to the basic character of the alkaloids, and the fact that they always contain nitrogen, it was suggested that they were connected with ammonia in some manner, and if so, that they might be readily broken down by distilling them with potassium hydroxid or caustic potash. They were regarded as derived or at least connected with ammonia. Hoffman, who added much to our knowledge of the amine compounds, considered that they were of the ammonia type and were tertiary amines. In attempting to find some reaction characteristic of the amines, Gerhardt and others, heated the alkaloids with caustic potash but were unsuccessful in obtaining any results that showed that their basic character was due to this structural cause although some of them do possess some properties resembling the amines. Others more closely resemble the ammonium compounds. Products obtained by heating some of the alkaloids with potassium hydroxid and distilling the volatile products were found to be the same as were obtained from the destructive distillation of bones. Later 1834, a study of bone oil by Runge led to the separation of a pure compound which was shown to have the formula C_8H_7N known as pyridin. This was later shown to be a cyclic (hetero cyclic) compound, like benzene, one group "CH" being substituted by nitrogen, i. e., trivalent nitrogen. A number of alkaloids have been shown to be constructed on this nucleus by substituting various hydrocarbon groups and are known as the pyridin alkaloids.

Among the pyridin alkaloids and derivatives from them are nuxtin, conin, atropin, cocaine. The study of the alkaloids by noting the action of potassium hydroxid proved to be a fruitful one. In 1842, Gerhardt obtained a compound from the destructive distillation of quinine that was a new substance. This was named quinolin because of its origin from this alkaloid. It was later shown to be structurally composed of a benzene and a pyridin nucleus joined by two atoms of carbon in common, the further complexity is due to substituting in this nucleus. Most of the alkaloids are esters and are consequently quite readily separated into the two parts of such compounds namely the acid and basic parts. A study of these constituent parts gives the complete facts as to the structure of the original compound. The esters may be decomposed by acids, alkalis, and water.

The nucleus quinolin has been found in quinin, cinchonin, cinchonidin, strychnin, and brucin, and hence these compounds are known as the quinolin alkaloids.

THE ISO-QUINOLIN GROUP.

This group of alkaloids have a nucleus isomeric with quinolin known as the quinolin group. Like the quinolin group, iso-quinolin has the empirical formula, C_9H_7N . The difference between this base and quinolin, so far as structural constitution is concerned, appears in the position of the atom of nitrogen. Not

only is the fundamental nucleus different in the two classes of alkaloids, but groups that enter into these nuclei are different, thus leading to a large number of possible compounds. To this group of alkaloids belong especially the alkaloids of opium. For example, morphin, thebain, narcotin, narcein, papaverin, and codain. There are also hydrastin, hydrastrinum and berberin.

THE PURIN GROUP OF ALKALOIDS

These compounds are not properly included with the alkaloids but they belong with the purin compounds or xanthin bases. The nucleus (purin) or atomic framework, shown by Fischer to characterize the xanthin bases, such as uric acid, the xanthin derivatives, guamin and adenin, is found in caffeine, theobromin, theo-phyllin, compounds that are yet classed with the alkaloids. The structural and mutual relationship of these to each other and to the xanthin bases has been determined synthetically and hence their classification is not a doubtful question. The alkaloids are methyl xanthina. Caffein is a tri-methyl xanthin and theo-bromin and theo-phyllin are di-methyl xanthins, the latter being an isomer of theo-bromin. The alkaloid of tea is sometimes called thein although its identity with the caffeine of coffee has been recognized for a long time.

There are several alkaloids whose structural relations have not been determined. Among these are pilocarpin, colchicin, and physostigmin. For the description of these alkaloids see statement in this text under their appropriate heads.

PHYSIOLOGICAL ACTION OF ALKALOIDS AS DETERMINED BY THEIR STRUCTURAL COMPOSITION.

It is within comparatively recent time that pure alkaloids have been prepared and consequently that their physiological effects could be determined. Some of these like quinin were prepared in a fairly pure condition during the first half of the last century but most of them belong to a later period. Attention has been called to the fact that a single alkaloid is rarely produced by the plant and hence in the usual extracts from the plant there are several alkaloids with varying physiological effects. These constituents vary in amount according to a variety of conditions under which the plant may produce them. Accordingly the extracts of such plants will vary in the proportion of the alkaloid present. For example, in cinchona bark the amount of quinin may vary from 2 to 13 percent. The physiological effect of the cinchona extracts will be markedly different in such extremes of composition, i. e. in reference to the alkaloid quinin.

In 1869, Crum, Brown, and Frazer called attention to the relation of the structure of organic compounds to their physiological effects. They were studying the comparative action of strychnin and brucin and although their knowledge of the structure of these two compounds was not complete yet they were able to trace a relation in this case. They found that the presence of methyl or ethyl groups strongly affected the active properties of these compounds. For example, brucin is regarded as the dimethoxy-derivative of strychnin. The discovery that most alkaloids are built about three nuclei, namely pyridin, quinoim, and iso-quinolin added new zest to the study of the physiological effects of certain organic groups when substituted in organic compounds.

Quinoim is a strong antipyretic and antiseptic but produces other results

that are decidedly toxic. By introducing the organic group, methoxy CH_3O , a compound is produced from which several derivatives have been formed, which possess antipyretic properties that have led to their use in medicine. These are known as analgen, kavin, kauloin, and thallin. Quinin differs from cinchonin in containing the methoxy-group in place of a hydrogen atom. This addition produces a much stronger antipyretic. The antipyretic effect of the methoxy group is shown in some of the derivatives of anilin. For example, anilin is a highly poisonous compound although it has good antipyretic properties. The addition of a similar group to the one in question produces a compound known as acetanilid or antifebrin. The group added, CH_3CO , is known as acetyl. In introducing the methyl group into acetanilid its antipyretic properties are somewhat reduced, but its anti-neuralgic properties are increased. The addition of hydrogen to quinolin increases its toxic properties while the hydroxyl group increases the antifebrin results. The antipyretic properties of anipyrin are probably due to the two methyl groups present in the molecule. The ethyl group when added to organic compounds that are poisonous, in many cases at least, decrease their toxic character. The same is true of the acid or carboxyl group. When morphin is heated with hydrochloric acid, water is formed, as in hydrolysis, and a new compound known as apo-morphin. This treatment has developed two hydroxyl groups. This new compound is an excellent emetic. The introduction of two acetic acid groups changes morphin into a mildly acting sedative known as heroin. Finally the purin alkaloids owe their properties to the methyl groups introduced into xanthin i. e. as distinct from the xanthins.

The accumulation of facts in this direction is increasing rapidly and some generalization from them will do for medicine what antisepticism and anesthetics have done for surgery, and change it from an empirical to a scientific basis.

Blyth in his work on "Poisons, Their Effects and Detection" has classified the alkaloids and other organic poisons into the following groups.

ALKALOIDS.

First group, liquid volatile alkaloids, under which there are grouped the alkaloids of Conium (conium, conhydrin), tobacco (nicotin), puerine from *Duboisia Hopwoodii* belonging to the same family as tobacco, and spartein from the Common Broom (*Spartium scoparium*).

Second group, the opium group, contains the alkaloids from the Poppy the more important being morphin, thebain, codem, of the Morphin group, and papaveramin, codamin, laudanin, nartotin, papaverin of the Papaverin group.

Third group, the strychnin or tetanus-producing group of alkaloids. Under this head are included the alkaloids strychnin and brucin, derived from *Strychnos Nux vomica* of the family *Loganiaceae*, the alkaloids *aspidospermin*, *quebracho* obtained from the Quebracho (*Aspidosperma quebracho blanco*) of the family *Apocynaceae*, and at least four others, the alkaloids from the Pereira bark *Geissospermum Vellozo*, pectinin; the alkaloids from Geisemum or Carouba Jessamine (*Geisemum sempervirens*) of the family *Loganiaceae*, gelsemin and gelseminin, the cocaine alkaloids obtained from *Erythroxylon coca*, cocain, hygrin, etc.; the alkaloids from the roots of the European *Corydalis cava*, corydalin, corybulbin, and six others. The corydalin in large doses causes epileptiform convulsions.

Fourth group, alkaloids aconitin, atisin, and japaconitin of the aconite group, obtained from several species of *Aconitum*, as *Aconitum Napellus*, *A. heterophyllum*, etc.

Fifth group, the mydriatic group of alkaloids. The alkaloids of this group are atropin, obtained from *Atropa*, *Belladonna*, and *Datura*; hyoscyamin obtained from *Datura*, *Hyoscyamus*, *Scopolin carnioleica*, and *Dubouia* (Hyoscin pseudohyoscyamin being also obtained from these plants), scopolamin from some of the same plants as the preceding, solamin which is, however, regarded as a nitrogenized glucoside, obtained from various species of *Solanum*, solanidin with stronger basic properties than solamin, obtained from plants of the same family; cytisin obtained from *Laburnum* (*Cytisus Laburnum*) of the family *Leguminosae*, found also in quite a number of other plants of the same family.

Sixth group, the Veratrum alkaloids containing the alkaloids jervin, pseudo-jervin cevadin, etc., obtained from various species of *Veratrum* as *V. album*, *V. viride*, etc.

Seventh group physostigmin, the most important alkaloid of the group derived from the Calabar Bean (*Physostigma venenosum*), and calabarin.

Eighth group, containing pilocarpin, obtained from the leaves of jaborandi (*Pilocarpus pennatifolius*) and four other alkaloids, jaborin, pilocarpin, isopilocarpin, pilocarpin. Jaborandi belongs to the family *Rutaceae*.

Ninth group, taxin, obtained from the yew tree (*Taxus baccata*).

Tenth group, the curare alkaloids which are obtained from the Curare plants (*Strychnos toxifera* and *S. Castelnai*) of the family *Loganiaceae*. The alkaloids are tubo-curarin, curin, etc. Prolocurin obtained from the latter species is a slightly toxic substance.

Eleventh group, colchicin alkaloid, this alkaloid is obtained from the seeds and roots of the common meadow-saffron or Colchicum (*Colchicum autumnale*).

Twelfth group, muscarin from the *Amanita Muscaria* or Fly agaric.

GLUCOSIDES.

The glucosides widely distributed in plants are compounds of glucose and organic acids and are certainly of great importance in connection with the poisonous principles found in plants. They have been grouped by Blyth into A first group the *digitalis group*, consisting of digitalin, digitonin, and digitogenin, all found in the common fox glove (*Digitalis purpurea*), (2) second group of glucosides acting on the heart and containing antiarin obtained from *Antiaris toxicaria*, the upas tree; the helleborin and helleboretin found in *Helleborus niger*, *H. viridis*, *H. foetidus* and eonymin, a resinous substance found in Wahoo (*Euonymus atropurpureus*) which is a powerful heart poison, the third group containing thevetin, obtained from *Thevetia nerifolia*; atrophantin from *Strophanthus hispidus* of the Dogbane Family belong to this group of heart poisons, but it is not a glucoside and only partly crystallizable; scillarin from squill; adonidin from the root of *Adonis vernalis* of the Crowfoot Family; oleandrin from the Oleander; nerin also from the Oleander, sometimes called the Oleander digitalin and the poison of the Madagascar Ordeal plant (*Tonghinia venenifera*). The fourth group contains the digitalin-like apocynin from the common Dogbane, and other Apocynums; erythrophleuin, convallamarin, a glucoside from the Lily-of-the-valley; coronillin from *Coronilla*, and cheiranthin from *Cheiranthus*. These behave like the Digitalins.

GLUCO-ALKALOIDS.

The gluco-alkaloids represent a class of compounds intermediate in connection between the alkaloids and glucosides. The achillein found in Yarrow (*Achillea millefolium*) and solamin in various species of *Solanum* should be mentioned. The latter substance has, however, been referred to in connection with the alkaloids.

SAPONINS.

The Saponins have been treated fully in another connection. They are poisonous and when dissolved in water form solutions which froth. Of these mention may be made of saponin and senegin.

OTHER VEGETABLE POISONS.

A third division of poisonous substances includes those which cannot be readily classified and under this head is santonin, a lactone found in the heads of *Artemisia* sp. A second division of this group is mezereon obtained from *Daphne Mezereum*. A third group is ergot of rye, containing ergotin, secusmatxin, and other substances referred to at length in another connection.

PICROTOXIN, CICUTOXIN AND TOXINS.

Picrotoxin is the active principle of the Indian Berry (*Corculus indicus* or *Menispermum coccineus*), which contains the active principle picrotoxinin, picrotin and menispermnin.

Tutin, a non-nitrogenous glucoside, is obtained from *Corlaria sarmen-tosa* and other species. Another poison belonging to the picrotoxin class has been isolated from the Japanese *Illicium anisatum*, a member of the Magnolia Family. The plant is sometimes called the Japanese Star Anise. To this group may also be added cicutoxin obtained from the Cowbane, the oil of savin obtained from the common savin (*Juniperus sabina*), croton oil expressed from the seed of *Croton tiglium*, the toxicalbumins of castor oil seed and of Abrus; nitrogen from various species of lupines (*Lupinus luteus*, *L. hirsutus*, etc.); the toxic substances in the cotton seeds; toxic substances in various species of Lathyrus, the toxic substances in Arum; in the black bryony (*Tamus communis*); the toxicalbumin of the black locust, and the poisonous substances of the male shield fern.

ANIMAL AND PLANT TOXINS.

Another group of poisonous substances is included under the head of ptomaines and animal toxins. The word ptomaine is used in a rather indefinite way, and is open to objection, but the classification given by Blyth as animal toxin is also objectionable. Many of these toxins are the products of bacteria some of these poisons are, however, the products of higher plants, as toxin of the black locust, *Abrus*), etc. The groups given by Blyth are the Amines, under which head we have methylamin, found in the cultures of the Comma bacillus, and the trimethylamin, non-toxic, found in a variety of putrefying substances.

AMINES

The Amines are basic and originate from ammonia; they include the diamines, belonging to the amin series, which are formed in putrefactive substances. Of these we have neuridin in putrefying substances, cadaverin, found

in cultures of *Spirillum* and putrid animal matter; mydalein, guanidin. The cholin group includes neuridin, betanin, and muscarin. The muscarin has been referred to elsewhere. The neurin is intensely poisonous, and atropin is an antidote to neurin. Tetanus produces tetanus. Tetanotoxin from tetanus produces tremor paralysis and violent convulsions. Mydatoxin contained in putrid horse flesh is poisonous in large doses, causing lachrymation, diarrhoea, and convulsions. Tyrotoxicon, isolated by Vaughan from milk, is toxic. The suso toxin isolated from hog cholera is said to be quite toxic.

ORGANIC ACIDS.

The last group of poisonous substances includes the organic acids, the most important of which is oxalic acid. This is widely distributed both in the free state and in combination with lime soda and potash in the vegetable kingdom. It occurs in some species of the geranium, spinach, *Phytolacca decandra*, pie plant, *Rumex Acetosus* and in *Atropa Belladonna*, in connection with potash. In Russian Thistle and *Salicornea* it occurs in combination with sodium. In clover, apple twigs, begonia, and many other plants it occurs in the form of so-called compound aggregate crystals, or rosettes of calcium oxalate, in the onion and some other plants of simple crystals. In aroids and Virginia Creeper it appears in the form of needle shaped crystals, known as raphides, which are formed during the metabolism of the plant, the oxalic acid being set free and uniting with the lime in the plant to form calcium oxalate.

Oxalic acid is commonly used by dyers and calico-printers and also by curriers and harness makers for cleaning leather, to remove iron stains, to bleach straw, etc.

Several cases of poisoning have been attributed to the use of plants (like the sheep sorrel, oxalis, etc.), that contain large amounts of oxalates.

Quite a number of cases of poisoning from this acid are reported, especially in Europe. The smallest dose of oxalic acid known to have destroyed life, according to Dr. Taylor, is 60 grams. Oxalic acid acts upon the central nervous system. There is temporary loss of voice, burning in the throat, burning in the stomach, vomiting, especially bloody matter, pulse weak, locally it acts on the mucous tissues.

Part II

A Descriptive Manual of the Poisonous Plants of North America Chiefly of the United States



Commane—Very Poisonous After Vasey

KEY FOR THE PLANT KINGDOM

Organisms without chlorophyll, the vegetative body a naked mass of protoplasm with many nuclei, reproduction asexual, spores free or enclosed in sporangia, spores produce motile swarm spores or amoeboid bodies.

A. *Myxothallophyta*. 158

Cells generally with cell membrane, with one or more generations, sexual reproductions frequently absent, the fertilized spores when present with one cell, which later separates from the mother plant, or a several celled body resulting from the fertilization of the female fructifying body, which later develops into a new plant.

B. *Euthallophyta*. 160

Small unicellular organisms, never green but frequently of other colors, blue greens, etc.; reproduction asexual by fission; spores formed in the interior of the cell or by transformation of vegetative cells into the endospores or arthrospores; nuclei absent, but a so-called "Central body" occasionally present; coloring matter equally distributed.

I. *Schwamphyta*. 160

Unicellular organisms generally colorless at least never green; membrane, consisting of a chain-like substance, occasionally gelatinous, cells frequently with cilia; reproduction by fission; spores when present endospores or arthrospores.

Schizomycetozoa. 161

Unicellular organisms like the preceding, cells contain chlorophyll and phycocyanin consisting of blue, blue-green, violet, or reddish pigments; swarm spores absent.

Schizophyceae. 184

One-celled organisms with nucleus sharply differentiated protoplasmic body with a simple denser protoplasmic membrane, pseudopodia absent, motile during most of their existence, cilia 1 or more, and with 1 or 2 pulsating vacuoles; chromatophores occasionally absent; reproduction asexual by longitudinal division.

II. *Flagellatae*. 188

Plants occurring mostly in water, always with cell-membrane and nucleus; green or other colors (brown or red) mixed with the green

III. *Euphyceae*. 188

Small unicellular organisms occasionally forming chain-like colonies. Cells possess two long cilia which arise from a furrow in the ventral surface. Found mostly in the plankton of salt water

Peridinales. 188

Small one-celled organisms of brown color, the chlorophyll masked by diatom found in chromatophores; cell-wall consisting of silica with a girdle and fine lines; reproduction asexual, division parallel to the long axis of the organism, and the formation of auxospores and sexual, by the formation of auxospores by conjugation.

Bacillariales. 188

Chlorophyll green algae; membrane without silica, reproduction by division,

swarm cells absent, sexual reproduction by zygospores through the union of two equivalent gametes. (Aplanogametes). *Conjugatae* 189

Chlorophyll green plants, occurring singly or in colonies (threads or flattened bodies, , one or more nuclei, reproduction asexual by producing multilateral zoospores and nonmotile aplanospores; sexual by the copulation of zoogametes, or spermatozooids and oospheres; the spores produce a new plant directly or generally produce swarm spores. *Chlorophyceae* 190

Plants of brackish or fresh water, consisting of internodes, short whorls of cylindrical branches, cells nucleated, growth from an apical cell asexual reproduction by means of bulbils and vegetative threads; sexual reproduction by egg-cells and spermatozooids, the latter are spirally coiled in the cells of the antheridium, the egg cell is contained in a spirally coiled structure and after fertilization becomes an oospore. *Charales* 193

Brown algae, chlorophyll masked by a brown coloring matter phycophaein, reproduction sexual and asexual, swarm spores, sperm cells and egg cells marine, tetragonidia absent. *Phaeophyceae* 194

Brown algae, reproductive bodies without motion tetragonidia present *Dictyotales* 194

Red or violet algae, chromatophores contain chlorophyll and a red coloring matter (phycoerythrin and rhodophyll), reproduction sexual and asexual, mostly marine. *Rhodophyceae* 194

Parasitic or saprophytic plants with one or more cells, chlorophyll absent, with apical growth mycelium usually evanescent, reproduction sexual and asexual, generally the latter; asexual by the formation of zoospores, conidia or spores. **IV Eumycetes** 195

The vegetative body mostly 1-celled, tubular, asexual by the formation of spores or endospores sexual by the formation of zygospores. *Phycomycetes* 195

Copious nonseptate branched mycelium, asexual reproduction by endospores or chlamydospores; sexual by zygospores. *Zygomycetes* 195

Mycelium occasionally sparingly developed, tubular, asexual, reproduction by swarm spores or conidia, sexual by the formation of oospores. *Oomycetes* 204

Mycelium, many celled, reproduction asexual or sexual by union of nuclei, conidia borne on basidia, number various. *Basidiomycetes* 209

Mycelium many celled; reproduction sexual and asexual, the latter by conidia, pycnidia and spermogonia with spermatia, sexual spores in sacs known as ascospores called ascospores. *Ascomycetes* 247

Fungi whose spores are not in sacs, or consist of sterile mycelium. Forms like *Oidium*, *Oosonium*, or *Mycorrhiza* *Fungi Imperfecti* 281

Organisms consisting of a fungus and an alga. Spores either in sacs (*Ascolichenes*) or borne like toadstools (*Hymenolichenes*) *Lichenes* 307

Plants with stem, root and leaf, corinophyte or in some cases thalloid. Two generations, gametophyte and sporophyte, antheridium with sperm cells tube cell absent *C Embryophyta Zoidiogama* 308

Many celled differentiated structure frequently with leaves and stem or

thalloid in some cases. Male (antheridium) and female (archegonium) organs are produced. Asexual spores in spore cases which open at the top, in true mosses.
I *Bryophyta*. 308

Spores alike or unlike, microspores and macrospores developing into flat or irregular prothallia, these bear the reproductive organs, (antheridia and archegonia), flowers and seeds absent, usually a well developed vascular system.
II *Pteridophyta*. 312

Plants with a microsporangium (anther) containing the microspores (pollen grains) which develop a tubular body, the prothallium (pollen tube) a macrosporangium (ovule) containing the macrospore (embryo-sac) which develops into a minute prothallium, this remains enclosed in the macrosporangium, after the fertilization of the egg cell in the macrospore a seed develops, plants with flowers and usually well developed tissues, the epidermis, parenchyma and vascular (*Embryophyta Siphonogama*)
D. *Spermatophyta* 325

Ovules not enclosed in an ovary I *Gymnospermae* 325

Resinous trees or shrubs, wood with tracheids, tracheae usually absent, fruit a cone of dry or fleshy scales.
Coniferae 327

Ovules enclosed in an ovary II *Angiospermae* 332

Embryo with 1 cotyledon, stem without distinction into pith, wood and bark, endogenous, leaves usually parallel veined, flowers generally on the plan of 3.
1. *Monacotyledoneae* 332

Flowers generally small, unisexual, regular with persistent perianth 6 or 3 stamens, carpels free or rarely united fruit a berry, drupe or nut, embryo small; endosperm copious.
Principes 369

Palm-like plants with palm-like leaves, flowers naked or with thick leaves of perianth, carpels 2 or 4 with 2 or 4 placentae
Synanthae

Mostly fleshy herbs or thalloid floating plants inflorescence a fleshy spadix subtended by a spathe or naked or reduced to few or solitary flowers on the margin or back of a thalloid body
Spathiflorae 370

Herbs generally with narrow leaves, flowers usually complete, their parts mostly on the plan of 3, corolla regular or nearly so, ovary compound superior, endosperm mealy.
Farinosae 372

Mostly herbs flowers with a well developed perianth, usually regular and complete; usually on the plan of 3, ovary superior or inferior compound; endosperm fleshy or horny
Liliiflorae 374

Large herbs; flowers irregular; ovary inferior, compound composed of several united carpels, seeds generally arillate, frequently with perisperm and endosperm.
Scitamineae 390

Herbs, tropical species frequently epiphytes, flowers very irregular, or in one family regular, generally complete and perfect, parts of the perianth in 3's or 6's, ovary inferior, compound; seeds numerous, endosperm present or absent.
Microspermae. 392

Embryo usually with two cotyledons, stem usually with wood, pith and bark marked, usually exogenous, leaves mostly netted veined, flowers frequently on the plan of 5.
2. *Dicotyledoneae*. 395

Petals separate or distinct from each other or wanting, occasionally some

carpels united, in one division ovules with many megaspores. In most plants of the other division there is a single megaspore with synergidal and antipodal cells. (*Choripetalae* and *Apetalae*) *a. Archichlamydeae* 395

Tropical plants with monoecious flowers and branches with a longitudinal ridge in which the stomata are hidden. *Verticillatae* 395

Dicotyledonous herbs, petals and sepals absent, flowers small spicate with bractlets *Piperales* 396

Trees or shrubs, flowers small, in catkins, imperfect sepals and petals none leaves simple, fruit a many-seeded capsule, seeds with tufts of hairs at one end. *Salicales* 396

Trees or shrubs; leaves simple, flowers small, monoecious or dioecious, in catkins perianth absent, ovary 1-celled, style short stigmas 2; endosperm none *Myrtales* 397

Woody plants with simple leaves, flowers, staminate spicate pistillate solitary *Balanopsidales* 399

Shrubs or trees, flowers small, dioecious, in catkins leaves simple, alternate, petioled, perianth absent in staminate flowers; pistillate, subtended by bractlets ovary 1-celled; endosperm thin. *Leinerviales* 400

Trees with alternate, pinnately-compound leaves flowers monoecious, with bractlets, staminate in catkins, pistillate, solitary or several, ovule solitary, erect; fruit a drupe, indehiscent or dehiscent, with woody husk, seed large 2-4 lobed, endosperm none. *Juglandales* 400

Trees or shrubs flowers small, calyx usually present monoecious, or rarely dioecious, in catkins, pistillate flowers subtended by an involucre which becomes a bur or cup in fruit. *Fagales* 402

Shrubs, herbs or trees, calyx present but corolla absent, flowers small, not borne in catkins, monoecious, dioecious or polygamous; ovary 1-celled superior *Urticales* 404

Shrubs, trees or herbs with scattered leaves, flowers in spikes, racemes or panicles usually perfect, single carpel. *Proteales* 415

Herbs or shrubs, generally parasitic, calyx present, corolla absent, flowers perfect, or imperfect, a single inferior ovary; fruit various. *Santalales* 415

Generally vines or herbs, leaves cordate, or reniform corolla absent calyx inferior, tube wholly, or partly adnate to ovary flowers perfect *Aristolochiales* 416

Generally herbs, occasionally trees, shrubs or twining vines leaves simple, mostly entire flowers small, regular, perfect, dioecious, monoecious or polygamous petals absent, stamens 2-9, filaments filiform or subulate ovary superior one-celled ovule solitary, fruit an achene, endosperm mealy *Polygonales* 417

Generally herbs, occasionally shrubs, flowers perfect, corolla usually absent when present polypetalous, calyx present, ovary superior, embryo coiled curved or annular, albumen present *Centrospermae* 423

Herbs shrubs or trees calyx usually of separate sepals, corolla generally present, polypetalous, ovary superior carpels many usually separate stamens generally free and more numerous than sepals. *Ranales* 444

Generally herbs, flowers regular and perfect, petals generally separate, stamens free ovary compound, superior, free from calyx. *Rhoeadales* 479

Carnivorous plants, flower scapose, corolla with separate petals or nearly so sepals generally distinct, stamens free, ovary compound superior *Sarracenales* 497

Herbs, shrubs or trees, usually with petals which are separate, stamens generally perigynous or epigynous, sepals generally united or confluent with receptacle which is concave, carpels 1 or more distinct or united into a compound ovary. *Rosales* 498

Trees, shrubs, or herbs usually with petals which are separate, united in some or entirely wanting sepals mostly distinct stamens few or occasionally more than twice as many as the sepals, alternate or opposite with them ovary compound, superior *Geraniales* 574

Trees, herbs or shrubs, petals usually present and separate sepals generally distinct, stamens opposite usually fewer than sepals or as many occasionally more than twice as many, ovary superior, compound, ovules pericarpous *Yapindales* 604

Shrubs, small trees or occasionally vines leaves generally alternate, flowers, small, regular stamens as many as sepals or calyx lobes alternate or opposite with them, ovary compound superior, ovules erect *Rhamnales* 620

Trees, shrubs or herbs leaves simple, mostly alternate, flowers regular, usually perfect, sepals separate, or more or less united, petals separate, or wanting, stamens usually numerous ovary superior, compound disk inconspicuous or none. *Malvales* 621

Shrubs, trees or herbs flowers generally complete, perfect, and regular or irregular in some, sepals distinct, or more or less united, petals almost always present and distinct, stamens usually numerous, ovary compound, superior, placentae mostly parietal. *Parvetales* 627

Fleshy plants, leafless, or with small leaves, generally spiny flowers mostly solitary, sessile, regular, perfect and showy, calyx tube adnate to ovary limb many-lobed, petals numerous stamens numerous, ovary 1-celled ovules numerous, fruit a berry. *Opuntiales* 634

Shrubs, trees or herbs leaves simple petals usually present and distinct, calyx 4-5 lobed or entire and petals wanting in *Thymeleales*, superior or inferior, ovary 1 or more celled ovules 1 or numerous. *Myrtiflorae* 637

Herbs, shrubs or trees with petals, leaves of calyx usually 5, stamens 4 to 5, ovary epigynous, adnate to calyx, ovule, 1 in each cavity *Umbellales* 645

Petals partly or wholly united rarely separate or wanting, coherence variable in some cases; tubular or funnel-form (*Sympetalae* or *Gamopetalae*)
b. *Metachlamydeae* 664

Flowers complete, regular with lobed or distinct calyx, corolla cleft gamopetalous; stamens free from corolla; ovary compound *Fricates* 664

Mainly herbs; corolla gamopetalous calyx generally free from ovary, stamens borne on corolla, as many as its lobes, or twice as many or more. *Primulaes* 675

Trees or shrubs, alternate, simple leaves, flowers generally regular, calyx free from ovary, inferior, corolla gamopetalous or polypetalous, stamens borne on tube, at base of corolla. *Ebenales*. 679

Trees, shrubs, herbs or vines, generally with opposite leaves, flowers regular, corolla generally gamopetalous, or rarely polypetalous or wanting, stamens borne mostly in lower part of corolla, as many as lobes or fewer, a ternate ovaries 2 and distinct. *Conartias*. 670

Rarely trees, shrubs, generally herbs, corolla nearly always gamopetalous, regular or irregular, stamens adnate to corolla tube, ovary one, superior compound. *Tubiflorae*. 698

Herbs frequently acaulescent or caulescent with opposite or alternate leaves; flowers small, perfect, polygamous or monoecious, calyx 4-parted, corolla free, stamens 2 or only 1, ovary sessile, superior, 1-2-celled or falsely more celled, fruit a pyxis. *Plantaginales*. 730

Plants with gamopetalous corolla, stamens as many as corolla lobes and alternate with them, or occasionally fewer, or twice as many, ovary compound inferior, adnate to calyx tube or ovary 1 or more celled, ovules 1 or more in each cavity of ovary, leaves opposite or verticillate. *Rubiales*. 740

Herbs or rarely shrubs with gamopetalous corolla or occasionally petals separate, stamens as many as corolla lobes or fewer, anthers generally united, ovary inferior. *Companulatas*. 748

MYXOTHALLOPHYTA

Fungus-like organisms without chlorophyll, regarded by some as animals, intermediate, in some respects, between animals and plants and hence called Mycetozoa by Rostafinski. In their vegetative condition, they consist of naked masses of protoplasm with many nuclei, the mass of protoplasm being called the plasmodium which creeps about on the substrata changing in form and thrusting out processes called pseudopodia which may later coalesce. After



Fig. 28. Slime Mould (*Trichia torris*). a. Before germination. b, c, d. Different stages in germination. d, e. Amoeboid body with Bagellum. After DeBary.




Fig. 18. Various slime moulds. *a-f*. Club root of cabbage, *Plasmoidophora Brassicae*; *a*. Swollen root *b* Spore, *c*. Spore germinating *d*. Plasmodium *e*. Cells showing aggregated masses *f* Spores in cells. *g*. *Lycogala epidendron*, *h, j*. Plasmodium with branches, *i*. Spore, *k*. Spore germinating showing cilium, *m*. *Stemonitis*; 1. Stipe, 2. Columella, *o*. *Capillitium*, *p*. *Trichia decipiens* sp. Sporangia, *q*. Elater, *r*. Spore.

a shorter or longer period the protoplasm contracts forming little heaps which contain the spores, the parts of the reproductive body are called the *sporangium* or spore case, the *peridium* or the wall of the case, the *stipe* or stalk, the *columella* or central axis in the spore case, the *capillitium* or fine threads, and the *spores*. The spores after absorbing water, germinate by breaking the wall and move about by means of cilia; sexual reproduction is entirely absent.

The division Myxothallophyta includes three classes. *Acrasidae* without swarm cells, *Plasmodiophorales* of which the club root of Cabbage, *Plasmodiophora Brassicae* is an example (a very destructive parasite upon cabbage, turnip, etc., in Europe and the Eastern States), and *Mycogasteres* which contains a great many species and genera common on spent tan bark, rotten logs, and the ground. Of the third class *Stemonitis Physarum*, *Lycogala* and *Fuligo* are common genera. No species of this class is poisonous so far as known.

EUTHALLOPHYTA

Cells generally with cell membrane, with one or more generations, sexual reproduction frequently absent, the fertilized spores when present, with 1 cell which later separates from the mother plant, or a several-celled body resulting from the fertilization of the female fructifying body, which later develops into a new plant. This division includes such plants as bacteria, blue green algae, the green algae, rusts, smuts, mildews, moulds, puffballs, mushrooms and toadstools.

SCHIZOPHYTA

Small unicellular organisms, never green but frequently of other colors, blue greens, etc., reproduction asexual by fission, spores formed in the interior




Fig. 19 Schizophyta. Schizomycetes Bacteria. 1 and 2. *Bacillus subtilis* 3 and 4. *Bacillus anthracis*. 1, 3 and 4x1000. 1 3, and 4 after Fränkel and Pfeiffer. 2 after Migula.

of the cell by the transformation of vegetative cells into endospores, or by the transformation of ordinary vegetative cells into arthrospores, nuclei absent, but a so-called "central body" occasionally present, coloring matter equally distributed. This sub-division includes the Bacteria or Schizomycetes and the Blue-green Algae or Schizophyceae.




Fig. 20. Schizophyta. Schizomycetes. Bacteria with flagella. 1. *Planococcus citreus*. 2. *Pseudomonas pyocyanea*. 3. *Pseudomonas syzyganea*. 4. *Bacillus typhi*. 5. *Spirillum temma*. 6. *Spirillum rubrum*. Fig. 1-6x1000, all after Migula.

SCHIZOMYCETES

Schizomycetes is one of the two classes of the sub-division Schizophyta. All the members of this sub-division are characterized by having no known sexual method of reproduction, multiplying by means of simple fission or cell division. The bacteria are distinguished from the fungi, or Schizophyceae, by the absence of the blue-green coloring matter which is characteristic of these forms. The two sub-divisions approach each other very closely at some points particularly among the branched bacteria. The shape of the bacteria is used as the character in the separation of the families. They are either rod shaped, and unbranched, spherical, bent, or spiral and straight and branched, and with or without sheathing, covering, or membrane. Five families are distinguished by Migula. Some of these contain considerable numbers of bacteria important from their toxigenic properties.

Bacteria are among the smallest of living beings, some undoubtedly being so small that they cannot be seen with the highest powers of the microscope. Others are large enough so that they may be seen as minute specks by the naked eye. In other words, they vary from less than $1/10 \mu$ to 100μ . They may be arranged in the case of the rod-shaped forms or bacilli, either singly or in chains. The same is true of the spirilla, or spiral forms. The cocci or spherical forms may be single, in pairs, in regular mass of 4 and multiples of 4, in chains, in irregular clusters, or imbedded in gelatinous mass forming zoogloecae. Mulu

plication, as before stated, is by simple fission or cell division. Growth takes place very rapidly in many forms, some being capable of growing to their full size and dividing to form two individuals in twenty minutes to half an hour. Under favorable conditions, this rapidity of multiplication explains the importance of the results obtained, and the products formed from such small plants. Some forms are capable of moving or swimming about by means of whips or flagella placed on all sides or simply at the ends. Other forms move by a sinuous or snake-like bending of the body. Many forms are incapable of motion. Spores are formed by many species; they are called endospores when formed singly within the bacterial cell, and arthrospores when formed by increase in size of the cells of the filament or part of a filament and its splitting into a




Fig. 2. The root tubercle organism (*Rhizobium leguminosarum*). 1. General view of root showing tubercles. 2. Root hair and strand with enlargements at a and c. 3. Cross-section of root at b bacterial tubercle. 4. Cells of clover plant filled with the organism nucleus at a. 5. Rod and y shaped organisms from 3b more enlarged. 6. Single cell containing bacterioids. After Frank.

number of small cells. These spores serve, on account of their great resistance to desiccation, and other unfavorable conditions, to tide the organism over until suitable conditions once more obtain. Bacteria are universally distributed, abounding in the soil, in the water, and being present often in the air except at high altitudes. Normally, they are absent from the tissues of living animals and plants, but are to be looked for practically everywhere else. Their food requirements are as various as their habitats. Some require the most complex organic compounds, while others cannot live in the presence of such, but manufacture their own food from inorganic substances. Most bacteria lie between these two extremes. In respiration, some bacteria require oxygen or air, others will not develop in its presence. Most species require an abundance of moisture for their development, but many species will withstand a considerable amount of drying. Light inhibits the growth and in many cases destroys the bacteria. As to heat requirements, some live only in hot water, others will develop upon the surface of ice, some best at blood heat, while most develop between 15° and 22° C.



Fig. 22. Schizophyta. Schizomycetes. Nodule forming bacteria. *Rhizobium leguminosarum*. 1. Root tubercle of *Lupinus*. 2. Cross-section of nodule. 3. Cell showing bacteria x 500. 4. Bacteria x 1500. After Woronin and Fischer.

Bacteria are also important in connection with the decomposition of organic matter. The nitrifying bacteria in the soil change the complex albuminous substances into nitric acid. This uniting with a base forms nitrates. The tubercle bacteria like *Rhizobium leguminosarum* are in mutual relation with clover and other leguminous plants and are important in the acquisition of nitrogen. Some bacteria play an important part in the dairy industry, the aroma and flavor of butter being due to these. Some, like the red milk organism (*Bacillus prodigiosus*), produce bad and disagreeable odors or cause the milk to become viscid or colored. Vinegar is produced by the acetic acid bacillus (*Bacillus aceticus*). Some bacteria produce diseases of plants like Fire blight of apples (*Bacillus amylovorus*), Cabbage rot (*Pseudomonas campestris*), Sorghum Blight, Corn wilt, etc. Some bacteria produce diseases of insects like Foul brood of bees, Silk worm disease, etc.

BACTERIA; POISONOUS PROPERTIES. It is believed best to consider in a general way, the various poisonous principles which are developed by bacteria before the discussion of the specific organisms and their specific poisons. Inasmuch as bacteria play a very important part in nature in breaking down dead tissues of all kinds, destroying them and returning them to their elements, or forming simple compounds, it is to be expected that among the multitude of chemical substances which are developed, there would be some which would be harmful

when taken into the body of man or animal. In fact, such decomposition products are known and as most of them, probably all, are basic, containing nitrogen, they have been grouped with that general class of vegetable alkaloidal substances called *Protonins*. If other poisonous substances than *Protonins* are developed, they are not known at the present time. In addition to these poisonous substances that arise as decomposition, other poisonous substances are produced by certain bacteria which are strictly synthetic that is not produced by the breaking down of complex compounds into more complex forms. The exact chemical nature of these substances is not understood, the reasons for this being that they are extremely unstable, it being impossible to heat them without destroying, and they cannot be recognized by any known chemical means. They must be distinguished and differentiated, and often detected only by animal inoculation and experimentation. These soluble substances excreted by the bacteria are called toxins. The term toxin is rather an unfortunate choice, because it refers simply to their poisonous properties. In the broad sense any poisonous substance is a toxin, but in the sense in which it will be here used, toxin indicates specific bacterial poisons excreted into the medium in which the




Fig. 33. *Bacillus clausii* from a corn disease, also found in sewage. Supposed at one time to produce toxic substances to which was attributed the corn stalk disease. After Burrill.

Fig. 34. Sorghum Rust (*Bacillus Sorghis*). a Young plant infected with the organism. b Leaf and sheath. c Bacilli. Modified after Kellerman and Swingle.

organism is growing and producing upon inoculation, anti-toxins. In addition to the products above mentioned, many bacteria undoubtedly owe their poisonous or intoxicating qualities to the fact that the protoplasm of living matter of the organism is poisonous or contains poisonous substances which are not excreted into the surrounding medium. When bacteria of this type are allowed to grow in favorable culture media for a considerable length of time, there is a certain amount of self digestion or autolysis which takes place and these

poisonous contents of the cells are liberated and then go into solution. They may be liberated, also, by grinding the bacteria, and extracting with water. These poisonous protoplasmic substances have been called toxalbumins, but this term commits one to the supposition that all of this type of poisonous substances are protein in nature. This has been by no means proven, consequently the term endotoxins is to be preferred.

The following terms used in discussions of immunity will need defining.

An Antitoxin is a substance capable of neutralizing a toxin by combining with it and is produced in the animal body as a reaction to the introduction of a toxin in non lethal doses.

A Bacterial Agglutinin is a substance produced in the animal body as a reaction to the presence of certain bacteria or their products. When introduced into a suspension of the organism the agglutinin will cause the bacteria to clump (agglutinate) into groups.

A Bacteriogram is a substance produced in the animal body as a reaction to the presence of bacteria or their products which will destroy and dissolve the corresponding (homologous) organisms.

An Opsonin is a substance found in the blood serum which will unite with bacteria and render them positively chemotactic for the white blood cells. This preliminary union of opsonin and bacterium seems to be necessary before engulfment and destruction of the bacteria by the white blood cells can take place.

In discussing the specific effects and products of the various species of bacteria, there would be an advantage in grouping these bacteria, according to the substances produced, and their effect upon animals and man. However, that this portion may be in keeping with the remainder of the text, the specific effect will be discussed and noted under each organism, and the organisms put in their correct place in the genera of Migula's system of classification. In many instances, bacteria not closely related produce effects that are very similar, in some of these cases, the discussion will be under the first of that group reached, the remainder of the group will contain simply the reference to the form under which the discussion is given.

COCCACEAE

Organisms globose or spherical in a free state, not elongated in any direction before division into one, two, or three planes, when united in pairs or groups, sometimes flattened on the proximal sides, containing five genera, three of which are of importance from our point of view.

BACTERIACEAE

Cells cylindrical or oval, dividing only in one plane, rods straight rod-shaped, without sheath, either non-motile or motile by means of flagella, contains three genera.

SPIRILLACEAE

Cells cylindrical, dividing in one plane, not straight, being bent or spiral, and without sheath, contains four genera.

CHLAMYDOBACTERIACEAE

Cells cylindrical, dividing in only one plane, enclosed in a sheath, contains five genera.

BEGGIATOACEAE

Cells cylindrical, dividing in only one plane, destitute of the sheath, united into threads containing sulphur granules, usually motile by means of the undulating membrane. One genus only




Fig. 25 Chlamydoacteraceae and Beggiatoaceae. 1. *Beggiatoa alba*. a Sulfur granules $\times 800$. 2-4. *Cladothrix dichotoma*. 2. Carnobium $\times 350$. 3. Part of filament magnified 1000 times. 4. Cells with flagellae $\times 1000$. 5. *Crenothrix polyspora*. a young threads. b, threads separating into bodies. c, older threads with spores $\times 1000$. Fig. 1, 2, and 5 after Zopf, Fig. 3 and 4 after Migula.

MYXOBACTERIACEAE

In the vegetative stage these occur as swarming rod-shaped organisms held together by a gelatinous substance secreted by the cells, they show slow creeping movements. They form cysts in which the spores occur. This very peculiar




Fig. 26. Myxobacteriaceae *Chondromyces pedunculata*. 1. General form of fructification, cyst and pedice. 2. Single cyst. 3. Rods forming the mass in cyst. After Thaxter.

group first described by Thaxter lives on the dung of animals and in habit resembles the Slime Moulds or Myxomycetes.

COCCACEAE

Micrococcus pyogenes, Var. *albus*, Rosenbach

White-pus Coccus.

This organism, when grown upon artificial media, produces no pigment, otherwise it is identical with the following.

Micrococcus pyogenes, Var. *aureus*, Rosenbach

Golden-pus Coccus.

Infections produced. This organism, or the preceding, or both, are found very generally associated with, and usually as the cause of wound infection and suppuration in general. They are usually found in furuncles, abscesses, carbuncles, and other inflammatory processes affecting the surface of the body. When present under certain conditions in the blood or various internal organs, they cause pyemia, septacemia, osteomyelitis, inflammations of serous membranes such as pleuritis, peritonitis, ulcerative endocarditis, etc.

Pathogenesis. Infection with this micrococcus causes a marked hyperleucocytosis. Its presence in tissues is generally followed by the production of pus made up of serum, polymorphonuclear leucocytes, disintegrated tissue, and the bacteria. Ordinarily, the area of infection and inflammation is walled off by an infiltration of the surrounding tissues by these polymorphonuclear leucocytes, forming the so-called pyogenic membrane.

Poisonous properties. In 1894, Van der Velde discovered that sterile filtrates from cultures of *Micrococcus* contain an hemolysin, which he termed staphylo toxin. In 1901, Neisser & Wechsberg studied this hemolytic substance and gave it the name staphelolysin. This substance will cause the erythrocytes to dissolve whether within or without the body. A true toxin, leucocidin, is



Fig. 27. Pus organism (*Staphylococcus pyogenes aureus*) shown in irregular bunches. After Flügge of the organism.

Immunity. The resistance of the body may be heightened by immunization with pure cultures of the organism. Immunity is undoubtedly due in large part to the phagocytic activity of the leucocytes. It seemed that virulence of the organism has very little relation to the production of toxins by them, for some very virulent types produce very small quantities of toxin. There is some substance secreted which is positively chemotactic to the phagocytes. Immunization is not due to the production of bacteriolysins in the blood. Immune sera have been produced, but have not proven to be of any practical importance. Vaccination is held by Wright to raise the opsonic index of the blood, to this he attributes increased resistance. Agglutinins are present in normal sera in most cases, but systematic injection greatly increases the agglutinating power of the blood.

Micrococcus coprinus, Mohler and Wapbury

Disease produced. Takosis

Animals originally infected. Angora goats

Susceptible animals. Mouse, guinea pig, rabbit

Animals naturally immune. White and brown rat, chicken, dog, and sheep.

Pathogenesis. Anatomically characterized by emaciation and anemia, congested pneumonic areas in the lung, splenic atrophy, and induration, spleen often being attached to the diaphragm and neighboring organs by fibrous tissues. The mucous membranes exhibit necrosed areas of mucosa and bacteria are distributed through the blood, consequently may be isolated from any of the internal organs.

Poisonous properties. Microscopically, the lungs are found with many of the terminal bronchioles and alveolae filled with mucus and desquamated epithelium, in the liver hyperaemia with fatty degeneration or periphery of many of the acini, catarrhs, nephritis, localized areas of parenchymatous degeneration in

the heart, capsule of the spleen thickened and contracted. Small intestines with a local superficial or completed necrosis of the glandular areas. The blood examination shows polycythemia, leucocytosis, and in advanced stages general poikilocytosis. True toxins not produced.

Immunity Filtrates seem to have some immunizing power, probably either bactericidal or opsonic. The serum of immune animals has little or no immunizing power. Sterilized cultures and filtrates heated for thirty minutes at 60° C. lose their immunizing power.

Micrococcus meningitidis, (*Diplococcus intracellularis meningitidis* Weichselbaum)

Name of disease Epidemic cerebral spinal meningitis.

Animals infected naturally Man.

Animals susceptible None, except when injected into a cavity in large quantities. Subdural inoculations in most cases produce a meningitis which however, does not agree with that produced in man.

Pathogenesis Severe inflammation of the meninges of both brain and spinal column characterized by the production of considerable quantities of pus. The lumbar puncture and a microscopic examination of the pus will show the organism present in large numbers. Probably the organisms sometimes reach the blood stream, and secondary infections are produced in various parts of the body by metastasis. It is very probable that the organism is present in acute rhinitis, and that the infection of the brain and spinal cord is secondary.

Poisonous properties Lapierre extracted what he called a toxin with glycerine, from old culture.

Immunity Second attacks of the disease are very rare. Lapierre claims to have immunized animals with the toxin and with cultures produced a preventive curative serum from immunized animals. Davis states that there is developed a bactericidal property in the serum, and also agglutinins.

Micrococcus lanceolatus, Fraenke

Disease produced Acute infectious pneumonia.

Animals infected naturally Man (and domestic animals).

Animals susceptible Rabbit, guinea-pig, dog, and mouse.

Animals immune Chicken and pigeon.

Pathogenesis The lungs are most frequently the seat of infection. The infected portion passing through several stages: first that of congestion, in which the air cells become filled with blood serum and red corpuscles, the former coagulates, the tissues become liver-like in consistency; this is followed by a marked invasion of leucocytes; the contents of the air cells soften and are absorbed and discharged. In most cases of the disease blood infection probably occurs, consequently in many cases infection of various other organs. Pleuritis is most common, then pericarditis, and even generalized peritonitis, endocarditis, arthritis, meningitis, otitis media, conjunctivitis, osteomyelitis, and degenerations in various internal organs, particularly the kidneys and liver.

Poisonous properties Poisonous substances are produced in greater or smaller quantities in culture media, but no true soluble toxin. Presumably, there is present an endotoxin. The pneumotoxin seems to be toxic toward all the organs of the body.

Immunity Immunity is probably due largely to phagocytosis and it is

likewise probable therefore, that it is due in large part to the opsonin content of the blood. There are no bactericidal or anti-toxic substances formed in immune blood. The blood serum of an immunized individual exhibits an increased agglutinating power. Immune sera for the prevention of the disease have not proven a success.

Micrococcus tetragenus, Gaffky

Disease produced. Associated with the tubercle-bacillus, and probably of importance in complicating the pus infections.

Animals infected. Man and animals.

Animals susceptible. White mouse, and guinea-pig.

Animals immune. House-mouse, field mouse, dog, and rabbit.

Pathogenesis. Inoculation of white mouse causes fatal bacteremia. The organism is found in tubercular infections, and probably hastens the necrosis of infected tissue. In some cases it may be the primary infecting agent.

Poisonous properties. Not known.

Immunity. Not known.

Micrococcus catarrhalis, Seifert

Disease produced. Superficial inflammations of the respiratory tract and conjunctiva.

Animals infected. Man.

Animals susceptible. None of the laboratory animals are susceptible, except when the organism is introduced in very large quantities.

Pathogenesis. Probably the primary cause in some cases of conjunctivitis, bronchitis, and catarrh, and in general superficial inflammations of the respiratory passages.

Poisonous properties. Not known.

Immunity. Not known.

Micrococcus gonorrhoeae, (*Diplococcus* of Neisser)

Disease produced. Gonorrhea.

Animals infected. Man.

Animals susceptible. None of the laboratory animals are susceptible.

Pathogenesis. Producing a severe inflammation of the mucous membranes of the urethra accompanied by blennorrhoea. Secondary infection of fallopian tubes, ovaries, uterine, etc., may occur.

Poisonous properties. Not known.

Immunity. Probably some immunity is developed after infection, but is not lasting. No method of immunizing is known.

Streptococcus equi, Schütz

Disease produced. Strangles.

Animals susceptible. Horses, asses, and their hybrids, and mouse.

Pathogenesis. Producing a severe catarrh of the nasal mucosa, with a swelling of the sub-maxillary, and pharyngeal lymphatic glands, abscesses generally form in the latter. May terminate in pharyngitis in a purulent pneumonia or pleuritis. Sometimes cutaneous exanthemata. Metastatic abscesses may appear in various lymph glands.

Poisonous properties. Not fully studied, but probably the same as those of the next organism, *Streptococcus pyogenes*.

Immunity Not known.

Streptococcus pyogenes, Rosenbach (*Streptococcus erysipelatos* Fehleisen)

Disease produced. This organism in various forms of inflammation and septic inflammation in general, sometimes alone, sometimes associated with




Fig. 26. Pus organism held by some authors to cause rheumatic fever. *Streptococcus pyogenes*. In chains with pus cells $\times 800$. After Flüge.

Micrococcus pyogenes albus, and *aureus*. Its specific cause, in many instances of septicemia, pyemia, phlegmon, abscesses, boils, erysipelas, ulcerative endocarditis, periostitis, otitis, meningitis, pneumonia, lymphangitis, bronchitis, inflammation of the serous membranes, as pericarditis, pleuritis, peritonitis arthritis, enteritis, endometritis, tonsillitis, salpingitis, has been held by some authors to cause rheumatic fever and also scarlet fever. Probably many so-called terminal infections are produced by this organism.

Pathogenesis. The organism is one of the pyogenic forms reacting much as the *Micrococcus pyogenes*, as has been described.

Poisonous properties. In many strains of *Streptococcus pyogenes*, there is present an endotoxin. This is little understood, however, at the present time. It is found that this endotoxin varies greatly, in some cases none at all being found in virulent types. It is susceptible to heat, organisms killed by chloroform being more poisonous than those killed by heat. Virulent streptococci also produce an hemolytic toxin called streptococlysin. This is a true toxin. The blood in fatal cases of streptococcal septicemia is often laked. The toxin is destroyed at a temperature of 70° for two hours and by peptic digestion. Substances which kill the leucocytes are also present in certain strains and inhibit phagocytosis. It is very probable that the pathogenic character of this organism is not entirely explained by its known toxic properties.

Immunity. Immunity against infection of streptococcus is probably due largely to the presence of opsonins in the blood, and the consequent activity of the phagocytes. Sera of animals which have been immunized by inoculation of non virulent or killed cultures seems to have some protective effect. Such has not come into general use, however. Agglutinins are produced for most strains.

BACTERIACEAE

Bacillus suispestifer, Salmon & Smith

Disease produced. Hog cholera. Probably not the primary cause, but associated with some unknown ultra-microscopic organism.

Animals infected. Swine.

Animals susceptible. Rabbits.

Pathogenesis. Post mortem examination reveals numerous petechiae, ecchymoses, and extravasations of blood into various tissues. This latter is particularly evident beneath the serous membranes. The spleen is enlarged, soft, and engorged. In subacute cases, large intestinal ulcers are formed.

Poisonous properties. Novy gave the name susotoxin to a poison base which he discovered in pure cultures. This is probably not a specific poison of the organism, certainly not a toxic.

Immunity Vaccination with killed or attenuated cultures develops an immunity, while vaccination against this particular organism is successful, it is not of practical importance, because of the fact that this particular organism is not the primary cause of the disease. Agglutination is well marked, and has been used in diagnosis.

Bacillus piscidus agilis, Sieber

Disease produced. No specific name given.

Animals infected. Fish.

Animals susceptible Frogs, mice, rabbits, dogs, and guinea-pigs upon inoculation.

Pathogenesis. Disease is marked by shortness of breath, ureter apathy, and finally paralysis.

Poisonous properties. The filtrate of cultures is poisonous, also the distillate. Cadaverin, and other known ptomaines have been obtained from cultures.

Immunity

Bacillus coli, (*Bacterium coli commune* Escherich)

Disease produced. This organism is a normal inhabitant of the intestinal tract of man and animals but under certain conditions produces inflammation of the internal organs, such as cholecystitis, peritonitis, meningitis, cystitis, suppurative nephritis and even generalized septicemia.

Pathogenesis. This organism is not highly pathogenic under ordinary conditions, and when found in inflammatory processes, it is generally associated with other organisms. Undoubtedly, one of the reasons why this organism is found in various organs in post mortems, is the fact that it gains entrance into the blood just before death, producing the so-called agonal invasion.

Poisonous properties. No specific coli-toxin has been produced.

Immunity. Animals may be immunized by the injection of killed cultures. The immunization resulting from the formation of bacteriocal antibodies and agglutinins.

Bacillus enteritidis, Gaertner

Disease produced. Meat poisoning.

Animals infected. Man.

Animals susceptible Mouse, guinea pig, rabbit, pigeon, lamb, and goat.

Animals immune Dog, cat, rat, pigeon, and sparrow.

Pathogenesis. It is believed at the present time, that many of the so-called cases of ptomaine poisoning which have occurred in the past, and been recorded as such in medical literature, are due to the presence of this organism, and its soluble toxin. The organism has been isolated repeatedly from spleen of fatal cases. The disease is contracted by eating infected meat, and is characterized by vomiting, and violent diarrhoea, followed by collapse, head-ache, and not infrequently urticarial or herpetic eruptions. Anatomical findings are not specific. Meat undoubtedly is infected only when coming from animals sick with an intestinal or general infection before they were slaughtered.

Poisonous properties. The organism produces in the meat, a soluble heat-resistant toxin in considerable quantities. This toxin in its heat-resisting properties, differs from most toxins and also in the fact that no anti-toxins are produced. Sufficient toxin is usually present to give the first effects of the

disease. The organism itself may or may not gain entrance into the circulation or organs.

Immunity. As before stated, anti-toxins are not developed. Agglutinins, however, are developed.

Bacillus typhosus, Zopf

Disease produced. Typhoid fever

Animals infected. Man. Inoculation of experimental animals usually negative, except when injected in considerable quantities.




Fig. 29. Typhoid fever bacteria (*Bacillus typhosus*). Section of tissue showing bacteria clustered in center. After F. Wang.

Pathogenesis. The organism invades the solitary lymph nodes and Peyer's patches in the intestine and produces more or less necrosis and sloughing of tissue. By means of the lymphatic channels, the internal organs are all infected, particularly the spleen which becomes very much enlarged. The bacilli invade the blood, and hence the disease is a true bacteremia. When they lodge in bony tissues, arthritis, periosteitis and osteomyelitis may be produced.

Poisonous properties. No soluble toxin has been discovered, an endotoxin, however, is present and may be secured through self-digestion in cultures, or by grinding and extracting the bodies of the bacteria.

Immunity. No true anti-toxin serum has been produced inasmuch as no toxin has been discovered. The blood serum of typhoid patients agglutinates the bacteria. Vaccination with killed cultures produces an immunity which lasts probably in most cases several years. The blood serum of animals immunizes against typhoid bacilli, but is not used because it is quickly thrown out of the system when injected, and because it possesses very little curative quality.

Bacillus tetani, Flügge

Disease produced. Tetanus, or lock-jaw

Animals infected. Man, white mouse, rabbit, guinea pig, mouse, rat horse, to a less degree cattle and most other warm blooded mammals. Most birds, amphibians, and reptiles are immune




Fig. 30. Tetanus Bacillus. *Bacillus tetani*. Spores and rods. a. Vegetative stage. b. Spore stage. After Abbott.

Pathogenesis. There are no gross characteristic anatomic changes, but microscopic degenerative lesions may be found in the ganglionic cells. The disease is characterized usually by a tetanus or rigidity of muscles.

Poisonous properties. A tetanus toxin is produced in quantities in media. It is believed to contain two principles, the first of which, the more important, affects the nerves, and is called tetanospasmin, the second which is hemolytic in its action is called tetamolysin. The toxin is destroyed by gastric and pancreatic digestion. It has a strong affinity for

nervous tissue, in test tube, practically all of the toxin will become fixed. It is absorbed in the body by the motor ends of the nerves and passes through this by means of the axis cylinders to the ganglionic cells.

Immunity. Natural immunity is probably in part at least, phagocytic in nature, but the presence of a toxin in the blood or in the body causes the production of the anti-toxin. The serum of animals immunized by toxin injections contains quantities of this anti toxin, so that it is used in immunization against, and in the cure of tetanus.

Bacillus botulinus. Von Ermengin

Diseases produced. Botulism, or meat poisoning

Animals infected. Man, principally

Animals susceptible. Guinea pig.

Animals immune. Dog and cat



Fig. 31. An organism found in sausage. *Bacillus botulinus*. After Jordan.

Pathogenesis. The ingestion of meat containing bacillus botulinus is followed, in from a day to a day and a half, by salivation, ataxia, bulbar paralysis, and death in from 25 to 50 per cent of the cases. It produces degeneration of glandular organs and vascular endothelium and consequent hemorrhages.

Poisonous properties. The organism growing in meat produces a characteristic toxin and it is this toxin already formed which produces disease, and not the proliferation of the organism after gaining entrance to the body. It has been found in decomposed hams, and sausages. The toxin differs from that of diphtheria and tetanus in that it is not digested by the gastric juice. It has a special affinity for nervous tissues, but is not so selective as tetanus toxin.

Immunity. Immunization with the toxin results in the formation of an anti-toxic serum which may be used in immunizing against the disease, or in curing. However, this disease is so rare that it is of no commercial importance in this country.

Bacillus alvei, Chesire & Cheyne

Disease produced. European foul brood

Animals infected. Honey bees.

Animals immune. None of the higher animals contract the disease when the organism is inoculated.

Pathogenesis. Destroys the larva of the honey bee

Immunity Not known.

Bacillus larvae, White

Disease produced. American foul brood.

Animal infected. Honey bee

Animals immune. All higher animals

Pathogenesis. Destroys the larva of the honey bee

Poisonous properties. Not known.

Immunity Not known.

Bacillus anthracis symptomaticus, Kruse

Disease produced. Black leg, Quarter evil or symptomatic anthrax.

Animals commonly infected. Cattle

Animals susceptible. Guinea pig, hog, dog, and rabbit

Animals immune. Bird, horse, goat

Pathogenesis. Irregular emphysematous pustules and areas. Muscles contain dark areas with blood serum and gas bubbles.

Poisonous properties. Doubtful, probably an endotoxin.

Immunity. Established by vaccination with the *Bacillus* attenuated by exposure to heat and drying.

Bacillus oedematis, Zopf

Disease produced. Malignant oedema.

Animals infected. Horse, sheep, goat, mouse, guinea pig, rabbit, dog, pig, chicken, and pigeon

Animals immune. Cattle

Pathogenesis. There is little blood infection by the organism but a general emphysema of the sub-cutaneous tissues, the gas bubbles being usually very numerous. Any of the body tissues may be affected.



Fig. 30b. Malignant Oedema. *Bacillus oedematis malignantus*. A. From spleen of guinea pig. B. From lung of mouse. Both $\times 700$. After Koch.



Fig. 30c. Malignant Oedema. *Bacillus oedematis malignantus*. Spores and rods. After Abbott.

Poisonous properties. Not definitely known.

Immunity. Immunization may be affected by vaccination with attenuated cultures. Attenuation is arrived at by passage of the organism through white rats.

Bacillus marsepticus, Koch

Disease produced. Mouse septicemia.

Animals susceptible. Hog, rabbit, mouse, white rat, pigeon, and sparrow.

Animals immune. Horse, cow, ass, guinea pig, cat, chicken, and goose.

Pathogenesis. The organism has been isolated from poisonous meat. Inoculations produce a true bacteremia. Microscopic examination shows the organism to be present principally in the capillaries. The spleen is enlarged, but otherwise, the internal organs show no characteristic lesions.

Poisonous properties. Unknown.

Immunity. Vaccination with killed or attenuated cultures, immunizes, but the serum of immunized animals possesses little curative power.

Bacillus pathicus, Nocard

Disease produced. Epidemic pneumonia contracted from diseased parrots.

Animals infected naturally. Parrots and man.

Animals susceptible. White and gray mouse, pigeon, rabbit, and guinea pig.

Animals immune. The dog is partially immune.

Pathogenesis. The disease is a true bacteremia, being associated in man with pneumonia.

Poisonous properties. Not known.

Immunity.

Bacterium influenzae, Lehman & Neumann

Disease produced. Influenza in man.

Animals susceptible. Rabbit, and guinea pig.

Animals immune. Most of the other laboratory animals.

Pathogenesis. Produces purulent bronchitis, and pneumonia. Sometimes there is metastatic infection of other organs producing diseases such as endocarditis.

Poisonous properties. The toxin is intracellular, probably an enzyme.

Immunity. Vaccination and inoculation do not confer a lasting immunity, in fact, infection in many cases tends to predispose to the disease.

Bacterium canceri, Kruse

Disease produced. Chancroid, or soft chancre.

Animals infected. Man.

Animals susceptible. Some of the monkeys.

Animals immune. Other laboratory animals.

Pathogenesis. Disease produced appears first as a small red papule which becomes larger, and ulcerates. The inguinal and other lymph nodes enlarge and ulcerate. Primary infection most frequently upon the genitalia, other tissues not frequently involved.

Poisonous properties. Not known.

Immunity. Acquired.

Bacterium pneumoniae, Zopf

Disease produced. Pneumonia.

Animals infected. Man.

Animals susceptible. Guinea pig, and rabbit.

Pathogenesis. The organism, probably an avirulent type, is found in normal saliva. Under some conditions, either alone or with other organisms, produces pneumonia or metastatic endocarditis, otitis media, and tonsillitis.

Poisonous properties.

Immunity Not permanent, probably opsonic in nature

Bacterium cholerae, Kill

Disease produced. Chicken cholera, and rabbit septicemia

Animals infected. Chicken, pigeon, goose, duck, rabbit, and mouse.

Pathogenesis. Generalized septicemia with minute hemorrhages, peritonitis, or the formation of diphtheritic areas.

Poisonous properties. Not known

Immunity Agglutinins well developed and immunization with killed or attenuated cultures induces the formation of bacterolymins and probably opsonins. In cultural and pathogenic character, this organism is closely related to the four following, all being classed under the general group of organisms producing diseases known as pasteurellosis.

Bacterium suis, Migula

Animals infected. Hog.

Animals susceptible. Rabbit, guinea pig, and less so chicken and pigeon

See preceding.

Bacterium sanguinum, Moore

Disease produced. Infectious leukemia.

Animals infected. Chicken. See preceding.

Bacterium bovisepiticum, Kruse

Disease produced. Hemorrhagic septicemia

Animals infected. Cattle. See preceding

Bacterium avium, Moore

Disease produced. Roup.

Animals infected. Chicken.

Bacterium asthenes, Dawson

Disease produced. Asthenia or going weak

Animals infected. Chicken.

Animals susceptible. Guinea pigs, and albits

Pathogenesis. ?

Poisonous properties. Not known.

Immunity ?

Bacterium anthracis, Migula

Disease produced. Anthrax or splenic fever

Animals infected. Mouse, guinea pig, rabbit, sheep, cattle, man.

Animals immune. Carnivora.

Pathogenesis. Oedema at point of inoculation, spleen very much enlarged, pulpy, internal organs generally hyperæmic. The organism is to be found in

the blood in all parts of the body. Acute degenerative in parenchymatous organs.

Poisonous properties. Neither a soluble toxin nor an antigen has been demonstrated though there is abundant evidence in the tissues of intense intoxication.

Immunity. Immunity may be developed by the inoculation with cultures of the organism grown at high temperatures. The blood serum of animals thus immunized possesses some immunizing power. *Spores* are probably important




Fig. 162. A culture of *Claviceps purpurea* growing on bread. (From the collection in the University of Michigan Herbarium.)



Fig. 163. Spores of *Claviceps purpurea* in a culture medium. (From the collection in the University of Michigan Herbarium.)

Bacterium necrophorum, Flugge

Disease produced. Necrosis in various organs.

Animals infected. Calves, lambs, cattle, sheep, goats, antelope, reindeer, red deer, roe, horses, asses, hogs, kangaroos, rabbits, dogs, chickens, kite, guinea pigs, and on experiment mice and pigeons.

Pathogenesis. The local lesion is sharply marked off usually yellowish, or of a dull brown, of a yeasty consistency, and having a characteristic odor of



Fig. 301. *Bacillus* of Splenic Fever. *Bacillus anthracis*. Spore formation and spore germination. A From the spleen of a mouse after 24 hours culture in nutrient solution. B Germination of spores x 650. C. The same x 1650. After Koch.

old cheese and glue. Often produces a diphtheritic false membrane. It has been described as producing the following in various animals. Necrotic dermatitis, sheep-pox, abscesses in rabbits, necrobacillosis of the hoof, necrosis in digestive tract, stomatitis, vaginitis, metritis, foot-rot of cattle and sheep, necrotic omphalophlebitis in young animals, jointill, multiple necrosis in the liver, lungs, and avian diphtheria, etc.

Disease produced. Diphtheria.

Poisonous properties.

Immunity.

Bacterium diphtheriae, (Flügge)

Animals infected. Man.

Animals susceptible. Horse, guinea pig, rabbit, mouse.

Pathogenesis. Local and general phenomena caused by a soluble toxin and necrosis of mucous surfaces, and the underlying tissues, and false membrane often forms consisting of fibrin, leucocytes, etc. There is a mild leucocytosis, probably due in most cases to the co-ordinate activity of streptococci.

Poisonous properties. A true toxin is produced in the body and when grown in artificial media. It is this toxin which causes the characteristic clinical picture in diphtheria.

Immunity. Injection of non-fatal doses of diphtheria toxin results in the production of an abundance of anti-toxin in the blood. This blood serum has powerful curative and prophylactic properties, and has obtained wide usage in general practice.

Bacterium tuberculosis, (Koch)

Disease produced. Tuberculosis, consumption.

Animals infected. Probably no animal is immune to all of the varieties of this organism.

Pathogenesis. The disease generally runs a chronic course, affecting practically all of the organs in the body. In man, characteristic lesions are to be found in the lungs, in cattle generally on the peritoneum. The disease is characterized by the formation of numbers of nodules in the tissues, varying in size from a pin-point to that of an egg. At first, these are hard, and firm throughout, but as they grow larger generally caseate. Microscopically, these tubercles are characterized by the presence in the center of the so-called giant cells with numerous nuclei surrounded by concentric rings of epithelioid and lymphoid cells.



Fig. 30g. Tubercle organism, *Bacterium tuberculosis*. After Möller and Waehburn.

Poisonous properties. Not definitely or thoroughly understood, but probably an endotoxin. The killed bodies of the bacteria or the product of their growth in media when injected into animals affected by tuberculosis, cause a characteristic temperature reaction. The material thus injected into animals for diagnostic purposes, is called tuberculin, and is in general used in veterinary practice.

Immunity. Agglutinins are produced in infected animals. Recovery from the disease occurs in many cases doubtless from the waning in of the bacteria. Artificial immunity may be experimentally produced by increasing the opsonic content of the blood, but as yet no accepted method of immunization has been developed.

Bacterium mallei, (Löffler) Migula

Disease produced. Glanders, and farcy glanders.

Animals infected. Usually the horse and ass.

Animals susceptible. Man, goat, cat, hog, field mouse, wood mouse, rabbit, guinea pig, hedge hog.

Animals immune. Cow, house mouse, white mouse, and rat.

Pathogenesis. The lesions usually appear in the horse and ass in the form of discrete, sharply marked ulcers upon the mucous membrane of the nose. The ulcers once formed generally remain open and continue to discharge pus. The lymphatic glands are infected, and generally the submaxillary glands enlarge and ulcerate. Infection of the lungs often occurs through inhalation. Another type of the disease is characterized by infection of the sub-cutaneous lymph channels, which become enlarged and ulcerate, breaking through the skin at various points.

Poisonous properties. This organism produces some poisonous principle similar in a sense to that found in tuberculin and under the name of mallein is used in the diagnosis.

Immunity. Successful immunization of animals against glanders has not been accomplished.

Bacterium leprae, (Hansen) Lehmann Neumann

Disease produced. Leprosy

Animals infected. Man

Animals immune. Laboratory animals.

Pathogenesis. The organism is found present in leprosy tissues, almost filling the cells in many instances. Several types of leprosy are differentiated




Fig. 30b. *Mycobacterium leprae*. a. Section through "glanderous" tissue the small rods massed or single. b. Bact. stained with methylene blue. After Flagge

on the basis of the organ or tissues in the body infected. In many of its aspects, the disease resembles tuberculosis.

Poisonous properties. The organism has not been successfully cultivated. Poisons are unknown.

Immunity.

Bacterium pestis, Lehmann-Neumann

Disease produced. Bubonic plague.

Animals infected. Man.

Animals susceptible. Rat, guinea pig, monkey.

Pathogenesis. Infection usually cutaneous. The lymph gland becomes swollen, and hemorrhagic and undergo more or less extensive necrosis, generalized septicemia in many cases, pneumonia and hemorrhages in various mucous membranes especially in the stomach and endothelial surfaces, such as the pericardium and in various parenchymatous organs with extreme degeneration of the latter. Spleen swollen.

Poisonous properties. The toxic substance is obscure. The filtrates from young cultures usually show little, or no toxicity, the older, more. The toxic substance seems to be susceptible to heat and is present in cultures killed by chloroform.

Immunity. Immunity may be conferred by the injection of sterilized or attenuated cultures, and this vaccination is practiced in some Asiatic countries.

Microspira comma, Schroeter

Disease produced. Asiatic cholera.

Animals infected. Man.

Animals susceptible. Laboratory animals naturally are immune to the disease, but by preventing peristalsis, and neutralizing the acid gastric juice some experimenters have succeeded in producing the disease. The guinea pig is susceptible to intraperitoneal inoculation.




Fig. 101. Cholera organism. *Microspira comma*. From the margin of a drop of broth containing pure culture of the *Spirillum*. a. Long spiral filaments. All $\times 600$. After Flügge.

Pathogenesis. It is essentially an intestinal disease, the organism living within the intestines. The characteristic lesions are produced through the absorption of the poisonous matters there produced. Large and small intestines deeply congested, diarrhoea, Peyer's patches and glands swollen, eventually mucosal necrosis in part. Ulcers eventually form, though perforations are rare. The parenchymatous organs show marked signs of degeneration, the vascular system, the nervous and respiratory systems show no characteristic lesions.

Poisonous properties. The essential poison is intracellular, undoubtedly an endotoxin. It is found in the filtrate of old cultures and in solution of bacterial cells.

Immunity. Produced by considerable quantities of bacteriolysins and probably opsonins. Agglutinins are also produced. Vaccination with killed or attenuated cultures has proven fairly successful, but not the use of the blood serum of immunized individuals as a curative or a prophylactic agent.

SPIRILLACEAE

Spirochaeta pallida

Disease produced. Syphilis.

Animals infected. Man and ape.

Animals immune. Other animals.

Pathogenesis. Produces primary lesions in form of ulcers at the point of inoculation, second as gummata in the parenchymatous organs.

Poisonous properties. Not known.

Immunity. Not well understood.

Spirochaeta anserina, Sakharoff

Disease produced. Goose septicemia.

Animals infected. Goose

Animals susceptible

Pathogenesis. Producing septicemia

Immunity

Spirochaeta Obermeieri, Cohn

Disease produced. Relapsing fever

Animals infected. Man.

Animals immune. Mouse, rabbit, sheep, and hog.

Pathogenesis. Produces relapsing fever in man.

Immunity

CHLAMYDOBACTERIACEAE

Cladothrix botry (Bollinger)

The mass consists of several distinct zones of different elements: the central portion granular with small round bodies radiating out from this angled mass of thread-like bodies; the outer portion consisting of conspicuous club-shaped colonies. The organism is quite polymorphic. In cultures the threads are from 3-10 to 5-10 μ in thickness with flask shaped or bottle like expansions. The organism may be grown upon all the artificial media. The colonies appear as small gray dots with translucent, radiating filaments. If kept for a few days at 37° C they are opaque and nodular; later they show a whitish downy appearance. In blood serum the nodules are yellowish or bluish-red in color; on agar agar the color becomes brownish with age; on potato, reddish yellow and the white down makes its appearance early.

Distribution. Widely distributed both in Europe and North America.

Pathogenic properties. The organism was discovered by Langenbeck in 1845, but was not described until 1878 by Bollinger. Israel in 1874-78 called attention to the disease in man, and Rostrom in 1899 made a careful study of the disease. The disease is not common in man but cases have been described by Murphy and Ochsner and Senn of Chicago. Two of the cases described by Murphy began with tooth ache and swelling of the jaw.

The disease may be caused by direct inoculation of pus, but there is good reason to believe that not an infrequent source of infection is by means of barley and other grains. There is reason to believe that it occurs in nature as a saprophyte. The history of many cases reported in man seem to indicate




Fig. 30., Lumpy Jaw *Cladothrix botry* showing the radiating masses. At the left the club-shaped bodies and branches (After Fournier). At the right, one of the radiating masses less magnified. (After Fournier).

this kind of infection and it is a well-known fact that certain fields are the source of infection. Dr McFarland says

When inhaled, the organisms enter the deeper portions of the lung and cause a suppurative or broncho-pneumonia with adhesive inflammation of the contiguous pleura. After the formation of the pleuritic adhesions the disease may penetrate the newly formed tissue, extending to the chestwall, and ultimately form external sinuses or it may penetrate the diaphragm and invade the abdominal organs, causing interesting and characteristic lesions in the liver and other large viscera.

Another allied disease is the Mycetozoa, or Madura foot (*Cladotrix maduree*), which is found in India, especially in the province of Sindh, but occurs also in other parts of Asia, in Europe and northern Africa, and a few cases have even been reported in North America.

Another *Cladotrix farcinica*, found in Guadeloupe country, is characterized by a superficial lymphangitis and lymphadenitis extending to the trachea and axillary glands. The glands enlarge, suppurate and discharge a pus. The internal organs have a pseudo-tubercular appearance.

The organism consists of long delicate filaments, characterized by distinct branching, the old cultures are rich in spores. The organism has been cultivated in the usual media.

It is pathogenic for guinea pigs, cattle and sheep. The culture is virulent for some time.

The papers by Norard on the farcinica organism and a paper by Mugrove, Clegg and Fick on streptothrix should be consulted.

SCHIZOPHYCEAE

Unicellular organisms common in fresh and salt water. They contain blue, blue-green, violet or reddish pigments; swarms, spores absent, are common in fresh and salt water. Simple in structure, existing as a single cell or as a chain held together by a gelatinous envelope, or in small colonies. Chlorophyll and other pigments not in definite bodies but distributed throughout the cell contents or else forming a sheath which lines the cell wall. Reproduction occurs by simple division, some forms produce spores which are thick walled thus enabling the organism to live over unfavorable conditions, after a period of rest these spores germinate and again reproduce in the vegetative way by fission or division. Some of the more common, more or less injurious types found in water are *Oscillatoria*, *Anabaena*, *Clathrocystis* and *Nostoc*. Cells contain phycocyanin and chlorophyll, the latter not visible because of the former.

CHROOCCACEAE

Cells spherical, singly or collected in colonies surrounded by a copious covering of mucilage forming gelatinous colonies of various sizes. The genus *Merismopedia* consists of flat rectangular colonies. Cell-division occurs in two directions. The genus *Gleocapsa* has spherical cells united into colonies, the cell with a thick colorless, brown yellow or violet coat. Some of the species are common in fresh water.

Clathrocystis. Henfrey

This alga occurs in colonies which are at first solid, but later become perforated. The colonies are held together by a gelatinous matrix. The cell contents are blue green, or rose-purple in color. The species most commonly found is the *C. aeruginosa*, occurring not only in Europe, but very widely scattered in




Fig. 3. Schizophyta—Schizophyceae. Blue Green Algae. 1. *Nostoc sphaeroides* $\times 400$. 2. *Gloeocapsa sanguinea* $\times 400$. 3. *Nostoc sphaeroides*. In A pair of cells is 4. *Chamaesiphon confervicola* $\times 400$ at the right, a caetidium; a: the left germinating arthrospores. 5. *Rivularia minutula* $\times 200$. 6. *Anabaena macrosperma* $\times 100$ at the right a caetidium at the left, germinating arthrospores. 7. *Placostema Tomassinianum* $\times 200$. 8. Filaments of *Lyngbya aestuarii* c—central body, ch—chromatin bodies, $\times 100$. 9. *Lyngbya aestuarii* $\times 150$ at the right filaments with hormogonia. 10. In a figures, ap—spores, h—heterocyst. Fig. 1, 2, 3, after Louke. Fig. 4, 5, 7 after Haugwitz. Fig. 6, 8, after Nadeau, Fig. 9a, 6, 9, after Wettstein.

North America, especially common in ponds and the plankton of lakes. Other species are *C. roseo-persicina* and *C. fuliginosa* the former being especially common in ponds and ditches which contain a great deal of decaying vegetable matter. The latter species is now generally referred to the genus *C. leospirastrum*, and the *C. aeruginosa* to the genus *Microcystis*.

OSCILLATORIACEAE

Cells in filaments, apical cells disc shaped with sheaths variable, sometimes wanting heterocysts absent, form hormogonia. Common representatives, *Oscillatoria* and *Lyngbya* which at times are common in fresh water.

Oscillatoria, Vauch.

The plant consists of more than one cell forming a simple filament held together by a common but stout gelatinous sheath, the cells being packed to-

gether like a row of lozenges. A few of the cells fall out of the sheath forming what is called a hormogonium. This starts a new filament which has a characteristic movement hence the name *Oscillatoria*. *Oscillatoria* is common in hot springs sometimes also covering damp soil in greenhouses. It is mainly through decomposition that these algae become noxious.

NOSTOCACEAE

Cells spherical in unbranched chains frequently torulose, sheaths gelatinous frequently forming jelly-like masses, reproduction by hormogonia and spores. Many species of the order are troublesome in water supplies. Some species of the genus *Nostoc* are used as food.

Nostoc, Vaucher

Colonies in flexuose chains united in definite gelatinous investment, cells usually spherical or ellipsoidal, heterocysts terminal or intercalary, spores spherical or oblong. These algae are very common in the lakes in southern Minnesota, Northern Iowa, and elsewhere in the United States. By decomposition *Nostoc* produces disagreeable products. Dr. Arthur, some time ago, found *Nostoc* in quantities in the lakes of southern Minnesota and at one time it was supposed one species produced poisoning of cattle. Dr. Arthur however, did not attribute the poisoning to this alga. One of the species of this genus frequently found is *Nostoc verrucosum* common in both the Old and New World.

Anabaena, Bory

Filaments straight or curved, surrounded by a thin sheath united to form a filiculent mass, heterocyst and spores intercalary. This alga also forms filaments which are free or united in a mass. In the filaments occur the vegetative cells, the heterocysts, whose function is not known, and a spore which serves to start the organism again. This organism causes much annoyance in water, not only in North America but in Europe. Dr. Farlow some years ago referred to its injurious properties. It has also been frequently mentioned by Parker as contaminating water supplies in Massachusetts, others have noticed it in New York, and Dr. Trelease has found it in Madison, Wisconsin. The latter writer says in speaking of the Waterblooms and other algae:

After a warm spring, on my return to Madison, June 26, 1887, I observed a considerable quantity of putrid scum on the shore of Fourth Lake, but the south wind scattered it before specimens of it could be obtained. The succeeding fortnight was hot and after a couple of calm days, succeeding a strong wind from the north-west, the southern half of the lake was filled with suspended particles about a millimeter in diameter. These consisted exclusively of *Anabaena husarii*, already in full fruit, the spores were the customary *Sphaerocyga* arrangement. A collection made June 20th.

This alga is common in many of our northern lakes and is a frequent pest in water reservoirs, producing pig pen odors and bad taste of water.

RYTULARIACEAE

Filiform filaments attenuated from base to apex, heterocysts basal or rarely absent, sheath tubular, gelatinous, or membranous.

Gloeotrichia, J. Ag.

Free floating colonies solid when young but inflated and hollow when old, the filaments radiating from the centre outwards.

Gloetrachia Pisum, (Ag.) Thur.

It forms small green spherical bodies about 1 millimeter in diameter, floating at various depths in the water. It consists of a mass of tapering threads arranged radially in the gelatinous matrix. The apices of the threads protrude more or less, giving it a bristly appearance. The base of each filament contains a heterocyst and above it a slender cylindrical spore and beyond it the ordinary vegetative threads of the alga.

According to Dr. Arthur it is common in Minnesota. He found it common in Waterville, Lake Minnetonka, Lake Phelan in Minnesota and East Okoboji Lake in Iowa. It was thought by the people of Waterville, Minnesota, that this alga caused the death of cattle which drank the water. The history of these cases is recorded by Dr. Arthur as follows:

"That some of the animals had drunk of the water and some a few hours only before they died was positively known, and that a' had done so seemed from circumstances quite probable. After the most careful examination the only plausible hypothesis that could be advanced in account for the death of the animals was that the alga present possessed some toxic or other baneful properties sufficiently powerful to kill a cow in a half hour or more after drinking freely of it. The well established reputation of all the algae for innocuousness made this hypothesis appear from the very first extremely improbable, but for want of the slightest hint in any other direction it was thought worth while to bear it in mind and to investigate the matter further.

About the middle of June 1884, word was received that eight cattle had died on the shore of Lake Tetonka. I at once started for Waterville, arriving on the twentieth and found the algae was abundant here in 1883 but still making the water green some fifty feet or more out from the shore toward which the wind had been blowing several hours. Although the conditions were not the most favorable yet it seemed best to attempt a direct experiment by giving the animals water charged with algae. After much delay the services of Prof. M. Stalker, state veterinarian of Iowa and professor of veterinary science in the Iowa Agricultural College, were secured to conduct the experiment. A horse and calf were employed. On June 30th, Prof. Stalker, with the assistance of Prof. Edward H. Porter of the university of Minnesota, and in the presence of a number of Waterville men the tests, the writer being unable to remain. The animals had not been permitted to drink for some twenty-four hours previous, and were consequently thirsty enough to take a large amount of water well charged with the algae. No bad results of any sort followed.

The thorough and able manner in which the test was made leaves no reasonable doubt of the perfect harmlessness of the algae in a growing condition. I append this last clause, because the citizens of the place still believe that the algae are at the root of the trouble, and that the test did not show it because they were not made at the right stage of their occurrence. Although no sufficient study of the habits of this plant has yet been made to enable one to speak with certainty yet it does not appear from present data that in some other stage it would give different results, unless it be when decaying, when it turns brown or reddish brown and gives off a peculiar stench. At this time the microscope shows the cells of the algae to be swarming with bacteria. Whether these are other than the common and harmless bacteria of putrefaction it is at present impossible to say. The probabilities are, however, entirely against the hypothesis that the decaying algae or the accompanying bacteria have anything to do with the trouble.

We are therefore obliged to sum up the economic part of this investigation by stating that the death of the animals is probably not due to the suspected algae, and that no clue to the real cause has yet been obtained."

Dr. Arthur in a recent communication states that he has had no evidence so far that these algae are poisonous. That the death of these animals was probably due to bacteria found in the marshes.

OTHER BLUE GREEN ALGAE

Quite a number of additional genera are known to occur in our fresh waters, among them the *Gloeoopsis* with cells single or in groups surrounded

by a gelatinous envelope, cell contents bluish green, brownish or reddish, *Marismopodia*, with division in two directions, cells arranged in tabular groups of 4, 8, 16, 32, 64, 128, etc.; *Lyngbya*, with filaments enclosed singly in a sheath, branchless or occasionally branched, cell contents blue and granular

FLAGELLATAE

One-celled organisms with nucleus, sharply differentiated protoplasmic body, some with a simple membrane, pseudopodia absent. motile during most of their existence, cilia 1 or more and with 1 or 2 pulsating vacuoles, chromatophores occasionally absent, reproduction asexual by longitudinal division. The *Euglenales* contain the family *Euglenaceae*. The most common of these is the *Euglena viridis* which occurs in stagnant pools. Cells are elongated spindle-shaped, cilia 1 and with a red eye spot at one end. Water where these are abundant is not wholesome. The *Euglena* forms small sphaeroidal nearly colorless colonies, the central portion of the colony is a hollow space filled with mucilage and the ciliated cells are arranged around the periphery, vegetative multiplication occurs by simple fission and by zoogonidia. The *Euglena*, when occurring in large quantities produce a fishy, oily odor. The related *Synura* produces an odor of ripe cucumbers with a bitter and spicy taste.

EUPHYCEAE

Plants mostly occurring in water, always with a cell-membrane and nucleus. Green or other colors mixed with the green (brown or red). This includes all of the algae or thallophytes that contain chlorophyll which is, however, frequently masked because of other pigments like red and brown.

PERIDINIALES

Small 1-celled organisms of brown color, the chlorophyll masked by diatomin found mostly in the plankton of salt water.

BACILLARIALES

Small 1-celled organisms of brown color, the chlorophyll masked by diatomin of the chromatophores, cell-wall consisting of silica with a girdle and fine lines, reproduction asexual parallel to the long axis of the organism and the formation of auxospores and sexual auxospores by conjugation.

The diatoms are especially important in considering water supplies since they are widely distributed and at times very common. The diatom is like a pill box, made up of two parts, one fitting tightly within the other, the walls are strongly silicified and marked with fine lines, the cell contents are colored brown. Economically, these algae are of some importance as food for fishes, in manufacture of dynamite, and for polishing.

Dr. Moore, speaking of Diatoms in water says

There are only a few species which are known to give rise to serious trouble in water supplies, but these occur quite frequently and in great quantities. Sometimes the infected water has an odor variously described as resembling fish or geraniums, and the taste is disagreeable enough to render it quite unfit for use. This condition is often produced by *Asterionella*. In addition to this effect, however, diatoms are extremely troublesome when contained in water to be used for the manufacture of paper or for laundry purposes, because

of the greenish-brown coloring matter they contain, which stains articles coming in contact with it. Whipple has observed that the growth of diatoms seems to depend upon certain definite conditions of the water—that is, they do not develop when the bottom of the pond or reservoir is quiet but in spring and fall, when the rising or lowering temperature causes the water to circulate and a good supply of air and nitrates is obtained, the growth is most luxuriant. Thus, it is seen that temperature is only an indirect cause, and not one that need be taken into account by itself.




Fig. 32. Diatoms. *Pinnularia viridis*. 1 View showing markings in wall. c—central nodule p—polar nodules r—raphe 2 Side view of single cell with raphe 3 In process of division. Fig. 1 and 2 after Pfäfer; Fig. 3 after Wettstein.

CONJUGATAE

Chlorophyll green algae, membrane without silica reproduction, swarm cells absent sexual reproduction by forming zygospores through the union of aplanogametes.

DESMIDIACEAE

The desmids are green unicellular organisms represented by such genera as *Cosmarium* and are found mostly with other algae. They are not especially troublesome. Common genera are *Closterium*, *Cosmarium* and *Desmidium*.

The filaments of *Zygnema* are small consisting of a single series of cylindrical cells placed end to end, occasionally with a slight constriction at the points of junction. Each cell has two star-shaped chloroplasts, each one containing a large pyrenoid.

Spirogyra, Link

Spirogyra is a common alga everywhere in our fresh waters, especially in quiet waters as in ponds and ditches. The filaments are simple, occur in bright green masses, often several feet long. Cells cylindrical, variable in diameter and relative length. Wall smooth and slightly gelatinous, chlorophyll arranged in




Fig. 12a. Lower figure. two views of diatom in lateral view / upper view. Upper figure. a desmidean *Closterium*, chl—chloroplastid. Charlotte M. King.

1 or more spiral bands, depending upon the species, nucleus is generally situated in the central portion of the cell; in the chlorophyll band occur what are known as the pyrenoids, reproduction both asexual, by the simple cutting off of the end cell, and sexual, by conjugation through the union of two cells.

ZYGNEMATACEAE

Cells cylindrical, unbranched forming threads, chromatophores present, in masses or spiral bands. Formation of zygospores.

CHLOROPHYCEAE

Chlorophyll-green plants, occurring singly or in colonies, (threads or flattened bodies) with one or more nuclei, reproduction asexual by producing multilateral zoospores and aplanospores, sexual by the copulation of zoogametes, or spermatozoous and oospheres, the spores produce a new plant directly or generally form swarm spores. Water net (*Hydrodictyon reticulatum*) *Pediastrum* and *Scenedesmus* are common in fresh water, *Pleurococcus* is common on trunks of trees. The *Conserveles* contain the sea lettuce, *Ulva latissima*, which is used as food, the *Conserve* with slender green filaments common in fresh water, the *Cladophora fraxia* in fresh water. *Cladophora* is a common alga and is quite rough to the touch, and may therefore easily be distinguished from *Spiragira* or *Zygnema*. The frond of *Cladophora* is branched, with many-




Fig. 33 Fresh Water Green Algae. Conjugatae. Left hand figure 1—Zygospore. 2—Zygospore. Right hand figure 3—Zygospore. 4—Zygospore. 1, 2, 3 and 4, in different stages of conjugation. 4 Zygospore.




Fig. 33a Botrydium granulosum. Green Scum. 1. A single large zoospore. 2. Rhizoid process.

nucleated cells or in some instances, only 1 or 2, chloroplasts occur on the margins with a single pyrenoid in each piece of the reticulum. The *Siphonaceae* contain the *Vaucheria*, found in damp ground, the long unsegmented threads of which produce antheridia and oogonia, and the *Botrydium granulosum* found on damp ground.

VOLVOACEAE

Unicellular organisms or forming colonies, each cell with a single chromatophore, forms gametospores and oospores.

Pandorina, Bory

The algae of this genus are collected together in spherical or subspherical colonies known as caenobia. Each caenobium contains about 16 cells closely packed within a gelatinous envelope, the cells are pyramidal in shape and reach almost to the center of the spherical colony; each cell produces 2 spores.

Volvox, (L.) Ehrenb.

Volvox consists also of globose colonies known as caenobia, each consisting of a large number of small cells from 200 to 22000, arranged in a single layer within a gelatinous sheath, the caenobium is a hollow sphere the cells being connected by protoplasmic threads of varying stoutness, each cell has a distinct chloroplast, 2 or more contractile vacuoles and a number of cilia. Reproduction occurs through asexual methods or by fertilization; in fertilization, the sperm




Fig. 34. Chlorophyceae. Pinn. Scum. *Ponderosa Marum*. a. Floating caenobium. b and c. Zoospores. d. Zoospores conjugating. e. Process of conjugation complete. f. Zygospore. After Lurich.




Fig. 35. Bigdee Wrack. *Fucus vesiculosus*. a. Section through conceptacle containing oogonia. b. Single oogonium with egg cells. c. Egg cells escaping. d. Antheridium with sperm cells. e. Egg cells in process of fertilization. f. Germination of spores, stained brown. g. Single sperm cells. After Thuret.

cell coming from the antheridium, unites with the egg cell which is contained in the oogonium, the caenobium because of the cilia has a rolling motion. Several species are common like *V. globator* and *V. minor*.

Eudorina, Ehrenb

The caenobium is globose or subglobose rarely ellipsoid and normally consists of 32 cells arranged within the periphery of a copious mucilaginous mass each cell contains 1 or more pyrenoids, reproduction takes place as in *Pandorina*. *Eudorina* like *Pandorina* produces a faintly fishy odor.

CHARALES

Plants of brackish or fresh water, consisting of internodes short whorls of cylindrical branches cells nucleated, growth from an apical cell, asexual reproduction by means of bulbils and vegetative threads sexual reproduction by egg-cells and spermatozooids, the latter spirally coiled in the cells of the



Fig. 36. Bladderwort *Equisetum veruculosa*. Air spaces shown in light areas (1) conceptacles (2) containing reproductive bodies. After Loewen.

antheridium, the egg cell is contained in a spirally coiled oogonium and after fertilization becomes an oospore. The Stoneworts or Charas are common in brackish water, and though not injurious frequently stop up canals and fill ponds so that it becomes necessary to pull them out.

PHAEOPHYCEAE

Brown algae, chlorophyll marked by a brown coloring matter, *phycophaein*, reproduction sexual and asexual, swarm spores, sperm cells and egg cells marine; tetragonidia absent.

The Phaeosporae contain the *Laminariaceae*, the Devil's Apron, *Laminaria digitata*, and other species from which iodine and mannite are derived. The *Macrocystis pyrifera* is of great length. The Clycosporae contain the family *Fucaceae*, the common Bladderwrack (*Fucus vesiculosus*) from which iodine, bromine and soda are obtained. The Sargasso weed (*Sargassum bacciferum*) found in the Atlantic ocean is also abundant in the Sargasso Sea.

DICTYOTALES

Brown algae reproductive bodies without motion, tetragonidia present. This group contains a single order Dictyotaceae comprising a few genera.

RHODOPHYCEAE

Red or violet algae, chromatophores contain chlorophyll and red coloring matter (phycoerythrin and rhodophyll); reproduction sexual and asexual; mostly marine. The red sea weeds are divided into several classes and numerous orders. The sub-class *Florideae* contains most of the species. Food is obtained from several species and the carrageen is furnished by *Chondrus crispus*, agar agar is obtained from *Gracilaria lichenoides* found in the Indian Ocean. The *Gloiopeltis coliformis* and other species are used by the Japanese as food. Many of the species are pretty and are much gathered on the sea coast.




Fig. 37. Red Sea Weed, *Nemalion multifidum*. 1 Branch with carpogonium and antheridium. 2-4. Different stages of development. 5 *Leptolus mediterraneus* with antheridium, carpogonium and spores. a—antheridia, c and o—carpogonia, t—trichogyne, s—sperm cells, e—spores, f—fruit. After Thuret and Bornet.

EUMYCETES

Parasitic or saprophytic plants with one or more cells, chlorophyll absent with apical growth, mycelium usually evident, reproduction sexual and asexual, generally the latter, asexual by the formation of zoospores, conidia or spores

PHYCOMYCETES

Thallus generally of a single branched tubular thread, septa in connection with the reproductive bodies only, threads containing many nuclei, reproduction sexual and asexual, in the latter the spores generally in sporangia (*Mucor*), conidia in chains (*Aibugo*), or at the end of the hyphae (*Plasmopara*), reproduction sexual by copulation forming zygospores (*Mucor*) or oospores in *Plasmopara* and *Aibugo*.

ZYCOMYCETES

Parasites or saprophytes mycelium branched but septate or septa in connection with the formation of the reproductive bodies, reproduction sexual by endospores, microconidia, or chlamydospores. A group of fungi represented by the Fly Fungus (*Empusa*) and Common Black Mould (*Mucor*)

MUCORACEAE

Sporangia with columella, many spored, zygospores between the threads of the mycelium. Few species have the two sexes united on the same plant generally they are on the separate individuals. According to Blakeslee *Sporodina* contains both sexes (homosporangic, homosporic, homophytic and homothallic) *Phycomyces* is dioecious, the zygospores producing at germination but one kind of germ tube which gives rise to a sporangium containing both male and female spores, (homosporangic heterosporic, homophytic, heterothallic) *Mucor mucedo* has sexes separated on different individuals but two different kinds of germ tubes are formed by the germination of the zygospores, (heterosporangic, heterosporic, heterophytic, and heterothallic) *Zygorrhynchus* is heterogamic. The same author* has recently reviewed the literature.


About 85 species widely distributed. The *Phycomyces* was first found in oil kettles, and not infrequently in oil cakes. *Sporodina* are parasitic on larger fungi. *Pilobolus crystallinus* is common on horse manure, the conidiophore being enlarged. The sporangia look like "fly specks" on the wall. This fungus is not injurious.

Mucor (Mickah) Link. Mucor Mould

Mycelium creeping, conidiophores simple or branched, sporangia spherica or pear-shaped, columella well developed, wall of sporangium mucilaginous, in some cases chlamydospores, or forming small chains or "cysts", zygospores produced by the fertilization of two gametes.

A genus of wide distribution of 50 species. The life history of a common species, the *Mucor stolonifer* (*Rhizopus nigricans*) found on bread and decayed fruits is as follows. The gray felted mycelium spreads through the substratum, and on the surface small black bodies, the sporangia, are produced. The conidiophore arises from the felted mycelium and bears an enlarged spherical head, the sporangium, within which, occur the spores. On adding water

* Bot. Gazette 1909 418. Reprint.



to the specimen, the wall of the sporangium collapses and the end of the stalk, known as the columella, turns back, giving it something of the appearance of an umbrella. The columella, before it collapses, projects into the sporangium




Fig. 40. *Mucor* Runum. Conidiophore. 2. Gemmae. 1. Chlamydo-spores. All greatly magnified. 1 after Vuillemin. 2 after Calmette. 3 after Wehmer. (Modified by Charlotte M. Kang).

The spores germinate readily when placed in a moist atmosphere. In addition to the production of a sporangium a stalk may bend over and cause the further extension of the fungus by producing what is known as a stolon.

In some species small, round, or elongated spores are produced in the mycelium which are known as chlamydo-spores, and spread the fungus. In addition to the formation of spores in the sporangium, zygospores are produced in some species, two threads of the mycelium lie in proximity and nearly parallel, each produces a tube these meet, the walls are absorbed, and, just back of the meeting point, a cell is cut off. The contents from the old cells pass into the newly formed cell. We also observe that the cell of one arm is somewhat smaller than that of the other. This spore is a resting spore or zygospore. It lies dormant for a period, then germinates by forming directly a conidiophore with its sporangium containing the spores.

Prof. Blakelee has shown with reference to the fertilization of some of the species of *Mucor* that it requires a male and a female plant. In speaking of *Mucor mucedo* he says:

Mucor mucedo has the sexes separated on different individuals as in *Phycomyces*. But two different kinds of germ tubes are formed by the germination of its zygospores. While some

germ tubes are male and produce only male spores, others are female and produce only female spores in the germ sporangium. The sporophyte as well as the gametophyte, therefore is unisexual.

"Raggi," used in the manufacture of Arrack, contains *Mucor Oryzae* (*Rhizopus*), which transforms rice starch into dextrose, the latter being then fermented by yeasts forms blackish brown sporangia and has a pear-shaped columella. *Mucor Romanii* of Calmette is commonly grown in China where it is found on rice husks and is made from these into Chinese yeast. It changes rice starch into sugar and has been used to some extent for manufacture of alcoholic drinks.

M. racemosus, common in decaying fruit, produces alcoholic fermentation. *Mucor fungus* is parasitic on species of *Collybia*, *M. Methylotrophicus* was found




Fig. 41. Common Black Mould (*Mucor stolonifer* or *Rhizopus nigricans*). 1 Sporangium and method of spreading by stamens. 2. Same showing rhizoids, conidiophore, columella, sporangium and spore. 3. Zygospore, showing method of conjugation. 4. Zygospore germinating, h, conidiophore.

in the stomachs of bees, *M. nigricans* was found by Neumann (1892) and later by Artanet (1893) in the eye of poultry but Barthelet does not consider it pathogenic.

Mucor corymbifer, F. Cohn.

Delicate, white mycelium spreading over the surface of the substratum conidiophores appressed, spreading branched sporangia in umbellate clusters, the lower sporangia smaller than the upper the latter 70 μ diameter, wall




Fig. 42 *Mucor corymbifer* after the bursting of the sporangium. After Lichthe m.

colorless, smooth, collapsing, spores colorless, small $2 \times 3 \mu$, columella club-shaped, brownish frequently papillate.

Distribution. Probably tropical, found in tropical drugs, in Europe and the United States.

Pathogenic properties. Lichtheim recognized this species as pathogenic. It grows better at a temperature of blood. When introduced into the circulation of guinea pigs, it produces death in 48 to 72 hours. Mycelium is found in kidneys, spleen and Peyer's patches of intestines which are swollen and ulcerated. Huckel found the organism in the human ear. Dr. Wolffner in Dr. Trelease's laboratory in St. Louis, Mo., cultivated the organism from the human eye. The clinical record of this case was as follows. A farmer near St. Louis was cutting corn with an old fashioned corn knife. A small piece of corn stalk flew into his eye, later inflammation set in followed by inability to see. Dr. Wolffner found a film over the surface. This film was removed and later *Mucor corymbifer* developed from it. It has been frequently found in ulcerated portions of the lungs, intestines, nasal cavity, and in the auditory canal. Dogs are immune.

The earliest recorded case of mucor in pneumomycosis was made by Fürbringer, who had under observation three cases of a disease in two of which he found a *Mucor*. According to Dr. H. C. Plaut, the cases of otomycosis are not infrequent in India. According to Siebenmann it occurs in 5.1 percent in all diseases of the ear and males, especially farmers and gardeners, are more predisposed than females. According to Hatch and Row, ear mycosis is common in India, they having observed 22 cases in one month. The most common fungi found in the ear are *Verticillium graphis*, *Aspergillus fumigatus*, *A. niger*, and *A. flavus*. *A. nidulans* is somewhat rare, as is *Mucor septatus*. The *Aspergilli* will be treated more at length in another connection.




Fig. 43. Black Mould, *Mucor rhaizopodiformis*. Branched conidiophores, large columella and discharged spores around the same. After Lichtheim.

Mucor Trichus, Locet and Costantin

This differs from *M. corymbifer* in a few characteristics of sufficient importance to cause Locet and Costantin to consider it a distinct species, *M. Trichus* having larger spores which are $4\ \mu$ in diameter and sporangia $35\ \mu$ in diameter. It was isolated from epidermal scabs appearing on a horse affected with tinea, produced by *Trichophyton minimum*. The *M. Regnierii* described by the same authors is similar to the preceding.

Mucor rhaizopodiformis, F. Cohn

Mycelium at first snow-white then gray, conidiophore single or clustered, brownish, $125\ \mu$ long, small rhizoid processes, columella broad, constricted at the base, sporangia spherical, at maturity blackish, spores spherical colorless, $5-6\ \mu$ in diameter. Closely allied to *M. stolonifer* or *M. unequalis*.

Distribution. Not uncommon on bread in Europe.

Pathogenic properties. Pathogenic like the preceding. When introduced into the circulation of guinea pigs it produces inflammation and the tissues of the spleen, liver and intestines are found to contain the mycelium of the fungus. The animal becomes inactive, lies on its side and drops its head. Small masses of the mycelium may be found in the kidneys.

Mucor pusillus, Lindt.

Mycelium spreading, with numerous chlamydospores which are capable of germination, conidiophores generally branched, sporangia spherical, brownish, $30-40\ \mu$ in diameter, spores ellipsoidal or spherical, $5-8\ \mu$ long, $3-5\ \mu$ in diameter, columella pear-shaped, zygospores seldom produced, spherical $70-84\ \mu$ in diameter, roughened, chlamydospores abundantly produced, species capable of changing cane sugar into invert sugar, producing the ferment invertase.

Distribution, and hosts. Widely distributed in both Europe and North America, occurring on various decaying objects.




Fig. 44. *Mucor racemosus*. Fragment of mycelium, which has undergone conversion into chains of oidia $\times 120$. After Beefeld.




Fig. 45. Mould, *Mucor racemosus*. 1 Branched sporophore of undischarged sporangia $\times 80$. 2 Optical section of sporangium $\times 368$. After Beefeld.

Pathogenic properties. Said to be pathogenic for various birds but Pierre Savouré, after some extensive experiments thinks that it plays no part in disease. It was not pathogenic for rabbits and guinea pigs. Bollinger states that it occurs in the respiratory tract of birds where it produces mucormycosis. It has been observed in cutaneous lesions in cavalry horses in France, although culture did not yield this fungus but yielded a trichophyte instead.

Mucor ramosus, Lindt.

Mycelium spreading in the substratum, small, branched, at first white then becoming grayish white, sporangia black, spherical with marginal spines $60-90 \mu$ in diameter; columella ovate, light brown, 50μ wide, spores somewhat spherical, $3-3.5 \mu$ in diameter, colorless.

Distribution and Habitat. Found in Europe, not abundant.

Pathogenic properties. Pathogenic for birds. Grows only at a temperature of blood, minimum $20-25^{\circ} \text{C}$., maximum $50-58^{\circ}$, optimum 45° .

Mucor ramosus, Lindt.

Mycelium branching, abundant in the substratum and superficial conidiophores $5-15 \mu$ wide, sporangia blackish, membrane but slightly colored 70μ in diameter, columella rounded at the end or blunt, spores colorless, with delicate membrane, smooth, 3 to $4 \mu \times 5-6 \mu$. This resembles *M. corymbifer* except in the character of the spores.

Pathogenic properties. Pathogenic for guinea pigs, death occurring in 30 to 36 hours.

Mucor septatus, Siebermann.

Mycelium at first white, later grayish, sporangia light yellowish brown, sporangia small, columella colorless, conidiophore branched spores small $2.5\ \mu$ in diameter

Distribution and Habitat. Found in Europe

Pathogenic properties. Pathogenic for human beings found in the ear

Mucor equinus (Constantin and Lucet), Pammel

Mycelium branched, at first white or whitish, floccose with simple pedicels without rhizoid processes, erect or suberect, becoming fascicled, hyphae $8-12\ \mu$ in diameter, columella spherical or subspherical $20-50\ \mu$ in diameter, spore roundish or slightly angular smooth $4\ \mu$ in diameter, chlamydospores numerous especially at blood temperatures.

Distribution. First found in Europe

Pathogenic properties. Found in horses. Guinea pigs and rabbits inoculated peritoneally die on the 5th or 6th day

Mucor pusillus (Lucet and Constantin), Pammel

Mycelium spreading, branched, brownish fawn color producing stolons and rhizoids, sporangia-bearing peduncles branched, conidiophores 12 to $14\ \mu$ wide $1-2$ cm. long columella ovoid pyriform slightly brownish $7-30\ \mu$ high, sporangia 8 to $37\ \mu$, lateral sporangia similar but smaller. Grows readily in nutrient media. The rhizoids sink into the substance, the simple conidiophores rise from the rhizoids. Lucet and Constantin placed this species in a new genus *Rhizomucor*.

Pathogenic properties. It is essentially parasitic and was isolated from




Fig. 46. *Empusa sphaceliperma*. 1. Larva of Cabbage Butterfly. 2. Sectional view. 3. Conidiophores and conidia $\times 300$. 4. Conidium $\times 600$. 5. Mycelium with zygosporangia $\times 350$. 6. Single zygosporangium $\times 600$. 7.2. *Conidiobolus strictus*, found on the gelatinous fungi like the Jew's Ear (*Auricularia*). 7. Conidiophore $\times 80$. 8.9. Same, much enlarged. 9. Discharging conidium. 10-12. Fertilization and forming zygosporangium. After Brefeld.

the sputum of a tubercular patient. It is pathogenic for rabbits and guinea pigs when inoculated. Lucet and Costantin think that cases of mucormycosis are more frequent than generally supposed. Meyer seems to have made the first observation of a *Mucor* in animals having observed it in the lungs of a dog. Heisinger in 1821 found a *Mucor* in the lungs of a goose.




Fig. 47. Fly Fungus. *Empusa Muscarum*. 1. *Empusa* on fly surrounded by a halo. 2. Part of body of fly general fructifying part (c), conidia (c) and secondary conidia (sc) x 40. 3. Fully formed conidiophore with conidium (c) and capsule (c) x 300. 4. Tubular conidiophore projecting a conidium (c) surrounded by part of the plasma (g) of the conidiophore. 5. Conidium (c) with a secondary conidium (sc). 6. Conidium (c) forming a mycelium tube x 300. 7. Secondary conidium (c) germinating x 300. 8. Part of chitinous integument of fly with conidium (c) penetrating the integument x 500. 9. Fatty bodies of fly containing mycelium of parasite x 300. 10. Yeast like sprouting cells (c) from the fatty bodies of a fly x 500. After Brück.

Mycelium abundant, generally parasitic on living insects, multi-nucleate, non-septate or may become septate, asexual reproduction by means of conidia which are cut off from the end of the sporophore, conidium with one or many nuclei, conidia forcibly ejected; sexual reproduction by means of zygospore, azygospores without fertilization also frequent. One of the most common species of this family is the House Fly Fungus *Empusa Muscar*. *Empusa sphaerosperma* is found on the larvae of Cabbage butterfly, *F. Gryllis* is on the Rocky Mountain Locust and the *Macrospora cicadina* is found on the Cicada. *Basidiobolus ranarum* occurs on frog excrement.




Fig. 48. Peronosporaceae, White Rust. *Albago canad.* 1. Inflorescence of Shepherd's Purse with fungus. 2. Mycelium with haustoria (h) x 198. 3. Conidiophores and conidia (spores) in chains x 400. 4 and 5. Formation of zoospores in conidia x 400. 7. Oogonium (o) and antheridium (a) attached, mycelium shown below. 8. Zygospore with thick wall. 9. Germinating zygospore forming a zoosporangium. 10. Zoospore. 7 0x400. 6. Germinating Zoospore. After DeBary.

OOMYCETES

Mycelium occasionally sparingly developed, tubular asexual, reproduction by swarm spores or conidia; sexual by the formation of oospores in the *Peronosporaceae* and *Saprolegniaceae*.

Synchytrium has a much reduced mycelium. Sexual reproduction found only in some of the genera of the family *Chytridiaceae*. The non-septate mycelium is reduced to a single sac-shaped cell forming a kind of gall in the host plant. One species of *Synchytrium*, the *S. decipiens*, occurs on the Hog-pea (*Amphicarpea monoica*).

The family *Pythiaceae* contains a destructive parasite of seedlings, the *Pythium DeBaryanum* and the *P. proliferum* upon dead insects in water.

ALBUGINACEAE AND PERONOSPORACEAE

Mycelium generally well developed. Reproduction sexual and asexual in sexual reproduction oogonia and antheridia, asexual spores, conidia, or zoospores.

Many members of this group are destructive parasites to cultivated plants like the potato rot fungus (*Phytophthora infestans*), the onion mildew (*Peronospora Schiedensiana*), the lettuce mildew (*Bremia Lactucae*), the mildew of the sunflower (*Plasmopara Halstedii*), the Clover mildew (*Peronospora trifoliorum*) which may be injurious to animals, the millet mildew (*Sclerospora graminicola*) which may also be injurious. As a type of this family the downy mildew of the grape (*Plasmopara viticola*) may be taken. It appears during the early summer and continues till frost. Leaves, berries and stem are affected. The upper surface of the leaf shows yellow patches underneath a white frosty mould. A section through the leaf will show the mycelium vegetating between the cells. The mycelium gives rise to the fruiting branches of the fungus the conidiophores, which pass out through the stomata. The conidiophores are dichotomously branched, and at their ends bear the conidia. When these conidia are placed in water they begin to change, at the end of an hour they swell and the contents divide. According to Dr. Farlow "at the expiration of an hour and a quarter the segments had resolved themselves into a number of oval bodies" which before long succeeded in rupturing the cell wall, and making their escape from the mother cell. Each of these zoospores is provided with two cilia. In some, zoospores are not produced, but the whole mass passes out, which soon produces a tube. The zoospores produce germ tubes which probably pierce the leaf of the grape. The temperature most favorable for germination is between 25° and 35° C. Inoculation experiments with the grape vine mildew show that on the second day the disease appears. Sexual method of reproduction takes place later in the season and occurs in the leaf. A slight swelling appears at the ends of the branches of the mycelium, which is spherical in shape, the cell wall being thick and pale yellow in color. The whole rounded mass is called the oogonium. The central part is the oosphere. A small body is developed from another (or the same thread) which has a long-




Fig. 49. 1 Downy Mildew (*Peronospora calotheca*). Mycelium between the cells and sporangia. 2 Potato Rot Fungus (*Phytophthora infestans*). Conidiophores, conidia borne on the branches. 3 Single conidium forming zoosporengium and the zoospores. 4 Discharge of zoospores. 5 Single ciliated zoospore. 6 Oogonium and antheridium. 7 Oospore and antheridium of *Peronospora blattarum* $\times 590$. 8 Conidiophore and conidia of *Bipodophora entospora* found on leaves of *Brigadier* $\times 200$. 9 Germinating conidia of *Bremia Lactucae* (the Lettuce Downy Mildew). 10 Conidium of *Peronospora leptosperma* germinating $\times 300$. 1-7 9-10 after DeBary. 8 after Cornu.

side of the oogonium, the antheridium. This pierces the oogonium and the protoplasm of the antheridium passes into the oosphere.

In the species that have been studied like *Peronospora parasitica*, and *Albugo candida*, the oosphere or egg cell contains a single nucleus, situated about at the center, the remaining nuclei having passed into the peripheral layer of the protoplasm of the periplasm.

A single male nucleus passes from the antheridium into the egg cell and fuses with the nucleus of the egg cell. Numerous investigations on this line have been made by Stevens,* Bertese,† and Wager‡. It is probable that the course of reproduction is similar for other species.

In fertilization karyoknetic changes occur. The protoplasm surrounding the oosphere is used to build up the wall of the oospore. Germination of oospore probably takes place in the spring. In *Albugo* or *Cystopus* the conidia are borne in a moniliform chain.




Fig. 50. Downy Mildew of Clover (*Peronospora trifoliorum*). a. Conidiophore. c. Stem. b. Conidium common in Europe on Red Clover, may be injurious to animals. After Smith.

* Bot. Gaz. 28:149, 21, 77, 34-420.

† Jahrb. f. wiss. Bot. 31:159.

‡ Annals of Bot. 4:127, 14:263.



Fig. 51. At the left, leaf of Green Foxtail (*Setaria viridis*) containing the oospores of *Sclerotinia grisea*. a—single oospore at a. After Trelease. At the right, spike affected by the same fungus b spikelet enlarged. The figure c the aph oospores from Hungarian grass oog. Oospore. oos—oospore or oospore. The middle figure, Husted, the right hand Charlotte M. King

In some forms the oospores are rare, and in the potato rot fungus they have apparently not been found. In some members of this group oospores are formed without fertilization, (Parthenogenesis)

Dr G P Canton,* who has made a careful study of the Lima Bean Mildew (*Phytophthora Phaseoli*) and the potato rot fungus (*Ph. infestans*) has been unable to find that the mycelial thread of the antheridium had the same origin as the one which bears the oogonium. It is possible that the fertilization is accomplished in a manner similar to that given for some of the Mucors. He says in a discussion of the potato rot fungus, "All of these facts are now in favor, rather than against distinct mycelial strains (heterothallic forms) except the last, which might indicate a homothallic form, one which contains both antheridia on the same mycelium."

SAPROLEGNIACEAE

Hyphae, long branched, undivided, zoosporangia cylindrical oospores produced from sexual organs, terminal cells are cut off and converted into ether




Fig. 52 Saprolegniaceae. Water Mould. 1, 3. *Saprolegnia Thureti* x 100. 2. Zoosporangium before the discharge of spores. 3. Same with biciliated spores being discharged. 4. The large spherical body, an oogonium and many oospores. 5. *Dicellaesphaera* a—oogonium, b—antheridium. 6. *Aphanomyces stellatus* a—oogonium, b—antheridium x 300. 7. *Leptomyces* sp. 8. Young zoosporangium x 200. 9. Part of older zoosporangium with zoospores sp and cellulose grains (c) x 300. 9. Zoospores x 430. Fig. 13 after Thuret. 4-6 after DeBary. 7-9 after Pringsheim.

oogonia or antheridia. The oogonia may give rise to one or many oospheres or egg cells. The antheridia are tubular and spring from the hyphae below the oogonia. They apply themselves to the oogonia and send out fertilization tubes to the egg cells. The latter then develop into oospores. The asexual method occurs as follows. An examination of the young threads of *Saprolegnia* will show long filaments which in places are filled with granular protoplasm. Some of these threads are separated from the rest of thread by a cell-wall. Soon the protoplasm arranges itself into polygonal areas. When mature the sporangium breaks and the zoospores are discharged into the water. When emptied a new sporangium is formed by the filament growing up into the old one, or in some cases a branch buds out below the oogonium.

Recent investigations indicate that the egg cell contains numerous nuclei, but as a general thing they are all degenerate but one. The antheridia also contain many nuclei. According to the investigations of some, one male nucleus enters the egg cell and fuses with its nucleus.

Trow,* Davis,† and Kauffman‡ have thrown light upon the development of the reproductive body.

Saprolegnia. Nees von Esenbeck. Water mould

Delicate branching hyphae, zoosporangia open from a terminal pore, zoospores pear-shaped with 2 terminal cilia. About 11 species common on decaying objects in water.

Saprolegnia monaca (Fringsheim) De Bary

Zoosporangia cylindrical; antheridia usually in close proximity to the oogonia frequently originating from the same branch, oogonia from short lateral branches, oospore spherical 16 to 22 μ in diameter, germ tube formed in germination.

Distribution. Widely distributed in North America and Europe.

Pathogenic properties. Occurs on dead insects thrown into the water, parasitic on living fish and crayfish. Frequently troublesome in aquaria. The *S. Thureti*, *Dellary* and *Achlya prolifera* are found on sick fish and crayfish. Hoffmann in 1867 stated that fish in aquaria died under the influence of *Mucor succedo* and *Saprolegnia*.

It is doubtful, however whether the *Mucor* produced death.

Basidiomycetes

Conidiophores arise from a many-celled, well developed mycelium, hyphae either separate or forming masses, texture soft, powdery or leathery, the spores various, in the most common type, the basidiospore is borne on special structures known as basidia from which arise little bodies called sterigmata into which some of the protoplasm of the basidium passes. In one group the mycelium consists of septate, branched threads, at maturity nearly disappearing because of gelatinization, mycelium gives rise to chlamydospores formed endogenously, reproduction sexual and asexual, usually the latter, comprises the sub-classes, Hemibasidia and the Eubasidia.

* Annals of Botany. 18: 541.

† Bot. Gaz. 35: 233.

‡ Ann. of Bot. 22: 361.

Hemibasidii

Mycetium, local or wide-spread, hyaline, septate, branched, becoming compact and giving rise to endogenous spores, the chlamydospores, color varies, in germination the spores produce a promycelium of terminal or lateral sporidia. These may propagate by budding like the yeast plant. The families are the *Ustilaginaceae* and *Tilletiaceae*.

HEMIBASIDIALES

This includes two families mentioned in the Hemibasidia

USTILAGINACEAE, Schrot

Usually parasitic fungi in the tissues of living plants. Sori usually exposed, forming dusty masses. Spores germinate by means of the septate promycelium which gives rise to terminal or lateral sporidia. In some cases these multiply like the yeast plant, or else produce infection threads. The order contains about 300 species, with the following genera in North America: *Ustilago*, *Sphaerolotheca*, *Metanopachium*, *Contracta*, *Schisonella*, *Mykosyrinx*, *Sporosporium*, *Thecaphora*, *Tolyposporium*, *Tolyposporella*, *Testicularia*. Many plants of the order are destructive parasites occurring upon economic plants like millet, timothy, pink, etc., and one species, the *Ustilago esculenta* on *Zizania latifolia* is edible being used by the Japanese for food.

***Ustilago*, Pers. Smut**

Mycelium septate, branched, gelatinous, sori on various parts of the host at maturity dusty, usually dark colored. Spores single produced in the fertile threads of the mycelium, the latter entirely disappearing at maturity. Promycelium septate, sporidia terminal or lateral, producing infection threads, secondary spores formed in the manner of yeast in nutrient solutions. The largest genus of smuts. About 250 or 260 species. Many of them are destructive parasites on cultivated and wild plants.

The *Ustilago minima* occurs upon the porcupine grass, (*Stipa spartea*) and the *Ustilago bromivora* upon the brome grass.

***Ustilago Zeae*. (Beck) Ung. Corn Smut**

Sori in the female or staminate inflorescence, leaves and nodes usually forming irregular swellings of variable size, at first covered by a membrane consisting of the gelatinized threads and tissues of the plant, soon rupturing, which exposes the blackish or brownish spores, spores sub-globose or spherical, or irregular echinulate 8-11 μ or sometimes 15 μ long. Spores germinate readily under favorable conditions, spore consists of an outer wall, which is spiny and an inner more delicate, the endospore, the germ tube or promycelium as it is called normally bears lateral bodies, the sporidia, but under more favorable conditions of food these may branch and bear secondary conidia. If the nutrient material is not exhausted this process of budding may be continued for a long time. These spores may propagate in a decoction of manure. It will then be seen that these budding conidia may be a center of infection.

The conidia as well as the secondary conidia are blown about by the wind and under proper conditions cause the infection of the corn plant. Several years ago Mr. F. C. Stewart made some extended studies of the germination of corn smut in which it was shown that the thermal death point of smut



Fig. 53. Maize smut (*Ustilago zeae*). Ear affected (a). Bracts (b). Smut balls (c). Kernels (d). In staminate flowers. Fig. 54. Smut balls making its appearance at the nodes. (C. M. King). 2. Foxtail Smut (*U. neglecta*). 3. Covered Smut of Barley (*U. hordei*).




Fig. 53. Maize Smut (*Ustilago zeae*). Cells showing thread of mycelium passing from cell to cell. a thread shows through. (A) section of cut sheath and thread. 2. Corn Smut Spores in process of germination; each spore is sending out a tube with small lateral bodies. 3. The same spores germinating in nutrient solution spreading like yeast. From U. S. Dept. Agr. 4. Kernel Smut of Sorghum (*Sphaerostheca Sorghi*) on Sorghum.

spores is 15 min. 105° 106° C. in dry oven, and 52° C. when immersed in water, and that corn is unable to come through an inch of soil after 15 minutes treatment with water at 70.5° C., and in dry oven at 78° C. Brefeld found that smut spores produced an abundance of secondary conidia when they were germinated in sugar solution, but with us this has never been a very satisfactory method of propagating them as the cultures soon became infected with bacteria which materially check the progress of the germination of spores.

Distribution and Hosts. Corn smut is found from the Atlantic to the Pacific wherever corn is cultivated, also in other parts of the world. In addition to occurring upon corn, it is found upon teosinte.

Poisonous properties. It has been held by many that corn smut is injurious to cattle. This has been a common belief in some quarters. In some kinds of smut a small amount of ergotin is found. Kedzie reports the following composition: Moisture, 8.30 per cent, albuminoids, 13.06 per cent, carbohydrates, 25.60 per cent, cellulose 24.69 per cent, sugar, 4 per cent, fat, 1.35 per cent, ash, much sand, 25.5 per cent. Professor Kedzie was unable to find any poisonous alkaloids. In 1888, the United States department of agriculture employed Professor Gamgee to ascertain the cause of the cornstalk disease. Professor Gamgee records his experiences as follows and concludes that smut is not injurious.

One cow was fed thrice daily one and one-half pounds of cornmeal and three ounces of smut mixed with as much cut hay as she would eat. The second had no smut in her ration, but was fed the same amount of hay. The amount of smut given in each case was increased to six ounces. The cow fed on dry food lost flesh. Eight days later the dose of smut was increased to twelve ounces three times a day. The cow on the wet food gained in condition, the other one lost. In three weeks the two cows consumed the forty-two pounds of smut. They had a voracious appetite the whole time and the only indication of a peculiar diet was a very black color of the excrement and the loss of flesh by one animal, although liberally fed on nutritious diet, which, however, was given in a dry state. It is evident that smut is not a very active poison in combination with wholesome food, and especially if the animal is a lowed grain food and plenty of water to drink.

Prof. W. A. Henry, in his work on "Feeds and Feeding" speaking of the work done by the Bureau of Animal Industry, Clinton D. Smith and Gamgee says:

In experiments by the Bureau of Animal Industry, U. S. Department of Agriculture, Washington, corn smut was fed to heifers without ill effects. With a little trouble but one ending without disaster, it seems reasonable to conclude that corn smut is at least not a virulent poison, if indeed it is one in any sense of the word. It is probable that in the Wisconsin cases, where one cow died and the other was indisposed, the animals suffered because of eating too much highly nitrogenous material rather than anything poisonous. Worse results might have followed the feeding of the same volume of corn meal or cotton seed meal. It would seem that there is little or no danger from corn smut unless cattle consume a large quantity. This is possible where they are allowed to roam through corn fields and gather what they wish. There may be cases where animals seek out the smut and eat inordinately of it.

A few years ago Prof. Smith of the Michigan Agricultural College gave the results of some experiments with corn smut. Varying amounts of smut were fed to three grade Shorthorn cows and one grade Jersey. Two of the cows were started with two ounces a day and increased to eleven pounds. Two others were started with two ounces and increased to a pound. The test lasted forty-nine days. They appeared to relish the smut. It produced no signs of abortion in pregnant cows, the milk yield was normal. Prof. Smith concludes that the smut in corn fields is not likely to prove injurious.

Real states that under certain conditions smut is likely to be injurious to




Fig. 88a. 1. Sprouting Grass Smut (*Ustilago panicum multiseptata*) showing large swelling in upper part of plant. C. M. King. 2. Millet smut (*Ustilago Crameri*) a, spores b, glumes of millet grain filled with a powdery mass of spores. C. M. King. 3. Kernel of smut corn (*Ustilago Pacheri*) on maize. Spores as right. Below a sectional view of an affected kernel. Panuol and King.

cattle. The experiments made by Moore also indicate, as do those of Smith, that smut is not injurious.

Beginning on the morning of January 17, 1894, and continuing until noon of February 2 (sixteen and one-half days), the heifers were fed morning and evening from two to three quarts of a mixture of equal parts by weight of cut hay and a mixture of corn meal, middlings and wheat bran, and sixteen quarts of smut. No injurious effects were observed by Moore. It seems reasonable to conclude from these experiments that under proper conditions corn smut is not injurious. In our experience no cases have ever been reported to us where cattle were supposed to have died from eating corn smut.

Professors Veranus A. Moore and Theobald Smith after making an exhaustive investigation of the so-called corn stalk disease, came to the conclusion that "corn smut is probably not very poisonous, but when fed in considerable quantity no doubt produces injurious symptoms." Miquel in an old work on poisonous plants published in 1838 in Dutch regarded the smuts as poisonous.

Dr. Peters of the University of Nebraska, makes the following comments on the subject of corn smut:

At a Farmers' Institute at David City a gentleman stated that he had often heard his neighbors say, and he had also read the same in agricultural papers, that cornstalk disease was caused by corn smut. He had the opportunity to make the test for himself. He was compelled to clear the farm he rented of the smut. His son gathered the smutty stalks into a yard where two cows ate considerable of the smutty leaves. No bad results followed, as witnessed by the gentleman himself and the owner of the place.

Mr J J Jolliffe in the Drovers' Journal of February 7, 1902, says

I have never had any bad results from smut. I have watched the stock eat ears that seemed 75 percent smut, and they devoured them apparently with as much avidity as sound ears, never affecting the health or appetite in the least.

We have seen cattle die in fields where there was no smut whatever on the stalks and also on some fields that had previously been cleaned as good as one could clean them from the smut, the cattle died in spite of this precaution. Often the owner of the neighboring fields, in which large quantities of smut were found, did not believe this theory and allowed his cattle to run in the stalks promiscuously without suffering any loss.

At the Illinois Experiment Station about sixty pounds of corn smut were fed to a grade Jersey steer, with grain and hay as follows:

From December 13, 1889, to January 2, 1890, 120 pounds of grain, 105 pounds of hay, 40 pounds of smut, and 371 pounds of water were given. From January 2 to January 24, 1890, 176 pounds of grain, 133 pounds of hay, and 39 pounds of smut were fed and 368 pounds of water given. When the experiment was begun, December 13, 1889, the steer weighed 560 pounds. January 2, 1890, he weighed 551 pounds, and January 24, 555 pounds. No evidence of disease was discovered.

Dr Kiborne records two experiments to test the effects of corn smut. In the first case the smut used came from a field in which several animals had died within five days after they had been turned into it. Three two-year-old steers were fed exclusively on smut laden stalks and free smut mixed with a small quantity of a mixture of corn meal and wheat bran, for seven days without ill effects. He concludes "It is safe to say that these animals consumed a much greater quantity of smut than the animals which died in the fields." In the second case, two heifers were fed in addition to corn and hay, sixteen quarts of smut morning and evening for sixteen and a half days. This feeding continued for several months. The animals appeared healthy at the termination of the experiment.

Dr N S Mayo records the experience of a farmer near Manhattan, who gathered the smut from the field and placed it within an enclosure. The cattle broke into the enclosure one night where the smutty corn was thrown and ate all they wished, but no injurious effects were observed.

Lstilago avenae (Pers.) Jens. Oat Smut

Sori, found in the spikelets forming a dusty olive brown mass, usually destroying the whole of the inflorescence or only a part, the spore mass at first covered by a membrane which later breaks, thus allowing for the scattering of the spores, spores olive brown, lighter colored on one side, spherical to sub-spherical or somewhat angular, minutely roughened, 6-9 μ in length. Spores germinate readily in water, infection takes place at the time of germination of the oats.

Distribution and hosts. Found wherever oats (*Avena sativa*) is cultivated also on wild oats, (*A. fatua*).

Poisonous properties. Probably not any more injurious than corn smut. When present in large quantities it may produce a sore throat, because of irritation. This fact is mentioned by White. The following note from Dr White refers to another species found upon grass:

The inflammation affects almost exclusively the face and genitals. It begins upon the former with a violent itching in about twenty-four hours after contact with the seeds, which is followed by a uniform redness, especially marked about the nostrils, and swelling of the eyelids. The appearance of the patient strongly resembles that of erysipelas. Later small vesicles develop, terminating in permanent excoriations.

Upon the male genitals it begins also with itching, followed by general swelling, with intense redness of the scrotum, and later by vesicles filled with a yellow serum, terminating



Fig. 56. A. Oat Smut (*Ustilago avenae*). From U. S. Dept. of Agri. B. Tall Meadow Oat Smut (*Ustilago perennans*)

Fig. 56a. Porcupine Grass Smut (*Ustilago lepidotes*) affecting parts of inflorescence and culm. a, spores, said by Dr. White to be an irritant. C. M. King.

in persistent and very painful erosions. The penis is sometimes affected, producing an inflammatory phimosis.

Delicate skins are most easily affected, and a moist condition favors the action of the parasite. Similar effects were produced upon the skin of the rabbit, by applying to it after being shaved some of the fungus removed from the seeds.

"After reading the above, I wrote to Prof. W. G. Farlow of Cambridge, our authority in cryptogamic botany, with regard to the occurrence of this species in America, and received the following reply:"

Your information about the poisonous character of *Ustilago hypoduræ* is something quite new to me. I do not know of any reference to the subject in botanical books. The spores of *Ustilagines* are known to be at times irritants when they reach the air-passages, but they are not poisonous to handle. *U. hypoduræ*, a species whose characters are not very well marked, I may say, has been found in two places in this country. I found it at Wood's Hole Mass. on *Phragmites* (reed). It was found by Curtis in North Carolina on *Arundinaria*, the cane, and what is probably the same species occurs in Iowa on a species of *Stipa*. The fungus may be much more common in this country than is now supposed, as few persons have collected fungi of this order.

Ustilago maydis, the corn-smut, grows upon our maize, and *U. setatum* attacks several of our grains, wheat, oats, barley, and our grasses, but I have never heard of their producing any irritative action upon the skin."

Ustilago nuda (Jens.) Kell. & Sw. Barley Smut

Mycelium found in spikelets forming a dusty olive brown spore mass, protected by a thin membrane which soon becomes ruptured and allows for the dispersal of the spores. Spores lighter colored on one side, minutely roughened, spherical, subpherical or elongated 5-9 μ in length. Infection probably takes place at the time of flowering of barley.

Poisonous properties. Like the preceding, *Ustilago Hordei*, found upon barley, differs from the species described because of an adhering purple black spore mass covered permanently by the lower parts of the glumes.

Distribution and hosts. Found wherever barley is cultivated in Europe and North America.

Ustilago Triticæ (Pers.) Jens. Wheat Smut

Spore masses found in the spikelets of the inflorescence forming black or olive brown mass, usually destroying the entire floral parts, and later spores are scattered by the wind, spores usually spherical or nearly spherical or somewhat elongated, minutely roughened, 3-9 μ in length.

Distribution and hosts. Common upon wheat wherever cultivated in Europe, Australia and North America and South America.

Poisonous properties. Probably injurious like the preceding smuts.

Ustilago Paniculæ (Wallr.) Wurt.

Sori in spikelets infecting all the spikes, spore masses enclosed by glumes, with a rather firm membrane, soon ruptured, permitting the scattering of the spores, spores dark brown, usually spherical or ovoid, occasionally elongated, rather prominently echinulate, 10-14 μ in length.

Distribution. Widely distributed in Europe and North America. One of the most common smuts wherever foxtail grows.

Poisonous properties. It is supposed by some farmers to cause abortion but there is nothing to support this view. According to Professor Power it contains a small amount of ergotin. Possibly injurious like the other species of smut.




Fig. 57. Spore spores of various kinds. 1. Tall meadow oat spore (*Lathyrus perennans*). 2. Timothy spore (*L. strigosus*). 3. Rye spore (*L. coarctatus*). 4. Chestnut spore (*L. stiloglossus*). 5. Foxtail spore (*L. negundo*). 6. Mistle spore (*L. parva mellea*). 7. Sandbar spore (*L. cerat*). 8. 9. *Triticum* germinating. 10. Secondary spores from conidia. 11. 12. *Urocystis* spores germinating. 13. 14. Micro-um of same. A. Suckers or hyphae. B. Hyphae passing from one cell to another. B 14 after Wulff.

Mr. W. A. Kelty informs me that the smartweed smut (*Ustilago utriculosa*) often produces severe irritation of the hands when corn is husked.

TILLETIACEAE, Schröt

Mycelium becoming gelatinized in the tissues, the promycelium gives rise to the terminal cluster of elongated sporidia which fuse or do not fuse in pairs, producing secondary sporidia which may be alike or unlike, or the sporidia germinate directly into infection threads. About 150 species, of which *Tilletia* is the most important genus, aside from the two species described below one species, *T. secalis* is found upon rye, *T. hordei* upon barley, etc., *Neovossia lowensis* on *Phragmites communis*, *Urocystis occulta* on rye, *Urocystis agropyri* upon quack grass, and *Entyloma ranunculi* upon anemone.




Fig. 58. Wheat Bunt (*Tilletia foetens*). At the left, a beardless variety with bunt kernels. At the right, a bearded variety with bunt kernels. From U. S. Dept. Agel.

Tilletia, Tul. Bunt

Sori in various parts of the plant, usually in the ovaries, forming a dusty spore mass, spores 1-celled, formed singly in the ends of the mycelial threads, promycelium of germinating spore short with a terminal cluster of elongated sporidia. 53 species are reported for the genus.

Tilletia foetens (B. & C.) Trel. Stinking Smut or Bunt

Sori in the ovaries, wheat glumes more or less spreading; spores light to dark brown, oblong or sub-spherical, or spherical or sometimes irregular,

16-20 μ in diameter, bad odor just before maturity and retaining the odor even in stored grain.

Distribution and hosts. Common in eastern North America, also in Canada and Manitoba.

Poisonous properties. It produces a bad odor when it occurs in flour and also gives the same a dark color and makes it unsalable.

Tilletia Tritici (Bjerk.) Wint. Wheat Bunt

Sori in the ovaries of wheat ovate or oblong, glumes spreading, spores chiefly spherical or sub-spherical, 16-22 μ in diameter, light to dark brown with winged reticulations. Infection of this and the preceding smut occurs at the time of germination of wheat, hence all of the stalks growing from the single wheat kernel become infected, mycelium growing upward with the growth of the plant.

Distribution and hosts. Common upon wheat wherever cultivated. Reported as destructive and abundant in Michigan, Montana and Kansas.

Poisonous properties. Same as in the preceding species.

Conidophores with true basidia—reproduction generally asexual, sexual in some cases through the fusion of nuclei—spores cut off from the ends of the threads or borne on little sterigmata. The group is divided into two divisions according to the form of the basidia. *Protobasidiomycetes*, the rusts and gelatinous fungi, *Autobasidiomycetes*, toad stools, mushrooms, and puff balls.




Fig. 59. The Gelatinous Fungi. *Tremellineae*. 1. *Tremella isidiosa* on wood. 2. Cross section through hymenium. 3. Basidia. 4. Conidia. 5. Basidiospores, $\times 450$. 6. *Tremellodon gelatinosum*. 7. Basidia of the same $\times 560$. 1-3 after Brefeld. 4-5 after Mueller.

PROTOBASIDIOMYCETES

Parasitic or saprophytic plants, basidia with longitudinal or cross septa, mycelium septate, branched, either in the interior of the plant as parasites or ramifying the substratum spores various. The following types occur spermatia, aecidiospores, uredospores, teliospores, and sporidia the spermatia are always accompanied by other spores, generally with the aecidiospores the aecidiospores are 1-celled and occur in cups the uredospores are 1-celled, occur in a powdery mass, and germinate immediately, teliospores arise from the same mycelium that produces the uredospores, one or more cells, on germination they produce a promycelium that bears the sporidia. This group contains the following families *Endophyllaceae* with a fungus parasitic on the leaves of spurge, stonecrop and house-leek, *Melampsoraceae* including several important economic fungi *Pucciniaceae* containing a large number of genera, the *Auriculariaceae*, gelatinous fungi common on decaying wood.

MELAMPSORACEAE

Teliospora forming incrustations on the surface of leaves; uredosori powdery aecidia without pseudoperidium (Caecoma) or with well developed




Fig. 60. *Uredineae*. Teliospores of different genera germinating. By germination originate the promycelia which divide into cells, each of which produces a conidium. 1. *Uromyces Pabae* x 460. 2. *Triphragmium Ulmariae* x 370. 3. *Melampsora betulina* x 370. 4. *Phragmidium Rubi* x 370. t—Teliospore, sp—Conidium. After Tulaane.

pseudoperidium, uredospores 1-celled occurring singly or in groups with or without pseudoperidium; paraphyses present, teleutospores 1-4-celled, closely or loosely united in the plant underneath the epidermis, *Calyptospora Goepfertiana* occurs upon the huckleberry and blueberry, (*Vaccinium*) connected with the *Aecidium columnare*, a very troublesome parasite upon *Abies*. The *Melampsora populina* occurs on the cotton-wood, the cotton-wood rust forming red sori on the leaf of the cotton-wood, with waxy incrustations. The other troublesome parasite, the *Chrysomyxa Rhododendri* occurs upon *Rhododendrum*. There are about 100 species in the family.

Coleosporium. Lev

Teleutosori forming flat waxy masses in the leaf, teleutospores composed of several vertical cells enclosed in a thick transparent membrane, each cell germinates by a single undivided promycelium which produces at the end a single sporidium, uredosorus reddish or orange, powdery, spores spherical or sub-spherical, ovate, elliptical, oblong or cylindrical, produced in basipetal chains. A small genus of 30 species.

Coleosporium Solidaginis (Schw.) Thum. Golden Rod Rust

Uredosori rounded, soon pulverulent and scattered, orange spores in short chains, spherical, oblong, or sub-cylindrical spiny, 20-35 x 15-20 μ ; the teleutosori at first orange, becoming red, flat often confluent forming waxy crusts; Teleutospores cylindrical or somewhat clavate generally 4-celled 60-70x15x25 μ , occasionally longer.

Distribution and hosts. Found in various Compositae, notably *Solidago canadensis*, *S. serotina*, etc., *Vernonia noveboracensis* and *Sonchus*. Occurs in both Europe and America.

Poisonous properties. Suspected of being injurious to horses, possibly produces stomatitis. Referred to at length under Golden Rod

PECCINIACEAE

Teleutospores with a short or long pedicel, spores single or in groups; spores one or more celled; frequently interspersed with paraphyses, spore mass powdery or gelatinous, sporidia arising either from the promycelium or from a similar sterigma after segmentation of the spore contents, aecidia with or without pseudoperidia, uredospores 1-celled, arising from the conidiophores. Includes the *Gymnosporangium macrospora* which produces its aecidium stage on the apple and the teleuto stage on the red cedar, the cedar apple gall with its long gelatinous horn being characteristic, and various aecidia connected with various rusts. Many species have a well marked alternation of generation, an aecidium on one host and the uredo and teleutospores on another host. The *Hemileia vastatrix* produces the coffee leaf disease of Asia, *Phragmidium subcorticum*, the rose rust. This large family contains 1500 species.

Recent investigations on the subject of the fertilization and reproduction in the rusts have been made by Profs. Olive, Blackman, Christman, Holden and Harper, and others. These studies seem to indicate that fertilization occurs, but that this fertilization is not, as was thought by the older writers, to be compared with that which occurs in some of the Ascomycetes. The older view was that the spermogonia were male organs and form a strictly morphological standpoint comparable to the structures of like character found in that group of



Fig. 61 Uredosori of Tickle Grass Rust. (*Puccinia emaculata*).



Fig. 62a. Teleutosori of *Puccinia coronata* on leaf of oats. Sheath affected with *Puccinia graminis*.

plants. Clements arranges the rusts with the Ascomycetes. He considers that the apothecium is reduced.**

Puccinia, Pers. Rust

Teliospori flat, usually powdery masses, teliospores separate with pedicels, usually consisting of 2 cells, occasionally 1-celled, or sometimes more than 2 cells, the germ pore of the upper cell at the apex the lower cell with the germ pore placed laterally below the septum, the promycelium septate with several spores. The teliospores germinate immediately in some species, in others after a period of rest. About 700 species.

Puccinia graminis, Pers. Common Grass Rust

The aecidiospores generally circular thick swollen with reddish spots surrounding the infected area, yellow below, peridia cylindrical with whitish torn edges, spores subglobose smooth orange yellow 15-25 μ in diameter spermatogonia on the upper surface consisting of small black dots, uredosori orange red, linear but often confluent forming long lines, powdery masses spore elliptical, ovate or pyriform, echinulate orange yellow 25-34 x 15-20 μ , germ pores 2 above the center on each side teliosori persistent open, generally forming lines on the sheaths, stems and inflorescence teliospores fusiform clavate constricted in the middle generally smaller below the apex thickened, rounded or pointed smooth chestnut brown 15-20 x 35-65 μ , pedicels long and persistent.

The life history of common grass rust is as follows. The common rust produces three stages. One stage occurs in the barberry and is known as the cluster cup fungus. This stage makes its appearance in the northwest some time during the month of June in the latitude of Ames, a little before the middle of the month. An examination of an affected leaf will show small black specks on the upper surface surrounded by a yellow spot. This is known as the

* The Genera of Fungi 5-98

** Christman in his studies (Bot Gaz 39:267 Trans. Wis. Acad. Sci. 13:317 Bot. Gaz. 44:18) of the common rust (*Phragmidium species*) finds that the ends of the hyphae produce a terminal sterile cell and a lower fertile one. The fertile cells fuse in pairs with one another the cell wall is breaking down. The nuclei are side by side and divide two of the daughter nuclei remaining in the lower cell and two passing to the upper part of the dividing cell. Then the upper portion becomes separated by a transverse wall and becomes the first spore mother cell. These last two are approximately equal. Backman (Ann. Bot. 18:323 see also Ann. Bot. 19:11) in his studies of another species of the same genus, states that the fertile or fusion cell contains a larger nucleus and that the male cell is reduced, that the hyphae which gave rise to an aecidium first cut off a sterile cell and the cell below which at first only contains a single nucleus becomes binucleated because of the passage of the nucleus from an adjoining cell. Six regular divisions of the nuclei occur and finally we have a chain of spore mother cells each having a pair of nuclei.

Dr. Oliver (Annals of Bot. 22:331 Bull. S. Dak. Agr. Ex. St. 1193) who has made a most careful study of *Triphragmium ulmariae* comes to the conclusion that two fusing gametes as well as their nuclei are approximately equal, and that the two gametes differ somewhat in time of development. "That the apparently normal and regular occurrence at the base of certain young aecidia of one to many multinucleated cells, points to the necessity of a broader conception as to the mode of development of the aecidium-cup than that held by either Backman or Christman. While the part which these multinucleated cells take in the development of the aecidium is as yet somewhat obscure the evidence appears to point to the conclusion that they are sporophytic structures and that they result from the stimulated growth which followed sexual cell fusion. Should this prove true, it is obvious that the 'fusion cell' does not at once function as a 'basal cell', at the bottom of each spore-row, as was claimed by Christman for this type of Rust. Further the occurrence of occasional instances suggesting nuclear migrations undoubtedly of a pathological nature, between the multinucleated cells of *Puccinia praevalens* throws doubt on the idea as to the normal origin of the binucleated condition in the aecidium-cup by this means."

It may be of interest further to state that Holden and Harper (Trans. Wis. Acad. Sci. 14:63) have studied a species of *Cercosporium* and find that the fusion nucleus divides in a manner similar to that of higher plants.




Fig. 42. Some Rusts. A B. D. E. *Puccinia graminis*. A. Accidium or Cluster cup on Barberry, a aecidium, a spermogonium on upper surface. B. Uredo spores & one-celled teliospores two-celled. C. Germination of uredospores showing long tube. D. Connection of stem showing two-celled teliospores. E. A teliospore germinating with promycelial tube (p) and sporilla (sp). F. Two-celled teliospores of *Puccinia coronata* on oat leaf. G. Teliospores of *Phragmidium intermedium*. All much enlarged. A and C. after Lortsen, B. D. after Uellary. E. after Tulane.

spermogonial stage, the flask-shaped bodies are called spermogonia and contain the spermatia which do not germinate, their function is not known. A sweetish fluid, which attracts insects, is frequently found in connection with these. Directly opposite the flask-shaped bodies are small globular affairs, "cups," (aecidia), slightly irregular on the margins. Owing to their upward growth they rupture the epidermal cells, and finally the lining layer of cells of the cups also breaks, thus exposing a large number of 1-celled spores borne in chains. These spores arise from short stalks contained at the base of these cups; the cluster cup spores are known as aecidiospores and are transported by the wind and other agencies, and have the power to germinate soon after maturity. When the proper host—a grass, such as bent grass, oats or wheat—appears, the germ

tube enters by way of the stomata, or the so-called breathing pores. The germ tubes produced by the spore of *Aecidium berberidis* are simple or branched, and in fourteen days usually give rise to the uredo spores, which occur in definite spots called sori. These spots occur in great numbers along the veins of the leaves. Before breaking open, the tissues of the leaf are somewhat paler at those places. The nourishment afforded by the host causes a vigorous mycelium to form, which soon collects in places, pushes the epidermis out, and an orange-colored pustule is formed which is known as the uredosorus.

A section through a diseased sorus shows that an abundance of the vegetative mycelium grows between the cells of the plant, and in some cases haustoria penetrate them. This pustule contains a large number of 1-celled, round or elliptical, spiny, orange colored spores, the uredospores. The spores have two membranes, the outer exospore being provided with wart like projections, while the inner endospore is provided with several pores through which the germ tube appears. These spores germinate in from three to four hours and can thus start a general infection. These spores, carried by the wind, rain or insects to another part of the same or another plant, germinate, the germ tubes branch and spread over the surface, but the tube cannot enter the host—a grass of some kind, such as wheat, oats or barley—unless it reaches the opening of the stoma, since it cannot bore through the epidermal cells. A single sorus contains hundreds of spores, and as a single plant may contain hundreds of pustules, it can readily be seen that rust must become quite general.

The red rust stage is followed by the black rust stage, known as the telento stage. The sori are brownish black in color, and frequently occupy the same place that the uredo stage did. The spores are dark brown in color, two-celled and smooth, having attached to them a persistent stalk known as the pedicel. The telentospores do not germinate till the following spring, when each cell produces a germ tube, the promycelium bearing lateral spores, sporidia. These sporidia, when in contact with the barberry leaf enter by boring their way through the epidermal cells.

The barberry cluster cup fungus, and its connection with common grass rust. It is not absolutely necessary for the common grass rust to have its first stage on the barberry, yet experiment has shown beyond doubt that it does occur on that plant. The theory has been advanced that appearing in one of its stages on the barberry gives the parasite new vigor. It is not improbable that in some places the mycelium or vegetative part of the fungus may be perennial in the tissues of grasses, as it is with many other fungi, probably this is true in southern localities. Beyond question this rust produces spores during the entire year in our southern states, and on the approach of early spring gradually moves northward. It may also be mentioned that in the west this rust certainly does not appear before the cluster cup fungus on the barberry appears. It is usually eight or ten days later, and then appears to a limited extent only. Rust often appears where barberry does not occur within hundreds of miles. This was especially noticeable during the early history of grain culture in the northwest. Rust follows a general infection.

Distribution and hosts. This fungus has been found not only upon wheat but also upon several species of *Bromus*, *Triticum* and *Triticum spelta*. Its distribution cannot be given because in most cases the *P. rubigo-vera* included this as well as the *P. glumarum*. It has been intimated above that the uredospores make their appearance on young germinating plants in the fall, but it

appears that the uredo spores are not common the following spring. The investigations of the authors quoted here indicate that not in a single case was it possible to produce uredospores in the spring from those of the autumn.

H. I. Bolley, of Fargo, N. D., remarks in regard to several cluster cup fungi which occur on members of the Boraginaceae family.

Several necidia of unknown life history have been studied with reference to their relations to the red rust of *Puccinia rubigo-vera*, many infection tests being made upon young wheat and oat plants, all with negative results.

In this region *Oenothera biennis* bears very profusely an necidium, which, because of its date of appearance, was worthy of suspicion, but tests enough were made to remove this notion.

P. rubigo-vera as well as the common grass rust, is very destructive in England and Australia, but according to Wolf, is not so common in Germany. A few years ago Professor Arthur investigated the subject of wheat rust in Indiana and found that this species was much more destructive to wheat in that state than common grass rust. The same year, 1889, the writer found that this rust was much more common on wheat in Iowa. Carleton says he is confident that the orange-leaf rust (*P. rubigo-vera*) does very little if any damage to the grain in this country, that in all cases of serious damage to the grain by rust the black-stem rust (*P. graminis*) is the real cause. In 1907, the leaf rust was very destructive to spring wheat in Iowa.

Puccinia glumarum, Schmidt

Aecidium unknown, the uredosori occur along the veins. The diseased leaf is frequently of irregular contour, color orange yellow, spores spherical, or short, elliptical, spiny. Teleutosori, grayish, covered by the epidermis on the stalks and leaves, less frequently on the flowers. Sori divided into chambers,




Fig. 63. Covered Rust of Wheat (*Puccinia rubigo-vera*) from wild Barley, perhaps the same as *P. glumarum*.

surrounded by paraphyses. Spores with short pedicels, mostly club-shaped, unsymmetrical, apex somewhat truncate, or with one or two projections.

Distribution. Common in Europe and probably also in this country, has usually been referred to as *Puccinia rubigo-vera*. In European mycological works, the aecidium of this fungus is said to be very common on common speedwell (*Isithospermum arvense*), *Lichium vulgare*, and *Anchusa officinalis*.




Fig. 64. Forms of rust on cereals. A. Common wheat rust, *Puccinia graminis* on sheath of wheat, winter spores germinating. B. The same. sporidia sp. C. Epidermis under surface of leaf with sporidium, sp. and germ tube. s. penetrating the epidermis. D. Uredospore germinating after being in water 14 hours in E. *Puccinia rubigo-vera* germinating. F. *Puccinia graminis*. Both cells have germinated. s, sporidium germinating, magnified $\times 500$. G. Crowned rust (*P. coronata*) from oat leaf. G after Bolley, the remainder, DeBary.

Common speedwell is a very common weed in St. Louis and other parts of Missouri and southern Illinois, but so far as known, the aecidium has not been found on these weeds.

Puccinia dispersa. Eriks. & Hen

This species of rust is apparently very common in Europe. There are three different stages. The aecidium stage produces circular or elongated, somewhat swollen, spots on the leaves, petioles and stem of several members of the borage family. The spores are between 20 to 30 μ or 20 to 30 μ \times 19 to 22 μ in diameter. The telio spores long remain covered by the epidermis. The sori are chambered, surrounded by numerous brown paraphyses, spores are mostly club-shaped, unsymmetrical and 40 to 50 μ long.

Puccinia coronata. Cda. Crowned Rust

The aecidium produces round or elongated spots with elongated, conspicuous aecidia, the spores from 18 to 25 μ \times 14 to 19 μ , the uredosori are long, confluent mostly on the upper surface of the leaf, they are orange-colored, and are soon exposed, each pustule containing a large number of 1-celled subglobose, roughened spores which are spherical or short-elliptical. The uredospores are yellow 20-32 μ in diameter by 20-32 \times 20-24 μ . The teliospores remain covered by the epidermis, and in this respect they resemble the covered rust of wheat (*Puccinia glumarum*). They usually occur on both sides of the leaf. The spores are short-stalked, cuneate and more or less truncate above, crowned with several projecting horns.

Distribution and hosts. Common wherever oats is cultivated and in several of its forms it occurs upon cultivated grasses. This is a well known destructive rust of oats and several other grasses and has received considerable attention from early mycologists. Klebahn has recently described this rust under several distinct forms. The *P. coronata dactylidis* in a narrow sense includes the rust upon *Dactylis glomerata* or orchard grass, *Festuca sylvatica* with aecidia on *Rhamnus frangula* and *P. coronifera*.

Ericksson and Henning distribute these forms into *P. coronata* I and *P. coronata* II. Historically this rust is of considerable importance since Gmelin was familiar with this disease in 1791 and described it as *Aecidium Rhamni* on *Rhamnus*. The aecidium stage occurs on species of buckthorn (*Rhamnus*) especially (*R. cathartica* and *R. Frangula*). In Iowa an aecidium is frequently found on a native buckthorn (*R. lanceolata*), but its connection with this host has not been studied. The aecidium attacks not only the leaves, but occurs on mid-vein, petiole, pedicels and flowers. As a result of the attacks, distorted leaves and flowers are produced.

Puccinia sorghii. Schw. Maize Rust

Uredo and teliosori upon the leaves and bracts, the former small, light brown sori, soon rupturing the epidermis; teliosori dark brown, the uredospores are 1-celled, round or more elongated and spiny, the stalk is detached, the spores measure 23-38 \times 20-26 μ , teliosori are elongated dark brown or black being broadly elliptical and 2-celled, 30-52 \times 16-24 μ , the apex may be thickened and somewhat pointed. These spores preserve their vitality for some time, but are dormant through the winter. In the spring each cell may germinate by producing a tube, known as the promycelium, which bears lateral bodies

known as sporidia. According to Dr J C. Arthur, it is undoubtedly connected with an aecidium on *Oxalis corniculata*.

Distribution and hosts. Common wherever corn is cultivated and according to Carelton, also upon teosinte.




Fig. 54. Corn rust (*Puccinia Sorghi*) on corn. Winter spores.

Uromyces. Link. Clover Rust

Aecidiospores in cup-like bodies with an evident pseudoperidium; uredosori powdery; uredospores 1-celled with several evident germ pores, teleutosori powdery; teleutospores 1-celled, separate, pedicellate, apex with a single germ pore, sporidia flattened on one side. About 250 species widely distributed. Many of the species produce serious diseases of cultivated plants, as *Uromyces pisi* upon the pea, the alfalfa rust, (*U striatus*.) and the bean rust, (*U appendiculatus* (Pers. Lev.) There are many other species found upon our wild plants. Some of these, when they occur upon forage plants, may cause mycotic stomatitis.

Uromyces Trifolii. (Hedw.) Lev.

Aecidia in circular areas of pale colored spots, pseudoperidia short, cylindrical flattish, edges, whitish, torn, spores sub-globose or irregular, finely roughened, pale orange, 14-23 μ in diameter; uredosori pale brown, round, scattered, surrounded by the torn epidermis; spores round or ovate, roughened; 20-26x18-20 μ with 3 or 4 germ pores, color brown, teleutosori small round almost black; long covered by the epidermis, spores globose, elliptical or sub-pyriform occasionally with wart-like swellings on the summit 15-20x22-30 μ , small, dark brown in color, pedicels long.

Distribution. Widely distributed upon various clovers, especially red clover and the white clover. So abundant is this fungus at times that the plants are covered with the brown dusty material. Miss Howell reports it as very severe in the state of New York at times. The writer commenting on this fungus some years ago, said

The fungus did not occur until August and only on the "rowen" or "aftermath." Later it was found quite abundantly on the campus and College Farm. So severely did it attack some of the plants, especially the stems and leaves, that in touching the plants, the hands became covered with brown spores.



Fig. 67 Clover Rust. *Uromyces Trifolii*. (Hedw.) Lev. 1. Aecidium spores above two cluster cups in which the aecidiospores are found. 2. White clover leaf showing the distortions produced by the aecidium stage. 3. Red clover leaf showing clusters of uredo spores. 4. Uredo spores. 5. Tetrads of spores. 6. An uredo cluster more magnified than in 2. Figs. 1, 2 and 3 after Miss Howell. Remainder by Miss King.

How long the fungus has affected clover plants in this country and especially in Iowa is not known.

Poisonous properties Clover rust has been suspected of being injurious to cattle. Dr John R Mohler of the Bureau of Animal Industry writes as follows with reference to mycotic stomatitis:

Several attempts have been made by the writer to determine the exact cause and also to transmit the disease to other animals by direct inoculation, but with negative results. Suspicion, however, has been directed by various observers to the *Uromyces* and the red and black rusts that occur in clovers. These fungi cause very severe irritation of the lining membrane of the mouth, producing sometimes a catarrhal, at other times an aphthous, and occasionally an ulcerous stomatitis.

Considerable irritation of the nose and throat is experienced when rusty oats and wheat are threshed. Virchow records a case of severe inflammation of the nose of an old lady in which he found a great deal of *Puccinia graminis*



Fig. 68. 1. *Toxistoma* (C. M. F.) ex. Soc. *Hymenogaster* sp. 2. *Pleurocybe* *quercina*. 3. *Marasmius* *eximius*. 4. *Tricholoma* *eximius*. 5. *Clavaria* *angulata*. 6. *Tricholoma* *eximius*. 7. *Tricholoma* *eximius*. 8. *Pleurocybe* *quercina*. 9. *Tricholoma* *eximius*. 10. *Tricholoma* *eximius*. 11. *Tricholoma* *eximius*. 12. *Tricholoma* *eximius*. 13. *Tricholoma* *eximius*. 14. 6-9 after Welstead. 10-12 after Haape.

AUTOBASIDIOMYCETES

The basidia of the hymenium more or less club-shaped, undivided sterigmata usually 4, occasionally 2, 6, or 8, coming from the apex of the basidium. The *Dacryomycetinae* with long club-shaped basidia and two long sterigmata. Basidiospores large, spores divided before germination, includes the group *Dacryomycetinae*, and an unimportant group, the *Exobasidinea* or small gall parasites containing the *Exobasidium* which occurs upon the cranberry and blueberry. The third group, *Hymenomycetinae*, contains a number of poisonous plants and will be treated more in detail.

HYMENOMYCETINÆ

Mycelium of septate hyphae, loose or delicate in texture or made up into strands or hard masses. Hymenium at the time of spore formation free, the basidia form a definite layer or hymenium which may cover the whole surface of the fruiting body, or may be restricted to a definite portion. The fruit is made up of more or less closely compacted threads, hyphae grown together, or it may be delicate and somewhat ephemeral, the hymenium may be free or gymnocarpous or covered from the beginning. The covering is called the veil which consists of a layer of threads extending from the margin of the cap to the stem, or the veil may envelop the entire plant. The volva is an envelope which in the young stage completely covers the plant; at maturity it is left in the form of a cup at the base of the stem or distributed from the cap to the base of the stem. The annulus is a ring around the stem formed by




Fig. 69. Cross section of Bracket Fungus *Polyporus ignarius*. A. Fungus threads, hyphae between the pores. J. Hymenium surrounding the pores, a number of basidia with spores. After Luessen.




Fig. 69A. Mushroom (*Agaricus campestris*). To the left a matured plant and to the right a young plant. (Strasburger, Noll, Schenck and Schimper)

the inner or partial veil. Occasionally cystidia form in *Coprinus*, chlamydo-spores are seldom produced.

POLYPORACEAE

Hymenium usually below, porous, tubular, honey-combed, reticulate or of concentric plates, spores produced on the inner surface of the pores. A family consisting of 2300 species, of wide distribution. Some are edible like *Boletus* (*Boletus edulis* B. *scaber*) and others of this genus, as the *Fistulina* (*Fistulina hepatica*) also known as the vegetable beefsteak, and the Sulphur Polyporus (*Polyporus sulphureus*) when young. Several members are destructive wound parasites of trees. Among these are *Polystictus versicolor*, the common Bracket fungus (*Pomes applanatus*), and *Trametes radiciperda* found on the roots of conifers, and producing death. The dry rot fungus, *Merulius lacrymans*, is widely distributed and destructive to buildings.

Boletus. Dill. *Boletus*

Soft or fleshy, the stratum of the tubes on the lower surface of the cap easily separated. They are nearly all found growing on the ground and have the stem attached centrally to the cap. Quite a number of species are edible, some are bitter and some are poisonous. A small genus of 200 species found both in Europe and North America. The *Boletus edulis*, according to European authority, is one of the most desirable of edible fungi. Professor Atkinson lists this as one of the edible North American species. The *B. scaber*, also a North American species, is according to Professor Peck, first class, but several species are poisonous and bitter. The *B. luridus* is regarded as poisonous. The fact that a species turns blue when the plant is cut, should not be regarded as indicative of its poisonous qualities, for this is due to the oxidation process of the fat in contact with the air.

Boletus fellens. Bull

Pileus fleshy, convex above, glabrous or nearly so, grayish-brown, buff-brown, reddish-brown or tawny, flesh, white, taste bitter, tubes long, convex in the mass to mature plants, at first whitish, becoming pale flesh color, stem equal or tapering upwards, usually reticulated at the top only, rarely wholly reticulated, commonly a little shorter than the pileus, spores oblong-fusiform, pinkish, .0005 to .0007 inch long.

Distribution. Widely distributed in woods and open places, found upon decayed stumps.

Poisonous properties. Prof Peck says

The Bitter boletus takes its name from the bitter flavor which its flesh persistently maintains. It is a common species, and one easily recognized by its reticulated stem and flesh-colored tubes taken in connection with its bitter taste.

The cap is rather thick, dry and smooth, but quite variable in color. This is generally some shade of brown tinged with red or yellow. The flesh is white, but when cut or broken and exposed to the air it sometimes assumes a pinkish tint.

The mass of tubes is generally somewhat convex in the mature plant, though it may be plane in the young plant. This also sometimes assumes a pinkish stain when bruised.

The stem varies greatly in length and thickness, and is sometimes crooked and deformed. It is usually reticulated at the top only.

The taste of the flesh in this *Boletus*, as well as in many species of *Lactarius* and *Russula*, is an important aid in the specific identification. In testing fungi for this purpose care should be taken to select only fresh, sound specimens, and the part tasted should not be swallowed.

Mr Hurd states that this species is not poisonous. No amount of cooking according to this author, will destroy the bitter flavor.



Boletus satanas, Lenz

Pileus large, yellowish-brown on its upper surface; lower surface blood-red at first, later becoming orange red, stalk yellow to reddish-purple with red reticulate markings, spores brownish ovate. Rank and unpleasant taste.

Distribution. In Europe and North America.

Poisonous properties. Said to be extremely poisonous. The *B. luridus* along with several poisonous species is eaten in Northern Russia. Ford states that these species may occasionally be the cause of transient disturbances in man and may occasionally cause fatal intoxication.

AGARICACEAE

Pileus generally expanded, stipe generally with central attachment, or nearly so, lateral, or sessile, gills simple or branched or anastomosing usually on the lower surface, lamellae folded or veined, radiating from the point of attachment; lamellae bear the basidia which in turn bear the four spores or rarely two, cystidia often present. A large order separated chiefly by the color of the spores. The *Melanosporeae* have their spores brown, purplish brown or black, in the *Ochrosporeae* spores are yellowish brown or rusty brown, in the *Rhodosporeae* spores are rosy pink, in the *Leucosporeae*, spores are white, whitish or pale yellow. Many species of the family, like the cultivated mushroom, (*Agaricus campestris*), the field mushroom, (*Agaricus arvensis*), the shaggy-mane (*Coprinus comatus*, Fr.), *Lepiota procera*, and others, are edible. The *Rastus gongylophora* of Southern Brazil, is cultivated by the leaf cutting ants for food. No invariable rule can be laid down for the poisonous species. Many of the *Leucosporeae* are edible, but many are deadly poisonous. A few of the poisonous species are described later.

Amanita. Pers. *Amanita*

The young plants covered by a membrane which in the button stage is more or less free with the surface of the pileus; later when the stem elongates




Fig. 70. Part of the hymenium of one of the Agaricaceae. *a*. Sub hymenial layer. *b*. Basidium. *c*. Sterigmata. *d*. Spores from basidium. *e*. Paraphyses. *e*. Cystid. After Bonn text book.

the volva is ruptured, stipe fleshy; volva and annulus present

In some species the remains of the ruptured volva persist, forming a kind of cup or sheath. In others they occur in the form of small scales or warts on the cap.




Fig. 71 Fly Agaric or Fly Amanita (*Amanita muscaria*). a. Mature plant. b. Top view of cap with scales. From U. S. Dept. of Agr.

Amanita muscaria, L. Fly amanita. Fly agaric

Pileus nearly flat at maturity, warty slightly striate on the margin, yellow to orange red, cap 3-8 inches broad, gills white or nearly so stem 4-6 inches long, $\frac{1}{2}$ inch in thickness, cylindrical, hollow, bulbous thickened at the base, which is more or less scaly from the fragments of the ruptured volva spore broadly elliptical, white Dr Farlow gives the following excellent description of this fungus

The fly agaric (*Amanita muscaria*), so called because decoctions of it are used for killing flies, is in most places, at least in the northern and eastern parts of the country, a common species—often a good deal more abundant than the common mushroom. It is found during the summer along roadsides, on the borders of fields, and especially in groves of coniferous trees. It prefers a poor soil, of gravelly or sandy character, and occurs only exceptionally in the grassy pastures preferred by the common mushroom. It grows singly and not in groups, and attains a large size, being one of the most striking toadstools. It



differs from the common mushroom in having gills which are always white, never pink or purple, and in having a hollow stem which is bulbous at the base and clothed with irregular, fringed scales on all the lower part. The pileus varies in color from a brilliant yellow to orange and a deep red, the yellow and orange being more frequent than the red. The surface is polished and has scattered over it a larger or smaller number of prominent, angular, warty scales, which can be easily scraped off. The gills and stalk are white, and there is a large membranous collar, which hangs down from the upper part of the stem. The general appearance together with the color of the pileus and gills noted above, are such that it is difficult to conceive how anyone who has ever seen a common mushroom or read a description of one could mistake this fly agaric for the mushroom. Nevertheless, in the writer's experience, no fungus is so often collected by children on the supposition that it is the common mushroom, and it is to the fly agaric that recent cases of poisoning in Washington, D. C., were due.

Distribution. Widely distributed in Europe and North America. Professor Coville, in speaking of this species after the death of Count Achilles de Voechj, and Chung Yu Ting, says.

The fly amanita is one of the largest, headstrong, and most dangerous of our mushrooms, and is the one whose character has been the most fully studied of all the poisoning species. It is abundant about Washington in the fall, growing in pine woods, a favorite situation in these woods being the vicinity of abandoned hog beds. The specimens that caused the death of Count de Voechj came from a pine tract about a mile west of Fort Myer, between Ball's Crossroads, and Columbia Pike.

Poisonous properties. The chief active poisonous principle of the fly amanita is an alkaloid called *muscarin*, but other poisonous substances, the chemical nature of which is not yet fully known, also occur in the plant.

Professor Atkinson, in discussing the Toxicology of the species says as follows:

The substance, *Cholein*, is of wide occurrence in the animal and vegetable kingdoms. It has been isolated from *Amorpha canescens*, *A. vernalis*, *Boletus luridus*, and *Heliopsis scabra*. It is not very toxic, but on uniting with oxygen it passes over to *muscarin*. According to Robert the substance formed from *cholein* on the decay of the mushrooms containing it is not *muscarin* but a very closely related alkaloid, *neurin*. This transformation of a comparatively harmless alkaloid to an extremely deadly one simply by the partial decay of the plant in which the former is normally found, emphasizes very much the wisdom of rejecting for table use all specimens which are not entirely fresh. This advice applies to all kinds of mushrooms, and to worm-eaten and otherwise injured, as well as decayed ones. *Neurin* is almost identical in its physiological effects with *muscarin* which is described below.

Muscarin is the most important because the most dangerous alkaloid found in the mushrooms. It is most abundant in *Amorpha canescens*. It is also found in considerable quantity in *Amorpha vernalis*, and to some extent, but still very dangerous extent in *Boletus luridus* and *Russula emetica*. It is quite probably identical with *boletarin*, isolated from *Amorpha phalloides* by Boudier. *Muscarin* is an extremely violent poison, .001 to .005 of a gram (.06 grains) being a very dangerous dose for a man. Like other constituents of mushrooms, the amount of *muscarin* present varies very greatly with varying conditions of soil and climate. This, indeed, may account for the fact that *Boletus luridus* is regarded as an edible mushroom in certain parts of Europe, the environment being such that little or no *muscarin* is developed.

Cases of Mushroom poisoning are frequent in some countries. Gaillard estimated the number of deaths in France at about 100 cases. Among the Americans deaths are not so numerous, although Palmer of Boston, has found 33 cases with 4 deaths.

Inoko of Japan, reports 481 cases in 8 years. The peasants of the Caucasus prepare an intoxicating beverage from *Amorpha muscaria* from which many individuals die.

Muscarin acts on the nerve centers, but cases seldom terminate fatally.

Helvella esculenta owes its toxicity to *helvellic acid* $C_{10}H_{20}O_7$. Very few deaths have been reported in recent years.

According to Kobert, *Amanita muscaria* contains, besides *choin* and *muscarin*, a third alkaloid, *fungus atropin*, (*pils-atropin*), this alkaloid, like ordinary atropin, neutralizes to a greater or less extent the muscarin. The amount of *pils-atropin* present varies, as other constituents of mushrooms vary, with varying conditions of soil, climate, etc., and it may be that in those localities where the *Amanita muscaria* is used for food the conditions are favorable for a large production of *pils-atropin*, which neutralizes the muscarin, thus making the plant harmless. Be this as it may, *Amanita muscaria* is deadly as ordinarily found. It is undoubtedly used quite largely as food in parts of France and Russia, and it has been eaten repeatedly in certain localities in these countries without harm.

Ford suggests, on clinical grounds, that it may not be the only poison present because even when this drug is completely neutralized by its physiological antidote, atropin, the patient, who has eaten *Amanita muscaria*, sometimes dies.*

The alkaloid *muscarin*, a tasteless alkaline substance with a tobacco-like odor, causes the contraction of pupils, *amoufin* $C_8H_{11}NO_8$ is an isomer of *choin*, and yields muscarin with muric acid and cetraric acid, $C_{10}H_{18}O_8$. Muscarin has been obtained synthetically from *choin*. It does not, however, produce quite the same symptoms.

Amanita Frostiana. Peck

Pileus convex to expanded, bright orange or yellow, warty, sometimes nearly or quite smooth, striate on the margin; lamellae white or tinged with yellow; stem white or yellowish, stuffed, bearing a slight, sometimes evanescent annulus, bulbous at the base, the bulb slightly margined by the volva; spores globose; 7.5-10 in diameter. From the character of the poisons it is quite distinct from the *A. muscaria*.

Distribution. New York to North Carolina

Poisonous properties. Professors Peck and Atkinson both list it as poisonous. Ford found an hemolysin of low grade intensity. Heated extracts were without action upon animals. Schmiedeknecht found a poison

Amanita phalloides. Fr. Death Cup

Pileus smooth, fleshy, viscid, greenish, brown or olive to amber, cap 3-5 inches broad, frequently free from remnant of volva. lamellae white. stem 3-6 inches long, annulate, spores globose, white. Prof. Atkinson says "The presence or absence of these scales on the cap depends entirely on the way in which the volva ruptures. When there is a clean rupture at the apex, the pileus is free from scales, but if portions of the apex of the volva are torn away they are apt to remain on the cap.

Dr. Farlow gives the following excellent description of this fungus

It is rather common and grows singly in woods and on the borders of fields, rarely appearing in lawns, and is not preëminently an inhabitant of grassy pastures, like the mushroom. It prefers a damper and less sandy soil than that chosen by the fly agaric. The pileus is often a shining white, but may be of any shade, from a pale dull yellow to olive, and when wet is more shiny than the mushroom or the fly agaric. It has no distinct scales and only occasionally a few membranous patches on the pileus. The gills and stalk are

* Sci. 20: 97-108





Fig. 72. Death Cup (*Amanita phalloides*) one-half natural size. From U. S. Dept. Agr.




Fig. 71. Stinkhorn, *Phallus impudicus*. Common in some vineyards and fields.

white, and the latter has a large ring like the fly agaric, and is hollow or when young is loosely filled with cottony threads, which soon disappear. The base of the stalk differs from that of the fly agaric in being more bulbous and in having the upper part of the bulb bordered by a sac-like membrane, called the volva. The volva is often of considerable size, but more frequently it is reduced to a membranous rim. In this species the stalk is longer and slenderer in proportion to the diameter of the pileus than in either the fly agaric or the common mushroom, and is buried rather deep in the soil or dead leaves, so that it often happens that the bulb is broken off and left behind when the fungus is gathered.

The following differences between the edible and two poisonous species are noted by Dr. Farlow:

(1) The common mushroom has a pileus which is not covered with wart-like scales; gills which are brownish purple when mature, a nearly cylindrical stalk, which is not hot-

low, with a ring near the middle, and without a bulbous base sheathed by a membrane or by scales.

(3) The fly agaric has a pileus marked with prominent warts; gills always white; a stalk, with a large ring around the upper part, and hollow or cottony inside, but solid at the base, where it is bulbous and scaly.

(3) The deadly agaric has a pileus without distinct warts; gills which are always white, and a hollow stalk, with a large ring, and a prominent bulb at the base, whose upper margin is membranous or bag-like.

(4) Other minor points of difference are the different places in which these species grow and also the colors, which, although they vary in each case, are brilliant yellow or red in the fly agaric, white varying to pale olive in the deadly agaric, and white usually tinged with a little brown in the mushroom.

(5) A word should be said as to the size and proportions of the pileus and stalk in these three species. In the mushroom the pileus averages from 3 to 4 inches in breadth, and the stalk is generally shorter than the breadth of the pileus and comparatively stout. The pileus remains convex for a long time, and does not become quite flat-topped until old. The substance is firm and solid. In the fly agaric the pileus, at first oval and convex, soon becomes flat and attains a breadth of 6 to 8 inches and sometimes more. The stalk has a length equal to or slightly exceeding the breadth of the pileus, and is comparatively slender than in the common mushroom, but nevertheless rather stout. The substance is less firm than in the common mushroom.

(6) The pileus of the deadly agaric is thinner than that of the common mushroom and from being rather bell-shaped when young, becomes gradually flat-topped with the center a little raised. In breadth it is intermediate between the two preceding species. The stalk usually is longer than the breadth of the pileus, and the habit is slenderer than in the two preceding species. All three species are pleasant in the taste, which shows that one cannot infer that a species is not poisonous because the taste is agreeable. The fly agaric has scarcely any odor. The two other species have certain odors of their own, but they can not be described.

Distribution. Widely distributed in Europe and North America in woods, groves and pastures.

Poisonous properties Professor Peck says

The Poison *amanita* is very variable in the color of the cap, and yet is so definite in its structural characters that only the most careless observer would be likely to confuse it with any other species. There is, however, a sort of deceptive character about it. It is very neat and attractive in its appearance and looks as if it might be good enough to eat. This appearance is fortified by the absence of any decidedly unpleasant odor or taste, but let him who would eat it beware, for probably there is not a more poisonous or dangerous species in our mycological flora. To eat it is to invite death.

Professor Atkinson says

Since the *Amanita phalloides* occurs usually in woods, or along borders of woods, there is little danger of confounding it with edible mushrooms collected in lawns distant from the woods and in open fields. However it does occur in lawns bordering on woods, and in the summer of 1899 I found several of the white forms of this species in a lawn distant from the woods. This should cause beginners and those not thoroughly familiar with the appearance of the plant to be extremely cautious against eating mushrooms simply because they were not collected in or near the woods. Furthermore sometimes the white form of the deadly *amanita* possesses a faint tinge of pink in the gills, which might lead the novice to mistake it for common mushrooms. The bulb of the deadly *amanita* is usually inserted quite deep in the soil or leaf mold, and specimens are often picked leaving the very important character of the volva in the ground, and then the plant might easily be taken for the common mushroom, or more likely for the smooth *Leptota* (*Leptota nana*), which is entirely white, the gills only in age showing a faint pink tinge. It is very important therefore, that, until one has such familiarity with these plants that they are easily recognized in the absence of some of these characters, the stem should be carefully dug from the soil. In the case of the specimens of the deadly *amanita* growing in the lawn on the campus of Cornell University, the stems were sunk to three to four inches in the quite hard ground.

The exact chemical nature of *phalloin*, an extremely toxic substance, is not certainly known, but it is generally conceded to be of an albuminous nature. That it is an extremely

deadly poison is shown by the fact that .0015 grain per 2 lbs. weight of the animal is a fatal dose for cats and dogs. It is the active principle of the most deadly of all mushrooms, the *Amanita phalloides*, or death-cup fungus. We quote again from Mr. Chesnut's account of phalloin and its treatment: "The fundamental injury is not due, as in the case of muscarin, to a paralysis of the nerves controlling the action of the heart, but to a direct effect on the blood corpuscles. These are quickly dissolved by phalloin, the blood serum escaping from the blood vessels into the alimentary canal, and the whole system being rapidly drained of its vitality. No bad taste warns the victim, nor do the preliminary symptoms begin until nine to fourteen hours after the poisonous mushrooms are eaten. There is then considerable abdominal pain and there may be cramps in the legs and other nervous phenomena, such as convulsions, and even lockjaw or other kinds of tetanic spasms. The pulse is weak, the abdominal pain is rapidly followed by nausea, vomiting and extreme diarrhoea, the intestinal discharges assuming the 'rice-water' condition characteristic of cholera. The later symptoms are persistently maintained, generally without loss of consciousness, until death ensues, which happens in from two to four days. There is no known antidote by which the effects of phalloin can be counteracted. The undigested material, if not already vomited, should, however, be removed from the stomach and intestines by methods similar to those given for cases of poisoning by *Amanita muscaria*."

Prof. Chesnut, writing in regard to the poisonous effect of this species, says that

The phalloin spoken of is one of the toxalbumins, an extremely virulent poison found in poisonous animals especially the rattlesnake. These toxalbumins are allied to those found in diphtheria and other diseases produced by bacteria.

Other species reported as poisonous or probably poisonous are *A. floccoccephala*, and *A. cothurnata*. Ford has shown that *A. spreata*, and *A. virosa*, *A. strobiliformis*, *A. chlorinasma*, *A. radicata*, *A. porphyria*, and *A. rubescens* are poisonous. The *A. verna*, a small spring form of *A. phalloides*, is also very poisonous.

Ford reports nearly 200 deaths since 1900 from this fungus in France, Germany Italy, and England. Ford gives the pathological changes described by Maschke to be as follows

1. Lack of post mortem rigidity
2. Widening of the pupils.
3. Failure of blood to coagulate and a cherry-red color
4. Ecchymoses and hemorrhages in the serous membranes and parenchymatous organs.
5. Dilation of the bladder with urine

Studor, Sabin and Schören found extensive necrotic and fatty changes in liver, kidney, heart and voluntary muscles. The amount of fat in the liver is nearly as great as in phosphorous poisoning.

Clinical symptoms. Often latent period of from 6 to 12 hours during which the victims remain quite well. They are suddenly seized with terrible abdominal pain, excessive vomiting and thirst. Diarrhoea may set in with mucous bloody stools, or there may be constipation. The paroxysm of pain may be so severe as to result in a peculiar hippocratic facies. The patients rapidly lose strength. In 3 to 4 days in children and 6 to 8 days in adults, coma develops, from which the patients cannot be aroused. Cyanosis and lowered temperature precedes the fatal exit. Ocular symptoms and convulsions do not ordinarily occur, but convulsions may be present on a terminal event. The mortality varies from 60 to 100 per cent.

Kobert obtained from *A. phalloides* a substance with marked hemolytic action, the dried extract dissolving ox blood 1 to 125,000. To this extract he gave the name *phalloin*, which he considered a *toxalbumin*.

Later this author* found what he thought was a poisonous alkaloid, "that all typical forms contained an alcohol soluble poison," that phalloin was occasionally absent.

Ford found that the extract of the fungus is a powerful hemolytic agent and quickly destroys the erythrocytes of guinea pig, fowl, pigeon, dog, goat, and man. This takes place at 37 degrees C. slower at lower temperatures. The corpuscles of sheep, beef, and swine are resistant. Raw and boiled milk act as an antidote — they are antihemolytins. Animals may be immunized by using non-lethal doses.

Since the above has been written there have come to hand several recent papers by Dr. W. W. Ford,¹ who states that he found muscarin in several "yellow Amanitas" found in New York and Massachusetts. The aqueous extract of *Amanita muscaria* first agglutinated and then slowly dissolved blood corpuscles. The agglutinin was heat resistant. The extracts produced hemolysis. The agglutinin is a glucoside. The *Amanita solitaria* also contains an agglutinin. The *Amanita frostiana*² contains a moderately hemolytic substance and free from resistant toxin and muscarin. By the same author the poisonous nature of a number of species is reported as follows. The *A. phalloides* produces a chronic intoxication in guinea pig, the animal dying in twenty-five days. The lesion is typical for amanita toxin. It is hemolytic for rabbit's corpuscles, in a dilution of 1:20. The poison from *A. tyrota* has a hemolytic strength of 1:200 in two hours and in dilution 1:100 at the end of 24 hours, and when heated to 60° C. kills guinea pigs in twenty-four hours, with signs of acute intoxication. The *A. sprete* contains hemolysin and toxin but in rather a low degree. It should be classed with the deadly poisonous mushrooms. The *A. porphyria*, *A. strobiliformis*, *A. radicata* and *A. chlorocoma* are all poisonous and contain a heat resistant substance which induces in animals a chronic intoxication, the *A. vittadini* and *A. rubescens* should also be included according to Robert.³

Dr. Ford in speaking of the poison in *A. phalloides* says:

"In a series of investigations published from the Johns Hopkins University it has now been shown that *Amanita phalloides* contain two poisons which for the sake of clearness we speak of as the amanita-toxin.⁴ The hemolysin is probably the same hemolytic substance which Robert had in his preparation of phalloin and the toxin is possibly identical with Robert's second poison. The hemolysin was found in every specimen of *Amanita phalloides* which has thus far been examined, and when obtained from the fresh plant is the most powerful hemolytic of vegetable origin known. Drs. Abel and Ford⁵ have shown that all enzymatic proteolysis can be removed from this substance by acetyl acetate in alkaline solutions and by freshly prepared metaphosphoric acid, and when thus freed from pepsin it continues to act upon blood corpuscles and gives the reaction of a glucoside containing a pentose. We have recently⁶ developed a method for the isolation and purification of this glucoside which has an activity of 1:300,000 in the pure state. Since its insensitiveness to heat and the digestive ferments the hemolysin is precluded from playing any important role in human intoxication. We are inclined to believe that the amanita-toxin is the active principle, and Schreiner and I⁷ have shown that this poison can be isolated by certain well defined methods. It also is one of the powerful organic poisons, four tenths of a milligram killing

* Sixteenth. Naturforschenden Gesellschaft Zurich 1909 26. Statement from Prof. Ford.

¹ The distribution of Poison in Amanita. The Jour. of Pharm. and Expt. Therapeutics, 275:264.

Notes on the Amanita-Toxin. Ford and Prouty. The Jour. of Pharm. and Expt. Therapeutics, 319.

² Jour. Inf. Dis. 4:437.

³ Lehrbuch des. Intoxikationen. Ed. 2. 617.

⁴ Ford Jour. Expt. Med. 8:437 May 26 1909.

⁵ Abel and Ford Jour. Biol. Chem. 11:73 Jan 1907.

⁶ Abel and Ford Arch. f. Exp. Path. u. Pharmacol. Supplement-Band. Schmeideberg Verlag 1908.

⁷ Schmeideberg and Ford Jour. Biol. Chem. 3:279 Sept 1907.

a guinea-pig in twenty-four hours. The amanita-toxin contains no proteid, does not respond to any alkaloidal reagents, and on fusion with potassium hydrate gives off iodol and pyrol. At first thought to be a conjugate sulphate, I have recently found in association with Mr. Prouty that this opinion is incorrect. We hope to ascertain the more exact characterization of this poison shortly."

Lepiota. Fr. *Lepiota* Toadstool

Plant with fleshy stem which can easily be separated from the cap gills usually free from the stem, in some species the top of the cap breaks from the scales which adhere, volva absent. A small genus widely distributed. Some species are edible, *L. procera* is said to be excellent as food.

L. Morgani. Pk.

A large fleshy plant, sometimes a foot across the cap, with a thick stout stem and a ring removed a little distance from the gills, the pileus when fully expanded, whitish, with dark scales, the spores and gills greenish.

Distribution. From Ohio southward and westward in grassy places, sometimes forming large fairy rings.

Poisonous properties. This plant is quite harmless to some people but to others it causes very unpleasant symptoms. It should be eaten with caution.

Russula. Pers. *Russula*

Cap red, purple, violet, pink, blue, yellow, or green; pileus fleshy, convex readily expanded and at length depressed, stem brittle, stout and smooth spongy within and confluent with the cap.

Russula emetica. Fr.

Pileus fleshy, quite viscid, expanded, polished, shining, oval or bell-shaped when young, rose-red to yellow or even purple, margin furrowed flesh white, gills free, equal broad, distinct and white, stems stout, solid, or occasionally spongy; spores spherical.

Distribution. Widely distributed in North America. Found in pastures and under trees. Readily distinguished by viscid cap and color. Mr. Hand states that it is easily recognized by its acid taste and free gills.

Poisonous properties. Mr. McIlvaine says that he has repeatedly eaten them and referred to a number of others who have also eaten them without any bad results, but Hand thinks that their acid taste is against their use or rather cautions their use. Prof. Ford states that they cause profound gastrointestinal disturbances, such as attacks of vomiting and diarrhoea, recovery only after thorough emptying of the stomach.

Volvaria. Fr. *Volvaria*

Universal veil forming a perfect volva, separate from outer part of the pileus, stem readily separated from pileus, gills free, at first white, then pink, and then reddish, and soft.

Volvaria bombycina. (Pers.) Fr.

This plant has a silky lustre; pileus is from 6 to 8 inches broad, globose, becoming bell-shaped, convex and somewhat umbonate, flesh white, gills crowd-

ed and flesh colored, stem is 6 to 8 inches long, tapering upward, spores rosy, smooth in masses and elliptical; volva large and somewhat membranaceous.

Poisonous properties. According to many authorities, this plant is edible, and it is likely that this and many other species can be eaten without serious trouble, although Gillot,* states that several species of this genus have caused death when eaten, though nothing is known of poisonous principle.

Inocybe Fr

In the genus *Inocybe* there is a universal veil which is fibrillose in character, and more or less closely joined with the cuticle of the pileus, and the surface of the pileus is therefore marked with fibrils or is more or less scaly. Sometimes the margin of the pileus possesses remnants of a veil which is quite prominent in a few species. The gills are adnate, or sinuate, rarely decurrent, and in one species they are free. It is thus seen that the species vary widely, and there may be, after a careful study of the species, grounds for the separation of the species into several genera. One of the most remarkable species is *Inocybe echinata* Roth. This plant is covered with a universal veil of a sooty color and powdery in nature. The gills are reddish purple, and the stem is of the same color the spores on white paper of a faint purplish red color.

Inocybe infida

This is slightly larger than *Panacolus papilionaceus*, with semiorbicular cap surmounted by a prominent nipple, which is dark reddish-brown, while the rest of the upper surface is light tawny brown. The upper surface also differs from that of the non-poisonous kind in being silky-scaly and shining. The lower surface differs in being much lighter, pale yellowish instead of brownish-black, and the spore print is about the color of oak wood.

Poisonous properties. Dr William A. Murrill has recently contributed an account on the poisoning from *Inocybe infida*, a plant which closely resembles the *Panacolus papilionaceus*. It appears that Dr Deming of West Chester, who poisoned himself and other members of the family, describes the following symptoms: The fungi were gathered in the morning just before dinner. They were stewed and served on toast at one o'clock, he ate about half a slice of toast with mushrooms, drank some tea, and ate one-half a stuffed egg, with lettuce and mayonnaise dressing and after dinner smoked one-half a cigarette. Soon after he began to feel "queer," then there followed a fullness in the head and a rapid heart action as if he had taken nitroglycerin, this was followed by a sweat, his clothing becoming wet, and at the same time there was no nausea or prostration, his mind became a little but confused. He then washed out the stomach, took castor oil and before the oil operated there was pressure and almost pain in the lower bowel. By evening he was as well as ever except somewhat exhausted. It appears that four other persons were affected with disagreeable symptoms from the eating of the mushroom.

Dr Deming says "In my case the beating of the heart, fullness of the head and sweating were very marked, though I ate about half as much as the others."

Dr Murrill says that there is nothing to suggest an irritating poison and that it is probably not narcotic.

* Etude medicale sur l'empoisonnement par les champignons. Lyon, 1900.

PHALLINEAE

Mycelium consists of branched strands matted together, from this is produced an oval body consisting of an outer wall, the peridium, and an inner peridium, between the two is a layer of gelatinous material, the outer portion of the oval body forms the volva, the central portion pushes through the peridium with a long cellular stalk, the upper one bearing the cap-shaped gleba, the spores are brown on club-shaped basidia, surrounded by a mucilaginous material giving off an offensive odor. This sub-order contains the *Clathraceae* and *Phallaceae*. The *Phallus impudicus* and the *Mutinus caninus* have been regarded as suspicious.

CLATHRACEAE

Receptacle latticed or irregularly branched, gleba enclosed by the receptacle. The following genera of this order are known to occur in the United States, chiefly in the southern states *Clathrus*, *Phallogaster*, *Symbium*, and *Anthurus*. Dr. Farlow* is authority for the report from Gerald McCarthy to the effect that in North Carolina hogs had been killed by eating *Clathrus columnatus* which a correspondent, Mr. G. W. Lawrence found growing in oak woods near Fayetteville. The animals died within twelve or fifteen hours after eating the fungus. According to Gillet, hogs are poisoned by these and by Phalloideae.

PHALLACEAE

Receptacle tubular or cylindrical with an external gleba. The common Stinkhorn *Phallus impudicus* has a thick hollow stalk of whitish color perforated with pores the upper part is honey-combed, resembling the moral. During the early stages, an egg shaped body may be seen coming from a mass of white mycelium. The egg-shaped body is more or less mucilaginous and contains the stalk and gleba, the latter becomes exposed later. Flies, attracted by the carrion-like odor and mucilaginous material of the gleba, scatter the spores and, apparently, are not poisoned. The fungus, however, is usually regarded as poisonous as are several related genera and species such as *Mutinus caninus*. The common Stinkhorn (*P. impudicus*) was formerly used as a salve in gout.

HYMENOGASTRINEAE

This contains the family Hymenogastaceae. The sub-order Lycoperdineae includes two families, *Tylostomataceae* and *Lycoperdaceae*.

LYCOPERDACEAE

Fructing bodies globular, oval or pear-shaped, solid and fleshy, often of great size, before maturity, a dense white mass of homogenous hyphae occurs, the fruit is surrounded by a peridium, in some cases double, the interior is made up of branched threads called the capillitium, containing the spores, fruiting bodies break open in various ways at maturity. This group contains several interesting families. Many of the *Lycoperdaceae* are well known, among these are the Earth-star (*Geaster*), the *Lycoperdon giganteum* and

* Farlow, W. G. Poisonous Nature of *Clathrus columnatus*. Bot. Gaz. 15:45-56. See also Hallett, B. D., Rept. N. J. Agri. Exp. Sta., 1894:417.




Fig. 74. 1. *Puffball* and their Allies. *Cantharellus*. 1. *Greater Amethyst*. 2. *Outer peridium* of 1. 3. *Section* of 1. 4. *Section* of 1. 5. *Section* of 1. 6. *Section* of 1. 7. *Section* of 1. 8. *Section* of 1. 9. *Section* of 1. 10. *Section* of 1. 11. *Section* of 1. 12. *Section* of 1. 13. *Section* of 1. 14. *Section* of 1.

Dr. Miquel lists the *Lycoperdon Bovista* as poisonous, thus and *Lycoperdon cyathiforme* are edible when fresh, but poisonous when the plants are mature.

NIDULARIACEAE

This is allied to the above and contains the Bird's Nest fungus. (*Crucidarium vulgare*), which occurs upon wood and manure, and the *Cyathus striatus*. The false truffle (*Scleroderma vulgare*) belongs to an allied order.

PHOTIDACEAE

Mycelium many celled, branched, reproduction both sexual and asexual, spores known as ascospores, limited in number. Arranged in two divisions, the *Hemiasci* and the *Eusci*.

Hemiasci

Parasites or saprophytes, reproduction generally asexual in fertilization, the contents of the antheridium and the oögonium fuse.

HEMIASCALES

An unimportant group with three orders, *Ascoideaceae*, *Protomyces*, upon the members of the carrot family.




Fig. 15. Fertilization of *Pyrenopeziza confusa*. 1. Three oögonia (o) with fertilizing processes (f). 2—Antheridia. 2. Oögonium after fertilization, with numerous nuclei. 3. Part of fruiting body, the ascogonium forming hyphae (ar), (a) antheridium, (o) oögonium. 1-3 greatly magnified. After Harper.

The family *Monascaceae* contains one fungus which has been found in mouldy corn and siage in Iowa, the *Monascus purpureus* Went. It is related to the *M. heterosporus* (Hartz) Shroter, which was found by Hartz in a soap factory. The coloring matter from *M. purpureus*, known as "ang-quac," is used in Eastern Asia as a pigment, being produced by the growth of the fungus on rice. The fungus consists of a mass of septate hyphae, producing conidia and perithecia with numerous asci, the ascospores are from 5-6.5 μ , in diameter. The

details of the structure of this fungus have been given by Olive,¹ Barker,² and Ikeno³ and in a paper to be published by Dr Buchanan. Dr Buchanan found this species in spoiled corn silage, which was responsible for the death of several horses in Iowa. This species possibly has been the cause of the disease, this fungus occurring only where air had access to the silage. The fungus found by Hartz produces a mycelium similar to the preceding with thick-walled swellings and color white or carmine red, conidia ellipsoidal, spherical, obovate, of two kinds, the smaller 2.5-3 μ to 7-8 μ , occurring in chains or singly, the larger occurring singly 9-11 μ in diameter, and arising from lateral branches, sporangia from short lateral branches are spherical 40-53 μ in diameter, many spored the sporangia are surrounded by branched hyphae, ascospores spherical or oval, colorless 4-5 μ in diameter, conidia and hyphae contain a carmine red pigment *physomycin*.




Fig. 75A. Corn Silage fungus (*Monascus purpureus*). 1, 2, 3, Conidiophores with conidia. 4, germinating conidium, 5, sterile hypha covering of perithecium sending out branches. these are sometimes tipped with conidia. 6, optical section of mature perithecium, spores still within asci. Found in corn silage by Dr Buchanan.

Eusci

Asci with definite number of spores, usually 2, 4, 8, 16, 32, seldom, but occasionally ∞ -celled.

EUASCALES

Contains the yeast plant, peach curl, plum pocket, ergot blue mould, powdery mildews, etc

PROTOASCINEAE

Asci single, in one group, without distinctive development of the mycelium, in the other with a distinctive mycelium bearing the asci with their spores.

¹ Annals of Bot. 17:167, pl. 12 & 13.

² Bot. Gazette. 39:56.

³ Ber. deutsch. Bot. Gesellsch. 12:259.

SACCHAROMYCETACEAE.

Vegetative cells single or in small groups, mycelium usually not evident, reproduction, by budding; ascospores, usually 4, produced in the cell, occasionally 8, 3, 5, or seldom 1.

The Saccharomycetes are fungi important in the process of fermentation. It is only in recent years that any parasitic species has been recognized. Metchnikoff, in 1884, found a parasitic yeast *Monospora bicuspidata* in Daphnids. Raum and Neumayer in 1891 declared yeasts were pathogenic. Busse, 1894, demonstrated that certain yeasts were pathogenic. Tokishige about the same time observed a yeast pathogenic for horses. Sanfelice isolated from the cancerous-like growth of an ox a *Saccharomyces* which was pathogenic for guinea pigs. The same author found another species in pigeons. Lydia Rabinowitsch studied 50 species of yeasts, of which 7 proved to be pathogenic. In 1895 Prof. Curtis found the second case of *Saccharomyces* in a young man clinically the disease resembled a myxosarcoma. It is doubtful whether these forms are true yeasts. Some of these appear to be Hyphomycetes rather than Saccharomycetes. I have therefore discussed these under the form genus *Oidium*.




Fig. 76. Yeast, *Saccharomyces mycoderma*. A. Process of germination. B. Mycelium budding in a weak nutrient solution. C. (a) Yeast-like form budding b) long cells.

Saccharomyces. Meyen. Yeast

Vegetative cells spherical, ellipsoidal, oval or pear-shaped, occasionally elongated mycelial like; asci spherical, ellipsoidal or cylindrical with 18 ascospores 1-celled spherical or ellipsoidal. About 40 species. The *S. apiculatus*, Rees, is important in the fermentation of fruit. The *S. ellipsoideus* causes the fermentation of wine. The *S. mycoderma*, Rees, forms a white mass on cider wine, cucumbers, etc., and prepares the way for the acetic acid fermentation. The *S. kefir*, Beyerinck, along with *Bacillus acids-lactici* and other bacteria is found in Kefir grains. *S. glutinus* Fres., the pink yeast, is found growing on nutrient media in laboratories.

Saccharomyces cerevisiae, Meyen. Common yeast

Vegetative cells, spherical or oval, 8-10x8-12 μ singly, or in several budding chains with one or more vacuoles, asci spherical or short elliptical 11-14, generally with 4 ascospores tetradform. It produces a white growth on gelatine and potato, does not liquify the gelatine, causes fermentation of grape sugar maltose and cane sugar. The biology of the fermentation of beer is as follows: Barley, which is ordinarily used for this purpose, is allowed to germinate; during the process of germination the starch, by means of diastase is converted into sugar, the sugar being afterwards removed with the water, this sugary fluid is then placed in large vats in dark rooms at a comparatively low temperature; the yeast plant is added and fermentation starts.

The fermentation of sugar is due to an enzyme found in the yeast plant, to which Buchner has given the name of Zymase. This enzyme breaks the sugar into alcohol and carbon dioxide. It is not necessary to have the living organism present to produce this fermentation, as a quantity of the yeast extract mixed with the solution of fermentible sugar will produce at the end of some days a small amount of alcohol. The enzyme decomposes very rapidly. Reynolds Green, in his book on fermentation, says

From these researches it appears certain that the production of alcohol whether in the presence or absence of oxygen is brought about by the activity of an enzyme. Its secretion by the cells of yeast attends the ordinary nutritive processes as well as the abnormal decompositions set up by incipient asphyxiation. The latter condition induces its formation in other parts of plants. The absence of oxygen stimulates the protoplasm of the cells to secrete it, the ultimate effect of its appearance being the liberation of energy as already stated.

Distribution. Widely distributed.

Poisonous properties. The chemical composition of alcohol is C_2H_5OH .

Different alcoholic drinks contain different percentages of alcohol. Ale and beers contain from 4 to 8 per cent together with bitters and malt extract, cider from 5 to 9 per cent sherry from 15 to 20 per cent.




Fig. 77 Yeast *Saccharomyces cerevisiae*. Ascospores in cells. Spores at f. Magn. 1000. After Hansen.




Fig. 7B. Yeast. *Saccharomyces cerevisiae*. 1. Single cell with vacuoles. 2. Cell budding $\times 1500$. 3. Numerous daughter cells $\times 1000$. 4. Cell with sporangia $\times 1200$. 5. *S. ellipsoideus*. 1-4 Modified from Luersen and Rees. 5 after Hansen.

Alcohol in its action is a germicide and when applied to the raw surface or wounds it is a stimulant and local anæsthetic, while in concentrated form it is an irritant and even caustic. When placed upon surfaces of the broken skin it causes cooling and contraction of the superficial blood vessels. When absorbed it hardens the tissues. Internally it causes a secretion of saliva and the heart is stimulated by the irritative action of alcohol. In large amounts it destroys the peptic ferment. Dr Winslow says:

Alcohol is essentially a heart stimulant and the most valuable one we possess. It makes the heart beat more forcibly and rapidly, and also increases blood pressure despite the fact that normally, alcohol causes dilation of the arterioles. In weakened bodily conditions, with vascular relaxation, alcohol may increase vascular tonicity. The heart and blood vessels are paralyzed by poisonous doses of alcohol and blood tension falls tremendously.

The local effect of alcohol upon the peripheral nerves resembles the action after absorption upon the system generally. The nervous system is affected in nearly the same order and manner as by anæsthetics, and the same stages may be observed. The stages include the stimulant, depressant and paralytic. The law of dissolution is demonstrated by alcohol as the more highly organized centres and those more recently developed in the process of evolution are first to succumb, and in following out this order the medulla, the first of the higher centres to be developed, is the last to be influenced by the drug. In accordance with this law the cerebrum is first acted upon. The period of excitement is brief and is due in a considerable degree to the increased cerebral circulation and flushing of the brain. It is essential to emphasize the fact that by far the most apparent and decided action of alcohol is one of depression upon the nervous system as a whole. The stimulating influence of alcohol upon the spinal centres is more marked in the lower animals than in man because the brain is proportionately small and poorly developed in the former. The primary stimulating effect of alcohol is shown in man by increased mental activity and apparent brilliancy, but acute reasoning and judgment are not enhanced, and in many cases there is almost immediate mental confusion and drowsiness induced.

This small order contains parasitic and saprophytic species. The asci are without perithecium, except in *Gymnoascus* and *Ctenomyces* where there is a rudimentary perithecium. The *Taphrinae* are undoubtedly related to the yeasts and by some are placed in one order known as *Gymnoasceae*, being represented by *Gymnoascus*. The *Gymnoasceae* exclusive of *Taphrinae* are sometimes placed with the *Plectasceae*, a fungus occurring on the dung of horses and producing simple-fruited organs, which consist of short-branched filaments arising either from a single hypha in which a cell is cut off, or several, one being spirally wound about the other. This becomes the ascus, which contains the ascospores. In *Eremascus* the ascus-producing part resembles certain zygosporae. In *Ctenomyces* the ascus is surrounded by simple torulose hyphae, representing a rudimentary perithecium. The *Lidawella spinosa* described by Matruchot and Dassonville is allied to *Gymnoascus*. It produces numerous ovoid short stalked asci with 8 ovate colorless ascospores. Parasitic on dog. The life history of parasitic members of this order may be represented by *Erasacus pruni*. This fungus grows on the fruit of various species of the genus *Prunus*, producing in plums what is known as plum pockets or bladder plums. The fungus, when fully developed, consists mainly of a single layer of palisade-like asci, which produce their branching mycelium in the parenchyma of the affected part, and later develop between the outer walls of the epidermal cells and cuticle. Here




Fig. 79. *Eremascae*. 1. Plum pocket (*Erasacus Pruni*) on *Prunus Padus*. a. Normal fruit, b. abnormal fruit. 2. *E. Anzincensis* on alder (*Alnus incana*), series enlarged. 3-5. *E. alutatorum*, 3. Surface view alder leaves showing hyphae (h) between cuticle and remainder of epidermal cell. 4. Formation of asci (ar). 5. Ripe asci with ascospores x 100. 1-2 after Wettstein, 3-5 after Sadebeck.

they grow and spread out to the surface, forming a single layer of cells, each cell swells, the cuticle becomes ruptured and a palisade-like layer of asci is formed. There are eight ascospores in each ascus which escape by means of an opening at the tip. The ascospores of *Taphrina* frequently germinate in the ascus, budding like yeast and in this budding condition they produce a small amount of alcohol. Another troublesome species is the Peach Curl (*Exoascus deformans*) which occurs on the young leaves of peaches. The *E. Cerasi* is another destructive species producing the "Witches Broom" of the cherry *Prunus Cerasus*. The *E. Betulinus* produces the "Witches Broom" in the Birches.

PEZIZINEAE, HELVELLINEAE, TUBERINEAE, PHACIDIINEAE

Mycelium well developed, asci borne upon large fruiting bodies and a continuous hymenium consisting of the asci, sterile threads, the paraphyses. Contains a number of common cup fungi like the *Peziza*, *Morchella esculenta*, and *Helvella*. A *Sclerotinia* produces a disease on red mangolds, beans, and hemp, *Sclerotinia* also occurs upon clover and the common brown plum rot (*Sclerotinia*). *Helvella suspecta* with a reddish brown pileus and a dirty flesh colored stalk is suspected of being poisonous. It has a nauseous, sweetish taste, and produces helvellic acid, a hemolytic, or blood destroying substance. The *Gyromitra esculenta* also produces helvellic acid and is regarded as poisonous. It owes its toxicity to the blood making properties. The *Helvellas*, *Morels*, and *Sclerotinia* belong to the family, *Helveliaceae*. *Tuber* produces tuber like bodies found in the soil. The organism is parasitic on trees. The fruiting bodies are enclosed by a peridium which consists of corrugated, smooth, or wart-like excrescences. The hyphae are compact. The ascospores occur in winding passages in the interior. Some of the *Helvellas* and *Morels* are edible. The truffle (*Tuber aestivum*) of the family *Tuberaceae* are edible.

The family *Phacidiaceae*, contains one important parasite of the alfalfa, the *Phacidium medicaginis*. The diseased leaves turn yellow and soon fall. The yellow leaf, or in some cases the green leaves, contain the small blackish or brownish specks usually upon the upper side of the leaf, the injury extend




Fig. 80. Enlarged plant branches. *Exoascus communis* on *Prunus maritima*, projecting beyond the mass are the asci, some of which contain the spherical ascospores. After Atkinson.




Fig 81 Tubercular Truffles. 1. *Tuber ericetum*, part of an ericid truffle, showing hyphae, asci, and ascospores, greatly magnified. 2. *T. acanthum* in log body. 3. *T. brunnescens* section of truffle. 4. Ascospore of *T. Magnusii*. 5. 3, 5, after Tulane. 6. after Wolcott.

ing to the lower surface. The spot contains a small pustule called the apothecium, which is cup-shaped. This cup-shaped body contains the asci (sacs in which the small spores occur, the ascospores with the asci—the two slender threads are known as paraphyses. This fungus is a serious parasitic disease of the alfalfa. The fungus does much injury to the fodder and it is not improbable that at times may be injurious to animals consuming the fodder.




Fig 82 Foliar spot. a. Enlarged leaf showing spots. b. Single spot enlarged. c. Ascus with ascospores, paraphyses rising from mycelium.




Fig. 83. Section through apothecium found on leaf (c. asc. asci and mycelium, Combe.




Fig. 84. Common Blue Mould (*Penicillium glaucum*). 1 Conidiophore, spores in chains. 2. Sclerotium or hard compact mass of fungus (threads hyphae) with asc. and ascospores. Asc. and ascospores shown above. Brefeld.

PLECTASCINEAE

Generally saprophytic fungi with a well developed mycelium, either buried in the sub-stratum or superficial. Reproduction sexual or asexual, asci either borne directly on the mycelium or in closed fruiting bodies, called perithecia.

ASPERGILLACEAE

Peridium thick, perithecia small, the sexual reproduction may be seen from the development as it occurs in the Blue Mould (*Penicillium*)

Penicillium, Link

A branched septate mycelium, conidiophores with septa, numerous branches near the apex, contains small flask-shaped sterigmata, spores borne in chains, conidiophores sometimes in bundles, as in the old *Coremium* asci develop in poorly-lighted places in a sclerotium-like body

Penicillium glaucum. Link. Blue Mould

At first a white mycelium spreads over the surface or through the sub-stratum, the mycelium, through an enzyme action, undoubtedly dissolves the starch, raised masses are formed on the surface, which consist of masses of mycelium thread strands, the strands send out lateral branches from the end of which a whorl of short branches appears, which give rise to one or more whorls from the ultimate branches a chain of small spores is produced, the last one on the chain being the oldest

The ascospores have not been found in corn, but occur in poorly lighted places and are produced in the absence of oxygen. The spores produced in chains germinate when the required amount of moisture and heat is present, so that unlimited numbers of generations may proceed from a single spore. These spores also preserve their vitality for a considerable length of time.

Brefeld has shown that they will germinate though kept in a dry place for several years. The organism grows at various temperatures, from near the freezing point to a considerable heat. It also resists antiseptics. It is one of the most troublesome fungi in stored fruit

Penicillium glaucum is an organism which contains diastase, maltase, emulsin and a ferment which inverts cane sugar. *Calcium oxalate* is deposited in the perithecia. Under certain conditions mannite is said to be produced. When the *Penicillium glaucum* occurs in grape must it delays fermentation

Distribution. The common blue mould is widely distributed in nature and is contained in a large number of the spores which drop in on fruits and decaying bodies and there germinate and produce fruiting bodies

Poisonous properties. This fungus certainly is not pathogenic. It is widely distributed on decaying fruit, it has been suspected, in several instances, of being poisonous, but there is no evidence to support the supposition that this is the case or that it produces toxic substances. Under certain conditions it may possibly produce mycotic stomatitis. It has been found in sputum, nasal secretions, and in the stomach, but these cases are without special significance.

Penicillium marneffii, Siebertmann

Mycelium at first white, flocculent, changing to blackish green when spores are formed, conidiophores slender, branching, bearing a chain of spores from 2-3 μ in diameter.




Fig. 85. Green Mould. (*Aspergillus glaucus*) on the left, *A. repens* on the right, both with conidia in chains, and conidiophores. After Siebenmann.

Distribution. Found in Europe.

Pathogenic properties. Found by Siebenmann in the ear

Aspergillus. Micheli

Parasites or saprophytes with branched septate mycelium, reproduction sexual or asexual, in the asexual, conidiophores enlarged at the end, the enlarged portion bearing numerous small sterigmata, or these bearing smaller sterigmata the conidia borne in chains, occasionally sclerotia form perithecia small with asci and ascospores. The ascigerous stage of a few only is known. The life history of the common herbarium mould was first worked out by DeBary. A little known *A. sulphureus* is said to cause muscular contractions, and tubercular bodies.

Aspergillus glaucus. (L.) Link

Mycelium on or in the substratum forming a bluish green growth, conidia spherical or somewhat elliptical, slightly roughened, 6-15 μ in diameter, borne in chains attached to a short simple sterigmata, perithecia form little yellow masses, each ascus has 8 colorless biconvex ascospores 8-10 μ in diameter.

The life history of this fungus is as follows. This species is common in stored grain and hay. The mycelium spreads over the surface and through the substratum it enters the kernel because of the dissolving action of an enzyme produced by the mycelium. From this mycelium erect threads (conidiophores or sporophores) arise which are enlarged at the end. From the enlarged portion of the conidiophores numerous small and radiating stalks (sterigmata) are produced, each bearing a chain of spores, the end spores of the chain being the older. These spores germinate under favorable conditions of moisture and heat, and again give rise to the same stage. In addition to this, the conidial stage, a second kind of reproductive body occurs. This is produced by the coiling of a branch of the mycelium having several turns. Two or three slender branches grow from the base. One of these grows more rapidly and connects with the top of the spiral coil formed first. The contents of those last formed




Fig 86. Common *Aspergillus*. 1. General appearance showing long conidiophore and sterigmata on end. 2. Perithecia with one ascus and ascospores. 3. Contents from an unripe perithecium. 4. A small part of a mycelium with conidiophore c and spore-bearing sterigmata, young sacogonium s, s. All after DeBary except 1.




Fig. 87. Moldy maize kernels. 1. *Aspergillus* (*Sterigmatocystis*). 2. *Aspergillus*. 3. *Rhizopus*. 4. *Pencillium*. Charlotte M. King.




Fig. 87a. Mycotic stomatitis caused from eating mouldy hay and parasitic fungi on forage plants. (U. S. Dept. Agr.)

unite with the spiral known as the ascogonium. After fertilization a perithecium is produced which contains the asci, each ascus being surrounded by a delicate wall and containing eight biconvex ascospores.

Aspergillus forms diastase and is capable of changing starch into dextrin and maltose.

Distribution. Widely distributed in nature on mouldy hay, corn and other grains.

Poisonous properties. The organism is not pathogenic but probably develops a poisonous substance which may produce disturbance. Dr. Law mentions a serious case, epizootic cerebro-spinal meningitis, in Pennsylvania, due to the feeding of mouldy timothy hay which was badly fermented. In Cairo, Egypt, 6,000 horses and mules perished from the same cause. Michener attributes this disease to foods undergoing fermentation due to toxic fungi. Williams, of Idaho, thought also that the fermentation of alfalfa, timothy and wild grass hay produced the disease. Dr. Law says:

In all probability as we learn more of the true pathology of the disease, we shall come to recognize not one, but several toxic principles and several different affections, each with its characteristic phenomena in the somewhat indefinite affection still known as cerebro-spinal meningitis.

It occurs in horses, sheep, oxen, goats, and dogs, preferably attacking the young which have not become immuned to the toxic substance. It occurs most commonly in winter and spring when animals shed the coat. Dr. Mayo, who investigated this trouble in Kansas, says that a disease known as "staggers," "mad staggers," or as he has termed it, enzootic cerebritis, is caused by feeding corn which is attacked by *Aspergillus gloxus*. The spores of the fungus gain entrance to the circulation, and find lodgment in the kidneys and liver. He supports his conclusions by experiments made by him on a guinea pig and a

young colt. He also quotes Kaufmann, who was successful in producing a disease with *Penicillium glaucum* and *Aspergillus glaucus*. There is considerable loss in many states from cerebro-spinal meningitis. In many parts of the country this is attributed, as I have said before, to mouldy corn. Dr. Biting, of the Indiana Agricultural Experiment Station, made an investigation of this question and concludes that mouldy corn is not responsible for this disease. Upon an examination of mouldy corn he found several moulds and a bacterium. To test the poisonous properties of these, two horses were injected under the skin with five cubic centimeters. Later, larger amounts were given, and each animal was induced to eat as much as five pounds of the infected meal per day. One of the moulds as well as the bacterium gave negative results, the *Fusarium* produced a redness of the gums and some salivation. In no case did cerebro-spinal meningitis result.

The results of the experiment show that inoculations with culture of the bacteria and moulds were ineffective. Eating of the mushes containing pure culture showed that only in the case of a growth of a species of *Fusarium* did any intestinal disturbance follow, and that in one case the feeding of the rotted grain produced considerable intestinal disturbance and some nervous symptoms, but that the disturbance was light in the other.

Grawitz succeeded in producing infection by adapting the digestive tract of the animals to an alkaline medium.

Roberts and Biting say in regard to this trouble in Indiana:

It affects horses, cattle and sheep, but the cause is not known. This disease is reported in stables in the fall and winter. The reports indicate that about an equal number of horses and cattle become affected, but that they rarely become affected at the same time. The horses and cattle kept in the same barn and fed the same kind of food will not become diseased at the same time. Most of the cases occur while feeding ensilage or shredded fodder and thus it has come to be called ensilage disease and shredded fodder disease. The character of the food, however, is only an incident, for cases may occur when other spoiled or fermented foods are present, or when only the best foods are used. The disease is often ascribed to mouldy and rancid corn, but our experiments with such foods and pure cultures of moulds from such foods were negative. Bad ventilation is also ascribed as a cause.

In regard to Mycotic Stomatitis of cattle which they attributed to moulds:

The particular organism causing the disease, if there be one, has not been described. It seems probable that the disease is due to more than one form of fungi which may be present on the pasture. The animals affected are cattle of all ages above 4 months. The disease is not contagious, but usually affects a number of animals in a given herd, and always while in pasture. The fact of a number of animals being affected is due to similar exposure and not to infection spreading from one animal to another. Attempts at direct inoculation have not been successful. The disease occurs in some localities every year, and in others seemingly under special climatic influences. I know a locality where it may be developed at any time by permitting cattle to graze along the roadside. The disease is much more prevalent on permanent blue grass than upon smooth pasture and is of rare occurrence upon pastures used in a crop rotation. The disease develops in pastures allowed to grow for some little time without being used. It is particularly liable to develop a few days after a good rainfall succeeding a dry period.

The symptoms are inability to graze, saliva dribbling from the mouth and frequent visitations to the watering trough, holding the mouth in the water as though it were burned. The animals appear to be hungry but cannot eat. The mouth is red and lips, gums and tongue swell. Blisters form and these soon give way to ulcers that may remain distinct or run together. In some of the aggravated forms the ulcers may unite so that when the crust comes off, it will make a cast of a lip or the whole end of the tongue. The crusts are usually from the size of a dime to that of a quarter. The tongue may swell to such an extent as to protrude from the mouth and the animal be unable to draw it inside. The muzzle may be increased one-half in size.

Dr Craig, of the same state, reports somewhat similar experience

Captain F. Smith, in his manual of Veterinary Hygiene, refers to the injuries from moulds, especially *Penicillium* and *Aspergillus*, calling attention to the brittleness of hay caused by fungi, and that the spores produce irritation to the respiratory passages. He states further that oats and bran have produced diabetes, paralysis, and subsequent death in horses. He refers to the case mentioned by Professor Varnell in which the horse died in three days from eating moldy oats. Professor Gamgee calls attention to the disease in France and Scotland in the years 1854 and 1856, due to horses feeding on grass which had become wet and musty. The animals suffered cerebral derangement, producing stomach staggers, so-called by English writers

Aspergillus Oryzae. Ahlburg

Rabbits inoculated showed convulsive symptoms. tubular feci occurred in the intestines. The *Aspergillus Oryzae* forms maltose and diastase and in Eastern Asia plays an important part in the manufacture of "sake" or rice beer, which has been a national drink of the Chinese for centuries.

Aspergillus malignum. (Lindt.)

Mycelium bluish gray, conidiophores short, the end pear-shaped, 22-24 μ wide, sterigmata branching, conidia in chains 3-4 μ in diameter, perithecia 40-60 μ , ascospores 6-8.

Distribution. Found in Europe.

Pathogenic properties. Grows best at the higher temperatures. Found by Lindt in the human ear




Fig. 48b. Pale Mould (*Aspergillus flavus*). Showing conidiophore and spores attached in chains. After Siebenmann.




Fig. 48. *Aspergillus Oryzae* on rice.
1. Conidiophore, sterigmata and conidia
2. Young conidiophore. Modified by Charlotte H. King after drawing by

Aspergillus flavus. Link

Hyphae arachnoid, white, the fertile erect, slightly caespitose, conidia 5-7 μ in diameter, small, globose, vari-colored, slightly wart-like, collected about the white sub-globose, wart-like apex; apex finally becoming yellowish, sclerotium very small, dark.

Aspergillus fumigatus. Fresenius

Forms greenish or bluish gray masses on the surface of the substratum, conidiophores short with a semi-spherical mass 8-20 μ in diameter. Sterigmata bear the spherical conidia 2.5-3 μ in diameter, which are at first bluish green and later brown. Sclerotia unknown. Grows best at a temperature of 37-40° C.

Distribution. Widely distributed.




Fig. 89. Section of kidney of rabbit showing mycelium of an *Aspergillus*. After Geawitz.

Pathogenic properties. It has been known for some time that several species of *Aspergillus* are pathogenic for animals. In 1815 Mayer and Emmert found the fungus in the lungs of a jay. In 1826 it was reported in the long bones of a white stork by Heusinger, and numerous other cases in birds like the flamingo, duck, chicken, ostrich, and turkey, have been reported especially in Europe. Kuhn, in 1893, furnished quite conclusive evidence that certain species of *Aspergillus* can produce necrosis and disease. Chantemesse, at the tenth International Congress in Berlin, called attention to a disease of pigeons resembling tuberculosis which he said was produced by an *Aspergillus*. Saxer attributed mycosis to an *Aspergillus*, and, according to Sticker, the disease may appear sporadic and endemic, the latter to persons who feed pigeons and to the hair combers in Paris. It is spontaneous in horses, cattle, dogs, and birds, and is sometimes quite epidemic in birds. The form of the disease when it occurs in the lung is called Bronchopneumomycosis, it appears that various species of *Aspergilli* also occur in connection with otomycosis, and occasionally in the nose or the eye. A very complete history is given by Drs

Mohler and Buckley in the report of the Bureau of Animal Industry. Hughes Bennett reported a case in the sputum of a tubercular patient and in 1847 Shuyter reported definitely on the *Aspergillus* in the lungs of a human being. Virchow in 1856 reported several cases. In 1879, Leber first described a purulent keratitis due to aspergillus infection. Drs. Mohler and Buckley in referring to the observations on pneumomycosis, say

Dieulafoy, Chantemesse, and Wedal, reported their observations and studies of pneumomycosis as it occurs in a certain class of men in Paris. These men feed thousands of young pigeons daily by taking into their mouths a mixture of grain and water which they force into the mouths of the birds much in the same way that the old pigeons feed their young. It had been a matter of common observation that these men were sufferers from a severe pulmonary disorder, but when their sputum was examined, instead of finding tubercle bacilli, only the threads of mycelia were detected. This observation was subsequently confirmed by Renon and other investigators. Until this time it had been held that the presence of fungi in the lung tissue was of secondary importance, but these observations dispelled further argument. Experiments on animals in which they were made to inhale the spores, were successful in producing the disease; thus it was that the natural infection was proved.

Renon, who made an exhaustive study of the subject, concludes as follows concerning aspergillosis:

1. That aspergillosis is a spontaneous disease affecting the bronch and lungs of birds and man, and creating in the animals a generalised affection similar to hemorrhagic septicaemia; that it develops in eggs in incubation and may contaminate the embryos notwithstanding there-in.

2. The disease may be transmitted experimentally. The natural and cultural characters of the fungus and the lesion it provokes are truly specific. In its pathogenic action it bears a strong resemblance to tubercle bacillus.

3. In man it develops upon the corner of skin, but has its particular evolution in the respiratory apparatus, creating pulmonary mycosis, resembling tuberculosis, and pulmonary gangrene but without the fetid odor. It may coexist with tuberculosis. Occasionally it is fatal after the formation of cavities in the lungs. It may invade the bronchial apparatus alone, causing membranous bronchitis of special form and of long duration.

4. In all its manifestations *Aspergillus fumigatus* may play a primary or secondary role in both man and animals. It is not, therefore, a simple saprophyte, but a true parasite.

Renon points out the relation of the occupation of man to his contracting the disease. When animals and men are kept where the mould is common, as in hair assorting establishments where rye is used to disentangle the hair, they become affected with the disease. The handling of dusty grain and feeds may lead to infection from *Aspergillus*. Saxer also went into historical details in giving his experiments with mycosis in man. In 1837 *Aspergillus* was observed by Ravolta in the pharyngeal abscess of a horse. Gott observed it in an auricular catarrh of a dog. Pech observed mycotic pneumonia in seven horses, where they had been fed mouldy hay. Several cases where the *Aspergillus* occurred in the trachea of cows have also been reported. Pearson and Ravenel record a case of pneumomycosis of the lung of a cow.

Infection takes place generally by the inhalation of the spores. The spores germinate in the bronchial branches, develop a mycelium and produce conidia-phores and spores on the surface. Drs. Mohler and Buckley, calling attention to the various aspergilli which have been found, say

Numerous experiments have been tried with the various fungi, especially in relation to the best temperatures for their development and fructification, and it has been found that, although a few are able to germinate in the bronchioles, the *Aspergillus fumigatus* is about the only one which develops a vigorous growth there and fructifies, the temperature of the human body seeming to be quite suitable for this species. Most of the other molds develop at a much lower temperature and are therefore usually harmless even if introduced

into the lungs. But for the *Aspergillus fumigatus* the lungs act as a veritable propagating house, furnishing a moist, nutrient and open which to grow and a congenial, warm, moist atmosphere with a sufficient amount of oxygen for its demand to come to complete maturity and for fructification to take place. When the fungous growth is localized in the bronchial mucous membrane, the condition is known as bronchomycosis. It may be that the tissues are able to forestall entrance into their substance and finally the fungi die and recovery takes place. In birds the growth may extend to the air sacs; this condition is then called cytomycosis. Cases of cytomycosis are very rare; and when it does occur, emaciation of the birds is the predominating symptom. When the lung tissue itself is the seat of invasion, the term pneumomycosis is applied. Invasion of the lung tissue by the mycelium is the occasion for an intense inflammatory disturbance with positive chemotaxis. However this tissue reaction seems to offer the most trifling barrier to the parasitic encroachment in such weakly subjects as birds. Generali states that delicate breeds of pigeons are noticeably susceptible to this disease.

In regard to the symptoms in birds, he says:

The birds become listless, mope, and do not follow the rest of the flock. When made to run they soon become exhausted and fall and have great difficulty in breathing. Even when disturbed they appear very weak and gasp for breath, extending their heads and making movements as if choking. There is a great thirst, but a diminution or complete loss of appetite. The birds become rapidly emaciated; the wings are pendant, the eyelids droop, comb and wattle become quite pale, and a general dejected appearance follows. Usually there is an intense diarrhea which weakens the bird very much. In the experimental disease the diarrhea is an accompaniment just as in that of a spontaneous development. The plumage is said to appear ruffled, and the respirations become croupy, even when the disease has not advanced very far. Later they are more rapid and a rattling noise can be heard. In the final stages suffocation is threatened.

When the air sacs are affected very few symptoms manifest themselves, though emaciation is marked. As in any similar condition of the lungs, fever is high, and symptoms that would be manifested in pneumonia of fowls would, of course, show here. There is more or less catarrh of the trachea and bronchi, and if these alone were diseased there would probably be nothing to attract notice other than symptoms of bronchitis. Bleeding from the nostrils has been observed in man and in animals, and it may be that this would also occasionally be seen in birds. If the air spaces in the bones become affected, lameness with swelling of the joints may result. The duration of the disease is quite variable and death may take place in from one to eight weeks from asphyxia or marasmus. Duration depends a great deal upon the portion of the respiratory apparatus that is affected. If the aspergillar nodules were localized in the mouth, as it is sometimes in pigeons, or in the bones or air sacs, the duration of the disease would, of course, be much longer than if in the bronchi or lung substance.

The pathological lesions are as follows:

The actinomycotic masses are noteworthy. The fungus may frequently become localized in kidneys, and muscles of heart.

The microscopic examination of these organs disclosed a picture simulating the gross appearance of an advanced case of pulmonary tuberculosis, with the exception that the bronchial tubes were almost completely plugged with a greenish velvety membranous lining.

In the bronchial divisions not wholly occluded by the croupous exudate are seen the characteristic aspergillar fruitheads in various stages of development, from that of a slight bulging end of the hyphae to those giving off their spores. Included within this avascular exudate are quite a few leucocytes and red blood cells, but their presence is by no means constant. The bronchial mucosa is often eroded and the lining epithelium replaced by a fibrinous coagula or by a membranous material composed of matted mycelia, threads from which hyphae extend into the air space forming spore-bearing fruitheads, owing to the presence of oxygen.

In animals in which the disease was experimentally induced by the injection of the spores into the blood vessels or into the lung substance, solitary venous resembling tubercular formations were quite noticeable in the lung tissues, and in these an occasional giant cell was discovered. In the lungs of a chicken which was inoculated directly into the lung substance, an acute solitary pseudo-tuberculosis was produced, accompanied by intense hemorrhages into the interstitial tissues, as was also the case in intravenous inoculations. In these tubercular nodules penetrating filaments could be made out, but the spores could

not be purely demonstrated, or at least differentiated from other cellular elements. Often the bronchial ramifications were the seat of hemorrhage, in which a noteworthy increase in the number of leucocytes could be observed.

In large rabbits the pathological lesions appeared to be as follows:

Rabbit No. 1006 failed to show any marked symptoms for the first two weeks after inoculation. It then began to lose weight, and on the twenty-fifth day was chloroformed. The postmortem examination showed an involvement of the liver, spleen, kidneys, and abdominal serous membranes, as in the preceding rabbit, but to a less extent. The organs of the thoracic cavity were apparently normal.

The optimum temperature of growth for the fungus is from 35°-40° C. Ceni and Beata in their investigations isolated a toxin from two species of *Aspergillus*, the *A. flavus* and *A. fumigatus*. Dogs inoculated intra-abdominally with large doses died within a few hours, showing tetanic symptoms and general hyperemia of all the organs. This work has not, however, been confirmed. Drs. Mohier and Buckley did not succeed in producing serious symptoms with the filtered product when injected into rabbits.

The *Aspergilli* also produce disease of the eye but, according to Plant, this disease is not of frequent occurrence; he discusses several cases under the head of keratomycosis. One case described by Leber is as follows: A farmer forty-five years of age, while threshing had the misfortune to have some chaff of oats thrown into his eye. The sclerotic coat became inflamed, followed by healing and total leucoma (leucom). Another case is cited where a pear was thrown against the eye of a farmer, and another case of a fifty-three year old patient, a miller by profession, who had a slight fever, his right eye becoming inflamed. The conjunctiva had the appearance of trachoma. The sclerotic coat was clouded and the surface of the eye brittle consisting of threads of fungi. Focha, who investigated this case, determined that the fungus was *Aspergillus fumigatus*. *Aspergillus fumigatus* has also been observed in the nasal cavities where it produces necrosis and a disagreeable odor.

In a review of a paper by E. Bowlin and L. Gauthier* the following statements are made with reference to the Toxin found in *Aspergillus fumigatus*:

From a study of this fungus in cultures and in experimental animals it was found that *Aspergillus fumigatus* produces a toxin which may be rightly contrasted with the toxins of bacteria. For the formation of this toxin in cultures it is necessary to have a mixture of protein, especially of the peptone type, and some carbohydrate, especially glucose, saccharose, maltose, or dextrin. The reaction of the toxin must be either neutral or alkaline. The effects of the toxin are chiefly observed in the nervous system and are produced more or less rapidly by the method of inoculation. The symptoms of poisoning from the toxin are muscular convulsions resembling tetanus and leading to death within a few hours if the animal does not recover. The rabbit and dog are very susceptible to the toxin, while the guinea pig, cat, mouse, and white rat are more refractory. The dog and cat are naturally immune to the spores of *A. fumigatus*, but are quite susceptible to the toxin produced by the fungus.

Treatment. To prevent the disease do not feed mouldy grain or fodder. Separate at once all diseased animals from the healthy. Use only thoroughly clean dishes, the troughs and boxes should be cleaned with formaldehyde.

Very little can be done in the way of treatment in the case of birds. Mohier and Buckley say that if a large number of birds are affected at one time, or if those affected are very valuable, treatment may be tried in the form of medicated vapors, such as those generated from wood tar or sulphur. A small quantity of wood tar is put in a pint of water and stirred with a red-hot iron. The person doing the fumigating should remain in the room and immediately remove any

* The Ann. Ent. Soc. Amer., 28 (1935) No. 3, of the Experiment Station Record.

birds that are overcome by the vapors. Burning sulphur or vapors of formalin may be tried in like manner. Hydrogen peroxid, solutions of potassium iodid or hyposulphite of soda may be used as intratracheal injections, and in case of local nodules in the mouth or nostrils the tincture of iodine may be applied to them with beneficial results.

It appears from the investigations with reference to kerato-mycosis, that infection generally occurs through the medium of feed, straw or something that is thrown forcibly into the eye. According to Plaut the simplest and surest method of dealing with the disease is to use a 2% solution of salicylic acid, three times daily, but inhalation of an atmosphere containing iodine is recommended by some of the German investigators, or the inhalation of ethereal oils. Immunity cannot be obtained by beginning with the injection of small quantities of spores and increasing the dose. Dogs are not immune against *aspergilli*. Mice are immune.

Aspergillus niger. Van Tieghem

An abundant mycelium in the substratum and on the surface becoming blackish, conidiophores long, sterigmata branched, conidia $3\frac{1}{2}$ - $4\frac{1}{2}$ μ in diameter, roughened, spherical or cylindrical sclerotia. The fungus contains diastase, invertase, and emulsin, it breaks up tannin into gallic acid and glucose, and converts sugar into oxalic acid.

Pathogenic properties. This fungus has been found both in the lungs and the ear, although less pathogenic than the preceding species.

Aspergillus subfuscus. Olen-Gade

Mycelium olive yellow or brownish when mature, in and on the substratum, conidiophores short, club-shaped, spores spherical, colorless.

Distribution. Found in Europe, closely resembles *A. fumigatus*.

Pathogenic properties. Pathogenic, but less so than the *A. fumigatus* or *A. niger*.

Aspergillus nidulans. (Eidam.)

The mycelium forms greenish masses; later the mass assumes a reddish color; conidiophores 0.6-8 millimeters long and 8-10 μ across, colorless, branched, sterigmata consist of a basal branching cell and two or more branches, each branch containing from 20 to 30 conidia, perithecia yellowish, 0.2-3 millimeters in diameter, ascospores 8.

Distribution. Found in Europe.

Pathogenic properties. The disease appears on the second day after inoculation in guinea pigs and death occurs in 60 hours. Kidneys are enlarged and show small white dots. White masses also occur in the peritoneum. It is pathogenic for cattle and man, and is occasionally found in the human ear.

PYRENOMYCETINEÆ

An important division of the fungi, containing about 10,000 species, many of which are troublesome parasites on cultivated plants. The mycelium is composed of delicate distinct hyphae or of closely coherent threads, frequently forming a pseudo-parenchymatous tissue, hymenium enclosed in a subglobose envelope called a perithecium, or with an opening at the apex, which is often



Fig. 89a. *Aspergillus nidulans*. 1. Conidiophore. 2. Branch of mycelium with asc and ascospores, magnified. 3. Asc. All greatly magnified. (After Tadm.)

prolonged to form a short tube or beak, numerous transparent asci arise from the base of the perithecium, these contain the ascospores; between the asci slender filiform bodies, called the paraphyses. Polymorphic fungi with conidia, spermogonia, and pycnidia, supposed to be connected with the ascigerous stage. The formation of the ascospores is in some cases presented by the development of sexual organs in which genuine fertilization occurs. The reproduction can be illustrated by the manner in which it occurs in the powdery mildew of the lilac, *Microsphaera Alm*.

The mycelium spreads over the surface of the lilac leaf the fungus draws its nourishment from its host by means of haustoria which penetrate the epidermal cells, the mycelium produces erect branches which bear these spores in a moniform chain the end spore being the oldest, these summer spores germinate immediately and propagate the fungus, later two hyphae cross and there arises an ova. cell, the oogonium, which is separated from the hypha by a cell-wall, at the base, from the same hypha springs a longer and thinner cell, also cut off by a cell-wall this cell is above the oogonium, and is known as the antheridium, from the base of the oogonium other cells arise which soon enclose it finally a brown perithecium is formed which bears dichotomously branched appendages, the perithecium contains the asci, in which are found the ascospores, which germinate, probably, in the spring. The accompanying figure after Harper illustrates the development.

Another type of one of the Sphaeriaceae, the *Gibbellina cerealis*, is common on stems of wheat where it produces at first a grayish brown circular spot, the mycelium frequently encircling the stem. The conidia are oval, the perithecia are numerous.




Fig 90. Stem Blight (*Gibberella cerealis*), one of the *Sphaeriaceae*. a, general appearance. b, asci with ascospores and paraphyses. c, stroma, mycelium, and perithecia. After Cavara.

PERISPORIALES

Perithecia spherical, closed, or with the ostium obscure, coriaceous or brittle carbonaceous, opening irregularly, generally without stroma, but mostly seated on a well developed, superficial mycelium. This division includes the order *Erysibaceae*.

ERYSIBACEAE

Superficial mycelium, branching, septate, closely adhering to the surface by means of the haustoria, asci arising from the base of the perithecium, delicate, thin-walled, colorless, oblong, obovate or suborbicular, stalked, usually containing from 2-8 ascospores, perithecium spherical with appendages, without ostium conidia (*Oidium*) simple, colorless, cylindrical, oval or ovate, borne one above the other on septate, colorless hyphae. Contains many important

parasitic fungi, like the powdery mildew of the grape (*Uncinula spiralis*), mildew of alac (*Microsphaera Alni*), mildew of sunflower (*Erysiphe Cichoracearum*), mildew of cherry (*Podosphaera tridactyla*)

Erysiphe. (Hedw.)

Perithecium containing several asci, appendages with simple threads, similar to and frequently interwoven with the mycelium. A small genus of 20 species of wide distribution.

Erysiphe communis. (Wallr.)

Amphigenous, mycelium abundant, persistent, or sometimes evanescent, perithecia variable in size and reticulate, appendages variable in length, often long; asci 4-8 or more, ascospores 4-8.

Distribution Found on a large variety of different hosts but common on plants of the order of Leguminosae, especially the forage plants like the pea (*Vicia sativa*), bean, clover and other members of the clovers.




Fig. 91. Powdery Mildews. 1-3. *Sphaerotheca Castagnei* on Hop. 1. Part of leaf of hop with perithecia shown in the form of dots. 2. Perithecia with tortuous appendages (ap) x 175. 3. Asci with spores within the ascospores x 380. 4. Powdery Mildew on Cherry (*Podosphaera tridactyla*), conidiophore bearing conidia (c). 5-7. *Microsphaera microsphaerula*. 5. On leaf. 6. Perithecia, greatly magnified. 7. Ascus and ascospores. 1 after Wettstein. 2-4 after Tulane. 5 after Landau. 6-7 after Winter.




Fig. 92. Powdery Mildew, *Sphaerotheca Castagnei*. 1. Ugonium (u) and antheridium (a). 2. Separation of antheridium cell. 3. Fertilization and formation of additional cells. 4-8. Further development of cells. At greatly magnified. After Harper.




Fig. 93. Powdery Mildew of Grass (*Erysiphe graminis*). A. Oidium stage and mycelium m. B. Perithecia with appendages and mycelium m. C. Perithecia with asci and ascospores. After Frank.




Fig. 94 Powdery Mildew of Bluegrass (*Erysiphe graminis*) Oidium stage, leaves at the right magnified, the one above more highly, showing the powdery substance. (Charlotte M. King)

Poisonous properties. The Veterinarians of Europe ascribe to these mildews a form of stomatitis.

Erysiphe graminis D C

Amphigenous, often epiphyllous, mycelium dense, felt-like, persistent white or gray, sometimes tinted brown, perithecium immersed in the mycelium, few and scattered, large, about 225 μ , in diameter, asci 16-25 μ oblong or oval, stalked, ascospores 8 or rarely 4, appendages rather short

Distribution and Hosts Found on many different grasses like blue grass (*Poa pratensis*), fowl meadow grass (*Poa serotina*), occasionally also on wheat or orchard grass. The following rather popular account treats of this disease as it is common in the west

Every one who has had occasion to walk through a blue grass meadow after a rain, especially in damp and shaded places close to the ground, must have noticed a white mealy covering on the blades of many of the leaves. The Germans have called this mehlthau (literally translated meal dew), which is certainly very expressive of its appearance. An examination with a microscope will show that this white substance is composed of spores and a mycelium. The mycelium is cobwebby and spreads over the surface, but does not penetrate the leaf. In numerous places erect branches are produced, these bear numerous spores. This stage was formerly called *Oidium monilioides* being named *Oidium* because the spores resemble an egg, although the resemblance is not marked in all cases of *Oidium*, the species was called *monilioides* because it was necklace like, referring to the manner in which the spores are borne. Worthington G. Smith states that the spores are so small that it would take about a million to cover a square inch.

In a powdery mildew occurring on the squirrel-tail grass, and supposed to be the same fungus, these spores are also capable of immediate germination. On blue grass the fungus frequently does not produce perithecia but ends its existence with the formation of conidia. It produces perithecia abundantly on wheat in Iowa.

These conidia or summer spores germinate, under favorable conditions, in from ten to sixteen hours. The temperature most favorable for germination is from 17-26° C. In a powdery mildew occurring on the squirrel-tail grass, and supposed to be the same fungus, these spores are also capable of immediate germination.

Under favorable conditions, especially moisture and damp weather, the fungus spreads rapidly. The leaf of grass affected by this fungus soon dries, and when the affected plants are disturbed, small clouds of dust arise especially in shady places. The perfect stage of the fungus is not of common occurrence, though if careful search is made in the fall, small black specks may be seen these are the perithecia and contain the asci and ascospores. It is the resting stage or winter condition of the fungus. The writer found the perfect fungus abundant on *Poa Wolfii* in Colorado, and Carver found it abundant on blue grass near Ames one season. The spores of the *Oidium* stage do not retain their power of germination very long, but the ascospores contained in the perithecium germinate the following spring, and when the tube comes in contact with the proper host the mycelium spreads over the surface of the leaf and causes the mealy appearance.

Poisonous properties. This species is abundant and often causes serious trouble it certainly renders the hay nearly worthless to be fed to animals. It often, no doubt, gives rise to a stomatitis such as is described for other fungi.

HYPOCREALES

Perithecia spherical or ellipsoidal, with an ostiolum, stroma when present variously colored, reddish, yellow, never black or hard.




Fig. 98. Various species of *Cordyceps*. 1. *C. ophioglossoides*. 2. *C. militaris*. a Stroma on a caterpillar (c). 3. Stroma on a fruiting form of *Biophomycet granulatus*. 4. Avespore $\times 200$. 5. Conidiophore $\times 150$. 6. Conidia of *C. ophioglossoides*. 7. *C. cinerea* on a beetle (c). 8. *C. varians* on a caterpillar (c). a in all figures sterile, b fertile part of the stroma. 1 and 6 after Lindau. 3-5 after Brefeld.

HYPOCREACEAE

Simple or compound, perithecia somewhat coriaceous, never black, bright colored, opening by a subcentral ostiolum, stroma soft, waxy, or occasionally cottony. A very numerous family containing many species. Contains the genera *Nectria*, of 250 species, some being parasitic upon trees, the *Gibberella* and the *Hypocrea* upon barks of trees, etc., *Cordyceps*, parasitic upon various insects, *C. militaris* being found upon *Lepidoptera*, the conidial stage of which is *Isaria farinosa*, the *C. Ravenelii* upon the larvae of the June beetle; *Polystigma rubrum*, parasitic upon the plum, *Epichloe typhina*, the so-called Cat-tail fungus found upon various species of grass, especially timothy and orchard grass. Contains also the *Gibberella Sanbornii*, a parasite on wheat, which is a stage of *Fusarium roseum* described later in this work.




Fig. 96.—Normal ovary of rye. Fig. 2—Same invaded by *Claviceps*. Fig. 3—Cross-section of ovary showing mycelium and spores of sphacelial stage. The round bodies are summer spores. Fig. 4—Sclerotium stage. Fig. 5—Sclerotium stage. Fig. 6—General view of sphacelial stage. Fig. 7—Development of ergot in spring. Fig. 8—Cross-section of globular head showing flask shaped perithecia. Fig. 9—Asci. A single perithecium showing elongated bodies in the center. Fig. 10—A single ascus with filiform ascospores protruding. These spores (reproductive bodies) germinate and infect the young ovary of rye. After Tuissanc. U. S. Dept. Agrl.

Claviceps, Tul. Ergot.

Stroma erect, consisting of a sterile stem, subglobose, fertile head from a subcylindrical, black, hard sclerotium; perithecium immersed in the stroma, flask shaped, asci, clavate-cylindrical; ascospores, filiform, colorless.

Claviceps purpurea, (Fr.) Tul

Sclerotium variable in length from $\frac{1}{8}$ to 1 inch long or more, long cylindrical, generally somewhat curved, wrinkled, purplish on the outside white within, usually several fruiting bodies from the same sclerotium, heads spherical, tuberculose, borne on short flexuose stems, asci narrow, linear, 8-spored, ascospores filiform, continuous, attenuated toward the end, 50-76 μ long

Ergot is a stage of a minute parasitic fungus, although its true nature was not known by early writers, it is mentioned by many of them. Lonicer about the middle of the sixteenth century, mentions its specific use. Thalus applied the name of "ad sistendum sanguineum"

Bauhin used the name of *Serole furfuracea*. De Candolle called it *Sclerotium clavus*. Although other names have been applied to it, the credit of working out the life history belongs to Tulasne, one of the most eminent of French mycologists

There are still many persons who believe that ergot is a degenerate kernel of rye or wheat, but the researches of Tulasne and other mycologists have laid at rest many of the vague theories concerning it. The black purple, or dark gray spurs found in the flowers of rye, wheat, and other grasses are simply one stage of a parasitic fungus, known as *Claviceps purpurea*. These spurs consist of a compact mass of threads known as the sclerotium stage; it was formerly called *Sclerotium clavus*.

No changes occur in ergot while it remains in the head, but the following spring, when laid on damp earth, it produces at different points small, roundish patches which are somewhat elevated. Soon a small white head appears which elongates, becoming stalked, and bearing a globular head at the tip. These heads change from a grayish yellow to a pinkish color. A cross section shows that the central portion is made up of closely woven hyphae or fungus threads while the edge contains a number of flask-shaped bodies, the perithecia, in which are found elongated bodies known as asci, each ascus contains eight filiform spores, the ascospores. The ascospores germinate and when coming in contact with a very young ovary the mycelium penetrates the delicate walls of the ovary and gradually displaces it. It is quite easy to trace out its life history by placing the ergot in damp sand and allowing it to remain over winter

The first indication of ergot in the summer is the formation of the so-called honey-dew a sweetish and rather disagreeable fluid, which is eagerly sought by flies and other insects which feed upon it. This fluid contains a large number of small spores so that insects can readily carry the fungus from a diseased ovary to one not diseased. These spores germinate immediately. This stage is called the sphacelia, and formerly was held to be a distinct fungus. In this stage the mass which has replaced the ovary is soft, but as it becomes older it hardens, ultimately a hard and compact mass, the ergot, is formed.

Distribution and hosts. Found on a large number of host plants. Rye is more subject to it than any of the other cultivated cereals. The largest specimens are usually produced on isolated specimens of rye coming up in fields. It seldom happens that all of the ovaries are affected. Wheat, especially winter wheat, is subject to the disease. The officinal ergot is usually obtained from rye. In Europe it has been reported on oats. Mr. C. W. Warburton found it on the same host in Iowa, in 1909.

Of our native wild grasses, wild ryes (*Elymus robustus* & *virginicus*, *E. striatus*, *E. canadensis*, *Aspella hystrix*) are most subject to the disease. Most cases of ergotism in the United States undoubtedly result from the ergot on various species of *Elymus*, in Iowa on the *Elymus robustus*, which is a common plant everywhere. *Agropyron occidentale*, a grass not uncommon in northwestern Iowa, and Quack Grass (*Agropyron repens*), are also much subject to its attacks. Scarcely a head of the Western Wheat Grass cultivated on the college farm could be found which did not have some ergot. This may be for the same reason that it occurs most abundantly on rye, namely, that the grasses occurred in isolated places. In some pastures, timothy (*Phleum pratense*) is much subject to the attack of *Claviceps purpurea*. Thus in an old pasture in Wisconsin I observed a large percentage of timothy which contained many heads which were ergotized. Blue grass (*Poa pratensis*), *Poa annua*, *Calamagrostis canadensis*, *Agrostis alba*, *Lycerna fistula*, and many others, in some seasons and localities, are diseased. Unusually large specimens sometimes occur on Wild Rice (*Zizania*) in Iowa.

It may be possible that some of the forms of ergot on grass may be referred to other species. Habited states, however, that ergot on *Elymus robustus* is *Claviceps purpurea*. The *Hordeum jubatum* contained apparently the same species, with some minor differences but these were due to the nature of the host. *Claviceps microcephala* (Wallr.) Tul., occurs on *Phragmites C. setulosa* (Quell.) Sacc. with yellow stroma on *Poa* and *C. pusilla* Ces. on *Andropogon ischaemum*.

Poisonous properties. The subject of ergot and ergotism is one of considerable importance to stockmen in many parts of the country. Scarcely a year passes without some complaints being received by the state veterinarians of the injurious effects of ergot. The writer receives several complaints of this kind every year. But the cases of ergotism today are not nearly so frequent as they were 40 or 50 years ago. We will, therefore, append here a short history of the disease.

Epidemics of ergotism have, without doubt, been correctly referred, before the tenth century. Wood states that epidemics of ergotism or chronic ergot poisoning have been recorded from time to time since the days of Galen (130-200 A. D.) and of Caesar (B. C. 100-44). From the ninth to the thirteenth century epidemics were frequent in France, and in the twelfth in Spain. They were first called plagues but later received special names. In 1596 Hesse and adjoining provinces were visited by this plague which was attributed to the presence of ergot in grain. In the epidemic in Silesia in 1722, the king of Prussia ordered an exchange of sound rye for the affected grain. Freiburg was visited in 1702, Switzerland in 1715-16, Saxony in 1716, and other districts of Germany in 1717, 1736, 1741-42. France was visited in 1650, 1670, and 1674. From 1765 to 1769 it was abundant in Sweden in rye

and barley. Linnaeus attributed it to the grain of *Raphanus raphanistrum*, which occurred in France in 1816, in Lorraine and Burgundy; it was especially fatal to the poorer inhabitants.

It has been observed that these epidemics follow a rainy season. Fleming states that in 1041 when the weather was so unpropitious, tempests, rains, and inundations occurring, many cattle perished from the disease. "In 1098, after inundations and heavy fogs, there was a general epizootic among cattle in Germany. In the same year ergotism appeared in the human species."

Dr. Randall, in 1849, called attention to a disease in New York, in which the involved parts were finally invariably affected with dry gangrene. He states that in the severe climate of New York farmers allow their cattle to winter in fields on blue grass (*Poa pratensis*) which is rich in ergot. A disease known as "hoof-ail" was correctly ascribed to ergot by James Mease, of Philadelphia, prior to 1838. The disease was quite severe in Orange county, New York in 1820. It was minutely described by Arnell. In 1857, the disease was quite severe in Portage county Ohio. A committee appointed by the Farmers' Association of Edinburg reported that the disease was due to ergot contained in the hay eaten by cattle. In recent years, epizootics of ergotism have been reported by Law in New York, Stalker in Iowa, and Faville in Colorado. In 1894, a very serious outbreak occurred in Kansas which was at first diagnosed as "foot-and-mouth disease." Dr. Salmon found, upon examining samples of hay from various localities in the state that these contained considerable quantities of wild rye (*Lymus virginicus*, var. *submuticus*) which in turn contained a large amount of ergot in one case 12 per cent and in another 10 per cent being found. From this he estimated that 5-6 per cent of the entire weight of the plant must have been ergot and that a twenty-pound ration of hay would contain four ounces of ergot.

Dr. Harshberger has called attention to an outbreak of ergotism from the use of ergotized red top, the fungus being common on red top throughout the United States and being one of the most common impurities in red top seed.

The ergot contains the substance leucin and the non nitrogenous substance ergotine, which according to the earlier investigations was regarded as the active principle and as an alkaloid. According to Wenzell ergot contains the two alkaloids, *erbohn* and *ergotin* $C_{25}H_{32}N_2O_8$, an amorphous, alkaline, feebly bitter substance. But according to the later investigations these substances are identical. Tanret isolated the crystallizable alkaloid *ergotinin* $C_{25}H_{30}N_2O_8$, this is a crystalline, slightly bitter substance, subsequently Robert found that this substance would not produce the action accredited to it and attributed its action to *ergotinic acid* and the alkaloid *cornutin*. The more recent investigation of Jacobs attributes the poisonous action to *chrysotoxin* an amorphous glucosidal acid. *Secounotoxin* is a compound of *sphaerolotoxin* and *iscalin* $C_{29}H_{38}N_2O_4$, accompanied by the harmless substance, *ergochrysin*. According to Robert *cornutin* is an alkaloid having a specific action on the uterus, causing it to contract, *sphaeric acid*, a non-crystallizable and non-nitrogenous substance which causes the poisoning and gangrene, *ergotinic acid*, a nitrogenous glucoside without action on the uterus and narcotic in its effects. Besides these substances it contains others, prominent among them being a sugar called mycose, which is also present in other fungi. Ergot stimulates the involuntary muscles of the stomach and the intestines, it causes a constriction of the arter-

loses and veins throughout the body with an increase of blood pressure. In toxic doses it paralyzes both the vasomotor centers and the heart muscle.

It appears from the experiments of Dale¹ and Barger and Carr² that *cornutin* does not occur as such in ergot but is an artificial decomposition. Taret discovered the first well defined crystalline alkaloid which he called *ergotinin*. The *secalin* of Jacobus³ is identical with ergotinin. Barger and Carr separate a second alkaloid which can be recognized chemically to this they gave the name of *ergotoxin*, $C_{28}H_{41}O_8N_3$. This substance is of great physiological potency. According to Dale⁴ ergotoxin produces in doses of a few milligrams "not only the characteristic reaction of ergot described by him, but also gangrene of the Cock's-comb, and other ergot effects described by Robert and others to *sphacelic acid*."

According to Cronyn and Henderson⁵ *ergotoxin* is a highly active alkaloid and has the properties of ergot most desired in medicine. It brings on long enduring vaso-constriction, increases uterine movements when injected in-



Fig. 98.—Effects of Ergotism: Hoofs of cattle showing flesh sloughing away. (Salmon.)

travenously and the same to a less extent when injected subcutaneously, but when given per os has very little action.

The toxicology of ergot is well described by Dr Winslow as follows:

Enormous single doses are required to poison animals or man. When as much as two drachms of ergot to the pound, live weight, are given to dogs, death is not constant. Three ounces, however, have proved fatal to small dogs. Acute poisoning is characterized by vomiting (in dogs), profuse salivation, dilation of the pupils, rapid breathing and frequent pulse. The animal cries out, has convulsive twitchings, staggering gait, paraplegia, intense thirst, and coma, terminating in death. Horses, cattle, and sheep are unaffected by any ordinary quantity of the drug.

Chronic poisoning or ergotism rarely occurs in animals owing to continuous ingestion of ergotized grains. It is characterized by gastro-intestinal indigestion, with nausea, vomiting, colic, diarrhoea or constipation, and abortion ensues in pregnant animals. In addition to gastro-intestinal irritation the symptoms naturally assume two forms: 1. The gangrenous form; 2. the spasmodic form. In the first variety of ergotism there are coldness and anesthesia of the extremities, including the feet, ears, and tail of quadrupeds, the comb, tongue, and beak of birds, followed by the appearance of passive congestion, blebs, and dry gangrene in the vicinity of these parts. The hoofs and beaks often drop off.

¹ Jour. Phys. 34 163, 1906.

² The alkaloids of Ergot. Jour. Chem. Soc. 91 337, 1907. These writers give a full literature on the subject.

³ Arch. Expt. Path. Par. 39 184.

⁴ Jour. Phys. 34 163.

⁵ Jour. Pharma. and Expt. Therapeutics. Aug. 1909.

Death ensues from general exhaustion. In the spasmodic form are seen tonic contraction of the flexor tendons of the limbs and anaesthesia of the extremities, muscular trembling and general tetanic spasm, with opisthotonos, convulsions and delirium. Death also occurs from ashenia.

Grünfeld fed various animals with sphacelic acid in food. In the cocks, gangrene soon appeared affecting the comb, next the wattles, tongue, lungs, and crop. In hogs, the ears became gangrenous and fell off. Horses and cows fed upon grains containing ergot lose their hoofs, ears, and tails. The cortin, according to Robert, acts through the nerve centers. Microscopic examination of the abdominal and thoracic regions shows a toxic polyneuritis.

Dr McNeil in describing the disease says

Ergot stimulates the nerve centers that cause the contraction of the small blood vessels supplying the different parts of the body and cause one of the two forms of ergotism, namely, a nervous form, and a gangrenous form.

Nervous Ergotism In this form the contraction of the blood vessels of the brain produces dullness and depression. The animal also suffers from gastro-intestinal catarrh, refuses food, and gradually passes into a condition of general wasting. The nervous form, however, may assume an entirely different aspect and the animal dies suddenly in delirium or spasm, or gradually from paralysis.

Gangrenous Ergotism: In this common form the checking of the blood, resulting from the contraction of the small blood vessels, causes a loss of a part or of all the limb below the knee or hock, the tail, or the ears. This form of the disease may manifest itself by the formation of ulcers at the top of the hoof or between the toes, and a toe may be lost or the entire hoof shed. The affected part dries, a small furrow or line of separation appears, completely surrounding the limb, dividing the living from the dead mummified tissue.

DOTHIDEALES

Perithecia reduced, asci arising from the stroma and not separable from it, stroma present, not fleshy, black or dark colored ostiolum present

DOTHIDEACEAE

Stroma pulvinate, elongated, black or nearly black, coriaceous perithecia inseparable from the stroma, asci 4-8 spored, hyaline, yellowish or brown.

Phyllachora, Nitschke.

Stroma variable, elliptical, oblong or lanceolate, covered by the epidermis, black, roughened, ascospores ovate, elliptical, or oblong, mostly hyaline. About 200 species

Phyllachora Trifolii, (Pers.) Eckl.

Stroma on the lower surface of the leaf, gregarious, collected in small, elongated groups extending along the nerves of the leaf black subglobose, prominent, often confluent, ascospores elliptical, hyaline continuous, 10-20 μ . In the early part of the season small whitish or pale brown spots appear on the leaf, which contains the mycelium of the fungus. Dr Trelease says

This fruits on the lower surface, producing numerous tufts of necklace-shaped threads, each of which ends in a 2-celled, egg-shaped conidia-spore. These tufts of threads, which, like the spores, are of a deep brown color, are packed so closely together as to completely cover the spots, though under a hand lens they can be distinguished as separate panicles. To the naked eye they appear dead-black. Later in the season similar spots are occupied by small, coal-black fruits that contain stylospores. Winter spores, produced in asci, are not known. The conidial form of this fungus is especially common on white clover, though both forms are at times found abundantly on red clover and other species.

The *Polythrax* is common on red clover and is one of the numerous species which may be injurious to cattle.



Fig. 99. Black Spot of Grasses (*Phyllachora graminis*). A. Cross section of leaf through a black mass of the fungus. B. Perithecia. C. An ascus with ascospores. a, b. Spots in grass leaf caused by the fungus. After Frank and Trelease.

Phyllachora graminis (Pers.) Fekl.

Stroma scattered or confluent, penetrating the leaf and more or less prominent on both sides, covered by a black and shining epidermis, roughened; ostiols obscure, asci short, stalked, cylindrical, $75-40 \times 7-8 \mu$, ascospores 8, paraphyses present.

Phyllachora graminis, occurs on many cultivated and wild grasses; other species occur on clover and other leguminous plants. This parasitic fungus disease causes blackish spots on the lower or both surfaces of the leaf. The fungus causing these black spots on grasses has been called the black spot disease.

During August, and especially later, the coal black spots along the veins are especially prominent, they are considerably less than one-eighth of an inch in length and width and occur on both surfaces of the leaf, but are more abundant on the upper. These black spots are composed of dense mycelium, which in the green leaves bears numerous small spores which serve to propagate the fungus in the summer. In dead leaves, small perithecia are found, which contain numerous elongated bodies, the asci, within which are found eight small colorless spores, known as ascospores. These latter carry the fungus over winter.

Distribution and Hosts. Widely distributed in both Europe and North America, very common upon Quack Grass, Wild Rye, Bottle Grass, Panic Grass, etc.

Poisonous properties. The genus *Phyllachora* is abundant at times and is associated with stomatitis.

SPHAERIALS

Perithecia generally with a distinct ostiolum, of various consistency, not reddish or membranous, brown or blackish, stroma when present dark colored outside and whitish within. Contains the families *Sordariaceae*, found upon decaying plants and substances, *Chaetomiaceae*, with superficial perithecia, gen-

erally with short ostiolum and an apical tuft of hairs or bristles; one species *Chaetomium charitum* common on paper. *Sphaeriaceae*, with membranaceous perithecia, apex perforated with a simple pore, contains a large number of parasitic fungi like the strawberry rust or spot disease (*Sphaerella Fragariae*), and the spot disease of the currant (*Cercospora angulata*).

FUNGI IMPERFECTI

The fungi included in this group are simply form genera, many of the species belong to the *Pyrenomyces*, some belong to the *Phycomyces*, and some to *Hymenomyces*. In this connection we shall describe a few only which may cause trouble in forage.

Helminthosporium gramineum, Rabh. Yellow Leaf Disease of Barley

Spots in parallel rows, causing the leaves to become marked with yellow lines of pale green color; mycelium of the tissue colorless, conidiophores brownish on the surface, spores large 3-6-celled.

Distribution. Widely distributed in Europe and North America on barley.

Helminthosporium turcicum, Pass. Leaf Browning of corn

Spots sharply limited, conidiophores brownish elongated, bearing several brown spores. Widely distributed in Europe on corn, and also in North America.

Helminthosporium inconspicuum, E. & E.

Leaves dead and discolored, discoloration sometimes interrupted by spots of various sizes; conidiophores brown with several-celled conidia.




Fig. 100. Yellow Leaf Disease of Barley (*Helminthosporium gramineum*). a. Hypha arising from short cells. c. Conidium and to the left a cluster of conidiophores. d. Mycelium.




Fig. 101 Spores of Yellow Leaf Disease of Barley (*Helminthosporium graminum*). a. and b. Spores germinating. c. Conidiophore. d. Leaf browning of Corn (*Helminthosporium turcicum*). Spores and conidiophore to the left. To the right, conidiophore pushing through stomata.

Distribution and hosts. On corn, widely distributed in North America.

Poisonous properties. All of these fungi may be regarded as injurious, possibly producing stomatitis.

Scolecotrichum graminis, Fockel

Elongated brown or purplish-brown spots, the centers of which are gray or whitish and contain minute black dots; these small dark spots contain the tufts of brown fungus threads, which make their way out through the stomata. The hyphae are sometimes septate and the spores are usually borne at the end or occasionally in a lateral position, these fruiting hyphae bear small, smoky-brown, two-celled spores, the cells of the leaf become much altered, because the colorless threads of the fungus permeate them. On barley the disease is marked by brown or purplish-brown spots which appear on the leaf transversely.

Distribution. Widely distributed in Europe and North America.

Poisonous properties. May possibly produce mycotic stomatitis.




Fig. 101a. Spot Disease of Orchard Grass (*Syntherisma graminis*). Cross section of leaf, general fruiting layer of fungus with conidiophores and conidia. A, spores germinating. After Trelease.



General appearance of fungus on leaf of Orchard grass. After Trelease.

Polydesmus Mont. Rape Fungus

Sterile hyphae repent, fertile erect, simple or branched septate colorless conidia, interstitial filiform, concatenate, fusiform or clavate, many septate and opaque.

Polydesmus exilis Kuhn. Rape Fungus

Forming minute, punctate, elongated dark brown spots, conidia elongated or somewhat clavate, narrowed upwardly, 18-12 septate; the septa but slightly constricted, olive-brown in color. The conidia are 120-140 by 14-16 μ , conidiophores short, straight or slightly irregular, septate, making their way through the stomata.

This fungus is widely distributed on rape and cabbage and has been referred to as *Alternaria brassicae*. It is, however, thought to be a distinct fungus.

Poisonous properties. In Europe this fungus has long been associated with mycotic-stomatitis of cattle, but mycotic-stomatitis may be produced as indicated elsewhere, by other molds and fungi. This disease is characterized by inflammation and ulceration of the mucous membranes of the mouth. Salivation is a prominent symptom, the feet become swollen and sore. Dr. Mohler says

"Superficial erosions of the skin, particularly of the muzzle, and of the teats and udders of cows, may also be present, with some elevation of temperature and emaciation."

The disease is not serious and in many cases recovery occurs. But where treatment is not resorted to the disease may prove fatal, death occurring in from 6-8 days. Dr. Mohler states that in serious outbreaks it is about 0.5%.

The treatment should consist of first removing the herd from the infected pasture or inclosure containing the fungus. They should be fed on good wholesome soft nutritious food, plenty of cold water should be given. Dr Mohler recommends dissolving 2 heaping tablespoonfuls of borax or 1 tablespoonful of potassium chlorate in each of the first two buckets of water taken during the day. If the animals permit the mouth should be swabbed out with some antiseptic wash, such as weak carbolic acid or creolin solution, or permanganate of potash, or hydrogen peroxid. Mohler recommends that range cattle can be treated by the use of medicated salt.

"This salt may be prepared by pouring 4 ounces of crude carbolic acid upon 12 quarts of ordinary barrel salt, after which they are thoroughly mixed. The lesions of the feet should be treated with a 2 per cent solution of carbolic acid or of creolin, while the fissures and other lesions of the skin will be benefited by the application of carbolic vaseline or zinc ointment. If the animals are treated in this manner and carefully fed the disease will rapidly disappear."

Cladosporium herbarum (Pers.) Link

This fungus and its allies are very common upon oats sometimes very destructive. It attacks all parts of the plant, but is especially common in the heads. The mycelium of the fungus grows not only on the surface of the plant but also in the interior, the conidiophores and spores are olive green, the former pass through the opening of the stomata or break through the epidermis, the spores are 1- to 2-celled, borne on the end or on short lateral branches and are extremely variable in shape and size.

The general effect of the disease is to cause the kernels to shrivel. The disease, as recorded by Cobb, occurs rather destructively on oats. Professor Peck records the occurrence of a *Cladosporium* on oats, which he describes as a new species, the *Fusicladium destruens*. He says in regard to oats

"The foliage of the plants presented a singular admixture of green, dead-brown and reddish hues, strongly suggestive of that of a 'rust-struck' field."

Peck thinks this fungus inhabits the leaves of some of our northern grasses and has escaped from them to oat fields. Giltay reports that plants are infected in the same way as in some of the grain smuts, the spores being carried over with the seed, and that the disease can be prevented by treatment with hot water. A species of *Cladosporium* commonly affects the kernels of maize and is at times quite troublesome.

Septoria Fr

Perithecia imbedded in the tissues of the plant, appearing as small blackish or brownish spots, conidia generally multicellular and colorless, produced from short conidiophores. A genus containing numerous species of wide distribution. Many of them like the *Septoria* on the black currant and gooseberry, and the blackberry leaf spot, *Septoria rubi*, are troublesome parasitic fungi of cultivated plants. All of these fungi irritate the mucous membranes when found in abundance in the leaf.

Septoria graminum, Detm.

Spots at first yellow, then reddish-brown and finally whitish, perithecia blackish or brownish-black, spores 50 to 60 μ long and 1.5 to 2 μ wide, numerous, usually 2-celled.



Fig. 102 A. Leaf of Chest, showing numerous small specks, the perithecia of *Septoria Bromi*, the spores in the perithecia shown at 2.

In a somewhat extended account of this disease Cobb states that the entire plant is not always involved. The fungus is variable, its character depending upon the host which it attacks. On *Poa annua* the leaf is mainly involved and in many cases is totally destroyed. Cavara states that the spots on the leaves are small, elliptical, red or yellow, or the latter may be entirely absent. The injury it does to young plants is very great, in some cases their total destruction has been observed.

Janczewski who has studied the life history of *Septoria graminum* states that this represents the pycnidial stage of *Leptosphaeria tritici* and that the conidial form is the *Cladosporium herbarum*. We have not found the *Septoria* in Iowa though the *Cladosporium* is common.

The *Septoria tritici* Desm. is closely related to the above and should perhaps be regarded as nothing more than a variable form of *S. graminum*. The spots it produces are at first yellow, then reddish-brown, and finally whitish. The spores are 50-60 μ long and 1-5 to 2 μ wide and usually divided. A *Septoria* on the glumes of wheat in Ohio has been reported by Selby.

Several other species of *Septoria* are allied to the above species, one, the *Septoria bromi* Sacc. is common in Iowa on *Bromus secalinus*.

Diplodia, Fr

Perithecia bursting out sub-cutaneously, sub-carbonaceous, papillate according to type, spores ellipsoidal, ovoid or oblong, 1-celled, fuscous, perforated, basidia rod-like, simple hyaline. From the original genus have been separated five genera as follows. Species with superficial perithecia *Diplodiscia*; with hirsute perithecia *Chaetodiplodia*, with clustered perithecia *Botryodiplodia*; with mucilaginous spores *Macrodiplodia*, with hyaline spores *Diplodina*.



Fig. 103. Spores of *Diplodia Zeae*. 2. Young spores on the conidiophores with sporophores attached. 3. Germinating spores. 4. Dark swollen hyphae of *Diplodia*.

Diplodia Zeae (Schw.) Lév

Pycnidia black and spherical to pyriform, those forming on the husk or stalk developing within the tissues and breaking through at maturity, the greater number of pycnidia, however, occur between the kernels and are situated in a stroma. Conidia dark brown, cylindrical to elliptical, obtuse, straight or usually slightly curved and 1-septate, one to several oil drops in each cell, 5 μ in diameter, spores germinate in 18-24 hours in 3 per cent glucose agar at 26° C, in somewhat longer time when grown on corn agar, germ tube arises from near distal end at each spore.*

Distribution. A serious parasitic disease generally found where corn is cultivated, particularly in Illinois,* Iowa, and Nebraska.

Poisonous properties. This fungus is widely distributed in ears of corn and may be responsible for forage poisoning.

* Heald F. D., Wilcox, E. M., and Pool, V. W. The Life-history and Parasitism of *Diplodia Zeae* (Schw.) Lév.

* Burrill, T. J. and Barrett, J. T. Ear Rots of Corn. Bull. II. Agr. Exp. Sta. 133-65 109. 22 pl.




Fig. 103A. Cross-section of the peridium of *Diplodia Zeae* on a corn kernel showing sporophores, conidia and mycelium. After Burrill and Harrett

Although the fungus has been cultivated, the toxic substance has not been isolated. The writer fed mouldy corn meal to cats and rabbits. It produced injurious effects in kittens, three of these animals fed with mouldy corn meal and milk died from the effects. Unfortunately, in this case different moulds, *Aspergillus glaucus*, *Fusarium*, and *Diplodia*, were used.

Dr. Erwin F. Smith and Florence Hedges write as follows of this fungus:

"It is also worthy of inquiry whether this fungus may not be the cause of the so-called 'cornstalk disease' prevalent among cattle in the west. It is also possible that to *Diplodia* should be referred the great numbers of deaths of negroes in the south during past three years (1905-1909) from the so-called pellagra, following the consumption of mouldy corn-meal, and mouldy hominy. This fungus is also the cause of mouldy corn in Italy. The only other fungi we have reason for suspecting in this connection are species of *Aspergillus*."*

There are striking similarities between the so-called forage poisoning of cattle and the Pellagra disease in Italy and they are probably referable to some of the fungi found in corn. Dr. Miquel[†] in 1838 suggested that a *Mucor* was the cause of Pellagra.

Fusarium, Lk

Mycelium spreading, more or less effuse; conidia spindle-shaped or sickle-like, many-celled at maturity; conidiophores branching; conidia borne at the apex. A genus of numerous species, many of which are of uncertain affinity, usually found on dead organic matter but several are known to produce diseases of cultivated plants, like *F. Lycopersici*, Sacc. which produces the "Sleeping Disease" of tomatoes, the mycelium occurring in the vessels of the roots and causing a wilting. The *Fusarium luvensis*, Briest, produces a mar-digomma, or foot-rot, of orange and lemon trees. The *Fusarium vasinfectum*, Atk., produces a disease of cotton, known as "frenching." The cotton wilt is caused by a species of *Fusarium* and the perfect form of this fungus according to E. F. Smith is *Necosmospora*.

* *Diplodia* Disease of Maize (Suspected cause of Pellagra). Science 30 60-61

† Die Noord-Nederlandsche vergiftige Gewassen. 43 Amsterdam 1839



Fig. 104. Wheat Scab (*Gibberella Saubinetii*). perfect form of *Fusarium roseum* L. 1. Wheat affected with wheat scab, upper portion destroyed. 2. Glumes covered with perithecia. 4. Perithecia. 3. Asci from perithecia with ascospores, one of these enlarged at 5. 7. Conidiophore and spores grown in agar. After Selby

Fusarium roseum, Link

Mycelium whitish or varying from yellow to orange, appearing at the time when the grain begins to turn, the head, or part of it has a whitish appearance and the chaff is glued together conidiophores branched, spores terminal or lateral, crescent shaped at first, 1-celled, finally 2 or more celled; color of the conidia white or in masses orange or pink. According to Saccardo the ascigerous stage is the *Gibberella Saubinetii* (Mont) Sacc with gregarious perithecia, coriaceous, or somewhat membranaceous, somewhat blackish in color asci oblong, lanceolate, ascospores fusiform, 3-celled. Definite cultural experiments have not been made in this country to determine the relation of this fungus to the *F. heterosporum*.

According to Burrill and Barrett* several forms of *Fusarium* occur on corn. Saccardo in a letter to the writer identified the common Iowa *Fusarium* on corn as *F. heterosporum*.

The *Fusarium heterosporum* Nees, is common in parts of Germany, and Tubeuf quotes Frank as stating that the destruction of rye is total in some

* Bull. Ill. Agrl. Exp. Sta. 133.




Fig. 105. *Fusarium*. 5. Macroconidia of *Fusarium* with the felty mass of mycelium. Produces a deep pink color. 6. Mycelium. 7. Corroded starch grains. 8. Conidiophores or sporophores. 9. Microconidia and macroconidia of another corn *Fusarium* frequently infecting isolated grains. 10. Mycelium of the same. 11. Microconidia and macroconidia of another *Fusarium* on corn, which produces a dense felty mass extending between the kernels to the cob. 12. A spore-producing hyphae in prune juice culture. 13. Germinating spores of one of the species. 14. Hyphal branches of the same, with microconidia and macroconidia. After Durill and Barrett.



Fig. 106. Moulds and bacteria from corn. 1 and 3. *Fusarium Aetiosporum*. 1. Mycelium. 3. Conidia. 2 and 6. Other moulds. 4 and 5. Bacteria. After Pammel and King.

places, the fungus investing the whole kernel. Rotrup mentions it as destructive to germinating barley. It also occurs upon ergotized rye and is regarded by some mycologists as distinct from *Fusarium culmorum*. It is probable that the various species of *Fusarium* infesting cereals should be referred to one species.

Poisonous properties. Whether this fungus is responsible for the disease referred to by Dr Mayo and other veterinarians, has not been definitely determined. It is true that experiments made by Dr Bunting, this writer, and others, show that no doubt the *Fusarium* fed in considerable quantities to cats and dogs has had an injurious effect. Cats did not relish milk in which this material had been placed. If nothing more *Fusarium* may be looked upon as producing stomatitis. Prof Sheldon refers this fungus to *Fusarium moniliforme*. In the diseased horses reported by Dr Peters the horses would lose their hair and hoofs and were said to be alkali-fed. Cattle and hogs were likewise said to lose their hair. Feeding experiments conducted on hogs with this corn as well as with pure cultures reproduced the symptoms in experimental animals. In this connection the statement of Dr Law's is of interest.

Fodders affected with cryptogams or bacterial ferments are undoubtedly at times the cause of encephalitis. Veterinary records furnish many instances of wide spread attacks of stomach sickness, abdominal vertigo, and cerebro-spinal meningitis in wet seasons, when the fodders have been harvested in poor condition or when from inundation or accidental exposure they have become permeated by cryptogams and microbes. Among comparatively recent accounts of this are those of Martin and Varnell (musty oats), Lombard, Depra, Erbe, Falck, and Twell (mould), Bouley and Barthelémy (musty fodder), and Ray (fermented potatoes). One of the most extended local outbreaks of cerebro-spinal congestion I have ever seen, occurred in the pit mines of the Wabigoon coal mines, which led on Canadian hay which had been soaked with rain in transit and had undergone extensive fermentation. It should be noted that there were the attendant factors of overwork, in anticipation of a strike, and a Sunday's holiday above ground in a bright summer sunshine.

The experimental administration of moulds, yeasts and microbes, have in the great majority of cases led to little or no evil result (Gangwe, Mayo, Dinwiddie, etc.) and there is a strong tendency to discredit the pathogenic action of these agents in reported outbreaks. The safer conclusion perhaps would be, to recognize the fact that they are not equally pathogenic under all conditions of their growth and administration. The oft-recurring epizootics of brain disease in connection with wide spread poisoning of the fodders in remote and recent times, probably imply that cryptogams or microbes and their products, plus some condition not yet fully understood, are efficient co-operative factors. If we can discover this as yet unknown factor and demonstrate that it operates with equal power in the absence of cryptogams and ferments, as in their presence, it will be logical to pronounce these latter as non-pathogenic under all circumstances. Until then cryptogams and bacteria must be held as probable factors.

In recognizing how much cryptogams and bacteria vary under different conditions of life, and what various products they elaborate at different stages of their growth we can theoretically explain the absence of the disease at one time and its presence at another under what seem to be identical circumstances, as also the variety of symptoms shown in different outbreaks. While this explanation cannot be said to be absolutely proved, it is not antagonistic to the facts in many of the best observed outbreaks, and may serve as a hypothetical working theory until actual demonstration can be furnished. The affection suggests a nervous system introduced from without, rather than a disease due to a germ propagated in the system.

In all probability as we learn more of the true pathology of the disease, we shall come to recognize not one but several toxic principles, and several different affections each with its characteristic phenomena in the somewhat red state affection still known as cerebro-spinal meningitis.

The malady has been described in horses, oxen, sheep, goats and dogs, attacking by preference the young, which is not yet inured to the unknown poison, and by preference in winter and spring, the periods of close stabling, dry feeding and shedding of the coat.

Dr. R. A. Craig of Indiana reports as follows in regard to interesting experiments made at the Indiana Station

In January four sacks of spoiled, mouldy corn were gathered from a stalk field adjoining a field in which cattle had developed cornstalk disease. A healthy heifer weighing three hundred and fifty pounds was fed four to five pounds (twelve to seventeen ears) of this corn twice a day. In addition mowage was fed. On the afternoon of the sixth day of the test the heifer appeared weak, went down in the stall and was heaped up twice in the afternoon. When down she struggled some, and when helped up "shivered as if cold. In the evening she was still trembling and appeared weak. The weakness disappeared the following day. A few days later a slight twitching of the body muscles was noticed. The feeding test extended over a period of sixteen days. Her appetite remained good throughout the test.

Dr. Craig adds the following

During the fall and early winter of 1898-'99, Biting reported losses in horses and cattle, supposed to have been due to feeding on spoiled corn. By feeding corn meal that was inoculated with a pure culture of a mold (*Fusarium* sp.) made from the spoiled corn, he produced salivation and redness of the gums of the two horses used in the experiment. Later spoiled corn was fed. On the fifth day one horse showed a slight salivation, colicky pains and diarrhoea. On the seventh day noticeable trembling and moving about and stupor. For two days the animal stood with the head pressed against the wall. A quick recovery followed and the nervous disease from which horses were reported as dying did not develop. The second horse showed nothing more than a slight irritation to the mouth.

Because of the close resemblance between toxic poisoning from sorghum and the symptoms of corn stalk disease, Price deemed it advisable to examine cornstalks for the substances which produce prussic acid in plants. Samples of stalks from fields in which cattle had died were obtained. In these samples he discovered an enzyme which had the property of decomposing a glucoside (amygdalin) and thereby poison as a result of enzyme was found. However no glucoside capable of forming this poison as a result of enzyme action was found. The results were not regarded as conclusive as only a few samples were examined, and the failure to discover a suitable glucoside did not prove its absence in the corn plant, or in other plants in the field.

Dr. Peters says in regard to the feeding of moldy corn to horses as follows

Numerous reports have been received from stock owners of a disease which they call cornstalk disease or spinal meningitis which affects horses in the stalks and also some which have not been in the stalks.

This disease is very rapid in its course. For this reason it is sometimes difficult to see animals alive or in the beginning stages of the disease. In the later stages the animals are usually in such violent excitement that the symptoms have to be studied from a distance. One peculiar feature about the disease is that it comes on without warning, often attacking an animal while at work. One of the first symptoms noticed is the refusal of feed. Some have observed an excessive thirst and a difficulty in swallowing. The head is drooped in a very peculiar manner denoting stiffness. The eyes become very dull and later almost totally blind. This is usually followed by delirium and death. When a horse becomes affected in the stall it sometimes presses its head against the manger or wall and as this symptom increases in violence it is not uncommon to find the stall and manger demolished. Another peculiarity of the disease is that just before the animal becomes violent, one can cross its legs and the animal will remain in the position semi-conscious for some time.

This disease has been attributed to many causes. It is practically conceded at this time that it is due to a fungus found on the food administered. Feeding experiments with mouldy corn at our Station and other Stations, have proven that mouldy corn is capable of producing this disease. In March, 1901 a quart etc of mouldy corn, which was taken from cribs of a farmer at Graf, Nebr., who had lost a number of horses with this disease, was fed to four horses. These horses were fed exclusively on this corn with a small quantity of good hay and on April 2 two of the horses were found to be affected, the symptoms being

the same as those in animals that died in various parts of the state. The black horse, John, was found in the afternoon of April 2 to be swaying in his gait. He refused feed, had the peculiar dullness of the eyes, and when his limbs were crossed he remained in that position entirely motionless for some time. The next morning the animal was very much worse and at noon he was killed so that an autopsy could be held. The post-mortem examination showed that all organs were practically normal except the brain which was much softer than normal. I will quote the description of the post mortem as given by Dr. Butler, which is as follows: "On removal of the brain the superior surface of the right cerebra, hemisphere was noticed to be slightly flattened over the anterior half. Palpation revealed a soft spot at this place. An incision through the apparently sound gray matter revealed what Mayo described as a serous abscess in which floated flocculi of broken down brain substance, which presented the appearance, as one stockman said, of a mixture of vinegar and curdled milk. This portion of softened and broken-down white brain substance is in no sense a serous abscess. The line of demarcation between the broken down and the healthy brain substance was not clearly marked, but surrounding the completely broken-down portion of a zone probably half an inch thick that was softer than normal and of a slightly yellow color. The liquid in the cavity, and in which floated portions of soft and partially broken-down brain substance, was slightly yellow, but in no instance was clotted blood or any other microscopic evidence of a hemorrhage to be found."

Dr. Butler and Dr. Mayo conducted an experiment with some mouldy corn from a farmer who lost four registered Percheron horses at Wakefield, Kansas. Four hundred pounds of the worst of this corn and fifty pounds of the chaff and screenings were sent to the Agricultural College in Manhattan, Kansas, and a feeding experiment was started with two colts, twenty-three months old. The experiment began on July 16, when each colt received $1\frac{1}{2}$ kilos twice daily. On July 22, they were fed $1\frac{1}{2}$ kilos of corn and cob meal twice daily. On the 26th of July $1\frac{1}{2}$ kilos of the damaged corn, well ground, cob and all. This was continued until August 19. One colt died August 21. Another experiment was conducted with a two year old colt, but fed with mouldy corn and good prairie hay. The temperature of the animal varied from 101-102° F. This colt died on July 26, the feeding experiment having




Fig 107 Dermal mycosis associated with Sarcoptic mange caused by *Pseudomonas equinum*, conidia and mycelium. 2-6. Conidia (macroconidia) in various stages of development. 4. Germinating. After Melvin and Mohler

begun on June 30. It was observed that after three weeks there were no notable changes except a gam in flesh.

It seems to me that there can be no question from these experiments that mouldy corn is dangerous to feed to animals.

Oxen, sheep, and dogs are also affected with a form of meningitis, due to mouldy conditions. Of course, it should be stated that in the above description by Dr. Law, no special mould fungus is referred to.

Fusarium equinum, Norgaard. Itch Disease of Horses

Mycelium immersed, septate and branched, conidia in cultures sickle-shaped segmented, in hair sacs and sebaceous glands spindle-shaped or crescent-shaped bodies.

Drs. Melvin and Mohler supplement the above characters as follows:

The *Fusarium* possesses three forms of spores, the microconidia small and oval, non-septate or two celled, the macroconidia large falcate, with sharp lanceolate ends, 3-5 septate, forming many aerial threads. 35-55 μ long 2 $\frac{1}{2}$ -4 $\frac{1}{2}$ μ wide the chlamydospores oval or oblong, thin walled, densely granular. 8-12 μ in diameter. The macroconidia occur during the later stage of growth. On culture media there is a white growth which becomes slightly colored. The most favorable medium is potato and sterilized bread but it grows well in agar, glucose, or saccharine agar.

Distribution. North America, California to Idaho.

Pathogenic properties. In December, 1901 Victor A. Norgaard contributed to Science an account of a disease affecting horses, said to be produced by a fungus in which he suggested that, pending investigation, the name *Fusarium equinum* nov. spec. be given. The following is an abstract of the article in question.

An epidemic skin disease appeared among the horses on the Umatilla Indian Reservation, Pendleton, Oregon, upwards of sixty percent out of six thousand horses having been affected. The disease manifested itself through severe itching and loss of hair over a most the entire body. Many of the animals died of starvation. An examination of samples of the skin was made in the Pathological Division of the Bureau of Animal Industry and the presence of *Sarcoptes equi* observed. However these parasites were not present in sufficient numbers to account for the almost complete alopecia, and examination of samples almost entirely denuded of hair failed to show their presence. Microscopic examination of sections of the skin stained with borax blue showed the presence of large half-moon spindle-shaped bodies, deeply stained, in the hair sacs and sebaceous glands. Further culture produced from one to five circular colonies of a fungus which grew rapidly and assumed a salmon pink color. Cover-glass preparations made from these colonies contained numerous sickle-shaped segmented spores, characteristic of *Fusarium*. Of the twenty-five known varieties of this fungus, according to Dr. Erwin F. Smith, hitherto none has been known to be pathogenic to animals.

Drs. A. D. Melvin and J. R. Mohler have given a somewhat more extended account of this form of dermatomycosis. They found present with the disease the *Sarcoptes scabiei*. In 1901 the disease appeared in a very aggravated form, some 2,500 animals were diseased out of 6,000 animals on the Umatilla Indian reservation. It is supposed that this disease was introduced from California in 1902 from trailed horses. It appears that the fungus apparently enters the hair follicles, penetrates between the cells of the epidermis or abrasion of the skin and involves the surrounding tissue, causing irritation, followed by pruritis, the animal attempting to rub itself against anything with which it comes in contact. When the scurf is rubbed off by the finger nail there is left in its place a red moist denuded surface. It affects almost the entire body except the knees and hocks. The crusts are of gray color at first but turn darker. When the tissue

is examined microscopically, the spores are found to be abundant in the hair follicle, the fungus causing the hair to drop out. The inflammatory process spreads in the sebaceous glands causing a suppression of the excretion with the formation of crusts and scab. Occasionally *Sarcoptes* were found but Drs. Melvin and Mohler do not believe that they were the principal cause of the disease, although when present the animal parasite may aggravate the trouble. All ages and breeds of horses are susceptible as are both sexes. The animal stands around the rubbing posts all day and finally dies.

Rabbits, dogs, guinea pigs are immune. Experiments with horses were not successful, which these writers think may be because the right stage of the fungus was not used for inoculation experiments.

Treatment The authors recommend dynamo oil and sulphur in the proportion of one pound of the latter to a gallon of oil. Coal tar sheep dips have also been used.




Fig. 109. Fungus of mouse (*Oospora porriginis*) from a culture. a. Mycelial threads. b. A single thread more highly magnified. After Flügge.

Oospora, Wallr

Fungus with small tufts spreading or pulvinate, mucedinous, loose, or somewhat compact, fertile hyphae, short, with few branches, conidia transparent, usually in chains, globose or ovoid, hyaline or slightly colored.

Oospora porriginis (Mont. and Berk.) Sacc. *Achorion Schonleinii*, Remak.

Favus, *Tinea favosa*. Honeycomb Ringworm

Mycetium flexuose, simple, branched, or forked, continuous conidia, ovoid, triangular or somewhat culical, varying 3-6 μ in diameter. mycelium in masses with granular protoplasm occasionally branched at the end, the ends swollen, club-shaped, branches of the mycelium with lateral branches, spores oval, round or angular, 3-8 μ long and 3-4 μ wide, single or in chains. The threads of the fungus are readily detected in the bulbs and the shafts of hairs when sodium nitrate or potassium hydroxid is added, but at a distance of two inches the fungus cannot be detected. Sections of the nail stained also show threads

of the fungus. This fungus has been cultivated in ordinary agar or by Kral's method. It grows well at higher temperatures, and in 24-48 hours the fungus threads appear. The spores germinate at 35° C in 14 hours, and in 24 hours a fine mycelium appears in the air. In nutrient media like potato, gelatin and agar, chlamydospores and yellowish bodies appear. The organism requires higher temperatures for its best development, the optimum is 35° C. The organism from some of the lower animals, however, grows at lower temperatures.

According to Walsch the best development of the fungus in the hair is near the upper end of the root, from here it extends upward or downward, the mycelium may be exfoliular or on the surface of the hair.

The favus of man does not differ especially from that of animals except in color, and in the shorter duration of the disease in animals.

Distribution. Widely distributed in Europe and North America, but, according to Hyde and Monigomery, less common in the United States, Austria, and England than in France, Scotland and Poland.

Pathogenic properties. This form of dermatomycosis known as Favus was discovered by Schönlein in 1839. In the middle ages, it was known as *Tinea*, meaning a moth or worm. Previous to the discovery of the organism by Schönlein, various troubles were classed as favus.

Heusinger suspected the fungus nature of the disease as early as 1826. Remak, in 1845 cultivated the organism upon apple and transmitted it to his arm. He named the fungus *Achorion Schönleini* and in medical literature it is frequently referred to by this name.

The favus organism of mice was discovered in 1850 by Bennett and recognized by subsequent investigators like Schroeder and Simon. Favus of cats, guinea pigs, and dogs was recognized by Saint Cyr. Gerlach found it in birds and Gruby three years after the discovery of the organism by Schönlein, found it in hairs on the heads of children and the hairs of the beard. In recent times, various views have been expressed with reference to the nature of favus and trichophytosis, it being held that these diseases are produced by different fungi. Pick in 1887, Walsch in 1896, and others considered the fungus to be polymorphic. Quincke distinguished three varieties but pathologists are not agreed on this point. Plant in his discussion of the parasitic fungi divides them into the following groups: the Favus and the Trichophytic groups. According to this view, the favus of man and that of animals are regarded as distinct fungi.

The disease in man generally appears where hairs occur but may appear also in other parts of the body as the eyes, nails, etc., seldom becoming general. Favus of the nails is called Favus onychomycosis. In lower animals the disease may occur on the head, nose, ears, back and more frequently, is generalized. Formerly this disease was common among the poorer classes. Today, it is not common in France, Holland, Scandinavia, Germany, England, Switzerland, Japan or America. On the other hand, it is rather frequent in Russia, Scotland, Italy, Spain, Asia, Austria and Egypt, young individuals being more susceptible to it than are older persons, probably acquiring the disease by contact. The disease is spread through spores of the fungus. The mycelium itself, according to Grawitz, is not capable of spreading the disease of animals. In man it is recognized by the development of minute yellowish or reddish points, delicate vesicles may surround these spots. Later, the fungus may develop a

erect the skin is very disagreeable being compared to the skin of man or the skin of cats.

Mice are most susceptible to the disease, all young mice since they come in contact with mice which have taken them up by means of the skin and finally guinea pigs.

According to the recent investigations of Frank (1891) three different types of fungi were isolated from mice and guinea pigs. The first type was the *Trichophyton* which has been found in the skin of mice and guinea pigs. The second type was the *Microsporum* which has been found in the skin of mice and guinea pigs. The third type was the *Trichophyton* which has been found in the skin of mice and guinea pigs. These may possibly be regarded as the most common causes.




Fig. 1. A plant stem heavily infested with a dark, irregular, and extensive growth of a fungus, likely a Trichophyton or Microsporum, showing the characteristic lesions of a fungal infection.

Sporotrichum, Link

Hyphae, branching irregularly and repeatedly, septate or continuous, usually equally procumbent, conidia acrogenous at the apices of the main and lateral branches, usually solitary beneath, ovoid or subglobose. This genus differs from *Botrytis*, especially in all the hyphae being procumbent and the conidia subsolitary, from *Trichosporum*, in never being dark colored. Very many species, imperfectly described by older writers, show mere forms, or mycelia.

Sporotrichum Furfur, Rob

Plaque yellow or yellowish brown to dark brown or brownish red spots varying in size from that of a lentil to that of a hand, either smooth or shining or dull, exfoliating. Found on the breast, stomach, or back. Never upon the hands, feet, seldom on the face. Slender hyphae 3-4 μ wide 7-13 μ long, variable as to length and thickness. The spores are clustered resembling a drop. On potato a characteristic growth of yellowish, orange red brown, blackish or greenish color. Old culture is grayish, brownish or violet color. In 3-4 days a whitish gelatinous mass forms, which in 1-4 weeks covers the whole surface. Conidia areidium-like surrounded by thick hyphae, occur in scales, 4-7 μ spherical. In cultures budding occurs.

Distribution. Common in some localities in Europe and America.

Pathogenic properties. Feht in 1840 observed that most of the inhabitants in a Swiss village were infected through cattle. Bazin in 1853 observed that many cavalry men were infected through horses. Papa in 1840 observed that this disease was frequently transmitted to men. In cities it chiefly occurs in cats and dogs, and through these it is conveyed to men.

It is especially common in people with tender skins and in tubercular patients, and is more common with women than men. In 1846, Fehstedt dis-




Fig. 110. *Sporotrichum Furfur*. After Kaposi.

covered the cause while Robin named the fungus *Macrosporon Furfur* K  bner, in 1866, made the first inoculation experiments.

Grawitz, in 1876, first cultivated the organism. His work was followed by that of such other investigators as Sehlen in 1890 and Koltjar in 1892, who succeeded in transmitting the cultivated fungus to guinea pigs and gave it the name *Oidium minimum*. It has also been named *Oidium Furfur*. Vuillemin also cultivated the fungus to which he gave the name *Malassezia Furfur*. Pure cultures of the organism grown on potato were transmitted to man.

Treatment. The best treatment, as recommended by Hyde and Montgomery, is a hot bath, the skin being rubbed with soap, following this the skin is bathed with clean water and sponged with a solution of sodium hyposulfite, 1 drachm (4) to the ounce (30)




Fig. 111. Erythrasma Fungus. *Sporotrichum minutissimum*. After Hyde and Montgomery

Sporotrichum minutissimum (Burckhardt) Paumel Erythrasma

It begins as small brown or reddish patches which become confluent, these spots may become as large as the palm of a hand and occur in the axial region, the scales contain Leptothrox-like threads which are branched and septate conidia small, round or angular. It grows well on agar agar, glycerine agar, gelatine, potatoes, and in blood serum. In nutrient media branching, septate hyphae 0.8-1.3 μ in thickness and 5-15 μ in length are seen, the short hyphae break up into numerous spores.

This disease was first observed by Burckhardt in 1869 and since has been observed by others. It occurs in the form of roundish or punctiform patches, sharply contrasted with the adjacent tissues. The younger areas are livid red while the older are yellowish or brownish.

A so-called Dhobie itch of the Philippine Islands is in part caused by this fungus. Though Hyde and Montgomery state that there are two other types of infection known by this name, one is trichophyton and one of bacterial origin.

Sporotrichum giganteum, (Unna) Pammel

Spores are free or in chains in nutrient medium 1.5-7 μ in diameter, the oval 4.5x3-6 μ , yeast-like budding resembling *Oidium lactis* also occurs; ectospores as well as chlamydospores present, the latter 8-12 μ in diameter with strongly refringent bodies in cultures ray like, the rays consisting of hyphae and spores, in liquid media only hyphae with ectospores. In the hair, knot-like bodies are formed with spores and hyphae embedded in mucilage.

The Colombia disease was described as *Trichosporon giganteum* Unna. It is a polymorphic fungus and the *T. ovoides*, Behrend is included but by some is regarded as different. The knotty masses of hair are less thick, and the spores are oval in shape, gelatine not liquefied. The superficial colonies resemble *Oidium lactis*. Chlamydospore 4-12 μ in diameter.

Other species of *Microsporon* have been described like *M. canis* in dogs, *M. tigris*, the *M. equi*, in horses and colts and another species in cattle. In calves a similar form occurs. The sheep are said to have the disease on the neck and breast, it also occurs in hogs, goats, and birds. These forms are said to differ slightly clinically. Pus is formed in follicles, and the hair is especially prone to drop out. A bad smelling liquid of a reddish color occurs. In lesions large spores, the ectospores, occur. The Onium chain like spores occur chiefly in the roots of the hairs.

Pathogenic properties. The disease is especially common in animals in which the skin is naturally fine, thin, and dry, and covered with hair sparsely, more common in the Arabian Barb, English racer, and American trotter of nervous organization than in heavier draft breeds. Old horses are more subject than young ones. Extended desquamation, excessive production of epidermal scales without any elevation of the skin, scurfy products may be found in patches scattered over the body, generalized or circumscribed as to the head, ears, crest and tail the hair may be pulled out with great ease. In cattle it occurs on neck and develops in connection with anaemia, spoiled fodder, and constitutional predisposition. Affects especially the head, neck, and back of dogs gorged with dainties and those becoming aged. The affected parts are covered with a floury or bran-like product lying upon a dry surface, the affection being usually limited to certain areas more or less destitute of hair. In the cat it may affect the whole dorsal aspect of the body being associated with extreme electrical susceptibility, the hair when touched, collecting in tufts. The scaly product is abundant.

Animals are said to spread trichophytic fungi, which supposition is important from a hygienic point of view. Since the disease sometimes occurs in school children, separate hooks for clothing and separate towels when bathing the hands and faces are recommended. A one per cent solution of bicloride of mercury will kill the fungus.

The disease was first described by Osorio in 1846, and was then thought to be confined to Colombia where it was called Driedra (Stone), but later was found in Europe, and Vuillemin reported it from Paris in 1902. Desenne of

Paris discovered the hyphae of the fungus and Malcolm Morris the spore-like bodies. Behrend in 1890 succeeded in cultivating the fungus.

Sporotrichum tonsurans. Barber's Itch

Hyphae slender 1.5-2 μ in diameter, straight, undulated, dichotomous, septate or non-septate, penetrate the hair follicles forming a matted mycelium. small pustules and scabs, in places devoid of hairs it forms red, scaly spots, discs and circles. The fungus is found between the uppermost layers of the epidermis just beneath the corneous stratum, the conidia are small, spherical or elliptical sharply defined 2-3 μ in diameter; the spore masses surround the root of the hair and are frequently densely and closely arranged like beads, in culture media like agar a many-rayed fungus occurs, the color varies with the medium, yellow, Bismark brown, cherry red, violet, rose, brown, blackish brown gelatine liquefied, spores swell after a few hours and produce 12 germ tubes from a single spore, mycelium with occasional swellings, ectospores formed in 60-96 hours, also small, branched air hyphae, small conidia 1.5-3 μ in diameter borne on short lateral branches. The Botrytis-like spores rise on the long thin, curved air hyphae. Oidium budding does not occur in nutrient media. The spores retain their vitality for six months but exposure to 45° C for a few hours will kill the organisms, it is sensitive to sunlight and common disinfectants. The fungus is polymorphic, one form having been classed, by Sabourne, with Botrytis. The large-spored trichophyte found on the scalp germinates at 37° C in a few hours but at room temperature, a much longer time is required.

Conidia 5 μ in diameter an abundant mycelium with dust-like growths, and, in three days, oidium-like spores, as well as ectospores, are produced on the potato, the disease being known as *Tinea Sycosis*.

The *T. circumscripta* produces, in animals, cherry patches each with a raised border and scales, and is also found on the head, arms, and neck, of man. *T. disseminata* produces small red pustules.

To *T. tonsurans*, also, is attributed *Picsema marginatum* which Kobner, while making a study of trichophytic fungi in 1864, recognized as a trichophyte. He also determined that the fungus on the nails, described in 1853 and 1855 by Baum and Meissner, was a trichophyte.




Fig. 112 Barber's Itch
(*Sporotrichum tonsurans*)
Filaments and spores. After Hyde
and Montgomery

Gerlach, in 1857-1859, demonstrated a trichophyte in bovine animals, and other investigators, later, recognized the case as *Herpes tonsurans*. *S. tonsurans* has also been described under *Trichophyton tonsurans* and as *Oidium tonsurans* Unna, in 1897, from twenty cultures described four species, *T. oidiophora* *T. erelmorphorum* *T. atractiophorum*, and *T. pterygodes*. It is probable that *Sporotrichum tonsurans* is a very variable species. Lindau places it under the genus *Oospora*, but it seems preferable to call it *Sporotrichum*.

Distribution. Occurs in Europe North and South America, Asia and Africa.

Pathogenic properties. In cattle, small, round, sharply defined spots occur which are covered with scabs and scales which project more or less above the skin and vary in size some being as large as the palm of the hand. Underneath the scales, is a purulent fluid with hollows that represent the empty follicle. In man as well as in animals, the hairs can be pulled out very readily. The eruption lasts from six to twelve weeks, outbreaks occurring from rubbing or scratching as a relief from the itching sensation that accompanies the eruption. In sucking calves it occurs chiefly about the mouth and is called "doughy mange" and, according to Hahn, is produced by the fungus *T. tonsurans*.

Most of the varieties can be transmitted to guinea pigs, cats and dogs and have even been transmitted to man during the process of sheep shearing. Healing takes place when the animal is inoculated subcutaneously. The large spored form, occurring on the scalp, forms pus and resembles moist eczema. Children take the disease from calves and by playing with cats and dogs.

Mycosis of the beard exists in two forms non infectious and infectious. In *Sycosis parasitica* the disease is accompanied by a severe inflammation of the hairy parts of the skin leading to infiltration and suppuration. Sabouraud classifies the parasite into a dry and a pus favus.

The *Leisma marginatum* supposedly caused by the same fungus was first described by Deyergie in 1854-1855. Berensprung having discovered the same fungus in 1855. It is slightly contagious, and more frequent in men than in women. Another form of the disease occurs in the mouth. In sheep the wool is felted and beneath it are bran like scabby parts, the fleece becoming very ragged in appearance. In poultry it shows itself by the loss of feathers. In horses, it occurs most often on the seat of the saddle. The spots vary in size, and the surrounding hair can be pulled out easily. In dogs, it affects the head and extremities. Usually the spots are round at first and sharply defined, later becoming hairless patches. occasionally they are dirty gray scabs.

Ringworm of the body or *Tinea circinata*, is characterized by the occurrence of one or more pea shaped or large circular reddish patches which are on about the same level as the integument and rarely 5-6 inches in diameter. In some forms there is itching. This *Trichophyton* was discovered by Gruby in 1844. This fungus can be readily recognized by making microscopic mounts. The mycelium is less branched and the threads are more slender than in the form previously described. The spores are like strings of beads.

The ringworm of the scalp, *Tinea tonsurans*, is a disease chiefly of children especially of those in schools. It differs from the preceding form in the fact that the fungus makes its way into the hair follicles. The patches are, at first, circumscribed, about the size of a small coin, covered or partly covered with roundish patches of slate gray color or a dirty yellow. The fungus is called *Microsporum adonis*.

Hyde and Montgomery state that there are at least two distinct and unrelated forms capable of producing ringworm, the *Microsporon adonis*, a small-spored fungus and the *Trichophyton*, or large-spored fungus. The *Microsporon* appears under the microscope in the form of a large number of round spores irregularly grouped or massed about the follicular portion of the hair. The mycelial threads are all within the hair proper while the spores terminate fine threads on the other surface of the shaft. The spores of *Trichophyton* vary greatly in size and are much larger than those of *Microsporon*. They are cuboidal, oval, or irregularly rounded. They occur in chains, up and down the hair or shaft. The mycelium is found without, never within, the hairs. The spores may be within (endothrix) or without (ectothrix).

Odium albicans. (Robin.) Rees Thrush

Forms a mould-like growth in the mouth of man and lower animals. Vegetative cells, yeast-like, spherical, elliptical, oval or cylindrical, 5-6 μ long, 4 μ wide, the elongated hyphae-like bodies variable in length conidia elliptical in chains grows well in nutrient media where it produces superficial, spherical, white, wax-like, granular colonies, varying color from reddish to white, chlamydospores in nutrient media and occasionally in the epithelial layer, it does not ferment lactose and saccharose, but ferments levulose and dextrose. According to Brebeck and Fischer there are two morphological forms of the organisms, a small oval and a large-spored form, however, this distinction is not generally recognized.

Distribution. Widely distributed in the United States, also in other countries, Germany, Austria, France, Italy, and Great Britain.

Pathogenic properties. John recognized the disease in 1816, while Buchner gave a somewhat detailed description in 1841. Langbeck and Berg discovered the fungus in 1839. It was thought by them that it was the cause of typhoid fever. Langenbeck demonstrated that the Fungus could be carried from a child, sick with the disease, to a healthy individual. Gruby in 1847, described the fungus under the name of *Aphaphyta*, placing it near the fungus *Sporotrichum*, while Robin, a French author on parasitic diseases, considered the fungus to be an *Odium*, naming it *Odium albicans*, a name frequently used by authors. Rees, however, placed it with the yeasts. *Monilia candida* is regarded by Plant and Landau as a synonym. Grawitz in 1877, made pure culture of the fungus and succeeded in producing the disease in guinea pigs. Klemperer found that when the fungus was inoculated into the circulatory system of guinea pigs, general mycosis resulted. Limossier and Roux (1889-1890) in their monograph, state that the mycelium occurs in the blood vessels of inoculated animals.

The fungus is very common in some sick chambers in regions where the disease is prevalent. It is most abundant in sucklings.

It occurs frequently in children of premature birth, and in weak children, the fungus is also found in aged persons, suffering from disability, it occurs chiefly upon the mucous membrane of the mouth, pharynx, and oesophagus, more rarely upon that of the stomach, intestine and vagina and upon the nipples of nursing women and bovine animals. It has also been found in the liver, kidneys and lungs. It penetrates the epithelium and even into the underlying, connective tissue. It is spontaneous in such animals as calves, birds, and

foals. The disease is fatal in many cases, some authors estimating the death rate as high as 22 per cent. Inoculated guinea pigs show a rise in temperature at first a lowering, accompanied by albuminuria, loss of flesh, and diarrhoea, death occurring in from 3-7 days. Immunity may be obtained in guinea pigs by beginning with small doses, and increasing these gradually to three times the strength. The product produced by the fungus is poisonous, 20-40 cc of the whole substance will kill a guinea pig weighing one kilogram.

Dr. Stuhr has contributed the following upon this subject:

Thrush is a mycosis of the mouth affecting children, calves, foals, and poultry, and is characterized by the formation of white patches upon the mucous membrane, which vary in size from points to large areas. It may involve the pharyngeal and laryngeal mucosae by extension. The disease is transmissible from man to animals. Young age, a weak constitution, gastric indigestion, uneasiness, milky and starchy diet predispose. Decaying food in the mouth offers a suitable place for the growth of the fungus.

Etiology. The specific cause of thrush is a vegetable parasite, *Oidium albicans* first described by Berg in 1840. It is one of the branching fungi closely related to the yeasts and grows readily on sour milk, in saccharine substances, on decayed wood, and on fresh cow manure. Calves fed milk from wooden pails which are not kept perfectly clean are particularly liable to contract the disease. The fungus descends into the epithelium and sometimes into the subjacent connective tissue, causing inflammatory infiltration and superficial necrosis.

Symptoms. The mucosa is diffusely red, swollen and tender and shows adherent white patches, varying in size, surrounded by a red inflammatory zone. When these white spots are rubbed off, shallow red ulcers are exposed. When the inflammation in the mouth is severe, or when the disease spreads to the pharynx and interferes with deglutition the prognosis may become serious. Usually however the disease is benign and yields readily to treatment.

Lesions. These are usually superficially located and rarely extend deeply into tissues. They begin with diffuse reddening of the mucous membrane and the formation of a somewhat shining, slimy, adhesive layer of grayish-white matter which is said to have an acid reaction. Later whitish dots appear upon the surface and gradually spread, sometimes coalescing. These whitish patches are false membranes composed of detached epithelial cells with a ramifying network of parasitic threads. The white color of the false membranes is markedly in contrast with the congested surrounding tissue. While the lesions are ordinarily restricted to the mouth they may involve the pharynx, oesophagus, (in chickens), larynx and even the stomach and intestines. Metastasis may occur and the fungus be carried to various parts of the body.

Treatment. This is aimed at the destruction of the fungus and for this purpose many substances have been recommended. The mouth should be cleansed at frequent intervals with solutions of borax, sodium hyposulphite, permanganate of potash, or chlorate of potash, etc. The system should be built up by feeding soft nutritious food, and the sanitary conditions should be improved.

Oidium hominis. (Russe.) Pammel. Blastomycosis

Cells spherical or ovoid, variable in culture. 8 μ in diameter with strongly refringent bodies in young cultures, nearly homogeneous and with oil drops, in old cultures, large cells with a thin membrane, culture, at first white, then grayish or yellowish, or yellowish-brown in plum cultures, in plum decoction

black, gas-producing in sugar medium, grows well in ordinary media but preferably in acid; grows best at high temperature. (An organism isolated by Curtis produced white colonies with oval, or club-shaped cells, frequently producing capsules.) Grows readily in nutrient media, does not form a pellicle on the surface in liquid media.

Descriptions of several so-called species of *Saccharomyces* causing Blastomycosis will be added to the above. In their development, blastomycotic fungi resemble true hyphal fungi rather than yeasts. It is convenient to discuss them here until their true relationship has been determined.

Distribution. Found in Europe and America.

Pathogenic properties. It was first isolated from the left tibia of a woman thirty-one years of age, the disease having first manifested itself by a purplish-red spot and swelling. An operation was performed, but it failed to relieve the trouble, new foci making their appearance after the operation and finally becoming general, being accompanied with pus formation. The patient died in 13 months, the lungs, kidneys, and spleen having become involved. There was no oedema, the organisms found in the lungs and kidneys being marked by small, nodular swellings.

Large amounts of culture, when inoculated into guinea pigs, dogs, and rabbits, produce the disease followed by death. In another case described by Curtis in 1895, the disease occurred in a young man. The organism isolated is pathogenic for rats, mice and dogs.




Fig. 111. Blastomycosis of the skin showing the elongated cells and budding forms. After Hyde and Montgomery.

Ziegler in his *General Pathology* summarizes our knowledge of the pathogenic properties of yeast as follows:

As parasites no importance has been attached to them until very recently, but the investigations of Busse, Buschke, Sarsfield, Curtis, and others have established the fact that there are also species of *Saccharomyces* of pathogenic importance. According to these observations, the pathogenic yeasts can multiply in different tissues, in the skin, peritoneum, lungs, and glandular organs, and can excite either purulent inflammations or proliferations of granulation tissue, which run a course similar to that of an infection with actinomycosis or tuberculosis. In inflammatory foci, the yeast cells are for the most part provided with a capsule. They may give rise to tumor-like swellings. Through degenerate changes, crescentic forms may develop from the oval yeast-cells.

In solutions containing sugar the blastomycetes form oval cells. Reproduction takes place through budding and constriction: on any portion of the parent cell there may develop an excrescence, which is constricted off after it reaches the size of the mother cell. Under

certain conditions the cells may grow out into threads, but in these threads no subsequent segmentation occurs, jointed threads arise through budding (Cienkowski, Grawitz). A dilute culture-medium favors the formation of threads.

Oidium granulomatogenes. (Sanfelice). Pammel. Blastomycosis

Forms nodular masses; grows in ordinary media; ferments sugar; uniform clouding of media, colonies white; the nodules consist of the fungus, giant epithelioid cells, causes a cheesy degeneration.

Distribution. Found in Europe.

Pathogenic properties. Pathogenic for hogs, occurs in the lungs, where it produces nodular masses.

Oidium lithogenes. (Sanfelice). Pammel. Blastomycosis

The fungus occurs in the cancerous-like growth of the lymphatics, is frequently surrounded by lime, on agar and gelatin forms white colonies, in stick culture the growth is needle-like, sugar is changed into alcohol and carbon dioxide. Fungus consists of spherical bodies.

Distribution. In Europe

Pathogenic properties. Pathogenic for guinea pigs, white rats, sheep, and cattle, producing nodular enlargements, frequently surrounded by a calcareous capsule.

W. W. Hamburger, in a recent number of the Journal of Infectious Diseases, refers to a morphological and biological study of blastomycosis as follows:

1. The strains of organisms appear nearly identical, so far as growth in test-tubes goes. A few minor differences are summed up below.

2. The organisms grow vigorously on the usual laboratory media, with perhaps a slightly more abundant growth on faintly acid glucose-agar.

3. Temperature is perhaps the most important factor in varying the gross and microscopic morphology, room temperature favors production of mycelia and aerial hyphae, incubator temperature inhibits production of hyphae and favors coherent, waxy yeast-like colonies (budding forms).

4. Those cultures which produce yeast-like growths at incubator temperature develop hyphae within 24 hours when withdrawn and placed at room temperature. Likewise the majority of yeast-like colonies will finally (in 17 to 30 days) show evidence of beginning hypha formation even if kept at 37 degrees C.

5. Glucose-agar slabs and broth form the most serviceable culture media if a limited variety is at hand. Duplicates should always be made to control differences in morphology at room and incubator temperature.

6. Four strains of organisms isolated from four cases of generalized blastomycosis appear identical.

7. Pronounced variations in the gross and microscopic morphology of the organisms are produced by variations in temperature. As a routine for purposes of study cultures should be grown at both room and incubator temperatures.

Distribution. Found both in Europe and North America

Pathogenic properties. Dr. Harris gives the following

Towards the lower animals pathogenic properties vary very much with the culture recently isolated cultures as a rule proving more virulent than older ones. Mice, guinea pigs, and dogs are most susceptible, succumbing after a variable time to a peritonitis inoculations whilst the white rat, rabbit, sheep, and horse are more refractory. In all, the lesions may be localized in the form of abscesses, or general infection may ensue where subcutaneous inoculation is practised.

Dr. E. R. Le Count and J. Myers discuss the case of systemic blastomycosis of a Polish laborer

The first noticeable departure consisted in a feeling of discomfort involving the chest on the right side extending through from front to back, later cutaneous lesions appeared,

located below the left ankle and extended down to the heel. He was obliged to stop work in December. The patient was emaciated, pale, anemic, and weak. Marked edema was present in the ankles, feet, face and arms. His nails were clubbed, inguinal adenopathy was noted. From the lesions blastomycotic fungus was isolated, the sputum also containing the organisms.

Eisendrath and Ormsby described the cultures as follows:

On March 22nd pus was removed from a subcutaneous abscess on the left forearm, which was inoculated on various media. Six days later growth was plainly visible, and after this time the cultures grew rapidly. These proved to be pure cultures of blastomycetes.

In the pus they occurred as circular forms and budding forms, having a double contour and the usual refractile capsule. On media the growth varied. It presented a moist, pasty surface on glycerin-agar with at times a wormy appearance or else presenting large folds and depressions. Microscopically, these cultures showed many oval and circular organisms, some budding ones, and much mycelial formation, the latter being both coarse and fine containing spores. Lateral conidia occurred. On glucose-agar the growth was more dry, white, and presented aerial hyphae and macroscopically there were fewer circular and budding organisms and more fine mycelia. On both glucose and glycerin-agar the media were penetrated to a considerable depth in a semi-circular manner.

Des. Le Count and J. Myers say, as follows:

The body was examined a few hours after death and the following anatomical diagnosis made: Blastomycotic bronchopneumonia blastomycosis of the peribronchial lymph nodes, of the pleura, the subpleural, and retropharyngeal tissue, the liver, the kidneys, the colon, the spinal column (dorsal vertebrae), the external spinal dura, the cerebellum, the left elbow, both knees and ankle joints, and of the skin and subcutaneous tissue with ulcerations, fistulae, and scars. Fibrous pleuritis. Passive hyperemia of liver and spleen. Severe atrophy of adipose tissue. Emaciation. Adenoma of thyroid and accessory spleen.

One notable feature of this case is the large conglomerate blastomycotic nodule in the cerebellum. In only one other case of systemic blastomycosis, that of Curtis, is there any record of changes in the nervous system, and the statements in that instance are purely clinical, death being due to meningitis. The reproduction by a process of sporulation demonstrable in the cerebellar lesion is likewise a new feature of the changes encountered in the lesions of this disease. The idea that in the nervous tissue the fungus may have found favorable or different conditions of nutrition, as an explanation for this method of multiplication, is opposed by the facts that the regions in which it was found were very minute, that it was not generally present in the cerebellar process, and the budding was commonly observed in the "abscesses" in the partitions between necrotic regions.

Highly interesting is the relationship between this case of blastomycosis and one of coccidioides disease described by Ophula. Up to the present two of the chief differences between blastomycosis and coccidioides granuloma have been the endosporulation observed in the tissues in the latter disease and its tendency to spread by the lymph channels. Although an widespread extension by the lymphatics was demonstrated in the case reported here, the extension to the tracheobronchial glands and in peribronchial lymph channels is unmistakable. The endosporulation in the cerebellum in part also resembles the methods of production described for the organisms of coccidioides granuloma. Taken together, these features in this instance of systemic blastomycosis are in accord with the belief expressed by Ophula of a close relationship of the organisms in the two diseases.

Ricketts, in an interesting monograph on "Oidiomycosis (Blastomycosis) of the Skin and its Fungi," gives the clinical history, cultural characters and histopathology of a large number of cases. The fungi are divided into three groups: 1) Blastomycetoid or yeast-like. (2) Oidium-like. (3) Hyphomycetoid. He says:

There are two histological forms of the fungus in the skin, the eosinophilous and the non-eosinophilous, the former being associated with the mould type of the organism. Aside from the infections considered in this communication, certain cases which have been described in the literature from time to time indicate that oidium-like organisms may cause other severe pathological conditions in man.

LICHEN

Symbiotic organisms consisting of higher fungi, chiefly of the class *Ascomycetes*, or rarely *Basidiomycetes*, the thallus consisting of algal cells enveloped by the mycelium of the fungus forming a felted mass. The algae are called gonidia and belong to the *Cyanophyceae* or *Chlorophyceae*. The reproductive bodies consist of spermogonia, which contain the spermatia. The asci contain the ascospores, and occur in apothecia. In the Basical lichens, spores are borne on basidia. Lichens are sometimes divided into fruticose, crustaceous, and foliaceous, but a more natural classification arranges the lichens into the *Basical lichens* and *Asci lichens*, with various families, such as the *Roccellaceae*, that contains the *Lunus*, *Roccella tinctoria*, the *Lecanoraceae*, containing the *Lecanora*, and the *Cladoniaceae* that contains the well known Reindeer Lichen, *Cladonia rangiferina*.




Fig. 115. Lichens, structure of Thallus and Apothecia. *Protosiphia mamillata* section through a part of the apothecium, a (at the right) below, Gonidia, a above, nucleus, b ascospores, c paraphyses, d, e, f. Section of thallus of *Idonia furcata* x 110. 1. Portion of thallus of *Stereocaulon vema*, new, a gonidia, b hyphae, c isolated gonidium (a with attached hyphae), d, e, f. *Synaldis rambourensis*, isolated gonidia, a with attached hyphae, (m). After Roemer.

EMBRYOPHYTA ZOIDIOGAMA

BRYOPHYTA

Seldom thalloid, generally with stem and leaves with well marked alternation of generations. They contain antheridia and archegonia similar to those of the ferns. The antheridia are stalked, ellipsoidal, spherical, or club-shaped, the sperm cells are biciliated, the archegonia flask-shaped, the ventral portion with a large center cell, the lower portion divided into an egg cell and ventral canal cell. At maturity the new canal cells become mucilaginous and disorganized.




Fig. 114. Spermatogonia of Lichen. *g.* Gonidia, fungal threads below. *sp.* Spermatia. Greatly magnified. After Tuckerm.




Fig. 145b. Lichens. 1 *Ochrolechia tartarea*. 2 *Rhizocarpon geographicum*. 3 *Lecanora subfusca*, on bark of tree. 4 *Calium*. 5 *Graphis scripta*. 6 *Lecanora ascalenta*. 7 No. 6 removed from substrate. 8 *Graphis scripta*. After Wettstein.

the mucilaginous material acting as a servant to attract the sperm cells. After fertilization occurs, spores develop. In addition to the sexual method of reproduction an asexual reproduction also occurs. The gametophytic and sporophytic stage are sharply differentiated.

In the development of the sexual generation the spore germinates giving rise to a tube which develops into a new plant, and is called protonema. There are two divisions, the common mosses, and the liverworts. The parts of the fruiting moss plant are as follows. The *calyptra* or membranous cap which covers the capsule and soon falls off, exposing the *operculum* which is a kind of lid, that is also thrown off. The *peristome* is developed within the operculum and contains teeth, between which the spores are discharged. The elaters in *Marchantia* are for the dissemination of the spores. The peristome differs in different genera, this affords a convenient means of classification. The spores are found in the capsule and running through the center is a slightly differentiated tissue, the columella. In *Funaria* the reproductive organs occur on different plants. The sexual organs are borne much like those of liverworts at the apex of the stem. The antheridia occur in a small rosette of leaves and are club-shaped, the upper part consisting of a single layer of large chlorophyll bearing cells in which small cubical masses occur, the biciliated sperm cells. The archegonia occur in young plants and closely resemble the archegonia of liverworts, except that they have a larger neck. The spores germinate by producing a protonema which early produces a rhizoid.

The liverworts and mosses are much more highly differentiated than any of the Thallophytes, being characterized by more or less differentiation into tissues. Their life history presents a well marked alteration of generations. The gametophyte is more conspicuous than the sporophyte the germinating spore produces the protonema, which consists of a branched filament, the cells containing the chloroplastids. The protonema is usually short-lived in the *Hepaticae* but in the true mosses is longer lived and may persist from year to year. The moss plant is attached to the soil by small unicellular root hairs, or by many curled filaments which, in mosses, are called rhizoids. The shoots of mosses bear lateral organs known as leaves. In *Polytrichum* and *Mnium* the leaf consists, essentially, of a single layer of cells except on the midrib. In the leafy-stemmed liverworts like *Frullania* two rows of lateral leaves occur. In *Marchantia* the leaves are rudimentary and occur on the under surface of the thaloid structure in the form of small scales. The small dots on the surface represent the stomata which are dome shaped structures consisting of a number of cells on each side. The stomata communicate with the photosynthetic system of the plant.

Bryophytes are divided into two classes, the liverworts — *Hepaticae* — and the Mosses — *Musci*, the latter represented by spagnum moss — *Polytrichum* *Bryum*, etc.

The mosses are distinguished from the thallophytes by their sexual reproduction, the antheridia or male organs are stalked, ellipsoidal or club-shaped, and enclose small cubical cells in which the ciliated sperm cells occur. These are ejected, float about in the water till the female reproductive organ the archegonium, is reached. This is a flask-shaped body containing a neck and an egg cell. At maturity the upper part of the canal cells become mucilaginous, the sperm cells pass down through the canal to the egg cell where fertilization




Fig. 16a. A Common Polypody (*Polypodium vulgare*) rh Rhizome. s Stipe. f Frond. r Rachis. a Part of frond with sori. c Sporangium. d Spores. B Moss *Mnium hornum*. a Inner peristome. b Outer peristome. c Capsule. d Spore. C Liverwort *Polytrichum commune*. rh Rhizoid. a Seta on stalk. c Capsule or sporangium. d Spore. D Common liverwort (*Marchantia polymorpha*). a Spores. b Enter. c Same as D. thallid body with female fruiting-body and cupules. F Same as D. magnified. r Rays with the spore cases containing the spores and elaters. p. Perigynium to the left and right archegonia, different stages. G Prothallium of fern with archegonia. h and anteridia. a. H Fern prothallium with young fern and root at r. I Prothallium of Moss (*Festuca hygrometrica*). r Rhizoids. b Buds. a Spores.

is brought about. After fertilization, the egg cell divides and gives rise to an embryo. The mosses differ from the ferns and their allies in a less differentiation of tissues and a slight development of shoot and root system. The vascular system and leafy shoots and roots are marked in the ferns.

PTERIDOPHYTA

Spores like or unlike microspores and megaspores developing into flat or irregular prothallia; these bear the reproductive organs (antheridia and archegonia), flowers and seeds absent, usually a well developed vascular system.

This sub division includes the class *Filicales* or ferns proper. The class *Equisetales* or horsetails, the *Lycopodiales* represented by the common club moss, (*Lycopodium*)




Fig. 1 Ab. Fern. A. Section through frond of *Aspidium Filix mas*, a leaf like body with vascular bundles and sporangia. B. Spore of *Aspidium Filix mas*. C. Spore of *Aspidium Filix mas*. D. Spore of *Aspidium Filix mas*. E. Spore of *Aspidium Filix mas*. F. Spore of *Aspidium Filix mas*. G. Spore of *Aspidium Filix mas*. H. Spore of *Aspidium Filix mas*. I. Spore of *Aspidium Filix mas*. After Luersten.

FILICALES

Leafy plants, fronds usually raised on a stipe, coming from a rootstock, leaves usually rolled up in the bud, circinate, spores all of one kind and size, produced in sporangia which occur on the back of the frond, these at maturity break open and discharge the minute spores, which develop prothalli that bear the antheridia and archegonia. The following sub-orders occur in the United States: the Ophioglossaceae represented by the common adder's tongue, *Ophioglossum vulgatum*, found in moist meadows, the Moonwort, *Botrychium Lunaria* and *B. Virginicum*, the Osmundaceae, large ferns with straight erect rootstocks, pinnate leaves, large globose sporangia with mere traces of a ring, the Royal fern, *Osmunda regalis*, Clayton's fern (*O. Claytonia*), the most common species in damp woods, and the Cinnamon Fern (*O. cinnamomea*), occurring in wet places, marshes, etc., the Filmy ferns Hymenophyllaceae, represented by the Bristle fern (*Trichomanes radicans*) Cyathaceae with such tree ferns as *Dicksonia*, Polypodiaceae, Common Brake, Maidenhair fern:



Fig. 117. Marsilea (*M. quadrifolia*), a young leaf, 2. Fruiting body, Bischoff.

Fig. 117a. Club moss (*Lycopodium obscurum*) 1. Plant with fertile shoots. 2. Scale and the sporangium. 3. Spores. After Wossnill.



Fig. 118a. Antheridium of fern with sperm cells. After Luerssen.




Fig. 119. *Pteris serrulata*. A single archegonium, canal, and neck cells mucilage protruding a 330. After Stranburger.




Fig. 120. Prothallus of Fern. Archegonia above, antheridia below among the hairs. After Luerssen.

Gleicheniaceae, some tropical ferns of few species; the Schizaceae represented by the Small Curly Grass (*Schizaea pusilla*), Climbing Fern, (*Lygodium palmatum*), sporangia ovoid or sessile provided with an apical ring, a family containing about 100 species; the Polypodiaceae the largest sub-order with 200 genera and 3000 species, Marsiliaceae containing *Marsilia* a common aquatic or semi-aquatic plant, represented by *Salvinia*, *Azolla*, also aquatic. The order *Marattiales*, contains the *Marattia*, tropical.

It is not at all strange that the ferns should be poisonous since Greshoff and others have reported the presence of hydrocyanic acid in these plants. Greshoff says the odor of oil of bitter almonds is especially intense in the young leaves of *Cystopteris fragilis* Bernh. and there is also a trace of HCN in the spores. He also calls attention to the presence of the same substance in the common brake (*Pteris aquilina*), and states that several tropical ferns namely *Davallia brasiliensis*, and other species are cyanogenetic, and that one fern, the *D. pentaphylla*, forms a large amount of this substance, especially the cultivated form *elegantissima*. Several species of the *Gleichenia* contain saponin.

POLYPODIACEAE

Perennial with horizontal erect, short or elongated rootstocks, leaves various, entire, pinnate, pinnatifid, or decomposed, vernation circinate (coiled), sori on the margins of the leaf or on the lower side, generally without an indusium (covering); sporangia with a vertical many celled incomplete ring, which on straightening out ruptures and discharges the spores. A few of the ferns, as Male Shield fern *Aspidium Filix-mas*, and the *A. marginale*, are used in medicine.

KEY FOR THE GENERA

Indusium absent	Polypodium
Indusium present, evident	
Sori marginal.	
Indusium with margin of frond rolled over	
Sporangia borne on a continuous marginal vein-like receptacle	
Stipe light colored	Pteris
Sporangia on the ends of the veins. Stipe black	Adiantum
Sori on back with special indusium covering the same.	
Sori linear or oblong.	Asplenium
Sori roundish on the back or rarely the apex of the vein.	
Stipe not articulated.	
Indusium flat or slightly convex or round reniform, fixed by the center, opening all round the margin	Aspidium
Indusium convex, fixed by a broad base, commonly reflexed as the sporangia ripen	Cystopteris
Indusium obscure, leaves closely rolled together with necklace-like segments	Onoclea

Polypodium. L. Polypody

Simple or pinnate fronds from horizontal rootstocks; stipes articulated to the rootstocks, sori (fruit dots) round, naked on the back of the frond in one or more rows each side of the midrib or scattered; indusium wanting. About 350 species, mostly tropical. The species in the Northern United States

are *P. vulgare*, and *P. incarnum*, the *P. vulgare* being more common northward. The *P. aureum* found in Florida is a large fern.

Polypodium vulgare L. Common Polypody

Creeping rootstocks covered with cinnamon-colored scales, stipes light colored, fronds 4-10 inches high, simple and deeply pinnatifid the divisions linear oblong, obtuse or somewhat acute obscurely toothed, sori large.

Distribution. Throughout North America, also Europe and Asia.

Poisonous properties. Used in catarrh and asthma. Supposed by some writers to be poisonous.

Adiantum L.

Sori marginal, borne on the under side of a transversely oblong, crescent-shaped or roundish, margin of the frond, the sporangia attached to the tips of the forking branched veins, stipe black, polished, leaves divided. About 80 species of wide distribution. The *A. Capillus-Veneris* in tropical and sub-tropical regions.

Adiantum pedatum. L. Maidenhair Fern

Root-stock slender, chaffy, stipe black, shining, dichotomously forked at the summit; pinnae arising from the upper axils of two branches of the stipe; pinnules short-stalked, numerous.



Fig. 121. Cultivated. Maidenhair fern (*Adiantum*)

Distribution. In moist woods from Nova Scotia to British Columbia and Alaska, California, Utah, Arkansas and Georgia, also found in Asia.

Medicinal properties. The Maidenhair Fern has a bitterish aromatic taste and was formerly much used as a demulcent, it is probably poisonous. The European *A. Capillus-Veneris* was used in catarrhal affections.

Pteris

Fronds once to twice pinnate, coming from a stout root stock, usually large plants. Sporangia in a continuous slender line occupying the entire margin of the fern frond and covered by the narrow edge which forms a continuous membranaceous indusium. *Pteris* and *Pteridium* are usually separated, about 100 species in the genus *Pteris*.

Pteris aquilina, L. Common Brake

Frond dull green, from 2-3 feet high, ternate at the summit of an erect stout stalk, variable in height from 1-6 feet, stipe coming from a black root-stock; the spreading branches twice pinnate, branches oblong-lanceolate.




Fig. 121a. Maiden fern (*Adiantum pedatum*). (Am. Agriculturist).

Fig. 121b. Brake (*Pteris aquilina*). The roots and stems of this plant are poisonous. The root-stocks contain a starch which is sometimes used as food. (Ada Hayden).

Distribution. Widely distributed in North America, from the Atlantic to the Pacific coast, also found in Europe. Probably the most common fern in the north, especially on the Pacific coast.

Poisonous properties. This plant has been reported as being an anthelmintic and also an astringent, it is suspected of being poisonous.

Asplenium, L. Spleenwort

Large or small ferns with entire, lobed, pinnate leaves, 2-3 times pinnate or pinnatifid, veins free, sori oblong or linear-oblique, straight or rarely curved, indusium straight or curved. About 30 species of wide distribution, several species cosmopolitan like 4 *Trichomanes*.

Asplenium Filix femina, (L.) Bernh. Common Spleenwort

Fronds 1-3 feet high, ovate-oblong or broadly lanceolate, twice pinnate; pinnules confluent on the secondary rachis oblong and doubly serrate or pinnately incised; sori short.

Distribution. Common in the north and especially in the dense woods, as far south as Missouri.



Fig. 122. *Asplenium Filix femina*. Spore bearing leaf 1/6 natural size. a. A single segment showing the under side $\times 10$. After Luermen.

Poisonous properties. The rhizome of the root is used in medicine although it is not official. It is supposed to possess properties the same as the Male Shield Fern.

Aspidium. Swartz. Wood Fern

Fronds with 1-3 pinnate leaves, free veins, sori borne on the back or rarely at the apex of the veins, indusium covering the sporangia, flat or flattish, scarious, orbicular and peltate at the center, or round kidney-shaped, opening all round the margin. About 200 species of wide distribution, common in the northern states.

Aspidium fragrans. (L.) Swartz

Fronds 4-12 inches high, glandular and aromatic, fragrant, rootstock stout, nearly erect, densely chaffy, as are the crowded stipes and rachis. Species found in Asia and Europe.

Aspidium Filix-mas. (L.) Swartz

Fronds large, 1-3 feet high; pinnæ linear-lanceolate, tapering from base to the apex, pinnules very obtuse, serrate at the apex and obscurely so at the sides, the basal uncisely lobed, sporangia nearer the midvein than the margin and usually confined to the lower half of each fertile pinnule.

Distribution. Native to Europe and found in rocky woods from Labrador to Alaska, northern Michigan, British Columbia, Greenland, Europe, Asia, and the Andes of South America.

Poisonous properties. It is suspected of being poisonous. The rootstocks have long been used in medicine as a vermifuge and contain the following substances—a fatty, green oil, traces of a volatile oil, resin, tannin, folic acid, $C_{80}H_{48}O_{12}$, flicin, aspidin, $C_{23}H_{31}O_7$, a fixed poisonous oil.

Dr. Winslow says

Large quantities of the drug cause hemorrhagic gastro-enteritis, tremors, weakness, stupor, coma, acute nephritis and cystitis. Six drachms of the oleoresin have proved fatal in man and sheep, five drachms in a medium-sized dog, and three ounces in a cow. *Aspidium* should never be given with oil which aids its absorption.

Aspidium marginale. (L.) Swartz

Much like the preceding, with evergreen fronds, small, thickish, ovate or oblong in outline, and from 1-3 feet high, pinnæ lanceolate acuminate, pinnules oblong or oblong-scythe-shaped, obtuse or pointed, entire or crenately-toothed, sori close to the margin.

Distribution. From Canada to Minnesota, Iowa, Missouri, Arkansas, and Georgia.

Poisonous properties. Probably has the same effect as the preceding species. Dr. Johnson says

Oleo-resin of male-fern is one of the best known remedies for tapeworm, and also one of the most efficient. Doubtless much of the disappointment experienced with it is attributable to inefficient preparations. Since, however, it has been demonstrated that *A. marginale* is quite as efficient and as this species is very abundant, there is now no good reason why reliable preparations should not be the rule rather than, as heretofore, the exception.

Cystopteris. Bernhardt. Bladder Fern

Fronds growing in tufts, 2-3 times pinnate, the lobes cut toothed, stipe slender, 2-4 pinnate leaves, sori roundish, borne on the back of the veins.



Fig. 123. Shield Fern (*Aspidium marginata*)
Said to contain flicic acid. Known to be poisonous.
(Ada Hayden).

Indusium delicate arched or level-like, attached by a broad base on the inner, partly under the sorus, opening free at the other side veins free. A small genus of 5 species, 2 common in the United States: *C. bulbifera*, long slender fronds bearing bulbils which propagate the plant *C. fragilis* with brittle stalk, the pinnae and pinnules ovate, lanceolate, irregularly pinnatifid or cut-toothed.

Onoclea. L. Sensitive Fern

Coarse ferns, creeping root stocks, fertile fronds erect, rigid with contracted pod-like or berry-like divisions, rolled up, sori roundish, imperfectly covered by a very delicate hood-shaped indusium attached to the base of the receptacle, when dry opening, allowing the spores to escape, sterile fronds foliaceous. A small genus of a few species.

Onoclea sensibilis. L. Sensitive Fern

Slender root stock with scattered fronds, sterile long stalked 2-15 inches high, triangular ovate, fertile fronds, contracted closely, bipinnate, pinnules rolled up into berry-like bodies.



Fig. 124. Flowering Fern (*Osmunda Claytoniana*). Reported as poisonous. (Ada Hayden)

Distribution. Moist meadows and thickets from Newfoundland to Florida and Minnesota.

Poisonous properties. Very abundant in hay from low meadows. May be injurious.

Onoclea Struthiopteris. (L.) Hoffman. Ostrich Fern

Fronds growing in a crown, root stocks stoloniferous, sterile, short stalked, 2-10 feet high, broadly lanceolate, pinnæ pinnatifid, veins free, the venlets simple; fertile frond shorter, pinnate with pod like or somewhat necklace-shaped pinnæ.

Distribution. Nova Scotia to Illinois, Wisconsin, Iowa and British Columbia, common also in Europe and Asia.

OSMUNDACEÆ

Large ferns, root stocks frequently stout and erect, leaves 1-2 pinnate, coiled in veneration, veins free, mostly forked, running to the margins of the pinnules or lobes, sporangia large, globose with mere traces of a ring, or none, borne on contracted pinnæ, on the lower surface of the pinnules. Three genera.

Osunda. L. Flowering Fern

Tall ferns growing in swamps or low ground. Fertile fronds much contracted and bearing on the margins the short pedicelled, naked sporangia on the margins of their radii like divisions without chlorophyll, sporangia thin, reticulated, opening by a longitudinal cleft into two halves, with a few thickened cells the rudiment of the ring.

There are six species in the North temperate regions, three species common in the North, the Cinnamon Fern (*O. cinnamomea*), clothed with rusty wool, the Royal Fern (*O. regalis*), a smooth pale green fern, 2-5 feet high, with 13-25 sterile pinnules, Clayton's Fern (*O. Claytoniana*), clothed with loose wool, but soon smooth; pinnae oblong lanceolate; some of the middle pinnae fertile. The *O. regalis* is used as a tonic and styptic. By some these ferns are regarded as injurious to stock.

EQUISETALES. Horsetails

Rush-like perennial plants, epidermis impregnated with silica; creeping root-stocks, stem generally hollow jointed, simple or branched, striated or grooved, provided with a double series of cavities and usually a large central one. branches verticillate leaves reduced to a sheath which is divided into teeth corresponding to the principal ridges of the stem stomata in furrows.




Fig. 125. Formation of Archegonia of *Osunda*. A. Early development seen from the surface. B. The same in vertical section. C, D. Farther development. E. Opened and closed neck. F. Neck in oblique section. G. Fertilization of the mature archegonium. a. Archegonium. b. neck of the same. c. central cell. d. egg. e. spermatozooids. f. neck canal cells. B-J $\times 240$. A. greatly magnified. After Luerssen.

sporangia 1-celled clustered underneath the shield-shaped scale of the cone, spores all alike, two thread-like elastic filaments (elaters) are attached to the base of the spore which roll around it when moist and spreading when ripe, prothallus green formed upon damp ground, usually dioecious. One order, *Equisetaceae*, and one genus, consisting of 40 species. Fossil horsetails numerous.

EQUISETACEAE

Equisetum, L. Horsetail

Perennia, jointed plants with creeping root-stocks, dull and blackish in color, often bearing tubers, roots in whorls from the nodes, stems usually erect, simple or branched, jointed cylindrical, the surface striated, the stomata occur in grooves either in rows or in bands, the nodes bearing a whorl of reduced leaves joined by their edges into cylindrical sheaths, the tips consist of persistent or deciduous teeth, branches when present in the form of whorls from the nodes, fruit consisting of a terminal cone containing the sporangia in which occur the greenish spores each spore provided with four hygroscopic bands, the elaters, spores produce two kind of prothalli, one male the other female, the male containing the antheridia, the female the archegonia. A small genus commonly called rushes or horsetails. Some ten species in eastern North America.

Equisetum arvense, L. Common Horsetail

Perennia, with annual stems, stomata scattered, fertile stems unbranched, destitute of chlorophyll, 4-10 inches high, soon perishing, sheaths distant, 8-12 toothed, the sterile slender 1-2 feet high, 10-14 furrowed producing simple or sparingly branched 4-angular teeth, herbaceous, triangular lanceolate.

Distribution. Abundant in sandy fields along roadsides and railroads, especially northward from Newfoundland to Virginia, California and Alaska. Also occurs in Europe and Asia.

Equisetum hyemale, L. Scouring Rush

Stems all alike slender rather stiff, evergreen, from 1½-4 feet high 8-34 grooved. Stem rarely producing branches which are usually short and sometimes fertile stomata arranged in rows, rough ridges with 2 indistinct lines of tubercles, the central cavity large, sheath rather long, cylindrical marked with a black girdle, their ridge obscurely carinate spikes persistent.

Equisetum hyemale L. var *robustum*, (A. Br.) A. A. Eaton

Stem perennial, tall and stout, 8-10 feet high, sometimes an inch thick, occasionally branched 20-48-grooved, the ridges roughened with lines of transversely-oblong tubercles sheaths rather short with a thick girdle at the base and a black limb ridges of sheaths carinate.

Distribution. In wet places, from Ohio, Iowa, to Louisiana, Mexico, California, and British Columbia, also in Asia.

Poisonous properties. The rushes have long been recognized in Europe as being injurious to horses, and there are records of their poisonous properties in American Agricultural Literature.

A writer in the American Agriculturist, many years ago, described accurately a disease which might be called equisetosis, and which was produced by poisoning from these rushes.

Mr H Lawrence of Spencer, Iowa, recently sent me a specimen of the last species mentioned above, writing that

"The cattle staggered and had the spours. One man lost 10 head of young animals."

Friedberger and Frohner state that the symptoms of poisoning by this rush are as follows

At first, excitement and anxiety the sensorium remaining unaffected later, uncertainty of movement, reeling and staggering at least, paralysis of hinder limbs, tumbling down, general paralysis, insensibility to external irritants, unconsciousness and coma. Pulse accelerated, appetite at first normal, but in course of time great disturbance of nutrition sugar in the urine. Course sometimes very acute death occurring in a few hours, but sometimes protracted (two to eight days), and at times chronic one to several weeks). In cattle, after excessive eating continuous diarrhoea becomes a prominent characteristic along with the paralytic symptoms, while, if the food be persisted with, cachexia and hydraemia, combined with weakness hurdering on paralysis, make the appearance. Autopsy reveals: hyperaemia, oedema, dropical effusions in the brain and spinal cord, especially on cerebellum, in cases of longer duration, hydraemia. Sometimes inflammatory changes in the mucous membrane of stomach and bowels. Therapeutic change of fodder purgatives and stimulants, especially camphor blisters along the spine.

Mr P J. O'Carra says of this plant

It has been found growing along roadsides and railroad tracks, but its occurrence in serious amounts is apparently confined to low moist meadows which are more or less sandy. As before indicated, it is confined mostly to the Missouri bottom. A thorough examination of several meadows in this region has shown that this plant often constitutes one-sixth or more of the bulk of the hay. There is no doubt that this plant causes a great deal of trouble, but to what extent is not known as many of the haystacks which were carefully examined contained the Rattle-box in considerable amount. Horses eating this hay suffered the combined effects of both poisonous plants.

Prof. Jones and Dr. Rich state

The first evidence of the trouble is more or less unthriftiness, the horse appearing thin and the muscles wasted. In from two to five weeks, according to the age of the horse and the manner of feeding, the animal begins to lose control of its muscles, sways and staggers like a drunken man, although its eye looks bright it ears well, and may even try to caper and play. After muscular symptoms become pronounced many cases refuse to lie down, standing until thrown down by disorderly muscular contractions. If it continues to eat the plant the horse in any case soon loses power to stand and goes down after which it becomes very nervous and struggles violently to get up but legs become more or less rigid, and at times all the muscles of the body seem convulsed. Even in this condition one well nursed patient lived two weeks. The horses are generally willing to eat although unable to rise, but become poor and tired from struggling finally dying from exhaustion. Life is much prolonged by turning from side to side three or four times in twenty-four hours thus preventing gravitational congestion of the lungs and kidneys. The pulse becomes slow until toward the end when it is rapid and weak. Temperature is below normal until the animal goes down, after which a fever develops in consequence of the nervous excitement and violent struggling. The extremities are usually cold, and in the winter horses suffering from *Equisetum* poisoning suffer severely from the cold, presumably because of diminished oxidation and consequent low body temperature. The visible lining membranes of mouth, nose, eye, etc., become pale.

Conditions Influencing the Effect of the Poison

Age of the horses. Young animals develop symptoms much more quickly and succumb to the *Equisetum* poisoning sooner than older ones. In one case under observation a mare eating *Equisetum* hay did not show symptoms until after four weeks, while her colt by her side developed typical symptoms of horsetail poisoning and died in ten days. None out of fourteen horses on one farm, all fed alike upon good hay, were bedded with waste hay containing large quantities of this weed. They ate this bedding freely and within three weeks all nine showed symptoms of poisoning, the remaining five bedded with straw kept perfectly well. The youngest a three-year-old, was down and died a few days later. The oldest an old brood mare, showed but slight symptoms, while the other seven of intermediate ages, all staggered and reeled, although they recovered.

Feed. Grain-fed horses resist the action of the poison much longer than those not grained. Horses seem to develop a depraved appetite for the weed. In the last mentioned



Fig. 129a. a. Scouring Rush (*Equisetum hyemale* var. *robustum*), b. Horsetail (*E. arvense*), fertile branch c. Sterile branch of *E. arvense*; d. *E. hyemale*. Said to be poisonous to horses. (C. M. King).

case, though all were fed good, clean timothy hay, they seemed to prefer the horsetail bedding, and even left their grain to eat it.

Condition of the plant.—We have no evidence that horses grazing upon the green plant are poisoned thereby. It may be that the plant is less poisonous in the early stages of its growth than when mature, or the laxative effect of the grass eaten with it may prevent the cumulative action of the poison. Moreover the plant rarely occurs in as large quantities in pastures as in meadows and apparently rarely need cause apprehension. If, however, it is abundant, close watch should be kept upon horses pasturing where it occurs that the animals may be removed at the earliest symptoms of trouble.

Treatment. In the way of treatment, the first and most important thing is to stop immediately the feeding of the *Equisetum* hay. Our practice further than this has been to give a purgative pill consisting of one ounce of Barbadoes aloes, one or two drachms of ginger, and sufficient English crown soap—soft soap—to make a ball or pill. This is put down the horse's throat at one dose, and following this we have usually given bran mashes night and morning until the digestive tract is entirely cleared of the poisonous plant. In case aloes cannot be easily obtained a quart of raw linseed oil will be very well. After the physic has operated a teaspoonful of powdered nux vomica is added to each grain feed, three times a day. This tends to relieve the muscular incoordination. When poison symptoms are severe and especially when staggering is very profound, slings should be used to support the animal for when once down it is very difficult to make it stand again even with the aid of slings. If, however, the above treatment is begun before the horse loses the power to stand and it can be kept on its feet, its life can be saved in practically all cases.

Steiner & Schröter in their work on the weeds of meadows mention this same plant and several other species as being injurious to stock, not only to horses, about which there is a difference of opinions, but to cattle. In the latter it produces diarrhoea. Cows become poor and the milk flow ceases or is checked.

That this disease is similar to one produced by mouldy corn is shown by the following quotation from Dr Peters:

It is also known that certain weeds commonly called horsetail have a faculty of producing a disease almost identical with this one. The experiments conducted by Dr Rich of the Vermont Station show that that weed is capable of producing similar symptoms.

EMBRYOPHYTA, (SIPHONOGAMA, OR SPERMATOPHYTA)

Plants producing seeds which contain an embryo with 1 or more cotyledons, a stem caudicle, a radicle, and a plumule these parts, occasionally not differentiated before germination, microspores, equivalent to pollen grains borne in microsporangia, ovules (macrosporangia) borne on a modified leaf called the carpel, containing 1 macrospore, equivalent to the embryo sac which develops the minute female prothallium, an archegonium, the egg cell in the embryo sac is fertilized by means of a sperm cell in the pollen tube the male prothallium generally but slightly developed. The Spermatophyta contain two main divisions based upon the character of the ovules.

GYMNOSPERMS

Ovules naked not enclosed in an ovary, attached to scales or wanting, pollen grains develop into the pollen tube, the male prothallium contains the sperm cell and fertilizes the egg cell in the ovule. The Gymnosperms are divided into six classes.

1 Cycadales. These include the *Cycas circinalis* well known in cultivation, an important plant of the tropics. The *C. media* of Australia produces rickets, a *Macrozamia* causes the same disease. Dr Stafford states that *C*



Fig. 126. (*Ephedra nevadensis*). A shrub in Southwestern United States.

circinnalis, known as the "Fadong" in the island of Guam, is poisonous, but the poisonous properties of the seeds are removed by soaking and repeatedly changing the water. He says also that the seeds when fresh are so poisonous that the water in which they are steeped is fatal to chickens. The group also includes the *Dioon* the seeds of which furnish a starch which is an article of food. The species of *Zamia*, a member of this group, are native to tropical Florida.

2. Bennettiales, A fossil group.

3. Cordaites, A fossil group.

4. Ginkgoales. These include the Ginkgoaceae, of which the *Ginkgo biloba* is well known and is frequently cultivated as an ornamental plant in the United States. Long avenues of these trees are planted in Washington. The fruit of the Ginkgo has a very disagreeable odor. The tree was common in the tertiary age.

5. The Coniferae.

6. Gnetales. This group is represented in the United States by *Ephedra*, shrubs with horse-tail like branches, small leaves and buckwheat-like seeds.

The *Welwitschia* of the above group is found upon stony ground in the tropical Old World.




Fig. 127. Hemlock (*Tsuga canadensis*). A common forest tree of Northern North America. Contains resin and the usual principles found in these resins. Said to be injurious.

CONIFERAE

Resinous trees or shrubs generally evergreen leaves, entire or scale-like, wood consists mostly of tracheids marked with large depressed disks tracheae only present near the pith and in the leaves, perianth none; flowers monoecious, stamens several together, subtended by a scale, anthers 2-7 celled, pollen grains frequently of three cells, one fertile and two inflated, ovules with two coats, borne solitary or together on the surface of a scale, straight or partly inverted fruit a cone, usually papery, but in some instances fleshy, sometimes berry-like seeds winged or wingless, endosperm abundant, fleshy or starchy, embryo straight and slender, cotyledons 2 or more. About 25 genera and between 275 and 300 species. They include the Podocarpus of the tropical regions the Taxus or Yew, the Norfolk Pine (*Araucaria excelsa* and *A. brasiliensis*) frequently cultivated, the White Cedar (*Cupressus Latsoniana*) of California, the *Cupressus nothofalcensis* of the northwest coast, the White Cedar (*C. thyoides*) which occurs in swamps, in the East, the genus *Picea* consisting of the spruces, Norway Spruce (*P. Engelmannii*), Tideland Spruce (*P. sitchensis*) one of the largest trees in Oregon and Washington; the Douglas Fir (*Pseudotsuga Douglasii*), one of the most valuable of the North American conifers, found in Washington, Oregon and California and in the Rocky Mountains, the Hemlock (*Tsuga canadensis*), abundant in North America, and the source of Canada or Hemlock Pitch, the bark containing an abundance of tannin, the leaves are said to be abortive, the *Tsuga heterophylla* of the Pacific coast which produces a valuable lumber, the Balsam Fir (*Abies balsamea*), which furnishes a kind of balsam that contains four acid resins and a volatile oil; the Black Fir (*Abies concolor*),

a large forest tree of the Pacific coast and the Rocky Mountains, the Sandarac tree (*Calistrus quadrivalvis*), which furnishes not only the sandarac gum used in making varnish, but also a dark-colored, fragrant wood capable of high polish and used in ornamental work, the Arbor Vitæ, or White Cedar (*Thuja occidentalis*), which contains *jenshor*, *thujone*, *thujin*, and the bitter glucoside *pinicrin*, its leaves being irritating to the skin, sometimes producing blisters, the Norway spruce (*Picea excelsa*), which contains resins and volatile oils and is the source of Burgundy pitch; the Black Spruce (*P. mariana*), from the young branches of which an essence is prepared that is used in the preparation of spruce beer, the White Spruce (*P. canadensis*), which, with the preceding species furnishes much of the wood pulp used in the manufacture of paper, and the Pines

Taxus. (Tourn.) L. Yew

Flowers generally dioecious or occasionally monoecious, axillary from scaly buds, sessile or nearly sessile, from small staminate catkins of a few scaly bracts 5-8 stamens, anthers 4-celled, fertile flowers solitary, erect subtended by a fleshy cup-shaped disk, fruit consisting of a fleshy disk which becomes cup-shaped and red and encloses the bony seed.

Distribution. About 6 species native of the north temperate regions. One upon the Pacific coast, *Taxus brevifolia*, is a tree. The European Yew (*Taxus baccata*), a well known poisonous plant, is frequently used for ornamental purposes in this country.

Taxus canadensis. Willd. American Yew

A low shrub, straggling over bushes, with linear leaves, green on both sides.

Distribution. In the woods from Newfoundland to New Jersey and Virginia, west to Wisconsin, Iowa and Minnesota.

Poisonous properties. One species is known to contain the alkaloid taxin, $C_{21}H_{33}NO_{10}$. Dr. Johnson says, concerning the poisonous properties of the American Yew

This plant, a variety, only, of the European yew, cannot be said to have, as yet, a place among medicines. It is believed, however, to possess poisonous properties, and is perhaps worthy of investigation. Regarding the poisonous properties of the berries, the author can state that he has eaten them without deleterious effect, but whether because the quantity was insufficient or not, is an open question. Cases of fatal poisoning from eating the berries of the European yew are on record, and therefore our variety is certainly open to suspicion.

Crestat refers to the poisonous nature of the yew as follows

The common yew, or ground yew of the northeastern United States is called poison hemlock in some places. The leaves of this shrub are probably poisonous to stock, as are those of the European yew. This species is more accessible to stock than are those of the western yew (*Taxus brevifolia*), which grows only in deep canyons.

Dr. Otto Lehmann* in his treatise on poisonous plants states that older naturalists regarded the yew as one of the most powerful of poisonous plants. Modern testimony is conflicting, but he regards the branches and leaves as poisonous for animals. Friedberger and Fröhner give the symptoms of poisoning from yew as follows: "Death may be sudden, resembling apoplexy, it may be preceded by staggering and convulsions, cases of long standing show gastro-enteritis. Give purgatives as remedies."

* Giftpflanzen. 121. Hamburg. 1882.

Pinus. (TOWN.) L.

Evergreen trees with short scale like leaves and longer leaves in bundles, the ordinary foliage leaves linear in bundles varying from 25, rarely 1, stamens in catkins, borne at bases of shoots, filaments short, anthers longitudinally dehiscent, pistillate bearing aments, solitary or clustered on the twigs of the preceding season consisting of numerous imbricated bracts, each with an ovule-bearing scale. Fruit a large cone. seeds 2 at the base of each scale, winged above. About 75 species of wide distribution. Of these the more important are White Pine (*P. Strobus*) one of the most valuable of North American Pines. Sugar Pine (*P. Lambertiana*) of California and Oregon. Austrian Pine (*P. Laricina*) furnishing Austrian turpentine. the Long leaved Pine (*P. palustris*) of the South, the most important source of turpentine which




Fig. 129. White Pine (*Pinus Strobus*). Branch bearing strobili. 2. Branch bearing pistillate flowers and young cones. 3. A cone enlarged. 4. Scales of pistillate flower, enlarged. 5. A young branch bearing young cones. 6. Fruiting branch with young cones. 7. Scale of cone with two seeds attached. 8. Seeds with wings attached. 9. Seeds, enlarged. 10. Seedling plant. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. One half natural size. (N. B. Green).

in this tree amounts to 70-80 per cent., resin, 15-30 per cent, volatile oil and some pinene $C_{10}H_{16}$, a very important constituent; *P. sylvestris* the source of Russian turpentine, *P. pinaster*, supplying the French turpentine, *P. heterophylla* and *P. echinata*, also turpentine trees; Loblolly Pine (*P. Taeda*) also containing pinene, the Western Yellow Pine (*P. ponderosa*) occurring from mountains of Colorado westward, a large tree 120 or more feet high, 4 or 5 feet in diameter branching widely spreading or drooping, bark light red, leaves in 3's or rarely 2's, cones stout, dense, heavy, ovoid-conical, each scale with short recurved prickles the variety *scopulorum* of the last named species, found in the front Rockies, a smaller tree with shorter leaves and Lodge Pole Pine (*Pinus contorta*), a tall straight tree, 80-120 feet high, and from 12 inches to 3 feet in diameter, with conical head, thin, light grayish brown bark, leaves 1-3 inches long, light green, rigid, often persistent cones. This last named species occurs from Colorado, Wyoming and South Dakota to the Pacific coast and is closely related to the Jack Pine (*P. Banksiana*)

Phenol and creosote oil are obtained from *Pinus palustris* and *P. Taeda*.

Poisonous properties. Cattle and sheep do not usually graze upon the leaves of conifers, but when forced to do so because of scarcity of fodder, sheep will eat the leaves, which may produce injurious symptoms. According to Friedberger and Frohner, plants containing turpentine are poisonous. Choibert in 1787, observed gastro-enteritis complicated by nephritis as a result of grazing on leaves of conifers. The first named authors find symptoms of haematuria, constipation, evacuation dry, and irritation of the kidneys.

Juniperus, (Toura.) L. Juniper

Flowers dioecious or monoecious, in lateral catkins, staminate catkins small, fertile catkins consisting of 3-6 fleshy scales, fruit appearing like a berry color of fruit bluish-black or blackish, frequently with white bloom seeds 1-3, wingless and bony. The *J. Oxycedrus* of the Mediterranean region produces "oil of cedar."

Juniperus communis. L. Common Juniper

A shrub or small tree with spreading or pendulous branches, leaves rigid, spreading berries dark blue. The variety *alpina* Gand is a low decumbent, or prostrate shrub with shorter, less-spreading leaves. It contains pinene and cadine. The oil and fruits are used in the manufacture of gin.

Distribution. From Nova Scotia to British Columbia, to Pennsylvania, Wisconsin, Michigan, Nebraska, and in the Rocky Mountains to New Mexico occurs also in Europe and Asia. The variety *alpina* is common in the mountains of New Mexico northward, not, however, as widely distributed as the species.

Juniperus virginiana. L. Red Cedar

A shrub or tree extremely valuable, frequently from 60-90 feet high pyramidal in form leaves scale-like, obtuse or acutish, dimorphic, the leaves of young plants being more or less flattened, spiny and awl-shaped, while those of the stem are scale-like and appressed, catkins terminal, berries on straight peduncles cones light blue or glaucous. The Platte Cedar (*J. scopulorum*) differs from the other in the development of the seeds.

Distribution. The Red Cedar occurs from New Brunswick to British

Columbia, south to Florida, Texas, New Mexico and Arizona, also the West Indies. The Platte Cedar occurs from Nebraska westward and is common in the foothills of the Rockies.

Poisonous properties. According to Dr Halsted it poisons goats which browse on it.



Fig. 129. Red Cedar (*Juniperus virginiana*). To the left a branch from an old tree to the right juvenile shoots, spiny. The plant is poisonous and injurious.

Juniperus occidentalis. Hooker

A shrub or small tree, with bark in shreds, leaves pale in color, closely appressed, obtuse or acutish, berries 4-5 lines in diameter

Distribution. Northwest along Pacific Coast.

The variety *monosperma*, Eng., shows stunted trees, frequently 2 or more feet in diameter, attaining an age of 1200-1300 years, leaves scale-like; berries




Fig. 130. Savin, Juniper (*Juniperus Sabina*). Fruiting branch. Known to be poisonous. (After Strasburger, Koll, Schenck and Schimper).

smaller than the above, frequently copper-colored; generally with one seed or sometimes more.

Distribution. Found from Colorado to Western Texas, Arizona, California, and Wyoming.

Several allied species have been described, among them *J. Knightii* by Nelson, and another by Sudworth, which occurs in the southwest.

Juniperus Sabina, L. Savin, Juniper, Swedish Juniper

A prostrate shrub with appressed leaves in pairs; margin slightly or indistinctly denticulate, berries on short recurved peduncles, 3-4 lines in diameter, 1-3 seeds. It contains the substance *sabinal*. It is official.

Distribution. Along the Atlantic coast, from Massachusetts westward to New York, Minnesota, Montana, and British Columbia, also in Europe and Asia.

Poisonous properties. The wood of Red Cedar is extensively used in the manufacture of lead pencils and was formerly also employed in making cigar boxes. The fruit of the common Low Juniper (*J. communis*) is used for flavoring gin. Red Cedar contains a fragrant volatile oil consisting of *cedrol* and *cedrene*. Cases of poisoning from this genus have been reported.

ARBOREALIA

Ovules enclosed in an ovary.

MONOCOTYLEDONEAE

Embryo with a single cotyledon, first leaves of germinating plantlet alternate, stems endogenous, consisting of an outer part, an inner mass of cells the parenchyma, and the bundles distributed through the mass, no distinction into pith, wood, and bark; leaves generally parallel veined, usually alternate and sheathing at the base, flowers generally on the plan of 3. This group of plants includes the palms, grasses, lilies, duckweeds, etc.

PANDANALES

Marsh plants, herbs or trees with linear leaves, flowers in spikes or heads, perianth of bristles or of chaffy scales, ovary 1-2 celled, endosperm mealy or fleshy. This order includes the Cat-tail (*Typha latifolia*), Screw pine (*Pandanus*) and the Bur-reed (*Sparganium*). The ripe fruit of the *Pandanus fragrans* is used as a relish in the Philippine Islands.

The Cat-tail is reported as poisonous. It is common across the continent and is found in swamps.

HELOBIAE

Aquatic or marsh herbs, leaves various, flowers perfect, monoecious or dioecious, perianth present or absent; stamens 1-numerous, carpels 1 or more, mostly distinct, endosperm none or little. This order includes the Pond Weeds (*Potamogeton*), of which there are many species, which float in the water and often give trouble in ponds of parks, fresh water eel grass (*Vallisneria spiralis*) water weed (*Elodea canadensis*), a troublesome weed in the canals of England and Europe. All of these plants are abundant in our fresh waters and afford food for crustaceans, which in turn are used as food for fish.



Fig. 131. Pond-weed (*Potamogeton natans*). 1. Apex of flowering shoot. 2. Flower viewed from above. 3. Flower viewed from side. 4. Diagram of flower (After Wossido).




Fig. 132. Pond weed (*Potamogeton*). Common in fresh water ponds.




Fig. 133. Bar-seed (*Sparganium*)



Fig. 134. Cat-tail (*Typha latifolia*). A common weed along shores of lakes and streams.

ALISMACEAE D. C. Water-Plantain Family

Aquatic or marsh herbs, generally with smooth, sheathing leaves, flowers perfect, monoecious or dioecious; sepals 3, persistent, petals 3, the larger, deciduous, imbricated in the bud, stamens 6 or more, anthers 2-celled, extrorse,



Fig. 135. Water Plantain (*Alisma Plantago*). A common marsh plant

pistils numerous or few, usually with a single ovule in each cell, fruit an achene, seeds small, erect. About 70 species of wide distribution in swamps. The Water Plantain (*Alisma Plantago*) of Europe and North America is common in the northern states. Several species of Arrowheads (*Sagittaria*) are used as food by the Indians and Chinese.

Sagittaria L. Arrowhead

Perennial with tuber-bearing root stocks and milky juice; basal leaves long-petioled, scape sheathed at the base, flowers monoecious or dioecious, borne near the ground in whorls, sepals persistent in pistillate flowers, reflexed or spreading, petals 3, white, deciduous, stamens indefinite, pistillate flowers with distinct ovaries, ovule solitary, fruit an achene in dense clusters, seed erect, curved.

Sagittaria Engelmanniana, J. G. Sm.

Perennial with stoloniferous roots; leaves very variable; scape 1-4 feet high, angled, lower whorl fertile, pedicel of fertile flowers, at least half the length of the sterile one, filaments smooth; achenes obovate with a long curved or horizontal beak.

Distribution. Across the continent and in Europe.

Poisonous properties. The tuberous stolons are eaten, if there is any poison contained in the raw state it is probably removed by methods of preparation for food.



Fig. 134. Arrowhead (*Sagittaria Regelmanniana*). Common in low grounds. Some species supposed to be poisonous. (After Miss G. E. Johnson, Rep. Mo. Bot. Garden).

GLUMIFLORAE

Endogenous plants mostly herbaceous, stems (culms) narrow or without leaves. Leaves usually narrow and elongated, entire or serrulate. Flowers small, generally perfect, in the axils of dry chaffy scales, called glumes, arranged in spikes or in panicles consisting of spikelets. 2 families, *Gramineae* and *Cyperaceae*.

Fruit a caryopsis	1 Gramineae
Fruit an achene	2 Cyperaceae

I GRAMINEAE. Grass Family

Fibrous-rooted annuals or perennials, rarely woody, generally with hollow stems. Alternate 2-ranked leaves, sheaths split or open on the side opposite the blade, flowers consisting of 2-ranked glumes, forming a 1 many-flowered spikelet, flowering glumes enclosing a small bract called the palea, stamens 1-6, usually 3, anthers versatile, 2-celled, stigma hairy.




Fig. 137 Corn (*Zea mays*). To the left pistillate flowers, g, ovary, s, stigma, to the right staminate flowers, s, stamens.




Fig. 137a. Rice-cut-grass (*Lernia tenuiflora*). The sharp edges of the leaf of this grass often cut the flesh of animals.



Fig. 137b. P. r. pine grass (*Spartina cynosuroides*). The sharp edges of this leaf cut like a knife often wound by animals.

A large order of about 3500 species, many of which are very important to man. Among them are the wheat, oats, rye, corn, wild rice, sorghum and sugar cane, the two latter being the sources of some of the sugar of commerce. Many grasses, also, are important forage plants, among which may be named blue grass, timothy, brome grass, and red top. Some grasses are used in medicine.

The Bamboo, native of the tropics, is valuable, being used not only for building purposes, but also in the manufacture of household furniture and in other ways.

Very few of the grasses have deleterious properties. A few, such as sleepy grass and millet, the latter of which is injurious to horses, are known to be poisonous. Some grasses, because of their stiff awns, penetrate the skin and even perforate the intestines, inflicting dangerous wounds. Needle grass and squirrel tail grass, or wild barley, are known to inflict injuries by lodging between the teeth, thus causing pus infection.




Fig. 137c. Spikelets of various grasses. 1. Common blue grass (*Poa pratensis*). 2. *Setaria*. 3. Spikelet of blue-stem grass. 4. Bristly grass (*Setaria verticillata*).




Fig. 137d. Scent glands of stink grass (*Eragrostis major*).

A great many grasses, because of their sharp edges on the leaf, inflict injuries by cutting the flesh. Of these we may mention the rice-cut-grass, (*Leersia*), and porcupine grass, (*Spartina cynosuroides*).

Holy grass (*Hierachloa odorata*) is sweet scented and contains coumarin. Indians use it to weave in baskets, mats, etc. Job's Tears, *Corn tachryma*, is used for rosaries.

KEY FOR GENERA OF GRAMINEAE

Spikelets jointed upon the rachilla below the glumes, 1-2 flowered.

Rachis bearded, spikelets spicate in pairs. 2 *Andropogon*

Rachis not bearded.

Pedicels bristly bearing 4. *Setaria*

Pedicels not bristly bearing.

Spikelets enclosed by a bur 5 *Cenchrus*

Spikelets plano-convex, not enclosed by a bur 3. *Paspalum*

Spikelets in pistillate flowers, borne on a cob 1. *Zea*

Spikelets not usually jointed above the persistent lower glumes

Spikelets 1-flowered.

Awn simple twisted 7 *Stipa*

Awn 3-pointed 6. *Aristida*

Spikelets more than 1-flowered.

Spikelets 2-several flowered, rachis often bearded, flowering glume with a twisted awn 8. *Avena*

Spikelets 1 or more-flowered with a zigzag jointed rachis, channeled

Spikelets solitary at the notches.

- Flowering glumes with the backs turned to the rachis 10 *Lolium*
 Flowering glumes with their sides turned to the rachis 11 *Agropyron*
 Spikelets 2-6 at each joint of the rachis 12 *Hordeum*
 Rachis not charneled 9 *Bromus*




FIG. 118. Spikelets of tall-meadow catgrass (*Archonatherum eximium*). 1 & 2 Staminate. 3. The lower flower with protruding styles upper flower with protruding stamens. The lower whorls are called sterile glumes. Each flower consists of a pair of flowering glumes, stamens and pistil.

1. *Zea. Mays. L.*

Spikelets unisexual, monoecious, the staminate 2-flowered in pairs, one sessile, the other pedicellate, arranged in terminal branches of a terminal panicle the pistillate 1-flowered, sessile crowded in several rows, along the much thickened continuous axis arising from the lower leaf-axis and closely enveloped by numerous large foliaceous bracts, glumes 4, awnless, those of the staminate spikelet acute, those of the pistillate very broad and obtuse or emarginate grain hard, only partially enclosed by the fruiting glumes. This well-known, tall, and striking annual grass has erect stems and broad leaves



Fig. 139 Vanilla grass (*Hierochloa odorata*). a. Spikelet with nearly equal lower glumes, b, with lower glumes removed, showing third and fourth scabrous glumes. c. pairs with stamens. d, pistil. (Dir. of Agric. U. S. Dept. of Agric.).

the terminal staminate inflorescence forms the "spindle" and the long projecting styles of the pistillate flowers constitute the "silk", the cob is formed by the union of the axes of several female spikes into a much thickened body.

The 1 or 2 species are of American origin, presenting many varieties in cultivation known as corn, Indian corn or maize (*Zea Mays*). Dr Sturtevant has arranged cultivated corn into the following groups:

- Pod-corn, *Zea tunicata*.
- Pop-corn, *Zea euvetia*.
- Flint-corn, *Zea indurata*.
- Dent-corn, *Zea indentata*.
- Soft corn, *Zea amylacea*.
- Sweet corn, *Zea saccharata*.
- Starchy sweet corn, *Zea amyloasaccharata*.

The so-called species and groups of Dr Sturtevant are hardly to be regarded as varieties. Some of the forms under conditions of culture and climate, revert to the original type.

A plant cultivated for so long a time by the Indians and civilized man has naturally given rise to diverse forms which we regard as nothing more than races of the very polymorphic species *Zea Mays*.

Some years ago Dr. Watson obtained from Moro Leon, through Prof. Duges, some corn which he considered a new species, calling it *Zea canina*. He says

The natural supposition was that we had here at least the original wild state of our cultivated maize. A careful comparison of the two, as thorough as the material at hand of the cultivated forms would permit, has led me first to doubt the probability of this, and now to consider the form in question a distinct species. The differences upon which this conclusion is based are in the habit of growth, the arrangement of the staminate spikelets, and the nervation of their glumes, the form of the glumes of the pistillate flowers, and the ready disarticulation of the ripened ear.

Dr. Harshberger, who is certainly a most careful observer, and who carried on some most interesting experiments on hybrids, considers our maize of hybrid origin and *Zea canina* is a hybrid of corn and *Euchlaena*. He says

Maize relates itself botanically to a native Mexican grass termed *Euchlaena mexicana*, and the fertile hybrids of this grass and maize are known producing a plant described by Watson as *Zea canina*. From the peculiar behavior of these hybrids, the writer has suggested that our cultivated maize is of hybrid origin, probably starting as a sport of teosinte, which then crossed itself with the normal ancestor producing our cultivated corn. This is speculative, but there cannot be any doubt that the close relationship of maize and teosinte points the way to the determination of the botanical characters of the original wild corn. Recently, Montgomery has suggested a theory as to the nature of the maize ear in which, in conclusion, he states that corn and teosinte may have had a common origin, and that in the process of evolution the cluster of pistillate spikes in teosinte were developed from the lateral branches of a tassal-like structure, while the corn ear developed from the central spike. It is probable that the progenitor of these plants was a large, much-branched grass, each branch being terminated by a tassal-like structure bearing hermaphrodite flowers.

Corn holds the first place in the list of crops produced in this country and North America produces four times as much as the remainder of the world. According to C. P. Hartley, Europe stands second, South America third, and Africa fourth. As a corn-producing country the United States has no rival. Argentina stands second, Hungary third, and Italy fourth. The average corn yields in four central states for five years, 1902-1906, were as follows:

State.	Bushels.
Illinois	342,115,835
Iowa	301,666,176
Nebraska	239,835,262
Missouri	210,082,426

Maize is one of the most important cereals of North America, being used as a food for man and stock, in the manufacture of starch and glucose, and in medicine the corn silk being used as a mild stimulant and diuretic. The oil from the embryo is a yellow viscid transparent liquid having a peculiar odor of corn meal. The silk contains maizenic acid.

Injurious properties. In many sections of the country where corn is grown and cattle allowed to feed on corn stalks, a disease occurs which has been called the corn stalk disease. This has been attributed to various causes such as corn smut, a bacterial disease, nitrate poisoning, bacterial poisoning and impaction of the stomach. Corn stalks are not easily digested and it is not to be wondered at that impaction should occur when cattle do not have access to plenty of water.

This same disease goes under other names, and it may be that there are several distinct types of diseases due to the feeding of corn stalks. Dr. Bitting describes a *Septicæmia hæmorrhagica* which is caused by an organism, the cocco-bacillus. The symptoms of this disease are as follows:

The symptoms depend upon the point of attack. If the respiratory system be attacked, there will be a rapid rise of temperature, difficult and rapid breathing, standing with the feet wide apart as in pneumonia, short coughing, the tongue protruded, and eyes prominent and congested. The animal will move only when urged to do so. The attack lasts for only a few hours. If the pneumonia be of less severe type the kidneys and bowels may show some affection before death.

If the bowels be the seat of attack, there will be bloating colic, noisy intestinal movement, straining and diarrhea. The bowel movements are soft, fluid and foul smelling, and may be blood stained. The urine will also be blood stained. If the infection take place from a superficial abrasion, the part will swell rapidly, become very large, be hot and painful, does not pit upon pressure, and does not crepitate. The swelling extends rapidly and if in the region of the neck, will cause suffocation. The course is short and generally fatal.

Dr. Bitting states that this disease must be differentiated from the corn stalk disease due to poisoning and that post mortem must be the means of separation in some instances. The only remedy is a change of pasture, as little can be done otherwise.

Recently much interest has been attached to the disease known as Pellagra, which has been treated elsewhere in this volume, but in this case it may be of interest to know that the disease has made its appearance in several of the southern states, notably Alabama and South Carolina, cases have also been reported in Maryland and Massachusetts, and a number of them in the Insane Hospital in Illinois. It is believed by the experts who have investigated this question that it is in some way associated with corn. For instance, Dr. Lavinder who investigated this disease with Assistant Surgeon-general Wyman cites the case of the disease on the Island of Corfu, where an epidemic followed when the people began to use an inferior imported quality of maize. Previous to this they had used their own maize which was carefully selected and prepared. There can be no question according to Lavinder that the introduction of maize collected in Spain, France, and Italy with unsanitary conditions and the use of poor maize greatly influence the spread of this disease in those countries. It is practically unknown in those countries where maize is not a staple article of food.

Dr. Arisberg of the U. S. Department of Agriculture, in a discussion of this disease before the American Society of Tropical Medicine, states that corn is one-fifth of the food of the Tennessee and Georgia mountaineers and one-third of the food of the negroes. Under the present conditions this corn is collected before maturity and often is shipped to distant points in poorly ventilated cars which makes it possible for moulds of different types to develop. Furthermore, the same person is authority for the statement that, in ten generations the fat content of corn has increased from less than 5% to 7 1/3% and that toxins are found to be related in quantity to the oil produced in the seed. Then, too, the weather conditions in this southern corn region have been extremely favorable in the last ten years for producing corn which would be immature and subject to moulds when transported.

Now it is a well known fact that for many years throughout the south they have had trouble with the so-called forage poisoning affecting live stock which had been fed corn, especially when mouldy. It seems to the writer that

there is some relation existing between Pellagra in man and forage poisoning in horses and cattle. Both are essentially produced by some toxic substance.

In this connection, the bulletin on the Grand Traverse or Lake Shore Disease, as investigated by C. D. Smith,* C. E. Marshall and Dr. Ward Giltner, is interesting.

2. *Andropogon*. (Royer) L. Beard Grass

Tall annual, or perennial grasses with spikelets in pairs upon each joint of the slender rachis, usually narrow leaves, terminal and axial racemes, one of them sterile, the other sessile, 1-flowered, and fertile, lower glume the larger, coriaceous and nerved, the second acute, stamens 13, grain free.

About 150 species widely distributed in tropical and temperate regions. Some of the species of the *Andropogon* L. are excellent grasses for forage purposes. Quite a number of them produce valuable oils like *Pamorusa* oil, obtained from *Andropogon Schoenanthus*, lemon grass oil from *Andropogon*



Fig. 140. Johnson-grass (*Andropogon halepensis*)
a, spikelet, c and d, glumes, e, f, g, parts of the flower.

* Sp. Buli, Mich. Agr. Exp. Sta. 58: 18.

citratus, and citronella oil from *A. Nardus*. The fibrous roots of the aromatic Cuscuta grass of India (*A. muricatus*), produce a substance used mainly as sachet powder, the fibres of the plant are used extensively by the natives in making mats.

Andropogon halepensis. Brot. Johnson Grass

A stout perennial with smooth, erect, simple culms, 3-5 feet high, and strong creeping rootstocks, panicle open, 6-12 inches long, the 3-5 flowered racemes clustered toward their extremities, outer glume coriaceous, second glume equaling the first and convex below, the third glume shorter than the outer ones, membranaceous, palea broadly oval, fourth glume ciliate awned; palea shorter than the glumes; nerves ciliate

A troublesome weed throughout the Southern States. First introduced as a forage plant.

Andropogon Sorghum. Brot. Sorghum

An annual with long, broad, flat leaves and ample terminal panicle. spikelets in pairs at the nodes, larger and rounder than in the preceding, rachis




Fig. 141 Sorghum (*Andropogon Sorghum*) 1 Kaffir corn 2 Jerusalem corn, 3, Amaranthogon. (Kansas State Board of Agri.)

not articulate, sessile spikelet with 4 scales, the outer hard and shining, the inner hyaline, the fourth scale on and subtending a small palea and perfect flower, or occasionally the palea wanting.

Sorghum is contained in a number of cultivated plants which are classified by Mr C. R. Ball under (1) Broom Corn, (2) Shailu, (3) Durra, (4) Sorghum and (5) Kaffir. Broom Corn used for the manufacture of brooms, is grown chiefly in the central Mississippi Valley, Kansas, Oklahoma, and the Panhandle of Texas. The Shailu, also known as Egyptian wheat, is culti-

vated extensively in India. The Durra have been cultivated for centuries in Egypt, and other countries of Africa, and in India for human and animal food. They are now cultivated in the United States, chiefly from Kansas to Texas. The Kaffir, native to eastern Africa from Abyssinia to Natal, was introduced into this country in 1875, and is grown chiefly in the semi arid regions for forage. Sorghum is grown largely for fodder. The pulpy juice contains cane sugar in variable amounts, and is used both for sugar and in the making of syrup, although the growing of sorghum for syrup seems to be on the decline. According to the census of 1890 the production was 24,000,000 gallons, in 1900, 17,000,000.

Poisonous properties Sorghum has long been recognized as poisonous. Mr C W Warburton in Bailey's Encyclopedia of Agriculture, says

Sorghum makes excellent pasture for hogs, but in many sections it must be pastured sparingly, if at all by sheep and cattle. After periods of extreme drought, or when growth is checked from other causes, the leaves of the sorghums often contain a large amount of prussic acid. A small quantity of this poison is fatal to stock, and death frequently results even after the sorghum is eaten. Natural growth seldom contains prussic acid in appreciable quantities, and it largely disappears on curing, so that cured sorghum may be fed with little danger. There is also some danger from hiccough cases and sheep should not be turned on sorghum pastures when hungry or when the plants are wet. With the exercise of care, however the crop can usually be pastured with safety. It should be at least two feet high before stock are turned on it, for cattle, sheep and horses it may be much more matured than for hogs.

Frosted cane is said to be especially injurious. Dr R H True, of the U S Dept of Agriculture, in commenting on the poisons from sorghum, says:

This office has from time to time received communications from stockmen, especially in the lower part of California, Arizona, and adjacent territory expressing a suspicion that the eating of the Johnson grass had caused the death of stock with rather sudden and violent symptoms. There has seemed to be little ground in poisonous-plant literature to support such an explanation. Last summer however convincing observations were reported from California by a stockman who had lost heavily, and a survey of the grass in question was obtained. The result of the study of this material was positive.

Mr A C Crawford, who investigated some cases, says

It has been noted that deaths in cattle frequently occur when on account of the failure of rain, the plants which have reached a certain size become stunted and withered. The toxic principle appears simultaneously over a wide area, but soon disappears if a rainfall occurs. The deaths of cattle have been attributed by some to an insect living upon the plant, and in Australia it is the belief that *Sorghum vulgare* which also yields hydrocyanic acid, becomes more poisonous when attacked by an insect during a drought. A similar observation has been made with *Sorghum vulgare* in the Sudan. Balfour found that one specimen of the plant which harbored aphids yielded more hydrocyanic acid than a second one without parasites. Penn has lately claimed that the deaths from Johnson grass in India were really cases of nitrate poisoning, as he found 25 per cent of nitrate of potassium in the stem of the plant and was able to produce somewhat similar symptoms in animals by feeding them this salt. Johnson grass is being introduced into Australia as a fodder plant, but as yet no reports of its poisonous action there have been noted by the writer.

Dr George H Glover of Colorado, also reports large loss of cattle in that state from eating Kaffir corn. Twenty-one head out of thirty-two cows died within an hour after first being placed upon the feed. Dr A. T Peters of Nebraska, investigated the subject of poisoning from sorghum and Kaffir corn. He states that in most cases where death of animals occurred, the animals did not regularly pasture upon sorghum, but broke into the fields from adjacent pastures or as they were being driven past fields of sorghum. The investigations proved that the animals did not die from bloat as had been re-

ported. Dr Peters speaks with authority as he examined the cases at first hand.

In response to one of these calls, I reached the farm of Mr Bert Foss, near Aurora, at 7.30 A. M., on August 3, 1901. Two days previous, fifteen head of his cattle had broken into a sorghum field, where they had remained twenty minutes. They were then driven into another field and were not seen again for several hours. When seen, three were sick, all of which died within a few hours. The symptoms were drowsiness, running at the eyes, twitching of the muscles, numbness of the limbs, staggering gait, inability to stand, involuntary passing of urine. On August 2d, two more cows broke through the fence and were on the sorghum field five minutes. One hour later, one of these animals, a four-year-old cow was very sick, but finally recovered.

We turned a small yearling steer on the sorghum at 8.10 A. M. August 3d. but he refused to eat any sorghum, and after thirty-five minutes, two more were turned into the sorghum, where they remained until 10.00 A. M. when only one, a small, red steer, had taken any sorghum, and he had eaten only a few leaves. They were then turned back with the herd. At 10.35 A. M. the small, red steer acted somewhat drowsy, but soon recovered.

At 11.00 A. M. we turned one red heifer and one yearling steer on the sorghum. The heifer was the only animal that ate any quantity and as subsequent examination showed, she ate only one and one-half pounds of green sorghum. At 11.10 this animal dropped to the ground. Upon examination it was found that she had stopped chewing her cud and there was a peculiar twitching of the muscles of the nose and head and also of the body. The animal was very dull. At 11.15 A. M. she was taken out of the sorghum field and allowed to lie in a stubble field. When lying down her head was turned toward the abdomen, preventing the symptoms shown by a horse having the colic. The eyes seemed dull and gave off a watery discharge. There was a partial paralysis of the tongue and great quantities of saliva ran from the mouth. The limbs and ears were cold. The pupils of the eyes dilated, pulse not perceptible, mucous membrane of the rectum protruding, involuntary discharge of urine and faeces. Upon pricking the animal with a knife on the lower limbs it showed no feeling. The animal was closely watched in the field by Mr Foss and myself and we observed that she did not take any weeds, but simply a small amount of sorghum, eating only the tops of the leaves. At 1.30 P. M. the animal was still lying on its right side and the muscles of the head were contracted and showed involuntary twitching. The limbs were paralyzed and the animal was unconscious. The mucous membrane of the mouth was of a salmon color. At 2.15 P. M. the animal was in great pain, and it was apparent that she would not recover. At the suggestion of Mr Foss the animal was killed in order to hold a post-mortem examination.

Post-Mortem Examination—Animal still warm. The bowels were opened and contents of paunch carefully noted. There was in all one pound and a half of sorghum leaves to be found in the paunch. No sourness of the contents. The same was immediately put up in Mason fruit jars with clean water and brought to the laboratory. The mucous membrane of the intestines normal, all other conditions of the animal normal.

In regard to the Colorado disease the following statement is made:

The cattle died on August 28th last. We lost 21 head out of 32 head which had been turned on the corn. Eleven head lived, but 4 of the 11 head had violent spasms, but recovered. The other 7 head were not affected. They were only on the corn 5 or 6 minutes. The first cow died in 15 minutes nearly all within an hour. One yearling lived over 6 hours. I gave it several doses of arsenic, thinking possibly one poison would counteract the other but it died in great agony. The cattle seemed to all go crazy at once, then stagger like a person intoxicated, fall in all directions, and die where they fell. I stuck all of them with a knife, the same as in a la fa blast but there wasn't any gas in them. The Kafir corn was planted on mud ground above irrigation. It was from 6 to 15 inches high and was burnt brown from the drought."

Shortly after the poisoning, Dr Glover visited the field and collected samples, which he generously placed at our disposal. These samples yielded prussic acid in greater amounts than any yet examined in Nebraska.

It appears more than probable that the sorghum plant under different cli-

matic conditions and different conditions of growth may produce varying amounts of prussic acid. I was told in Texas that the sorghum most poisonous to live stock is the second growth.

In regard to the chemistry of the subject, Dr. S. Avery says

In 1886 Berthelot and Andre ascribed the cause to excessive amounts of potassium nitrate (salt peter). Williams of the U. S. Department of Agriculture also suggests saltpeter as a cause of the trouble. Hilner has shown that the amount of nitrate in Nebraska fields was too small to produce fatal results. This writer suggests that the plant under certain conditions develops a highly poisonous chemical compound. Slade in the Annual Report of the Station for the present year (1907) put forward the theory that such a compound might be produced by the action of an enzyme upon a glucoside found in the plant through a process of abnormal growth. On June 27th of the present year the Chemical News of London contained an article on Cyanogenesis in Plants by Dunstan and Henry. This article, which finally confirms Slade's prediction, was not known to Mr. Slade or to the writer till October 10th. In brief, the English Chemists isolated from Egyptian *Sorghum vulgare* a glucoside capable of liberating prussic acid. In the meantime Mr. Slade had detected Prussic acid in (a) sorghum grown in western Nebraska, determined the per cent. and secured strong evidence in favor of the glucoside theory.

During the first two weeks in September the writer discovered that Prussic acid could be obtained from leaves of healthy sorghum in the fields about the Station. As the past season was abnormally wet, nearly all of the fields had made a vigorous growth. By distilling water from a sufficient quantity of leaves, determinable amounts of Prussic acid were evolved in all cases, though the amount was well below the dangerous. Of the common Nebraska forage plants, sorghum and Kaffir corn alone yield Prussic acid.

The substance *dhurrin* $C_4H_7NO_4$ occurs according to Dunstan and Henry in young plants of *A. Sorghum*. A glucoside resembling that found in almonds also occurs, it differs however but is capable of being converted into hydrocyanic acid, HCN. The investigations made at this station show that the prussic acid is not present as such, but that it is liberated from a glucoside, (1) by an enzyme in the plant as in the case of sorghum poisoning, and (2) by the action of boiling water on the plant. Glucosides of this sort are in themselves harmless and are dangerous only when they liberate prussic acid. The experiments mentioned above also showed that even dried plants may contain a very large amount of combined prussic acid. We should expect that such a fodder would be as fatal to stock after curing as when standing in the field. Experience, however, seems to prove the contrary. Enzymes rapidly become inactive when dried in the presence of protein substances, according to Dr. A. F. Woods, Chief of Division of Plant Pathology.

Antidotes. Prussic acid has a tendency to unite with certain carbohydrates, forming additional products. These compounds are much less poisonous than the free acid. Both glucose and milk sugar unite with Prussic acid to some extent even in dilute solutions. Aside from this action these carbohydrates retard the action of the enzyme in liberating Prussic acid. These facts suggest that, in case the animal is not in such a condition as to render medical treatment out of the question, the following may be effective.

A strong solution of glucose, which nearly every farmer has at hand in the form of "corn syrup" or molasses, may be administered.

Large quantities of milk have in a number of instances been administered apparently with good effect.

In all cases the animal should have as much fresh air as possible.

3. *Paspalum*, L. *Paspalum*

Spikelets spiked or sometimes racemed, in 2 to 4 rows on one side of the flattened or filiform rachis, awnless, 1-flowered; glumes 3, rarely only 2, 1 glume flowering, flower coriaceous, orbicular or ovate, stamens 3, spikes 1 or more at or toward the summit of an elongated peduncle.

Species about 160, chiefly in warm temperate regions in both hemispheres. In South America they constitute an important part of the plants of the Pampas. One species is used in medicine and several species are excellent forage plants for the South. One species is troublesome as a weed in the Southern States. The Kona Millet (*P. scrobiculatum*) known to be poisonous and injurious to animals and man in India is used during times of scarcity of food and causes poisoning. The seed, especially the testa and pericarp, contains a narcotic poison which causes delirium and vomiting. Cattle should not be allowed to feed on it when it is ripening.




Fig. 142. Korean Foxtail (*Setaria italica*.)

Fig. 143. Korean Foxtail Miller (*Setaria*)
U. S. Dept. Agr.

4. *Setaria*, Beauv

Spikelets jointed upon the pedicels, panicle densely racemed or spiked, surrounded at the base by a few or many persistent awn-like bristles, which rise below the articulation of the spikelet.

Species about 10, in temperate and tropical regions. Some species are used as food especially in China, Japan and India. Several are important forage plants, like the broom corn millet, and Hungarian grass. Three species are weedy in eastern North America.

Setaria italica, Beauv. Italian Millet or Hungarian Grass

A stout, erect, somewhat glaucous annual, 3-8 feet high, with broad leaves large, dense, compound, spiciform panicles 3-8 inches in length, nodes bearded, with short, appressed hairs, leaf-blades lanceolate, narrowed at the base, long-acuminate, 8-16 inches long, $\frac{1}{2}$ to $1\frac{1}{4}$ inches wide, scabrous. Panicles dense, cylindrical, $\frac{1}{2}$ to $1\frac{1}{2}$ inches in diameter; rachis densely villous. Aetæ 1-3, green or purplish, retroscly scabrous. Spikelets elliptical, strongly convex $1\frac{1}{2}$ to 2 lines long, obtuse, second and third glumes about equaling the flowering glume, 5-7-nerved; flowering glume glossy, nearly smooth. Widely cultivated. Quebec to Minnesota, south to Florida and Texas.

Setaria germanica, Beauv. German Millet

A caespitose annual, from 1-3 feet high, with narrow panicles, about $\frac{1}{2}$ inch in diameter and long, usually purple aetæ; some forms approaching *Setaria viridis*.

This form is usually regarded as only a variety of the Italian Millet, and is found in cultivation only or perhaps springing up from seed on land cultivated the preceding season. The German Millet differs from the Italian in having a more dense or compact, and usually erect panicle or "head." Widely cultivated in most parts of the world.

Poisonous Properties. Numerous complaints have been made from time to time with reference to poisoning from millet.

Dr Hinzbauch states in regard to this trouble that in the winter of 1891 and 1892 a disease commonly called millet disease was prevalent to a considerable extent in North Dakota and that this disease was attended by a death rate of 7-10 per cent. It received the name of millet disease from the fact that from 95 to 98 per cent of the animals that were affected had been fed on millet. He says

"When millet is fed in considerable quantities it stimulates the kidney to increased action. The urine is light colored and the bladder evacuated every two or three hours, large quantities of water being passed at each time. At the time the first symptoms of lameness were noticed, the kidneys had almost ceased to act."

And then he goes on to say:

"When the cause was kept up a sufficient length of time for the reaction to set in, the material which would under normal conditions be secreted by the kidneys was allowed to remain in the system and produce deleterious effects."

Apparently the condition of the millet had little to do with this action. In a later bulletin on the same subject Dr Hinzbauch reports a more extended investigation, giving considerable experimental data as well as urinary analyses

The post mortem examinations revealed some interesting facts. The cartilages on the ends of the long bones show deep furrows running in a direction parallel with the motion during flexion and extension.

Both grooves of the astragalus were partially denuded of cartilage, so that the corresponding elevations of the tibia which articulate in the grooves did not have cartilage interposed between them. The whole general appearance, instead of being of a white, glistening color, was of a dark, dull color bordering on brown. The fluid which escaped from the joint when opened, instead of being a yellow, amber color, was brown and contained red blood corpuscles, indicating that inflammation was present. The joint fluid was brownish black in color and contained red blood corpuscles.

In conclusion we would say that our experiments here have thoroughly demonstrated that millet, when used entirely as a coarse food, is injurious to horses (1) In producing an increased action of the kidneys. (2) In causing lameness and swelling of the joints. (3) In producing infarcts of the blood into the joints. (4) In destroying the texture of the bone, rendering it softer and less tenacious, so that contraction causes the ligaments and muscles to be torn loose. The experience of many farmers with whom I have talked confirms the above conclusion, and we could multiply case after case showing that the above conditions are the results of feeding millet.



Fig. 143. Sandbur (*Cenchrus tribuloides*).
spiny bar enclosing spikelets; b, section of
the same; c, lateral view of a spikelet. U. S.
Dept. Agri.

The North Dakota Station has published the results of further experiments on the subject of feeding millet. Two tests were made. In the first trial two geldings in good health were fed hay and grain for about two weeks. Millet was then substituted for hay for about ten days. These experiments confirmed those made previously.

Ladd has isolated a glucoside from the aqueous extract of millet hay, which, when fed in small quantities, gave the characteristic symptoms.

From the experiments made by Dr. Hinebaugh and others, it would appear that feeding millets alone as coarse fodder is injurious to horses. It produces an increased action of the kidneys and causes lameness and swelling of the joints. It causes an infusion of blood into the joints and destroys the texture of the bone, rendering it soft and less tenacious, so that the ligaments and muscles are easily torn loose.

5. *Cenchrus*, L. Sand Bur

Annual or perennial grasses, flat leaves, spikelets surrounded by a spiny involucre which becomes coriaceous and forms a deciduous, hard, rigid bur which falls away at maturity, glumes 4, the 2nd and 3rd membranaceous, the 4th hard, the palea enclosing the perfect flower, stamens 3; styles united below.

Species about 12 in tropical and warmer temperate regions. One widely distributed from Maine to New York, Florida, Texas, California and the Dakotas.

Cenchrus tribuloides, L. Sand Bur

An annual with erect culms a foot or more high, flat leaves about 6 inches long; burrs of the involucre with strong, barbed spines; 2-flowered.

Distribution. Common in sandy fields and waste places, a weed along railroads and in sandy soil.

Injurious Properties. This plant frequently inflicts mechanical injuries, entering the flesh and thus causing serious inflammation. This applies to man as well as to lower animals.

Aristida, L. Triple Awned Grass

Perennial or annual grasses; narrow, often involute leaves; spikelets narrow, 1-flowered, outer glumes unequal, often bristle pointed, flowering glume tipped with 3 awns, palea small, 2-nerved, stamens 3, styles distinct, grain free, linear enclosed in the scale, callus variable, often sharp-pointed and rigid. About 100 species in warmer regions of both hemispheres but of very little economic value, the majority being found in dry sterile soil, several species, like the Purple *Aristida*, however, are common in dry soils of the West. The latter is of little value for forage purposes. The awns of *Aristida hygrometrica* of Queensland bore into the skin of animals and occasionally reach the intestines, thus causing death.

None of our species produces serious trouble except, possibly, the Long-awned Poverty Grass.

Aristida tuberculosa, Nutt. Long Awned Poverty Grass

A rigid, much-branched annual, 12-18 inches high, with nearly simple panicles, 4-7 inches long, branches erect, rather distant, the lower in pairs, one short and few flowered, the other elongated and many flowered, empty glumes



Fig. 144a. Long-awned Poverty Grass (*Aristida subarcalosa*). a, Spikelet with lower glume, b, flowering glume with divergent long awns. (U. S. Dept. Agri.).



Fig. 144b. Short-awned Poverty Grass. (*A. brevicauda*). Occurs in sandy and gravelly soils. (U. S. Dept. Agri.)

nearly equal, 12 lines long, awn pointed, flowering glume about 10 lines long, twisted above to the division of the awns, and with a densely barbate sharp-pointed callus, awns nearly equal, divergent or reflexed, $1\frac{1}{2}$ 2 inches long, distinctly articulated with the glume.

Injurious properties The sharp pointed callus slightly injurious in the same manner as *Stipa*.

7 *Stipa*, L.

Perennial grasses with 1 flowered spikelets, flower falling away at maturity from the membranous, persistent, lower glumes, fertile glumes coriaceous, cylindrical, involute, and embracing the smaller palea and cylindrical grain a long twisted or spiral awn jointed with the apex, the base consisting of a beard and sharp pointed callus, stamens generally 3; stigmas plumose

About 100 species found in temperate and tropical regions. The Stipae often produce injurious effects upon animals.

Injurious properties. It has long been known that *Stipa capillata*, L., indigenous to Russia, and the *Stipa spartea*, Trin., and *S. avenacea*, L., native to North America, as well as *Arctida hygrometrica* Br., native of Queensland,




Fig. 145. Esparto Grass (*Stipa tenacissima*). Used for making paper, ropes and mats. It is not known whether this species, like the *S. umbrosa* and the *S. spartea*, acts like a narcotic on animals. (Baillon Diet.)

and *Heteropogon contortus*, L., native of New Caledonia, frequently bore into the skin and in estives of lower animals where they cause fatal inflammation and peritonitis. Prof. Blanchard in a recent number in "Archives de Parasitologie" calls attention to injurious properties of *Stipa Vescaana* which is found in Uruguay and other countries of South America. In this case the needles injure the eyes producing an intense keratitis often followed by inflammation of the cornea. Sheep become blind and they are unable to get food, hence

die from hunger and thirst. An instance is also recorded of a case where so many of these needles had accumulated among the feathers of an American Ostrich as to cause extensive ulceration which finally resulted in the death of the bird. The old world *S. nebrascensis* acts very much like our sleepy grass

Stipa comata, Trin. and Rupr. Western Stipa. Needle Grass

A rather stout, erect, caespitose perennial, $1\frac{1}{2}$ -4 feet high, with mostly involute leaves, loosely-flowered panicles, 8 to 12 inches long, spikelets with nearly equal, long-attenuate-pointed, empty glumes about 12 lines long, and thinly




Fig. 146. Western Stipa or Needle Grass (*Stipa comata*)

pubescent flowering glumes about 6 lines long, awn slender, $2\frac{1}{2}$ -3 inches long, strongly flexuose or variously curled and twisted. Distributed in western Iowa, Nebraska, Utah, Oregon, California and Arizona.

Stipa spartea, Trin. Porcupine Grass

A stout erect perennial, with simple culms 3 to 5 feet high, long, narrow leaves and contracted, few-flowered panicles, 4 to 8 inches long spikelets

larger, empty glumes subulate-pointed, 12 to 18 lines long, slightly unequal; flowering glume 8 to 10 lines long, including the barbed and very sharp-pointed stipe or callus, sparsely pubescent below and crowned with a few short hairs;



Fig. 149. Needle or Porcupine grass (*Sasa sporea*) a, a single spikelet; b, flower more highly magnified, with sharp pointed bearded callus. (Div. Agron. U. S. Dept. Agr.).

palea nearly as long as the glume; awn stout, 3 to 6 inches long, twisted below and twice geniculate above. June to August. Common on dry, gravelly roads and high prairies.

Distribution. North America. From Wisconsin, Illinois to Missouri, Kansas, Nebraska, Dakotas and Minnesota to New Mexico, Manitoba to British Columbia.

Injurious properties. Dr. M. Stalker says the fruits of the porcupine grass are a frequent source of inconvenience and injury to living animals.

In many of the northwestern counties of Iowa this grass grows in the greatest profusion, and during the latter part of June, the season for mowing and consequent falling of these spikes, they are the occasion of much annoyance and in some instances the death of domestic animals. Only such animals as are covered with wool or a

thick growth of long hair are seriously inconvenienced. Sheep suffer most. The spines readily find a lodgment in the wool, and after burrowing through it frequently penetrate the skin and bury themselves in the flesh. A large number of these barbs thus entering the tissues of the body produce an amount of irritation that is sometimes followed by death. I have seen large numbers of these imbedded in the skin and muscular tissues of shepherd dogs that were covered with a thick growth of soft hair. These mischievous animals frequently exhibit the greatest dread at being sent into the grass during the season of danger.

Professor Bessey in his inquiries into the structure and nature of this plant received several responses, one of which, from Professor King, formerly of the University of Wisconsin, was as follows:

In connection with the two notes relating to the fruit of the poison one grass, it may not be without interest to say that while engaged in geological work in Dakota, north of the Northern Pacific railroad, we were much annoyed by the fruit of this grass. Indeed, I found the only way to walk with comfort through this grass was to roll my pants above my knees and my socks down over my shoes. I also observed, on several occasions, these seeds planted two inches deep in the soil with the awns protruding from the ground. It is plain that with the point of one of these fruits once entered below the soil, the swelling and shrinking, due to varying amounts of moisture, would work the seeds directly into the ground.

The *Stipa comata*, or needle grass of the west, which is common throughout the Dakotas, and throughout west Dakota, Nebraska, Wyoming and Colorado, is common in prairie hay, and Prof Thomas A. Williams mentions that though a forage plant, and not cut until the needles have fallen so that the stock may not be injured, the fruit of this plant often injures stock to a considerable extent. During the past summer in Alberta, Canada, the writer suffered some inconvenience from the penetration of the fruit through the clothes.

Stipa robusta, Vasey. Sleepy Grass

A large grass from 4-6 feet high growing in dense clumps. Leaves involute setaceous, large, flattened, 12 feet long, panicle 1 1/2 ft long, spikelets 4-5 lines long, on short pedicels, empty glume nearly equal 3-5 lines long, variable in length up to 1 1/2 inches, slender flexuous, pale about 1/2 length of glume.

Distribution. From Colorado to Texas and Mexico.

Poisonous properties. This is the grass which is properly called Sleepy Grass and is poisonous. Dr Vasey says the variety in parts of Texas and Mexico is known as Sleepy Grass, so called for its intoxicating and narcotic effect upon horses or cattle which feed upon it.

In the west this species of grass has received the common appellation of sleepy grass. It has long been regarded by range people as poisonous. Dr Palmer, who found this grass in Coahuila, observed that it was poisonous to cattle, horses, and sheep, causing them temporary sleepiness. Later Dr Havard states that in 1898 he received from Dr M. E. Taylor, of Stanton, N. M. a grass with the following statement:

Perumbouts grows a grass the eating of which by horses will, within a few hours, produce profound sleepiness or stupor lasting twenty four or forty eight hours, when the animals rally and give no evidence of bad effects. It is known among cowboys as "sleepy grass" and dreaded by them on their "round ups" as their horses are liable to eat it and cannot then be kept up with the herds. The tradition is that horses that have once eaten of it will not touch it again.

To quote from Dr Havard

From the same gentleman I received a letter in 1890, in which he says: "Since I corresponded with Dr. Taylor it has been brought to my notice that cattle are affected in a similar way to horses, and that the curious properties which so affect animals are contained in the blades. Quite a number of our horses have been ill this spring

after having eaten it. It usually takes them about a week to recover, during which time they are unfit for work, and especially so during the first three days."

Captain Kingsbury, of the Sixth United States cavalry, under date of March, 1890, wrote me from Fort Stanton that the sleepy grass affected nearly all his horses at two camping places. It was hard work to make them walk.

The similarity of symptoms, whether observed in Coahuila or in New Mexico, is certainly remarkable, and furnishes strong evidence of the substantial accuracy of the observations as reported. It would seem, then, reasonably established that this plant possesses narcotic or sedative properties, affecting principally horses, but also cattle and probably other animals, that animals are not fond of it but eat it inadvertently or when under stress of hunger that cases of poisoning occur especially in the spring, when the radicle and lower blades first come up, and that the active principle resides in these blades, and perhaps only during that season.

8. *Avena*, L. Oats

Annual or perennial grasses, usually with flat leaves and paniced spikelets, spikelets 2, many-flowered, or rarely 1 flowered, lower flowers perfect, the upper staminate or imperfect, empty glume unequal, membranaceous and per-



Fig. 148. Wild Oats (*Avena fatua*). a, empty glumes b, flowering glumes. (L. S. Dept. Ag.)

sistent, flowering glume deciduous, generally bearing a twisted awn on the back between the two acute teeth at the apex, rachis and base of flower often bearded, stamens 3, style short and distinct, grain oblong, linear, grooved on one side invested by the palea. About 50 species in temperate regions. The cultivated Oats (*Avena sativa*) is the best known representative of the genus and has long been used for food for man and animals. Several native species produce good forage.

Avena fatua L. Wild Oats

An erect, glabrous annual, 3-5 feet high, with flat leaves and spreading panicles of large, nodding spikelets, spikelets 2 to 4-flowered, with empty glumes $\frac{1}{4}$ 1 inch long and pubescent, flowering glumes 6 to 9 lines long, awns nearly twice as long as the spikelets. Wild oats is highly esteemed as a forage plant on the Pacific Coast especially California.

Distribution. Native to Europe but now abundant in grain fields of the Rocky Mountains the Dakotas, Minnesota, and the Pacific Coast.

Injurious properties. Bezoars are sometimes produced by the common oat and Dr. Harz thinks it is a dangerous food material because it favors the development of these "hair balls." The barbed and awned seed of the wild oat may probably sometimes also lodge in the mouth and produce inflammation or other results of mechanical injuries.

Avena sativa, L. Common Oats

A well known erect annual, 2-4 feet high, with flat leaves and expanded panicles of rather large, pendulous, and, usually, 2-flowered spikelets. Lower florets sometimes awned.

Distribution. Widely cultivated in Europe, North America, Asia, and in all temperate regions. Commonly cultivated in Northern United States, Canada, and the Pacific Coast. The species is native to eastern temperate Europe and western Asia although the wild form has not been found. According to some authors, cultivated oats originated from wild oats *Avena fatua*. This is very doubtful.

Injurious properties. Harz reports the occurrence of phytobezoars in horses which had been fed oats straw. These bezoars in their origin and structure are similar to those occurring from feeding on cacti and the crimson clover referred to elsewhere.

9. *Bromus*, L.

Spikelets 3 to many flowered, panicle, glumes unequal, membranaceous, lower glume 1-5 nerved, flowering glume either convex on the back or compressed-keeled, 5-9-nerved, awned or bristle pointed from below to the groove of the oblong or linear grain, stamens 3, styles attached below at the apex of the ovary. Coarse grass with large spikelets at length drooping or pedicels thickened at the apex. About 40 species, of which Beal lists 27 as either native or introduced into the U. S.

Bromus tectorum, L. Awned Brome Grass

A slender, erect, leafy annual, 7-25 inches high, with narrow, softly pubescent leaves and open, nodding panicles, 3-7 inches long, spikelets each 5-8 flowered, with unequal, acuminate-pointed, hairy, empty glumes, and rough or



Fig. 149. Common Oats (*Avena sativa*), sometimes the cause of phytohemorrs in animals. (La Seed Co.)

hairy glumes 4-6 lines long, awns 6-8 lines long, blooming period from June to August. First introduced into the United States from Europe, it is without forage value, and, while not greatly troublesome in Iowa or eastward, has become a serious pest farther west, in Utah and Colorado.

Poisonous properties. This plant produces injuries similar to those caused by Squirrel-tail grass, the awned glumes working in under the teeth causing inflammation and suppuration. Animals eating this grass may lose their teeth as a consequence.



Fig. 150. Awned Bromegrass (*Bromus tectorum*). a. Sterile or outer glumes. b. Spikelet. U. S. Dept. Agr.

10. *Lolium*, L. Darnel and Rye Grasses

Annual or perennial grasses with flat leaves and terminal spike, spikelets many-flowered, solitary on each joint of the continuous rachis placed edge-wise, empty glumes except in the terminal spikelets, only one flowering glume, rounded on the back, 5-7 nerved, palet 2-keeled, stamens 3, grain adherent to the palet, 6 species, 2 more or less naturalized in the eastern states. Natives of the Old World. Two species, the Italian rye grass and the common rye grass, are valuable forage plants. Darnel is a troublesome, poisonous grass.

Lolium temulentum, L. Darnel, Poison Darnel

An annual with smooth stout culm, 2-3 feet high; leaves with scabrous sheaths and short ligule, spike 6-12 inches long, spikelets 5-7 flowered, empty glumes sharp pointed, as long as the spikelets; flowering glume awned or awnless. Commonly found in grain fields.



Fig. 151. Darnel (*Lolium temulentum*). b. Spikelet. a. Empty glume. L. S. Dept. Agr.

Distribution. Naturalized in eastern North America and abundantly so on the Pacific Coast.

Poisonous properties. It is a well known fact that a number of grasses are poisonous. It was well recognized by the ancients that darnel (*Lolium temulentum*) was poisonous, for it is written "But while men slept, his enemies came and sowed tares among the wheat."

Darnel when ground up with wheat and made into flour, is said to produce poisonous effects on the system, such as headache and drowsiness. This poisonous property is said to reside in a narcotic principle, *tolin*, a dirty white, amorphous, bitter substance yielding sugar and volatile acids, which, according to Hackel, "causes eruptions, trembling and confusion of sight in man and flesh-eating animals, and very strongly in rabbits, but it does not effect swine, horned cattle or ducks." Landlev states that the grain is of evil repute for intoxication in man, beast and birds, and brings on fatal convulsions. Haller

speaks of it as communicating these intoxicating properties to beer. It acts as a narcotic, acrid poison. Darnel meal was formerly recommended as a sedative poultice. In Tayler's work on poisons, the statement is made that the seeds whether in powder or in decoctions, have a local action on the alimentary canal and a remote action on the brain and nervous system. He states further that no instance is reported of its causing fatal injuries to man, and as much as three ounces of a paste of the seeds have been given to a dog without causing death. Then he goes on to cite the experience of Dr. Kingsley, in which several families, including about thirty persons, suffered severely from the effects of bread containing the flour of darnel seed. These persons had staggered about as if intoxicated. It is claimed by some investigators, however that this plant is not poisonous. One writer claims to have made bread from flour said




Fig. 152. At left, a hypha at base of seedling of "Darnel" (*Lolium temulentum*). At right, hyphae in the starch endosperm of a seed. A, hyphal layer of grain nucleus; B, starch cell; C, wall of starch cell; D, knot formation in an intercellular space. After Freeman, redrawn by Charlotte M. King.




Fig. 152. "Darnel" (*Lolium temulentum*). Section of outer part of a grain which has been in a germinating chamber 24 hrs. i, pale; k, pericarp; o, crushed integuments; s, outer row of nucellus cells. B, cavities with nucellus (probably old cell lumina); A, hyphae; C, nucellus; D, starch endosperm. After Freeman, redrawn by Charlotte M. King.

to contain considerable darnel and experienced no injurious effects. When mixed with flour and water the dough is foamy and narcotic in its action. There are other grasses which produce similar narcotic effects. Quite recently it has been claimed by several European investigators that the fruit of *Lolium temulentum* contains a poisonous fungus. Guerin states that the hyphae of a fungus constantly occur in the nucleus of the seed and the layer of the caryopsis lying between the aleurone layer and the hyaline portion of the grain or nucellus. He also thinks that the toxic action of the *Lolium* is due to this particular fungus hypha. The threads were also found in *L. arvense* and *L. baccatum*, but, as yet, have not been found in *L. italicum* and but once in *L.*

perenne, the fungus is allied to *Endoconidium temulentum*, which has been found on rye.

The *Lolium* fungus, according to Guerin, lives symbiotically in the maturing grain and is therefore not a parasite, but Freeman has observed that occasionally it is injurious although it is generally stimulating. Nestier who made an examination of *L. perenne*, *L. multiflorum*, *L. temulum*, and *L. festucaeum*, found nothing comparable to the fungus mycelium which occurred in *L. temulentum*. He also succeeded in demonstrating the presence of the mycelium of the fungus as indicated by Guerin. According to Nestier, the *Fusarium roseum* is identical with the fungus occurring in *L. temulentum* found by Guerin, but this has not been confirmed and seems very improbable. Hanaussek considered the fungus to be related to the smuts, but Freeman found no evidence of spore formation, the septa are infrequent and the intercellular course different from that for smuts. The subject has, in recent years, been investigated by Prof. Freeman who, in a general way, confirms the reports of previous investigations and says:

The probabilities of relationship with the ergot of *L. temulentum* are very interesting. The frequency of occurrence of ergots of *Lolium* in England is strangely coincident with that of the fungus in the grain, e. g. most abundant in *L. temulentum*, less so in *L. perenne* and exceedingly rare in *L. uilatum*.

It is certainly not one of the rusts and the Ustilaginaceae are the closest affinity, perhaps, the fungus is carried from one generation to another by the sterile mycelium, when the embryo of the grain pushes out during germination, the hyphae, being in the "seed" keep pace with its growth and can be detected in the growing point throughout the life of the plant. Prof. Freeman says:

The hyphae sometimes penetrate the aleurone layer at any point and invade the starch endosperm. There exists in the scutellum, at the base of the scutellum and at the lower end of the inner groove of the grain, a layer of hyphae which lies directly against the embryo, constituting an infective layer.

11 *Agropyron*, Gaertn. Quack or Wheat Grass

Annual or perennial grasses, with flat, or involute leaves. spikelets 3-many-flowered, compressed, 2-ranked, alternate on opposite sides of the solitary terminal spike, 1 at each joint, or, occasionally, all, or the lower in pairs, sessile, with the side against the axis; glumes transverse, nearly equal and opposite, lanceolate. flowering glumes rigid, rounded at the back, 5-7 nerved, pointed or awned from the tip, pale flattened, bristly, ciliate on the nerves, adherent to the grain.

About 40 species, in temperate regions. The root of quack grass is used in medicine. several species, like western wheat grass (*Agropyron occidentale*) and slender wheat grass (*A. tenerum*) are valuable for forage purposes. Quack and western wheat grasses are also good soil binders.

Agropyron repens, (L.) Beauv. Quack Grass

Perennial, 1-3 ft. high, from a creeping, jointed rootstock, sheaths usually smooth, scabrous, or pubescent above. spikes 3-10 inches long, erect, spikelets 4-8 flowered, empty glumes strongly 5-7 nerved near the apex, awnless or sometimes short awned.

Distribution. Widely naturalized, a good forage plant and also a bad weed. In eastern North America, it occurs in cultivated fields and by roadsides and is a troublesome weed.



Fig. 154. Quackgrass (*Agropyron repens*). The roots (rootstocks) used in medicine. The roots contain considerable of a nutritious carbohydrate. (C. H. King.)

Medicinal properties. Quack grass is not known to be poisonous. The ancients since the time of Pliny have used the drug in medicine and it was also used by the Germans in the 10th century. The root stock is official. Gerard ascribed to the root diuretic, lithontriptic virtues or properties. The root contains considerable sugar and a substance called tritidin, an amorphous, gummy substance easily transformed into sugar. It is found useful in the mucous discharge from the bladder. Quack grass and Western Wheat Grass frequently contain ergot.

12. *Hordeum*, L. Barley.

Annual or perennial grasses with flat leaves, cylindrical spikes, spikelets 1 flowered, with an awl-shaped rudiment on the inner side, 3 at each joint of the rachis of a terminal spike, the lateral ones usually imperfect or abortive and with a short stalk, empty glumes side by side in front of the spikelets, forming a kind of involucre, flowering glume and palea herbaceous, the former long and awned from the apex, stamens 3, styles very short grain usually oblong and adherent to the palea, spike often separating into joints.

About 20 species widely distributed in both hemispheres. Of the Barleys, the 2-rowed barley (*Hordeum distichum*) and the 4-rowed barley (*H. vulgare*) are well known in cultivation, being used for malting purposes and occasionally in medicine. The awns of cultivated barleys produce mechanical injuries to stock. Several members of the genus are very troublesome weeds.

Four-Rowed Barley. Annual, 2-3 feet high, smooth leaves linear-lanceolate, keeled, nearly smooth, sheaths striate, smooth, auricled at the throat; ligule very short, spikes 3-4 inches long, somewhat 4-sided rachis flattened, pubescent on the margins, spikelets with 1 perfect floret, empty glumes narrowly linear, pubescent, terminating in a slender awn, flowering glume 5-nerved, scabrous near the apex, long-awned, awn flattened, keeled, somewhat 3-nerved, serrulate on the margins.

The cereal is without doubt one of the most ancient of cultivated plants. It is supposed to have originated from *H. spontaneum* Koch which grows wild in Asia Minor and Caucasian countries to Persia and Beloochistan as well as in Syria and Palestine.

Hordeum jubatum, L. Squirreltail Grass

An annual or winter annual from 6 inches to 2 feet high producing fibrous roots, forming solid and compact bunches; leaves not unlike those of blue grass, but paler in color, from 2-4 inches in length, margins scabrous, flowers in dense spike from 2-4 inches long, pale green or purplish in color, consisting of a number of 1-flowered spikelets, 3 occurring at each joint 1 being perfect (bearing stamens and pistil), 2 others awl-shaped, and borne on short stalks, 1 sterile spikelet occurring on each side of the perfect flower which bears a long awn, at each joint will be found 6 empty, long awned glumes, spreading at maturity which give to the plant its bristly appearance, when mature, the spike breaks up into joints consisting of the rudimentary spikelets and a perfect flower so that each joint has 1 "seed," the number of seeds in the spike varying from 35 to 60. A single cluster of plants may therefore produce from 300 to 2000 mature seeds. The plant has a wonderful capacity for "stooling." From a single plant as many as forty spikes may be produced and the number often no doubt exceeds this.

Distribution. It is found in marshes, in moist sand along the sea shore, and near the northern lakes. Its present distribution is from Nova Scotia to New Brunswick along the Atlantic coast, Maine to Maryland and westward to the region of the Great Lakes, Minnesota, Saskatchewan and the Mackenzie river the Dakotas, Iowa, Nebraska and the Rocky Mountain region south to Texas, California and southern Mexico. It is also reported from Europe and Siberia.

Originally it was chiefly distributed in the Rocky Mountain region occur-

ring in the saline soils of the plains, the great lakes and along the seacoast extending far northward. Its extension eastward and westward has taken place in the more recent times.

Hordeum serotinum, Schreb. Little Barley

An erect annual from 4-10 inches high, more or less geniculate at the lower end, sheaths smooth or upper often inflated; leaf blade 1-3 inches long, spikes narrow, empty glumes rigid, those of the central spikelet scarcely lanceolate, all awn pointed flowering glume of the central spikelet awned or nearly so.

Distribution. Common and troublesome especially as far east as Missouri, Nebraska, British Columbia, and California.

Hordeum murinum, L. Wild Barley

An annual from 1-2 feet tall, erect or geniculate at the base leaves rough, spikes from 2½-5 inches long, spikelets usually in 3's, scales awned, the empty glumes awnlike and scabrous, the second scale of the lateral spikelets not ciliate, the flowering glumes scabrous at the apex, bearing an awn about 1½-1¾ inches long.

Common on the Pacific coast and the dry regions of Utah, New Mexico, Arizona, and occurring on ballast in the Eastern states.

Mechanical injuries It has long been known that the barbed awns of barley, wild barley and other plants act injuriously in a mechanical way. In the west this is especially true of wild barley (*Hordeum jubatum*).

Dr. S. H. Johnson, of Carroll states in the Carroll Herald that this grass, when found in hay and allowed to ripen, if in any quantity, is very injurious to horses' mouths. He says:

The small awns seem to work in and cause deep ulcerating sores, which form under the tongue and lips. The writer has seen a large number affected and made a careful examination, and found the awns deep in the flesh, where they had remained for three months or more. I have seen lips eaten completely through and tongues eaten almost off by the grass. As to cattle, I have seen some affected, but not to any extent because the mucous membranes are much thicker. The sooner the grass is eradicated the better.

Professor Nelson, who has carefully studied this question, says on the injury to stock:

The awned heads, when taken into the mouth, break up into numerous sections, scatter within the mouth and everywhere adhere to the mucous membrane, which soon becomes pierced with the long stiff awns. As the animal continues to feed, more awns are added, and those already present are pushed deeper into the flesh. Inflammation soon results and leaves the gums of the animal in condition to be more easily penetrated. The awns are particularly liable to be pushed down and alongside and between the teeth. As the swelling and festering progresses the awns are packed in tighter and pushed deeper and cause suppuration of the gums as well as ulceration of the jaw bones and the teeth. Through the absorption of the ulcerated sockets and roots the teeth become loosened and even drop out, but the animal, impelled by hunger, still endeavors to eat such hay as may be offered.

The above statements apply largely to *H. jubatum*, but are equally true of all other species given above.

2. CYPERACEAE. Sedge Family

Grass-like, or rush-like herbs. Culms slender, solid or rarely hollow frequently triangular, terete, quadrangular or flattened; roots fibrous and frequently creeping rhizomes, leaves narrow, sheathes closed; flowers perfect or imperfect




Fig. 155. Wild Barley (*Hordeum jubatum*). Produces mechanical injuries to animals. (U. S. Dept. Agri.)



Fig. 156. Common Little Barley (*Hordeum germinum*). Causes mechanical injuries. (Charlotte M. King.)

arranged in spikelets, 1 or 2 in the axil of each glume, spikelets, 1 many-flowered, scales 2 ranked, or spirally imbricated, persistent, or deciduous perianth free, composed of bristles, scales or rarely wanting, anthers 2-celled ovary 1-celled, ovule 1, erect style 2 or 3-cleft, endosperm mealy, embryo minute. A large family of comparatively few genera (65) and 3,000 species of wide distribution. *Carex* is found in colder regions, while *Cyperus* is in warmer

regions. About 600 species of *Cyperus*, 200 of *Scirpus*, 200 of *Rynchospora* and 1,000 of *Carex*. The Papyrus (*Cyperus Papyrus*) of Africa and Sicily was used by the ancients as writing material. Common rush (*Scirpus lacustris*) a cosmopolitan plant found in water and marshes, is used for making mats and baskets. The rhizome of *Carex arenaria* is used in medicine.



FIG. 157. Sedge (*Carex arenaria*). 1. Flower (on plant). 2. Staminate flower with glume. 3. Pistillate flower. 4. Pistil. 5. Back of pistillate flower. 6, 7. Staminate and pistillate flowers of *C. hystrix*. (After Weisbach.)

PRINCIPES

Woody or herbaceous plants with endogenous stems, flowers in spikes, generally on the plan of 3, free, regular or slightly irregular, stamens 3-9 or numerous but generally 6, ovaries, free, 1-7-celled usually, fruit dry or a

Fleshy drupe Contains the family *Palmeæ*, a large family of 1,000 species, of which the most important palms are as follows. The date palm (*Phoenix dactylifera*) of Asia and North Africa, now cultivated in warmer regions of Europe, California and Arizona, and an important article of commerce in North Africa. the *Corypha* which furnishes sago, fiber, and a seed which is used as a substitute for coffee, and the *Washingtonia* of Southern California, frequently cultivated. Vegetable wax is derived from *Copernicia cerifera*. The wine palms (*Raphia rufifera* and *R. pedunculata* of eastern Africa), furnish raphia fiber. The *Metroxylon Rumphii* of the South Sunda Islands furnishes sago. The betel-nut palm (*Areca Catechu*) is much used as a narcotic the poison derived from this being known as arecain, half a grain of which is sufficient to kill a rabbit in a few minutes. It acts upon the heart and influences respiration causing tetanic convulsions. It also causes a contraction of the pupil of the eye. It is used to some extent as a vermifuge and in India and the Islands of the Pacific it is applied as an external remedy. The nut contains the alkaloids, *arecolin*, *arecain*, *arecoidin*, and *gubarin* which are used as vermifuges for dogs. The orange colored fruit is about the size of a hen's egg. When the nut is wrapped in quicklime and used, it imparts a red color to the saliva, it injures the teeth, and eventually destroys them. The resinous exudation from dragon's blood (*Daemonorops Draco*) of the East Indies is used in the manufacture of paints and varnishes. The oil from the oil palm (*Elaeis guineensis*) of West Africa and eastern South America is an important article of commerce. The cocoa-nut palm (*Cocos nucifera*) in tropical countries, especially the Islands of the Pacific, is an important article of food. The milk is the endosperm. The juice in the nut before maturity is unwholesome, being strongly diuretic and likely to cause serious results when taken into the system. A fermented drink is made from the juice of the plant which causes obesity and premature old age. A fiber known as coir is made from the husk. Vegetable ivory (*Phytolophus macrocarpa*) of tropical countries is a well known article of commerce. "Tuba" or Philippine toddy is made from the sap of the flowering spadix of *Hypha fruticosa*. Toddy is also made from the juice of *Arenga pinnatifida*, a plant which also furnishes an almost imperishable fiber. The "Royal Palm" is the "Yagua" (*Roystonea borinquena*) of Porto Rico, the sheathing bases of the leaves of which are used in thatching and siding the houses of the poor. An oil is produced from the husk and nut-like seeds of the *Acrocomia* or corozo palm which is distributed through tropical America from Mexico and Cuba to Paraguay.

SPATHIFLORÆ

Mostly fleshy herbs with endogenous stems, or thalloid floating plants; flowers generally in a fleshy spadix subtended by a spathe or naked, or a few solitary flowers on the margin or back of the thalloid structure.

ARACEÆ Arum Family

Herbs with pungent juice, leaves with long, slender petioles and abounding in raphides. flowers borne in densely-flowered fleshy spadix subtended or enclosed by the spathe. rootstock tuberous, floral envelopes none or of 4-6 sepals, stamens 4-10 filaments short, anthers 2-celled, ovary 1 several-celled ovules 1-several in each cell, fruit a berry, seeds various, with 2 coats, the outer fleshy, endosperm abundant or none. About 900 species of wide distribution.

Many of the plants, as the skunk cabbage, *Symplocarpus foetidus* possess acrid and noxious qualities. This is a native herb which is acrid and has a disagreeable odor. The fleshy spadix of *Monstera deliciosa* of the Mexican




Fig. 158. Common European Arum, Cuckoo-pint, or Wake-robin (*Arum maculatum*). Leaf; spadix; longitudinal section of ovary; germination; longitudinal section of seed; embryo. (After Paget.)

Cordilleras is edible. The vegetable calomel (*Acorus Calamus*) is used in medicine and contains the bitter principle *acorum* and an alkaloid. The sweet calomel is poisonous, under some conditions, causing disturbed digestion, and, in severe cases, gastro-enteritis, persistent constipation, followed by diarrhoea and passage of blood in the feces.

The *Calla palustris*, a marsh plant, has acrid properties and is used in Lapland with bread. The bulbs of *Amorphophallus* are rich in starch and are edible. The *Richardia africana* is frequently cultivated and used as food, a starch being also made from it. The poisonous substances contained in it are removed on roasting and boiling. In some of the fruits of aroids, like *Arum italicum*, *saponin* has been found, also needle-like crystals of oxalate of lime. *A. maculatum* is poisonous and causes severe dermatitis, paralysis, and, in the

case of children, even death. The *Thomsonia napalensis* of India, according to Major Kirtikar, is an acrid poison but its deleterious properties may be removed by roasting. The arrow arum (*Peltandra virginica*) of eastern North America is an irritant.

Arisaema. Mart.

Perennial herbs with tuberous rootstock or corm, having acrid properties, leaves simple or compound, scape simple, spathe convolute, generally arched above, spadix with flowers near the base, floral envelopes none, flowers monoecious or dioecious, stamens 4, anthers 2-4-celled, pistillate flowers with a 2-celled ovary containing many ovules, fruit a globose, red berry, seeds with copious endosperm. About 50 species found in temperate climate.

Arisaema triphyllum. (L.) Schott. Indian Turnip

Corm turnip-shaped, farinaceous, leaves generally 2, divided into 3-foliate



leaflets, ovate, spadix mostly dioecious, club-shaped, much shorter than the arched spathe, which is green and purple striped, ovules 5-6, berries shining, forming a dense head. The dragon head (*A. Dracontium*) with solitary leaf pedately divided into 7-11 oblong lanceolate leaflets, and spadix tapering to a long slender point, is common in rich woods from Minnesota and Iowa, eastward.

Distribution. The Indian Turnip occurs in moist woods from Kansas and Minnesota to Nova Scotia and Florida.

Poisonous properties. The corm of Indian Turnip is so extremely acrid that a decoction made from it has been used to kill insects.

The family *Lemnaceae* is allied to the *Araceae*. It contains the Duckweeds, (*Lemna*).

FARINOSAE

Herbs with endogenous stems and mostly narrow leaves, flowers usually complete, parts usually in 3's or 6's, corolla regu-

Fig. 150. Indian Turnip (*Arisaema triphyllum*). A common plant of our woods. Corm contains an acrid poison.

lar or nearly so, ovary compound, superior, endosperm of the seed mealy. This series contains the *Xyridaceae* of which the yellow-eyed grass is an example, the *Eriocaulaceae*, of which the pipewort (*Eriocaulon septangulare*) at the Atlantic seashore is a good illustration, the pine-apple family (*Bromeliaceae*, of 350 species, in tropical and warmer regions, represented in the south by the Spanish moss (*Tillandsia usneoides*) which hangs in long festoons

from trees, and by the pine-apple (*Ananas sativus*), a well known fruit now cultivated extensively in Florida, from which has been isolated the enzyme, *bromelin*, a powerful ferment capable of rapidly digesting vegetable and animal albumen. It acts in the presence of either acid or alkaline carbonates and is related to *trypsin* and *pepsin*. In the same family is the pinguin (*Bromelia Pinguin*) or wild pine-apple, the slightly acid pulp of which is edible and the fiber valuable. The plant is armed with stout spines which made the passage of troops difficult in the late Spanish war.

In the same order are the Spiderworts belonging to the family *Commelinaceae*. The common blue spiderwort (*Tradescantia virginiana*) of sandy and



Fig. 161. Common Rush (*Juncus tenuis*). A weed with tough stems, along beaten paths and roadsides. (Charlotte M. King.)

gravelly soils, has mucilaginous stems, blue ephemeral flowers, and is common everywhere in eastern North America. Several species of *Tradescantia*, like the wandering Jew (*T. Zebryna*), are commonly cultivated. Another family of the order is the *Pontederiaceae*, containing the pickerel weed (*Pontederia cordata*) and the *Eichornia speciosa*, which is frequently cultivated in green-houses and has become a very troublesome weed in the rivers of Florida and elsewhere in warm countries.

LILIIFLORAE

Herbs or occasionally shrubs with endogenous stems and monocotyledonous seeds, perianth generally well developed, flowers generally regular and complete, their parts in 3's and 6's, ovary superior, or inferior compound; endosperm horny or fleshy. This series contains the family *Juncaceae*, called

rushes, some of which, like wire-grass (*Juncus tenuis*), are troublesome weeds. *Luzula* is common at high altitudes and in northern states.

The *Dioscoreaceae*, or Yam Family, contains but few species in the United States. To this belong the wild yam root (*Dioscorea villosa*) of our woods, the Japanese yam (*D. esuricata*) and the air potato (*D. bulbifera*) of Asia, sometimes cultivated in the Gulf States for its large tubers. Yam starch is obtained from several species, the most important of which are *D. alata*, *D. zativa*, *D. japonica*, and *D. esuricata*.

The family *Taccaceae* contains *Tacca pinnatifida*, the roots of which are the source of the Tacca starch of Tahiti

and the neighboring islands. The plant is grown also in Brazil and India.

FAMILIES OF LILIIFLORAE

Ovary mostly superior

Perianth segments distinct or partly united, the inner petal-like
fruit a capsule or berry.....

Liliaceae




Fig. 160. Flowers, fruit and leaves of Yam (*Dioscorea villosa*). A common plant in thickets.



Fig. 6. Wake Robin (*Trillium at-rolet*) Canada Lily (*Lilium Canadense*) The *Trilliums* are considered poisonous. (C. H. King).



А. Яковлев
сфотографирован в 1905 г.



Судовая. «Л. II. Парм»

Ovary inferior, at least in part

Stamens 3, opposite the inner segments. Hamodoraceae

Stamens 6.

Erect perennial herbs, flowers perfect Amaryllidaceae

Stamens 3, opposite the outer segments Iridaceae

I. LILIACEAE Lily Family

Herbs or rarely woody plants with regular, symmetrical flowers, perianth not glumaceous, 3 sepals, 3 petals, 6 stamens, ovary 3-celled, fruit a pod or berry, embryo enclosed in the hard albumen. A family of about 1,600 species, including among others, several ornamental plants like the lily, lily of the valley, and yucca, some medicinal plants like squill, aloe, and false hellebore, and several poisonous plants like death camas and colchicum, the latter native to Europe and Africa. The fatal poisonous nature of *Colchicum* was familiar to the ancients, it being known to contain several poisons, such as the alkaloid *colchicin* $C_{27}H_{22}NO_6$, an amorphous, yellowish white gum, chiefly an alkaline bitter substance, which, on boiling with acids, yields *colchicem* $C_7H_{11}NO_6$ and a yellowish green resin.

Animals that eat the plant suffer with acute gastro-enteritis, coma, staggering, weak pulse, and increased urination. The family also includes several economic plants like the onion (*Allium Porrum*), garlic (*Allium sativum*), chives (*A. Schoenoprasum*), shallot (*A. acrotonum*), hyacinth (*Hyacinthus orientalis*); New Zealand flax (*Phormium tenax*), native to New Zealand where it occupies much of the country, and is now used in large quantities for making ropes and mats, (*Yucca filamentosa*) and (*Y. angustifolia*), the former a well known plant of the South and the latter a well known plant of the West, both species frequently cultivated for ornamental purposes, a large number of other species of the genus *Yucca* being also found in the Southwest. The day lily (*Funkia subcordata*), several species of the tulip (*Tulipa*), and several species of *Lilium* are cultivated. Perhaps the most common in the old gardens is the tiger lily, (*L. tigrinum*). Several species of the aloes are common in cultivation in greenhouses. They are also medicinal, containing the substance *aloinum*, a neutral principle, which yields *barbatorin* $C_{17}H_{20}O$. Aloes are cathartic. The California or Mariposa lily belongs to the genus *Calochortus*. The asparagus (*Asparagus officinalis*) is cultivated and is a well known vegetable. The cultivated asparagus (*Asparagus medeoloides*) is a native to the Cape of Good Hope. The dragon-tree (*Cordylus terminalis*) is frequently cultivated. Some of the species of the latter like "Ti" of the Sandwich Islands are of economic importance. The roots of "Ti" contain a saccharine matter, from which the natives extract sugar. They also bake the roots and eat them. The remarkable dragon-tree of the Canaries is noted for its large circumference and comparatively low height. The Botany Bay resin (*Xanthorrhoea hastata*) is chiefly used as a shellac for making colored varnishes.

Yucca leaves contain salicylic acid. Several investigators have reported saponin in the roots of *Yucca filamentosa*, *Y. angustifolia*, and *Y. imperialis* contain the alkaloid *imperialin* $C_{24}H_{40}NO_4$. In the former, Dr. Helen Abbot Michael reports the presence of several resins, the amount varying from 8-10 per cent in the root. She regarded the saponin as a constructive glucoside which served to unite what are known as the Saponin groups. Saponin occurs in many different plants, especially in the *Sapotaceae*.



Fig. 162. *Colchicum* (*Colchicum autumnale*). Flowering plant and longitudinal section of stem and bulb. Faguet.



Fig. 163. Carrion plant or Death Carnation (*Zygadenus venenosus*). A western plant, extremely poisonous to sheep.

The fly poison (*Amianthus muscaetoxicum*) is a smooth plant with simple stems from base, broadly linear leaves, white flowers in simple racemes widely spreading perianth without claws or glands. Occurs from Long Island to Florida to Arkansas. It is a well known fly poison of the south. It is related to *Veratrum* and *Melanthrum*.

The *Bulbine bulbosa*, of Australia, is poisonous to cattle, sheep and horses, which, after eating it, display such symptoms as lying down, rolling continually, having scours and a mucous discharge from the nose. The tuberous herb, *Gloriosa superba*, of India, according to Major Kirtikar, is a violent emetic.

the roots of this, when eaten, produce death in four hours. It is said to contain the bitter principle *superbin* ($C_{22}H_{40}N_2O_{17}$), perhaps identical with *scillatoxin*. The leaves and roots of *Paris quadrifolia* of Europe have a bitter taste. The berries are said to poison chickens and to produce gastro-enteritis in man. The *Alos succotrina* contains from 4-10 per cent of a bitter principle *alosin*, also some *emodin*. Representatives of the genus *Scilla* and *Urginea* yield *scillipicrin*, *scillin*, and *scillatoxin*, the latter of which resembles *digitoxin*, the first of these acts upon the heart, where used as an emetic, it has proved fatal because of its irritant action on the intestines. The seeds of *Sabadilla officinalis* are used as a parasiticide. They contain *cevadil* $C_{61}H_{48}NO_3$, *cevadilin* $C_{64}H_{52}NO_3$, and are the principal source of *veratrin* $C_{32}H_{46}NO_2$, and the glucoside *scillium* or *scillitin*. According to Friedberger and Fröhner animals poisoned with "rat poison" (squill) had cerebral convulsions and erysipelas. The rhizome of Solomon's Seal (*Polygonatum giganteum*) has an acrid bitter taste.

Chlekerinchee (*Ornithogalum thyrsoides*) is reported by Dr. Lautard (1) to have been the cause of acute gastro-enteritis in horses in South Africa. The species *O. Murrai* may possibly be poisonous since it is allied to the above which an African veterinarian reports to be poisonous. The Star-of-Bethlehem (*O. umbellatum*) which is a pretty cultivated garden plant in the northern United States has become an escape in Kentucky and is regarded as a rather troublesome weed. The Tulip and *Prisallaria* are also poisonous.

GENERA OF LILIACEAE

Perianth bell-shaped, gamophyllous.

Fruit a berry..... 5. *Convallaria*

Perianth cleft or divided.

Fruit a berry..... 6. *Trillium*

Fruit a capsule.

With scarious bracts..... 4. *Allium*

Without scarious bracts

Roots bulbous..... 1. *Zygadenus*

Roots not bulbous.

Sepals with claws, free from the ovary..... 2. *Melanthium*

Sepals without claws..... 3. *Veratrum*

1. *Zygadenus*. Michx. Camas

Smooth, erect, perennial herbs from bulbs or rootstocks, leaves linear greenish or white flowers in panicles, stamens free from perianth segments capsule 3-lobed and 3-celled. A small genus of about 8 species, native to North America and Mexico.

Zygadenus venenosus Wats. Death Camas

A pale green, slender perennial, 6 inches to 1½ feet high, from small coated bulb, leaves rough, somewhat shorter than the stem, flowers borne in a raceme, yellowish or yellow, polygamous segments of the perianth ovate or elliptical, free from ovary, bearing a roundish gland, capsule much larger than the perianth.

(1) Am. Vet. Rev. 30:298.

Distribution. From South Dakota to Nebraska, Colorado, Utah, California, Montana and British Columbia.

Zygadenus chloranthus. Pursh. Smooth *Zygadenus*

A glaucous perennial 1-3 feet high, coming from an elongated bulb, leaves flat, flowers borne in racemes, few flowered, greenish; segments of the perianth oval or obovate, muted below and adnate to the base of the ovary, capsule longer than the perianth.

Distribution. Common especially northward in Iowa and Minnesota to Alaska, in the Rocky Mountains to New Mexico and east to Missouri. It may also be poisonous.

Poisonous properties. Mention of the poisonous nature of the various species of *Zygadenus* has frequently been made, especially by the early explorers, the poisonous bulbs encountered by them being referred to as poison camas or poison sago, so called to distinguish them from the edible *Quamash*, which is commonly called *kamas*. These species bear essentially the same name today, except that in some places they are also called *Lobelia*. The bulbs are apparently much more poisonous than the leaves, but if the ground is very dry sheep are less likely to pull them up than when the ground is moist. After rains, however or early in the spring it is possible that some of the bulbs may be pulled up and thus eaten by sheep. In Montana, according to Chesnut and Wilcox, large numbers of sheep are killed by eating death camas. These authors state that in one band two thousand were poisoned and one hundred of these died. In another band two hundred were poisoned and ninety died.

Prof. Hilman reports that the wild sago (*Z. paniculatus*) is probably responsible for the death of a considerable number of cattle in certain alkaline districts in Nevada. Dr. S. B. Nelson, in experimenting with this species had wholly negative results. He fed one pound of the plant in blossom and fruit to sheep. Dr. Wilcox and Prof. Chesnut made tests on rabbits and sheep with extracts and fresh plants, and in every instance obtained positive evidence of poisoning. In these instances the plants were not in flower. Prof. Chesnut says stock is poisoned while pasturing by eating the bulbs along with the leaves or the leaves alone, or by the seeds when present in hay, as they sometimes are. Stock, especially sheep, are usually killed by eating the plant before it has blossomed in the spring. Cases of poisoning are so common in Oregon and Nevada that the term "lobeliaed" has been used to indicate the result from this kind of poisoning.

According to Chesnut and Wilcox the symptoms of poisoning are remarkably uniform.

The first signs of poisoning are a certain uneasiness and irregularity in the movements of the sheep. These irregularities rapidly become more and more pronounced, accompanied by incoordination of the muscular movements, spasms and rapid breathing. Although sheep are highly excited under the influence of *Zygadenus* poisoning, the cerebral symptoms seldom constitute a condition of frenzy. It was readily observed that until a few minutes before death ewes were able to recognize their lambs, and indicate in other ways that they were not in any sense crazed. The later symptoms were those of complete motor paralysis, combined with an exceedingly rapid and sharp breathing and a frequent weak pulse. The duration of these different stages of poisoning varies to a considerable extent, and depends entirely upon the amount of death Camas which the sheep had eaten.



Postmortem examinations made show that in every instance the lungs were congested with blood, being a hepatized condition. There were no lesions in the membranes of the brain. In cases of adult sheep the effect upon the digestive organs was not marked. There were usually to be noticed an increased salivation and continued regurgitation through the mouth and nostrils. "Symptoms produced experimentally by feeding the death camas to sheep were the same as those characterizing natural poisoning by this plant.

The toxic substance has not been isolated. Chesnut and Wilcox observed that the ground material macerated with luke warm distilled water produced a substance that had a decided soapy feeling, and that the pure juice was distinctly irritating when left on the hands for several minutes. The physiological action of the *Veratrum* is somewhat similarly caused by the active poisonous principle in camas. It is probable that many of the *Melanthaceae* have similar properties. Dr. Wilcox recommends, in case of poisoning by death camas, the hypodermic injection of strychnin in 1/20, 1/10 and 1/5 grain doses, the hypodermic injection of atropin in 1/60 and 1/30 grain doses, and solutions of potassium permanganate and aluminum sulphate. From 5 to 10 grains of each of these compounds are dissolved in water and given as a drink to adult sheep. Hogs take the same doses as sheep, horses from 15 to 20 grains, and cattle from 30 to 50 grains. Occasionally the material is injected directly into the stomach, but ordinarily the more convenient method is to allow the animals to drink it. The substances *veratrin*, *sabadin* and *sabadinin* have been obtained from *Z. venenosus*.

2. *Melanthium*. L. Bunchflower

Perennial tall leafy herbs with a thick rootstock. leaves linear to oblanceolate, flowers on large panicles, monoecious or polygamous, greenish yellow; perianth of 6 widely spreading segments raised on slender claws free from the ovary, stamens shorter than the perianth; pistil with 3 styles; capsule 3-lobed and 3-celled. A small genus of 4 species, in eastern North America.

Melanthium virginicum. L. Common Bunchflower

Tall leafy stemmed plants 3-5 feet high, leaves linear, the lower sheathing, the upper similar and sessile, flowers in an ample panicle, fragrant, perianth of flat segments, greenish yellow, styles persistent, capsule 3-celled, 8-10 seeds in each cavity.

Distribution. In low meadows and prairies from New England to Iowa river basin to Minnesota to Texas and Florida.

Poisonous properties. Several correspondents in Iowa have attributed poisoning of horses to this plant. Several related plants of the *Melanthaceae* like *Zygadenus* and *Veratrum* are known to be poisonous.

Mr J. R. Campbell, of Blockton, Iowa, writes us the following

The specimens I sent you, and which you have identified as *Melanthium virginicum*, have been the reputed cause of a number of cases of poisoning here this summer. The veterinarian here pronounced it acetone poisoning as the symptoms are similar but he decided this weed caused it as it has been found present in every case. In the first cases that he noticed here nearly all the horses in the livery barn were attacked after partaking of hay which contained an abundance of the matured seed pods of this plant. None of the animals died. The liveryman then had his men pick out all the weed, and he has not been troubled since. Several cases have occurred at different places since then, all traceable to this weed.




Fig. 164. Bunch Flower (*Melantherum virginicum*). Common in low meadows Eastern Iowa and southward. Often mixed with hay and causes poisoning of horses. (Charlotta M. King.)

At the place where I obtained these specimens the owner said he had cut the meadow and fed hay off it for fifteen years and never had any trouble until this year. Hay cut last fall seems to contain the poison, seed heads were fully mature, meadow is now and wet.

The following are the symptoms as described by the veterinarian. Heart fast and very weak, respiration shallow and labored, great muscular weakness, retching, considerable stobbering, some sweating; temperature was normal. The effect lasted three or four hours, and the animal was stupid and lacked appetite for one or two days afterwards.

The disease stopped when a ration of hay containing none of this weed was fed. Since writing the above, Dr. Blanche, a veterinarian in Belle Plaine, this state, found that horses fed with hay containing this plant "became ill and acted as if they were crazy. The symptoms were much like those from aconite poisoning." These bunchflowers have long been used to poison flies, and Hyams, of North Carolina, says that they are poisonous to crows. The *M. latifolium* and *M. parviflorum* have similar properties. According to Chesnut the Indians of Mendocino County, California, use the soaproot or Yuki (*Chlorogalum pomeridianum*) to stupefy fish. This plant is closely related to *Melantherum*.



3. *Veratrum* (Tourn.) L.

Perennial herbs, leaves broad, clasping, veined, and plaited, flowers in large panicles, greenish, polygamous or monoecious, perianth in 6 parts, spreading, greenish or brownish, without glands or nearly so, and not clawed; stamens short and free, ovary with 3 persistent styles, capsule 3-lobed, 3-celled and several-seeded. A small genus of 10 or 11 species distributed in north temperate regions. One species used in medicine all are poisonous.

Veratrum viride, Ait. American White Hellebore

A stout, leafy perennial from 2-7 feet high, with fleshy root 1-3 inches long. Flowers in ample, dense, spike-like racemes, it blooms from May to July.

Distribution. Common in swamps and wet woods, especially in eastern North America, west to Wisconsin, south to the mountains of Georgia, and north to Alaska.

Veratrum californicum, Durand. California Hellebore

A stout perennial from 2-8 feet high, fleshy root, flowers in a large loose terminal panicle, perianth segments whitish with long and narrow floral leaves.

Distribution. Common in the mountains of California and the Rocky Mountains as far north as British Columbia, south to New Mexico.

Poisonous Properties. Prof. Chesnut says

Cases arise mainly from overdoes in medicine, but instances of accidental poisoning are reported for man and for various animals and birds. In one case all the members of a household were poisoned by eating the young leaves, which were mistaken for those of marsh marigold (*Calix palustris*) and prepared for food. Animals do not reject the plant which is acrid and burning in the fresh condition, but young animals sometimes eat it with fatal results. The roots are not often mistaken for those of edible plants, but being fleshy and especially rich in alkaloids, they are somewhat dangerous. The seeds have been eaten by chickens with fatal results. The general effect is very much like that of aconite (*Aconitum Napellus*) being directed chiefly against the action of the heart and spinal cord, both of which tends to paralyze.

The symptoms of the poison are burning in the throat with increased salivation, producing a weak pulse, labored respiration and profound prostration. The root was used by the Indians in making snuff. Dr. Halsted attributed deaths of human beings as well as of cattle in New Jersey to this plant.

The number of poisonous substances found in hellebore is quite large. Of these so-called *veratrin*, $C_{32}H_{48}NO_{11}$, of earlier writers, has an alkaline reaction, and a burning taste, it produces violent sneezing and dilates the pupil. However, later investigators have separated this into the following bases: the very toxic *cevadon* $C_{28}H_{48}O_9$, *veratridin*, $C_{37}H_{53}NO_{11}$ and *sabadilin* $C_{34}H_{58}NO_8$.

Veratrum album L., *V. lobelianum* Bernh., *V. viride*, Ait., also contain in addition to the bases named above, two other bases, *sabadin*, $C_{29}H_{51}NO_8$, and *sabadinin*, $C_{27}H_{45}NO_8$, and also the following substances: *servin* $C_{28}H_{47}NO_8$, a pure alkaloid *rubijervin* $C_{26}H_{42}NO_7$, *pseudojervin* $C_{29}H_{43}NO_7$, *protoveratrin*

$C_{22}H_{51}NO_{11}$, *protoveratridin* $C_{22}H_{43}NO_9$, and the latter glucoside *veratamarin*. *Verum* is a powerful depressant to the heart muscles and vaso motor centers; large doses therefore weaken the pulse. It depresses respiration and death occurs from asphyxia. Dr Winslow, in speaking of the toxicology, says

The symptoms exhibited in *Veratrum viride* poisoning are salivation, vomiting, or attempts at vomiting, purging, abdominal pain, muscular weakness, difficulty in progression, loss of power and general paralysis, muscular tremors and spasms, and, occasionally, convulsions. The pulse is unaltered in rate at first, but later becomes infrequent and compressible and finally rapid, threadlike and running. The respiration is shallow, the temperature is reduced, the skin is cold and clammy, there is semi-consciousness, loss of sight and death from asphyxia. Treatment should be pursued with cardiac and respiratory stimulants, as amy, nitrite (by inhalation), alcohol, strychnin and atropin (anale acid as a chemical antidote, opium to subdue pain, and demulcents to relieve local irritation of the digestive tract. Warm water should be given the smaller animals to wash out the stomach and to hasten vomition, and quietude should be enforced. In man, fatal poisoning is rare, since the drug is spontaneously vomited. The same would probably apply to dogs. Recovery has ensued in horses after injection of two ounces of *veratrum album* root.



Fig. 164a American White Hellebore (*Veratrum viride*). A poisonous plant of Eastern North America.

4. *Allium* (Tourn.) L. Onion, Garlic and Leek

Perennial bulbous plants, bulb solitary or clustered, leaves generally linear, a few lanceolate or oblong, stem simple or erect, flowers in umbels subtended by bracts perianth white, purple or pink, the parts distinct or united at the base often becoming dry, the 6 filaments awl shaped, ovary 3-celled or incompletely so; capsule with 1-2 black seeds in each cell. About 275 species of wide distribution contains a number of important economic plants, among them garlic (*A sativum*), garden leek (*A Porrum*), chives (*A Schoenoprasum*), shallot (*A ascalonicum*), onion (*A Cepa*), and the golden garlic (*A Moly*), cultivated for ornamental purposes.

Allium vineale, L. Field Garlic

A slender scape, naked from an ovoid membranaceous bulb, 1-3 feet high; terete and hollow leaves, channeled above, frequently densely bulbiferous, flowers greenish or purple.

Distribution. Common in meadows and wheat fields of Virginia, and eastern states from Connecticut to Virginia and Missouri. Naturalized from Europe.

Allium triococum Ait. Wild Leek

Scape naked, 4-12 inches high from an ovoid bulb, leaves fibrous articulated, leaves oblong, lanceolate or elliptical, few, appearing long before the flowers in spring, flowers in umbels numerous, greenish white, one ovule in each cavity, capsules strongly 3-lobed, seeds black and smooth.

Distribution. Common in the woods from western New England to Minnesota and Eastern and northern Iowa, especially in the low damp grounds.

Allium canadense L. Wild Garlic

Scape 1-2 feet high, coming from an ovoid bulb, the outer coats fibrous reticulated, leaves narrow linear; flowers in an umbel frequently with small bulbs; flowers pink or white.

Distribution. Common in meadows or low grounds in New England to Minnesota and Iowa, south to the Gulf.

Injurious Properties In parts of the country where these onions grow there is frequent complaint of milk taking the flavor of onions where cattle feed upon them. Chesnut and Wilcox do not mention any species of the genus *Allium*, except some of the species found in Montana, which may impart to milk a disagreeable flavor. Friedberger and Frohner state that onions produce slavering.

Prof. A. Lautard* has prepared an abstract of a report by Dr W W Goldsmith in the Journal of Comparative Pathology and Therapeutics upon onion poisoning in cattle. Briefly it is as follows

Loads of onions partly started to shoot and partly decayed, were unloaded in a meadow where nine head of cattle were grazing. After a week the cattle seemed sick

* Amer Vet. Review. 26:63

and one died, displaying the following symptoms: Increase onion odor tucking up of flanks, constipation in some, purging freely in others, one vomited abundantly another very ill, grunted, was much constipated, staggered in walking, was very tender in loins, temperature 103°, urine dark and smelling of onions. Treatment: Feeding with soft food and hay. Large doses of linseed oil. One animal that was very ill got also extract of belladonna and carbonate of soda. All but one of the animals recovered. At the autopsy of the dead one, the rumen was found inflated and also the bowels. Liver enlarged and of light color. Kidneys dark green and with offensive odor. Rumen contained large quantity of onions and grass. The whole carcass and organs smell of onions.

5. *Convallaria* L. Lily of the Valley

A low smooth herb with horizontal root-stocks, flowers white in a one sided raceme, stamen 6, ovary 3-celled, berry globose. A genus with one species.

Convallaria majalis, L. Lily of the Valley

A smooth perennial herb with horizontal root stocks and 2 or sometimes 3 oblong leaves; flowers in racemes, perianth bell shaped, white, 6-lobed, stamens 6, inserted on the base of the perianth, ovary 3-celled, 4-6 ovules in each cell,



Fig. 165. Lily-of-the-Valley (*Convallaria majalis*). A well known cultivated plant possessing poisonous properties similar to those of Foxglove U. S. Disp. Agr.




Fig. 166. False Agave (*Agave virgineus*). A plant of the Southern United States.

berry roundish, red and few-seeded. The species is native to Europe, Asia, and the Aleghanes, and is widely cultivated as an ornamental plant.

Poisonous properties. The plant contains two glucosides, one, *convallamarin* $C_{28}H_{44}O_{12}$, an extremely poisonous crystalline compound with a bitter sweet taste, with a physiological action on the heart like digitalin, a substance found in the common foxglove, and *convallarin* $C_{24}H_{40}O_{11}$, which is crystalline, has a sharp taste and is purgative in its action. Chesnut says "The emegative and purgative actions of the lily of the valley are quite marked." The action of the heart is infrequent and irregular, and finally death occurs from paralysis.

Trillium L. Birthwort

Herbs, with naked stem from a short, horizontal root stock, netted veined, simply whorled leaves, in 1 or 2 whorls, colored flowers, 3 green persistent sepals, 3 colored petals which wither with fruit, stamens 6, hypogynous linear adnate anthers on short filaments, sessile stigmas 3, ovary 3-celled, fruit a berry.

The principal species of the United States are the wake-robin (*T. sessile*) which flowers very early in the spring, is from 2-4 inches high and is common northward and eastward, the sessile-flowered wake-robin (*T. sessile*) which bears sessile dull purple flowers with narrow sepals and petals, and leaves that are often blotched and occurs from eastern Iowa southward, prairie wake-robin (*T. recurvatum*) of the west, which has dull purple petals but differs from the preceding in having narrow leaves, large white-flowered wake-robin (*T. grandiflorum*) which bears a large white flower raised on a peduncle later recurving from the erect, the flowers becoming purplish, and rounded, ovate, sessile leaves, and birthwort (*T. erectum*) much like *T. grandiflorum* except that the flowers are not so large and are unpleasantly scented. Both of the two last named are found in the eastern and central states.

Poisonous properties. *Trilliums* have long been considered poisonous. All species are emetic. Lindley states that the roots have a violent emetic action. The fruit should be regarded with suspicion.

HAEMODORACEAE Bloodwort Family

Perennia, herbs with fibrous roots, leaves, narrow, lanceolate and somewhat erect small perfect flowers which are woolly or scurfy on the outside; flowers in panicles, perianth 6-parted or 6-lobed adnate to the ovary, stamens 3, opposite the 3 inner segments of the perianth, stigmas 3, fruit a 3-valved capsule, seeds few or numerous. A small family of 9 genera and 35 species mostly native to Africa, Australia and tropic America.

Lachnanthes L. Red-Root

A stout herb with short rootstock red, fibrous, perennal root, leaves, equitant and sword shaped, crossed at the base and scattered on the stem, flowers, numerous, borne in a woolly, cymose panicle perianth, 6-parted, the outside segments smaller than the inner, stamens, 3, opposite the 3 inner divisions, pistil with 3-celled ovary few ovules in each cavity, seeds few flattened nearly orbicular, fixed by the middle. A species of a single genus native to southeastern North America and western India. (*Cyrtanthes*)

Lochneranthus fructuosus, (Walt.) Ellis.

A stout, tall herb with numerous yellow flowers, 6-parted perianth and few seeds.

Distribution. In salty swamps near the coast in southeastern Massachusetts, Rhode Island and New Jersey to Florida. This plant is commonly called the pink-root of the Atlantic coast.

Poisonous properties. Prof Chesnut says that throughout the South, white hogs are supposed to be particularly subject to the poison contained in this plant. Dr Halsted says "Throughout the southern states, this plant abounds and the preponderance of black over white-skinned hogs is claimed to be due to this paint-root. White hogs with free access to the plant are soon killed off, while black ones are not."

This is not the only case of the color of animals seeming to have an influence upon their distribution. Thus, white horses in Prussia, it is claimed, are injured by eating milkweed, while dark horses are not. In Sicily, there are black sheep only, as white ones are killed off by a species of St John's wort (*Hypericum*). "While the claim of the immunity of black pigs from the effects of paint-root seems to be a common belief, further investigation should be made before this should be asserted definitely as a fact."

Family *Amaryllidaceae* Amaryllis Family

Mostly perennial herbs with bulbs, rootstocks or corms. Scapose flowers regular or nearly so, perianth 6-parted or 6-lobed, the lobes or segments distinct, united below into a tube, adnate to the ovary, stamens 6, style single, capsules several, many seeded. About 800 species, chiefly native of tropical or warm regions, a few in temperate regions. Some well known representatives are daffodil, (*Narcissus Pseudo-Narcissus*), Polyanthus, (*A. Tazetta*), poet's Narcissus, (*N. poeticus*) producing intense gastro-enteritis. Jonquil (*N. Jonquilla*), snowdrop, (*Galenthus nivalis*); amaryllis, (*Amaryllis Belladonna*), tuberose, (*Polygonium tuberosum*), the latter widely cultivated, the American aloe or agave, the most common species in cultivation being the century plant (*Agave americana*) native to Mexico and Central America, the Mexican drink, pulque, being made from the sweet liquid obtained from this plant at the time of flowering. Several species are used for the manufacture of fibre, the best known being the sisal, (*Agave rigida*). The mauritius hemp, (*Furcraea gigantea*), is native to Mexico and has been introduced into Zanzibar. Many members of the family have acrid properties and some of them are poisonous. *Buphane disticha* is used by the Hottentots to poison their arrows. Poet's narcissus contains *pseudo-narcissin*. *Amaryllis Belladonna* contains *belladonna* and *Sprekelia formosissima* contains *amaryllin*, a belladonna-like alkaloid. The *Lycorn* species contain *lycorn*, an alkaloid with the formula $C_{32}H_{37}N_2O_4$, and a second alkaloid *lycanin*, $C_{34}H_{39}N_2O_2$. *Agave heteracantha* contains *agavesaponin*. Dr MacDougal states that the sharp pointed leaves of *Agave Schottii* often penetrate leggins and leather shoes inflicting painful injuries.

Zephyranthes. Herb

Smooth herb with coated bulbs, narrow leaves, flowers scapose, large erect, pink, white or purple, perianth funnel-form from a tubular base. The 6 divided petals are united below into a tube subtended by an entire or 2-cleft



Fig. 167 Saffron (*Crocus sativus*) The flowers furnish the saffron of commerce. (Faguet)

bract, ovary 3-celled, style long, filiform, 2-cleft at the summit, ovules numerous, capsules membranaceous, seeds flattened, blackish, small genus of 30 species, native to America.

Zephyranthes Atamasco (L.) Herb. Atamasco Lily

Leaves bright green and shiny from an ovoid bulb, scapes erect, bracts 2-cleft, perianth white, pinkish or light purple; segments shorter than the two stamens.

Distribution. In most places from eastern Virginia to Florida and Alabama.

Poisonous properties Prof. Chesnut, in speaking of this plant, says that the Atamasco of the southeastern United States is supposed by some persons to cause the disease in horses known as "staggers."



Fig. 168. Atamasco Lily (*Zephyranthes atamasco*). A plant of the southeastern United States, supposed to cause "staggers" in horses.



Fig. 169. Blue Flag (*Iris versicolor*). The rootstock is poisonous. The plant grows in low grounds. (After Johnson.)

Family IRIDACEAE. Iris Family

Perennial herbs, frequently with bulbs, corms or tubers. Leaves equitant, erect, 2-ranked. Perianth of 6 segments or 6-lobed, its tube adnate to the ovary; stamens 3, adnate to the ovary; anthers facing outward. Ovary inferior, mostly 3-celled, style 1 or 3-cleft, stigmas 3, opposite the three stamens, ovules generally numerous in each cell, embryo small, endosperm, fleshy, or horny.

About 1000 species, of wide distribution. Common native plants of the family are the blue flag (*Iris versicolor*), growing in low grounds of the North,

the Carolina blue flag (*I. carolina*) of the South, blue eyed grass (*Sisyrinchium angustifolium*). Many species of the exotic blue flags, Iris like the dwarf garden iris (*I. pumila*) and the common flower-de-luce (*I. germanica*), the common crocus or saffron (*Crocus vernus*) used for coloring, freesia (*Freesia refracta*), tritonia and gladiolus are cultivated for ornamental purposes. The orris root (*Iris florentina*, *I. pallida* and *I. germanica*) is an article of commerce used for perfume and tooth powders. It contains myristic acid. A substitute for saffron is obtained from the flowers of the South American saffron (*Crocasmia aurea*)



Fig. 169a. The petaloid bilobed stigma and stamen of Iris. (Kerner).

Iris (Tourn) L.

Herbs with creeping or horizontal root stocks, and erect stems with equitant leaves, flowers large, regular, panicled, perianth of 6 segments united below into a tube, the outer dilated, spreading or reflexed, the 3 inner, smaller, stamens inserted at the base of the outer perianth; ovary 3-celled fruit a capsule; seeds numerous. About 100 species in the North Temperate regions. The *Iris florentina* contains the glucoside *irigenin* $C_{15}H_{14}O_8$. This is derived from *irisin*.

Iris versicolor, L.

Root stock fleshy, stem roundish leaves erect, leaves shorter than the stem, flowers bluish, perianth deeply 6-parted, the 3 outer divisions reflexed, the 3 inner smaller, erect stamens distinct, covered by the petaloid stigmas.

Distribution. In marshes, thickets, and wet meadows from Newfoundland to Manitoba, south to Florida and Arkansas.

Poisonous properties and uses. The root contains the substance *irisin*, or *iridin*. The acrid resinous substance, *irisin*, acts powerfully upon the gastro-intestinal tract, liver and pancreas, causing a burning sensation and congestion. That the root is poisonous may be seen from the following statement made by Dr Rusby

Another rhizome whose acrid taste is likely to prevent ingestion in poisonous quantity, is that of the common *Iris versicolor* L. Still, because this is commonly known as the blue flag, there is some danger that it might be eaten in mistake for calamus, which is commonly known as sweet-flag. If so, it would prove seriously, if not fatally poisonous

as its well-known emetic-cathartic properties, even when used by drying and keeping, are powerful, and in a fresh state would be decidedly violent.

Dr Johnson says

Iris, in full doses, is an active emeto-cathartic, operating with violence, and producing considerable prostration. Its effects upon the liver appear to be analogous to those of podophyllum. In sick headache dependent upon indigestion, small doses, frequently repeated, often act most happily. It has been largely used by eclectic practitioners, and is highly esteemed by them as a hydragogue cathartic, an alterative, emagogue, vermifuge, and diuretic.

One case of poisoning has been recorded in this state. Other species of *Iris* of which we have quite a number in the U. S. must be looked upon with suspicion. Mention may be made here of the *Iris missouriensis* and *I. carolina*. The root stocks of our cultivated species like *I. pumila* and *I. sibirica* must be looked upon with suspicion. The South African *Homeria colina* naturalized in Australia, according to Maiden, is poisonous to cattle browsing on the plant.

Large herbs with endogenous stems and monocotyledonous seeds, flowers very irregular, ovary inferior, composed of several united carpels, seeds



Fig. 170. Ginger (*Zingiber officinale*)
a. Entire plant. b. flower. (Charlotte M. King, after Strasburger, Schenck, Noll and Schimper.)

with endosperm. This order contains the important family *Musaceae* in which is found the banana (*Musa sapientum*), well known as an article of commerce. It is extensively cultivated in the tropics and one of the most important food plants in all warm countries. The fruit is eaten fresh when ripe, a kind of flour is also made from it. *M. textilis* is an important fiber plant being the source of Manila hemp, large quantities of which are imported from the Philippines. The ravenala or traveler's palm also belongs to this family. It has an oily, edible, arillus which is bright blue. The family *Cannaceae* contains the Indian shot (*Canna indica*), frequently cultivated for ornamental purposes in this country, in tropical regions, however, a starch is made from the rhizome of this species and from *C. edulis*. *C. flaccida* is a native of the southern United States and has a pretty blue flower. The family *Marantaceae* contains the West Indian arrowroot (*Maranta arundinacea*). The family *Zingiberaceae* includes ginger (*Zingiber officinale*) which contains gingerol and is used as a condiment and stimulant. The ginger of commerce is derived from the fleshy rootstock, the plants grown in Jamaica being considered most valuable. These are cultivated in regions having an altitude of 2000 feet. Malabar cardamon (*Elettaria Cardamomum*), round cardamon (*Alpinia striata*), bastard cardamon (*Amomum xanthioides*), Bengal cardamon (*A. subulatum*) and Java cardamon (*A. maximum*) also belong to this order. The *Kasmpferia rotunda* of India, is a bulbous or tuberous rooted biennial which according to Major Kirkitar, causes profuse salivation and vomiting when administered internally. The rhizome of *K. Galanga* furnishes a perfume.

Arrowroot comes from *Curcuma leucorrhiza*, and turmeric from *Curcuma longa*. The tuber of the latter, when powdered, is used as a yellow dye-



Fig. 171. *Canna* (*Canna flaccida*). A native American *Canna*.

Fig. 172. Banana Fruit (*Musa sapientum*). A well known tropical fruit. W. S. Dodgson.

stuff, in making turmeric paper, as a condiment, especially in curry powder, and as an aromatic stimulant. The zedoary, (*C. Zedoaria*), is used in Himalaya India, where it is native, in place of turmeric. Galangal is the root of *Alpinia officinarum* which grows on the Chinese coast. Another species, *A. Galanga* is used on the island of Java.

HERBACEOUS

Herbs with endogenous stems, flowers very irregular or in a few cases regular, generally complete and perfect, and parts in 3's or 6's, ovary inferior compound, seeds small, numerous, without endosperm.

Family ORCHIDACEAE. Orchid Family

Perennial herbs with corms, bulbs or tuberous roots. perfect and irregular flowers, perianth of 6 divisions in 2 sets, the 3 outer similar in texture to the 3 inner petals, one of the 3 inner, different in form and is called the lip. In front of the lip is a column composed of a single stamen, or in *Cypripedium* of two stamens, and a rudiment of the third, pollen in 2 or 8 pear shaped sacs called pollinia which are united by little threads. Stamens variously united with the thick, fleshy style into a column, ovary 1-celled with many ovules on a three parietal placentae, capsule 1-celled, 3-valved, seeds numerous. A large order of about 5000 species of wide distribution, most abundant in the tropics. Many of the plants like the *Cypripedium*, *Angrecum* and the *Cattasetum* are cultivated for ornamental purposes.

The sap of commerce is obtained from the *Orchis masculata*. The flavoring material, vanilla, is obtained from *Vanilla planifolia* native to Mexico and widely distributed by cultivation; this plant contains from 1½ to 3 percent of vanillin $C_8H_8O_3$. Other species of the genus *Vanilla* also furnish vanilla but in smaller quantities, these are *V. Pompona*, *V. guianensis*, and *V. palmarum*. Vanillin is also made from coniferin and eugenol, and occurs in other orchids as *Spiranthes* and such plants as *Spiraea Ulmaria* and *Lupinus albus*. It is used for medicine.

Orchids contain some alkaloids, for example, *Phalanopsis amabilis* contains a tonic alkaloid, according to Boorsma, which is closely related to coniferin, $C_{18}H_{32}O_8$.

Cypripedium L.

Tufted roots, perennial, glandular, pubescent herbs. leaves large, many nerved, flowers solitary or few, sepals shiny, spreading 3 distinct or 2 of them united into one, under the lip. petals spreading, resembling the sepals, lip of large inflated sac, column declined with a fertile stamen on each side a sterile petaloid stamen above, which covers the summit of the style, pollen granular, stigma broad, obscurely 3-lobed, moist and roughish. About 40 species, mostly tropical.

Cypripedium parviflorum, Salisb var *pubescens* (Willd.) Knight. Yellow Lady Slipper

Perennial, with leafy stem, 2 feet high, pubescent, leaves oval, or elliptical, acute. sepals ovate, lanceolate, usually larger than the lip, yellowish or greenish, petals narrower, usually twisted, lip flattened laterally, pale yellow with purple lines.



Fig. 173.




Fig. 174.

Fig. 173. Small Yellow Lady Slipper (*Cypripedium parviflorum* var. *pubescens*). A beautiful flower of early summer seen in the woods of eastern Iowa. (M. King.)

Fig. 174. Glands of several species of Moccasin flower, which are said to contain the toxic substance. 1. Hair gland of *Cypripedium pubescens*. 2. Hair gland of *Cypripedium hirsutum* in water. 3. Hair gland of *Cypripedium pubescens* in water. (Charlotte M. King, after Nessler.)

Distribution. In woods and thickets, chiefly east of central Iowa and Minnesota to Nova Scotia; occasionally in Colorado, Nebraska and Alabama.

Cypripedium candidum Muhl. Small White Lady Slipper

A slightly pubescent perennial, leaves lance-oblong, acute; petals and sepals greenish purple spotted, sepals ovate-lanceolate, lips white striped with purple inside, flattened laterally, convex above.

Distribution. In bogs and meadows from New York to Minnesota, Iowa, Nebraska and Missouri.

Cypripedium hirsutum Mill. Showy Lady Slipper

A rather stout downy perennial 2 feet or more high, leaves ovate pointed, sepals round ovate or orbicular, longer than the petals, which are obovate, lip inflated, white, pink purple stripes.

Distribution. In woods and swamps from Nova Scotia, Ontario and Georgia west to Minnesota and Iowa.

Poisonous Properties. Dr. Babcock, many years ago, found that the several species of Lady's Slipper produced dermatitis. Years ago the writer heard of a case of poisoning where a young man carried a large bunch of



Fig. 175. Showy Lady Slipper (*Cypripedium hirsutum*). A plant thought to cause dermatitis. C. M. King.



Fig. 176. Flower of another orchid (*Habenaria*).

Showy Lady Slipper and became poisoned very much as if it were by poison Ivy. Prof. Chesnut in referring to the poisoning from these plants says:

The poisonous character of these plants was not suspected prior to 1875, when Prof. H. H. Babcock, of Chicago, who had annually been suffering, supposedly from recurrent attacks of ivy (*Rhus*) poisoning, discovered that the affection was most probably caused not by the ivy, but by the two species of Lady's Slipper named above (*C. parviflorum*, var. *pubescens* and *C. hirsutum*) instances were afterward reported, but the facts were not positively ascertained until 1894, when an investigation was made by Prof. D. T. MacDougal of the University of Minnesota. It was discovered that these plants are provided with glandular hairs which cover the surface of the stem and leaves and contain a poisonous oil which is especially abundant at the fruiting season. Its action on the skin is very similar to that of *Toxicodendron*, the active constituent of poison Ivy (*Rhus Toxicodendron*), but its exact chemical nature could not be ascertained on account of the small quantity obtainable. Experiments with the stem and leaves upon individuals showed that over half of them were affected by the first two species, and that the last was also poisonous, but in a minor degree. No accidental cases have been recorded against it. No specific antidote has been suggested.

Dr MacDougal* made a personal experiment with a mature specimen of *C. hirsutum* on which there were newly formed seed pods. This plant was broken off near the base of the stem and the leaves brushed lightly over the arm.

A slight tingling sensation was felt at the time, and, fourteen hours later, the arm was greatly swollen from the shoulder to the finger tips. The portion covered by the plant—covering an area of 50 sq. cm.—was violently inflamed and covered with macules, accompanied by the usual symptoms of dermatitis and constitutional disturbances. By treatment of the most approved kind, the arm was reduced to its normal size in ten days, but the effects were perceptible a month later.

Nestler discovered that the secretion contained in these hairs was a fatty acid readily soluble in alcohol and benzol and producing a mildly acid reaction. He also states that his results with *C. pubescens* were negative but that with *C. spectabile* (*C. hirsutum* Mill.) he secured positive results, producing a dermatitis, the action, however, not being so pronounced as that reported by MacDougal. He also found that, as stated above, the maximum poisonous effect was during the formation of seed capsules and that the poison was in the hairs of the plant as is the case in the Primrose. Nestler did not succeed in producing dermatitis with *C. parviflorum*, *C. acule*, *C. macranthum*, *C. montanum*, or *Calceolus*. As some of these species produce an abundance of raphides in the stem, it is evident that dermatitis is not caused by these crystals, but rather by a substance found in the stem. Dr MacDougal suggests that the raphides may serve the plant as a protection from animals.

Nestler also asserts that the *Cypripedium* may contain an additional substance *myelin* which Seng has found in Ginkgo seed, and Nestler himself observed in the fruit of *Capricorn annuum*. It is not a cardol.

From the root of *Cypripedium* a substance is obtained which is sometimes administered to children as a substitute for opium. It contains a bitter glucosidal principle.

Class, DICOTYLEDONEAE

Stem usually oxogenous with pith, wood and bark (endogenous in a few plants), the woods traversed by medullary rays, leaves usually pinnately or palmately netted-veined, embryo of the seed with 2 cotyledons or occasionally 1, parts of the flower usually in 5's, rarely in 3's or 6's.

Archichlamydeae

Petals separate and distinct from each other or wanting. Includes many plants classed as *Apetalae* and *Polypetalae*. In some orders, as *Leguminosae*, the lower petals are more or less united and joined at the base.

VERTICILLATAE

Contains a single family *Casuarinaceae* of 20 species, mostly Australian, with monoecious flowers. The *Casuarina equisetifolia* of the tropical Old World furnishes a hard wood known as iron wood and in Egypt the trees are used as a shelter belt for bananas.

* Minn. Bot. Studies. 1894 32-36.

1 Das Sekret der Drüsenhaare der gattung *Cypripedium* mit besonderer Berücksichtigung seiner hautreizenden Wirkung. Nestler Ber. der Deut. Bot. Gesell. 25:554-567




Fig. 177. Black Pepper Plant (*Piper nigrum*). 1 Part of shoot with young fruit. 2. Tip of fruit spike. (After Wauson.)

PIPERALES

Herbs with exogenous stem, with neither petals nor sepals; flowers in spikes, bracteolate. Largely tropical and includes the family *Saururaceae* or Lazarus's tail, the peppers, *Piperaceae*, including black pepper (*Piper nigrum*), a well-known condiment of the tropics containing the alkaloid *piperin* $C_{17}H_{19}NO_3$ and a volatile oil $C_{10}H_{18}$, cubeba (*P. cubeba*) containing *cubebin* $C_{19}H_{27}O_3$ and the oil of cubeba, kava kava (*P. methysticum*) native to the Pacific Islands, containing *methysticin* $C_{15}H_{15}O_8$, which is used to make stimulating drinks, *P. longum* of India, *P. chaba* of India and the Philippines, the betel pepper (*P. Betle*) of the Malay Islands, the berries of which are chewed with the Betel Nut and the Matico, or the Soldier's Herb (*P. angustifolium*) of South America, the hairy leaves of which are used as a styptic. The South American *Peperomias* are well-known greenhouse plants. Other species of peppers are used in medicine. The so-called "causimon" (*P. peltatum*) according to Mr. Combs, is a powerful diuretic. "Matico de Peru" (*P. angustifolium*) is an acriid, bitter plant containing a green volatile oil.

SALICALES

Trees or shrubs with simple flowers, imperfect catkins, perianth wanting, fruit a many-seeded capsule, seeds with a tuft of hair at one end. This series contains only one family the *Salicaceae*.

Salicaceae Willow Family

Diocious trees or shrubs, alternate stipulate leaves, the stipules often minute and soon falling, staminate and pistillate flowers borne in catkins, one to each bract, without calyx or corolla, staminate flowers with 1-numerous



Fig. 78. Weeping Willow (*Salix caprea*). 1. Flower bud. 2. Staminate flower. 3. Pistillate flower. 4. Fruit. 5. Fruit. 6. Summer branch. 7. Bud and leaf scar. 1, 2, 3, 4, 5, one half natural size. M. H. Clancy.

stamens subtended by a cup-shaped disk; pistillate flowers with a 1-celled ovary, stigmas 2-4, simple or 2-4 cleft; fruit a 1-celled and 2-4 valved pod bearing numerous seeds provided with long silky hairs. There are only two genera and about 200 species, found in temperate and Arctic regions. The bark of some species of the family is used in medicine because of its astringent properties. The willow contains the glucoside *salicin* $C_{11}H_{16}O_5$. Poplar contains *populin* $C_{10}H_{18}O_5$. The Balm of Gilead (*Populus candicans*) may cause blistering, and the European *P. balsamifera* causes colic.

Myricaceae. Sweet Gale Family

Monoecious or dioecious shrubs with alternate, coriaceous, aromatic leaves, flowers in short scaly catkins, staminate flowers with 2-16 but usually 4-8 stamens, ovary with 2-8 scales and 2 linear stigmas, fruit a small 1-celled




Fig. 179. Pistillate and staminate flower of willow. a. seer gland, b. scale.




Fig. 180. Leaves of Cottonwood (*Populus deltoides*). A well known native tree growing on the borders of streams throughout the U. S. east of the Rockies. (W. S. Dudgeon.)



Fig. 180a. Sweet Gale (*Myrica*). Common in the east.



drupe, the outer part frequently covered with wax. About 35 species of wide distribution. The sweet fern (*Myrica asplenifolia*) is sometimes weedy in sandy fields in the North, it contains an oil of strong, spicy, cinnamon-like odor, bayberry wax is derived from *M. cerifera*, common along the Atlantic coast and the Gulf of Mexico. The bark of *Myrica nagi* contains *myricetin* $C_{15}H_{10}O_8$. The leaves of *M. acris* are used in the preparation of bay rum.




Fig. 101 Balm of Gilead (*Populus canadensis*). 1. Flowering branch of staminate tree. 2. Same of pistillate tree. 3. Flowering branch. 4. Scale of staminate catkin, enlarged. 5. Scale of pistillate catkin, enlarged. 6. Scale without flower displayed, enlarged. 7. Mature fruit. 8. Seed, enlarged. 9. Longitudinal section of seed, enlarged. 10. Embryo, enlarged. 11. Twig or branch, showing buds. 1, 2, 3, 11, one-half natural size. M. H. Cheney

BALANOPSIDALES

Contain a single family, the *Balanopsidaceae* of New Caledonia, which in turn contains a single genus *Balanops* of 7 species.

LEITNERIALES

Shrubs or trees with entire, petioled, simple leaves, flowers in catkins, staminate flowers subtended by what appears to be a perianth, sepals 3-4, ovary 1-celled, style slender, endosperm thin. Only one family, the *Leitneriaceae*, which consists of a single genus *Leitneria*, with perhaps 2 species. *L. floridana* occurs in swamps in southern Missouri to Texas and Florida and produces a wood lighter than cork, probably the lightest wood known.

JUGLANDALES

Trees with alternate, pinnately compound leaves, flowers monoecious, bracteolate, the staminate in long drooping catkins, pistillate solitary or several together, staminate flowers of 3 many stamens with or without a perianth, per-



Fig. 8. Black Walnut (*Juglans nigra*). 1. Flowering branch. 2. Staminate flower, enlarged. 3. Perianth of staminate flower, enlarged. 4. Pistillate flower, enlarged. 5. Longitudinal section of pistillate flower, enlarged. 6. Pistillate flower, enlarged. 7. Longitudinal section of pistillate flower, enlarged. 8. Pistillate flower, enlarged. 9. Pistillate flower, enlarged. 10. Pistillate flower, enlarged. 11. Pistillate flower, enlarged. M. M. Carey.

anth adnate to the ovary, anthers erect, 2-celled; pistillate flowers usually 2-bracteolate, calyx 4-lobed or with petals, ovary inferior, 1-celled or incompletely 2-4 celled, ovule solitary, erect, styles 2, fruit generally a drupe, dehiscent or indehiscent, the involucre regarded by some as the calyx, encloses the nut, which is incompletely 2-4 celled, endosperm none, cotyledons corrugated, only One family *Juglandaceae*. The English Walnut (*Juglans regia*), from the Mediterranean region to the Himalayas is extensively cultivated in California, Spain, France, Italy, and other warm temperate countries, butternut (*Juglans cinerea*), from New Brunswick to North Dakota and Nebraska, produces a valuable wood which is, however, inferior to the black walnut (*Juglans nigra*), distributed from Massachusetts to Minnesota, Kansas and Texas but




Fig. 183 Cork Wood (*Ledneria floridana*). The wood of this plant is extremely light. It grows in swamps in Southern Missouri and Florida. (W. S. Dudgeon.)

the timber is becoming scarce. There is a popular impression that the black walnut is poisonous to vegetation growing under the trees. California Walnut is *J. californica*. The Japanese walnut (*J. Sieboldiana*) produces a large, thick-shelled nut. The bark of *Juglans* contains juglandic acid $C_{10}H_8O_6$. The *Pterocarya caucasica* is a native to the Trans-Caucasus.

The genus *Carya* is native to North America and yields valuable timber and nuts. The nut of the pecan (*Carya illinoensis*) is an important article of commerce in Texas and other southern states. The wood is also used. The shellbark hickory (*Carya ovata*) and the Missouri hickory (*C. laciniata*) supply valuable woods which are used in the manufacture of ax handles and for parts of wagons. The nut of the latter is large but like the preceding one and the

C. tomentosus has a hard shell. The bitternut (*C. cordeformis*) and pigment (*C. glabra*) have soft shells. The bark of hickories is used to flavor and color glucose to imitate maple sugar.

FAGALES

Monoecious or rarely dioecious trees or shrubs, with simple alternate leaves stipules deciduous; calyx usually present; corolla usually wanting; staminate flowers in catkins; stamens 4-30; pistillate flowers solitary, clustered in scaly catkins, ovary more or less 2-7 celled with 1-2 pendulous, straight ovules, all the ovules but one disappearing in fruit; axis becoming a husk or cup, embryo large, endosperm none. This is an important order, including the chestnut, oak, birches and alders. The bark of a few species of *Quercus*, because of its astringent qualities is used in medicine. The European filbert (*Corylus Avellana*)




Fig. 143. American Chestnut (*Castanea dentata*). 1. Flowering branch. 2. Staminate flower enlarged. 3. Diagram of pistillate flower cluster. 4. Pistillate flower in bud. 5. Longitudinal section of pistillate flower. 6. Perianth of fruiting branch. 7. Longitudinal section of nut. 8. Involucre spore. 9. End of young branchlet. 1, 4, 7 one-half natural size. M. M. Cheney.

and our haxe, not (*C. americana*) are articles of commerce. Both the American beech (*Fagus ferruginea*), and the European (*F. sylvatica*), are prized for their nuts, which contain a valuable oil. The chestnuts are well known in commerce. The American chestnut (*Castanea dentata*) is common in the United States. The European species (*C. sativa*), has long been cultivated in North America. The Japanese (*C. japonica*) is also cultivated. In southern Europe, starch and flour are made from the nuts of the Spanish chestnut (*C. sativa*). The wood of many species of oak is used for interior finishing and furniture, among these are white oak (*Quercus alba*), red oak (*Q. rubra*), pin oak (*Q. palustris*), English oak (*Q. Robur*) and the live oak (*Q. virginiana*) of the south. The bark of several species like the scarlet oak (*Q. coccinea*), English oak, and chestnut oak (*C. Muhlenbergii*) is used for tanning. Cork is obtained from the bark of *Q. Pseudo-suber*. The galls of *Q. laurifolia* are used for dyeing purposes. They are known as the Aleppo galls and contain from 60-70 per cent of tannic acid. The bark of the White Oak is used in medicine and is recognized as medicina in the United States Pharmacopoeia. The main constituent of oak bark is tannin which is used like other astringents.

The acorns of the Bektes, the evergreen oak of Europe (*Q. ilex*) are used for food. The oaks are well represented in eastern North America, several handsome species also occurring on the Pacific coast.

Formerly and perhaps even now, in some regions the acorns of the White Oak and the Bur Oak were dried, roasted, ground, and used very much as the coffee berry.

In Europe, various species of oaks cause sickness and death in hogs and cattle. Dr. Chevreul suggests that this might possibly be caused by the tannin or bitter principle contained therein.

In sections of the country, where oaks are common, hogs are allowed to run in the forests, the farmers considering that the acorns are fattening. In some parts of the south, it is believed that the mast of oaks makes excellent feed for hogs but is poisonous to cows, a small amount merely decreasing the flow of milk while a greater quantity causes death. It is also claimed that the "sweet mast," that of the white and bur oaks, is less poisonous than the "bitter mast" of mast of black, pin, red and cow oaks. Mr. R. B. Watson made some inquiries upon this subject for me among southern farmers and obtained evidence of four men, which differed slightly in detail but agreed in the conclusions that mast is poisonous to cattle but rather beneficial to horses and hogs. Some say that the coarse hulls or cups clog the digestive tract and cause unthriftiness, others that there is actually poison in the mast. That cattle are affected more seriously than hogs or horses may possibly be explained by the difference in the structure of the digestive organs.

In some localities tympany is said to be produced in cattle that browse on the leaves and bark which are very strongly astringent. The white oak contains about .10 per cent of tannin. *Q. laurifolia* contains in addition to tannic acid mentioned before, from 2-4 per cent of gallic acid. That other plants of the order are injurious has been indicated by Freilberger and Fröhner who state that the European Beech produces violent colic, tetanus, mania and fits of madness resembling those produced by strychnin, that the autopsy shows lesions resembling strychnin poisoning. They recommend giving tannin, morphia and chloral hydrate.



FIG. 284. Red Oak (*Quercus rubra*). 1. Flowering branch. 2. Staminate flower, enlarged. 3. Pistillate flower cluster, enlarged. 4. Fruiting branch, enlarged. 5. Acorn. 6. Cupule. 1, 5, natural size. M. M. Cheney.

The alders (*Alnus glutinosa*) of Europe are used as ornamental trees. Several trees of the genus, on the Pacific coast, are fair sized and produce good timber. The birches are valuable both as ornamental trees and for the excellent quality of their timber. The white birch (*Betula alba*) of Europe is frequently cultivated and birch tar is obtained from it. Cherry birch (*B. lenta*), and yellow birch (*B. lutea*) produce most valuable wood which is used for interior finishing. Paper birch (*B. alba* var. *papyrifera*) is used for making spoons and canoes. The black birch (*B. nigra*) is common along our streams. The common source of oil of wintergreen is the cherry birch. This oil resembles that obtained from *Gaultheria procumbens*.

URTICALES

Trees, shrubs, or herbs. flowers never borne in catkins. monoecious, dioecious, or polygamous, ovary 1 celled, superior. *Urticales* are divided into the

the elm, the elm, *Moraceae*, the mulberries, and *Urticaceae*, the nettles. The castor oil plant of Mexico, the bread nut tree of Central America (*Brassica* *integrifolia*) and the *Latex* of tropical America furnish rubber.



1. 185. Yellow Birch (*Betula lutea*). Flowering branch. 2. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 841. 842. 843. 844. 845. 846. 847. 848. 849. 850. 851. 852. 853. 854. 855. 856. 857. 858. 859. 860. 861. 862. 863. 864. 865. 866. 867. 868. 869. 870. 871. 872. 873. 874. 875. 876. 877. 878. 879. 880. 881. 882. 883. 884. 885. 886. 887. 888. 889. 890. 891. 892. 893. 894. 895. 896. 897. 898. 899. 900. 901. 902. 903. 904. 905. 906. 907. 908. 909. 910. 911. 912. 913. 914. 915. 916. 917. 918. 919. 920. 921. 922. 923. 924. 925. 926. 927. 928. 929. 930. 931. 932. 933. 934. 935. 936. 937. 938. 939. 940. 941. 942. 943. 944. 945. 946. 947. 948. 949. 950. 951. 952. 953. 954. 955. 956. 957. 958. 959. 960. 961. 962. 963. 964. 965. 966. 967. 968. 969. 970. 971. 972. 973. 974. 975. 976. 977. 978. 979. 980. 981. 982. 983. 984. 985. 986. 987. 988. 989. 990. 991. 992. 993. 994. 995. 996. 997. 998. 999. 1000.

The family *Urticaceae* contains the bread fruit (*Artocarpus integrifolia*) an important article of food for the natives of the Pacific Islands, and the tack fruit (*A. integrifolia*) the fleshy exocarps of which are, however, somewhat poisonous. Gumbo gum, a very good substitute for rubber is obtained from this genus. The tree belonging to this family is the coccoloba tree of the Americas produces small fruit which is used in Egypt for food. The common fig (*Ficus carica*) is the most valuable it includes the common and Smyrna fig of commerce containing 60 to 70 per cent of grape sugar. The India rubber tree (*F. elastica*) is the source of some of the India rubber. The bay



Fig. 186. Rubber Tree (*Castilleja elastica*). Staminate flowering branch. A commercial rubber tree of Central America. (After Paguet.)

yan trees (*P. Bengalensis*, *P. religiosa* and *P. altissima*) of the East Indies furnish she-lac. The mulberry (*Morus nigra*) is largely cultivated both for its fruit and for its wood, the latter being very durable for posts. The white mulberry (*M. alba*) is extensively planted, the leaves being used as food for the silk worm. The Osage orange (*Maclura pumila*) of Arkansas, Indian Territory and Texas produces a very durable wood used for posts and pulley blocks. The fruit is said to be poisonous. The wood of fustic (*M. tinctoria*) of the West Indies is used for many purposes. The bark of the paper mulberry (*Broussonetia papyrifera*) is made into paper, and in Japan is also made into cloth. The paper mulberry is cultivated in the South.

The Upas tree (*Anhars toxicaria*) contains anhydrous $C_{27}H_{42}O_{10} + H_2O$. This tree furnishes an arrow poison which the natives prepare from the plant. It is a semi-liquid greenish black substance. The poison acts on the brain and respiratory nerves, causing vomiting and loss of sensation.

Ramie grass cloth or China grass (*Boehmeria nivea*) produces a fine fibre but it is difficult to separate it from the bark and wood. The hop (*Humulus Lupulus*) is cultivated and is the source of the oil of hops which imparts an aromatic, bitter flavor to beer. It contains lupulin which is a tonic and slightly narcotic. The elms are commonly cultivated as shade trees. The best is the American




Fig. 187. Fig-tree (*Ficus*). 1. Flowering branch. 2. Pistillate flower cut through longitudinally. 3. Staminate flower. 4. Fig in longitudinal section. (After Wossido.)



Fig. 188. Flowers and leaves of the American Elm (*Ulmus americana*). 4. familiar tree furnishing an important commercial wood. (W. S. Dudgeon.)



Fig. 189. Deadly Opes Tree (*Anacardium toxicaria*). Flowering branch, portion of staminate flower, longitudinal section of pistillate flower. Alice Taguet.

elm (*Ulmus americana*) which supplies a wood used largely in the manufacture of chairs. The partly ornamental elms are rock elm (*U. racemosa*) and slippery elm (*U. fulva*). The bark of the latter is used in medicine. The rock elm is used in the manufacture of bicycles, the wood taking a pretty finish. One of the common elms of Europe, cultivated in the United States, is the *campestris*. The *U. montana* is the Scotch elm. The hackberry (*Celtis occidentalis*) is a valuable shade tree and furnishes pretty wood which is difficult to work. The *Chlorophora excelsa* is one of the best timber trees of West Africa.

URTICACEAE. Nettle Family

Herbs, trees, or shrubs with stipules, flowers monoecious or dioecious, rarely perfect calyx from the 12-celled ovary which forms a 1-seeded fruit, stamens as many as the lobes of the calyx or sometimes fewer, opposite, ex-albuminous or albuminous, when albuminous the radical points upward cotyledons broad. A small family of 500 species some of which, like *Borhmersa*, produce valuable fibers. The leaves of *Pilea pumila* are demulcent and are said to be valuable in Rhus poisoning.



Fig. 190 Slippery Elm (*Ulmus fulva*). Flowering branch. 2. Fruiting branch. 3. Winter branch with buds. 4. Same showing flower buds beginning to enlarge. 5. Summer branch. 6. Flower enlarged. 7. Longitudinal section of flower, enlarged. 8. Longitudinal section of pistil, enlarged. 9. Stamen, enlarged. 10. Cross section of ovary, enlarged. 11. Longitudinal section of fruit, enlarged. 12. Seed, enlarged. 13. Longitudinal section of seed, enlarged. 14. Embryo, enlarged. 1, 2, 3, 4, 5, 11, one-half natural size. (After M. M. Cheney, in Green's Forestry in Minnesota.)

Genera of *Urticaceae*

Herbs with stinging hairs.

Leaves opposite; flowers 4-parted

2. *Urtica*.

Leaves alternate, staminate flowers 5-parted

3. *Laportea*

Herbs or trees without stinging hairs.

Herbs, pistillate flowers spiked

1. *Cannabis*.

Trees, staminate flowers racemose

4. *Maclura*.1. *Cannabis*, Tourn. Hemp

Dioecious herbs with tough fiber to the inner bark, greenish flowers, sepals 5 in the staminate, 1 in the fertile flower, achene, crustaceous

Cannabis sativa, L.

Stem from 4-8 feet high with broad, divided leaves, the linear-anceolate segments sharply and closely serrate, greenish flowers with narrow staminate panicles and erect pistillate spikes, the sterile with 5 sepals and 5 stamens, fertile flowered spiked, with 1 sepal, fruit hard ovoid, achene oblong

Distribution. Native to Europe and Asia and in waste places from New Brunswick to Tennessee, Kansas and Minnesota



Fig. 191. Hemp (*Cannabis sativa*). Staminate and pistillate flowering branches, fruit, longitudinal section of fruit (After Faguet.)

Poisonous Properties. The resinous secretions of this plant possess very powerful medicinal properties which, however, are said not to be produced by the plant when grown in temperate climates.

Indian Hemp (*Cannabis indica*) is probably not essentially different from the common hemp and has been used in medicine for a long time. According to Dr. Houghton and Mr. Hamilton the American grown product is equal to the Indian Hemp.

The use of hemp seems to have spread through India, Persia and Arabia during the early middle ages. The Hashishin, a sect of the Moravians, killed a large number of the Crusaders during the 11th and 12th centuries by the use of hemp as an intoxicant. The drug is largely grown in India and Turkestan. The form of hemp commonly reached by commerce is called Bhang or Hashish and consists of dried leaves and small stalks frequently mixed with fruits. This is smoked in India with or without tobacco. Ganjah is obtained from the flowering shoots of the female plant or stalk, a stiff woody stem several inches long which is pruned to produce flowering branches. The tops of these are collected then pressed by being trodden by the feet. From this mass comes the drug known as ganjah. It grows in an altitude of six thousand feet. The other forms of the plant consumed in India are Bhang and Charras. Subjee or Bhang is used for smoking. The narcotic ingredient found in majun and charras is undried resin which is obtained by the natives who, when passing among plants wear rubber aprons to which the resin adheres, after which the product is scraped together. The principal constituents of hemp are resin and a volatile oil. The oil or amber colored substance has an oppressive hemp-like smell, and furnishes a resinous substance, cannabin which crystallizes in needles and acts like strychnin. Cannabinol, with intoxicating properties, is obtained from cannabin and is a product from the glands of *Cannabis*. Cannabin hybrid ($C_{18}H_{27}$) is a substance with the camphor-like odor, it is antispasmodic and soporific, and anodyne and a nerve stimulant. Dr. C. F. Millspaugh referring to the products of plants affording this oil concluded from experiments made, that this drug causes depression, epilepsy, vertigo, congestion, followed by cephalalgia, ear ache, toothache, dryness of mouth, throat, lips and lids, it produces nausea, vomiting after coffee, palpitation of the heart weakness of the limbs and dreaminess during sleep. It produces the same symptoms in animals.

The stem of hemp is used by the Mohammedans who smoke it in combination with other substances. They also smoke the non-uried leaves. It is intoxicating and restful to the smoker and alleviates pain, increases the appetite, causes sleep, and induces cheerfulness. It also produces violent coughing and nose bleed.

Hemp is most important in China, and other Asiatic countries, for the manufacture of cordage. The growing of hemp for the same purpose is also carried on to some extent in Nebraska and Kentucky. The seeds of the plant furnish food to birds.

2. *Urtica* (Tourn.) L. Nettle

Herbs with stinging hairs, flowers greenish, monoecious or rarely dioecious, clustered, staminate, with 4 stamens, fertile, with 4 sepals in pairs, fruit an erect, ovate, flattened acheme. A small genus of 30 species.

Urtica gracilis Ait.

A perennial from 2-6 feet high, sparingly bristly, leaves ovate, lanceolate with slender petioles, long, acuminate, sharply serrate, 3-5-nerved, the slender petioles sparingly bristly, flowers dioecious or with staminate and pistillate clusters. The stinging hairs of this and other species of the genus contain formic acid. A common weed in dry or moist ground along fence rows from Canada to British Columbia, Kansas and North Carolina.

Poisonous properties. The nettle and some other plants produce what is commonly called "urticaria" or nettle rash. It is an inflammatory disorder with a burning and itching sensation. It may come out in large or small patches, remaining for a few minutes or several hours and may disappear as abruptly. It usually leaves no trace behind. The nettle is supposed to contain an irritant toxic principle, *formic acid*, but recent studies seem to indicate that the urticaria is probably caused by one of the toxins.

The following species of the genus have urticating properties: *Urtica membranacea*, *U. spatulata* and *U. pilulifera*.



Fig. 192. Stinging Nettle (*Urtica urens*) (From Darlington's Weeds and Useful Plants.)

Urtica urens L. Small Stinging Nettle

An annual from 1-2 feet high, stem 4-angled, tough, branching with a few stinging virulent hairs; leaves elliptical or ovate, serrate or incised, with scattered stinging hairs, flowers loose or in racemose spikes, sepals 4 petals 4; fruit straight, ovate, flattened achene.

Distribution. From New Foundland to Florida and also on Pacific Coast.

Poisonous properties. This nettle has been used in medicine but it is not official. Formerly it was used for flagellation of the skin.

Urtica holosericea Nutt

A tall perennial with stinging hairs, leaves thick, oblong, ovate or ovate-lanceolate, flowers in open panicles.

Urtica dioica L. Stinging Nettle

An erect perennial, leaves and stems beset with stinging hairs, leaves thin, ovate, long petioled, acute or acuminate at the apex, cordate at the base, sharply serrate, flowers in large clusters, cymose-paniculate, often dioecious.

Distribution. Native to Europe but largely naturalized in North America from Atlantic coast to Minnesota and Missouri.

Poisonous properties. Poisonous like the preceding

3. *Laportea* Caudichaud. Wood Nettle.

Perennial herbs with stinging hairs; flowers monoecious or dioecious in loose cymes, the lower mostly sterile, staminate flowers with 5 imbricated sepals; 5 stamens and a rudimentary ovary, pistillate flowers with 4 unequal sepals; stigma elongate, awl-shaped; achene ovate flat, endosperm scant or obscure. About 25 species in warm countries.



Fig. 193. Common Nettle (*Urtica dioica*). Sometimes causes urticaria. (From Johnson's Med. Bot. of N. A.)

Laportea canadensis Gandichand. Wood Nettle

Perennial stem 2-3 feet high, leaves ovate, pointed tip, long-petioled, sharply serrate, fertile cymes divergent, achene smooth, as long as the calyx.

Distribution. In rich woods from Nova Scotia to Minnesota and Kansas and south to Florida.

Poisonous properties. It acts similarly to nettle, the poisonous action being even more pronounced. *L. crenulata*, *L. gigas*, and *L. simulans* also possess similar properties.

Maclura Nalt. Osage Orange

Tree with milky juice, leaves alternate, pinnately veined, stipules caducous; stout, axillary spines; flowers dioecious, staminate in loose, short racemes with 4-parted calyx and 4-stamens, pistillate, capitate with a 4-cleft calyx enclosing the sessile ovary and long exserted style; fruit an achene surrounded by a fleshy calyx, endosperm none; embryo curved, it contains a single species named *Taxylon* by Rafinesque.

Maclura pomifera (Raf.) Schneider Osage Orange

A tree 30-50 feet high; leaves ovate to oblong, lanceolate, pointed, mostly rounded at the base, green and shining, the syncarpous fruit is globose, yellowish green 2-4 inches in diameter, the wood is hard and tough and is used in the manufacture of wagons for paving, fencepost, etc. The tree is extensively planted as a hedge plant.



Fig. 194. Wood Nettle (*Laportea canadensis*). A common wood plant causing urticaria. (C. M. Kang.)

Distribution. In rich woods from Missouri to Kansas to Texas, widely cultivated in the north from southern Nebraska to southern Iowa, Illinois and eastward.

Poisonous properties. This species is listed as poisonous by Professor Bessey in Nebraska. Dr. Halsted notes that a friend of his while working in Osage Orange hedges suffered considerably because of inflammation following the piercing of the thorn. The writer had a similar experience.

Dr. Bessey says "The Osage Orange (*Mocura pennsylvanica*) which has been shown by Dr. Halsted to be more or less harmful as an external poison, is very commonly grown in the southern portion of the state, and it thus adds another to the plants to be avoided by some people. Although I am quite sensitive to some of the external poisons, I have myself never experienced any bad effects from handling the leaves or fruit of the Osage Orange."

PROTEALES

The proteales include one family, the *Proteaceae*, with nearly 1000 species, native to the tropics, mostly of the southern hemisphere.

SANTALALES

Herbs or shrubs generally parasitic; flowers solitary or clustered without corolla; calyx present, imperfect or perfect, pistil 1. Of the two families in the United States, the *Loranthaceae* contains the Southern mistletoe (*Phoradendron*) parasitic upon various deciduous trees like the oak and elm, the *Arceuthobium* of Europe, and the Rocky Mountains, species parasitic upon conifers, one also occurring on spruce trees in eastern North America. Hyams is authority for the statement that the berries of *Phoradendron flavescens* are poisonous to children. Several deaths have been attributed to them. *Santalaceae* contains the fragrant sandalwood (*Santalum album*) of the Indian-Malayan region which contains an oil used in medicine for venereal diseases and for




Fig. 195. Wild Ginger (*Asarum canadense*). Wild ginger is used in medicine. The roots are spicy fragrant, some plants related to it are poisonous. (W. S. Dudgeon.)

perfumes. The bastard toadflax (*Comandra umbellata*) of our northern woods is parasitic upon the roots of flowering plants. The family *Balanophoraceae* consists of chlorophyllless parasitic plants with twining or acaulescent stems, and is native in tropical woods and savannas of Java, India and Australia.

ARISTOLOCHIALES

Plants with twining or acaulescent stems, leaves cordate or reniform, flowers perfect; calyx inferior, the tube adnate to the ovary or partly so, corolla none, ovary generally 6-celled. There are only three families, one of which occurs in North America. The *Aristolochiaceae* includes the wild ginger of the North (*Asarum canadense*) which is more or less purgative and probably also to be regarded as suspicious, its rhizome furnishing the substance *asarin* and a volatile oil which is used in perfumery, the *A. europaeum*, listed by Lehmann as poisonous because of its purgative action and blistering proper-




Fig 196. Southern Mistletoe (*Phoradendron juavescens*). The berries of this plant are said to be poisonous. (W. S. Dodgeon.)

ties, the Dutchman's pipe (*Aristolochia macrophylla*) frequently cultivated and hardly as far north as Minnesota and Wisconsin, the gooseplant (*Aristolochia grandiflora*) of Brazil whose flowers emit an offensive odor, but in spite of this fact the plant is cultivated in greenhouses, Virginia snakeroot (*Aristolochia Serpentaria*) the root stock of which is used as a tonic and contains a volatile oil borneol a bitter poisonous principle aristolochin $C_{22}H_{22}N_2O_{11}$ and the alkaloid aristolochinin. The European (*A. Clematitis*) produces colic and other gastric disturbances and is listed among the pungent narcotic poisons. The substance *asarum* when heated is irritating.




Fig. 197 Dutchman's Pipe (*Aristolochia macrophylla*) Leaf and flower dehiscient fruit. (After A. Paget.)

Several species of the genus *Aristolochia* are used as antidotes against snake-bites and this use is clearly indicated in some of the specific names, as in Virginia snakeroot (*A. Serpentaria*). Other plants of the genus are said to be poisonous, *A. grandiflora* being an example of this. The Arabs use *A. sempervirens* and *A. indica* as snake poison antidotes. According to R. B. White, the Guaco (*A. mexicana*) is a cure for snake bites. Many other plants are used for the same purpose, several compounds being well-known remedies. Among the latter are *Liatris squarrosa*, *Cacalia tuberosa* and *Prenanthes alba*. Other plants having the same qualities belong to the families *Ranunculaceae*, *Orchidaceae*, *Violaceae*, *Polegaceae*, *Liliaceae*, *Umbelliferae*, *Filices* and *Palmae*. One has only to look through such works as the Robinson and Gray's Manual, or Britton's Manual, or various old medical works for the common names of plants with the word snake attached to them, to understand how prevalent was the belief that these plants were antidotes against the bite of venomous snakes.

POLYGONALES

Herbs, shrubs, or trees, often climbing vines, leaves alternate or occasionally opposite, jointed stems, flowers small, regular, dioecious, monoecious or polygamous, calyx 2-6 cleft or parted, inferior; stamens 2-9, inserted near the



Fig. 198. European *Aristolochia* (*Aristolochia Clematidis*) Flowering stem. Listed as a narcotic poison. (After Faguet.)

base, perit. with superior ovary, fruit an achene endosperm mealy. Contains a single family (Polygonaceae).

POLYGONACEAE. Buckwheat Family

Herbs, shrubs, or trees, often climbing, jointed stems, stipules in the form of sheaths, juice often acid or acid; leaves alternate or occasionally opposite; flowers small, regular, mostly perfect, calyx more or less persistent, ovary 1-celled, bearing 2-3 styles or stigmas and a single erect ovule; fruit an achene, 3-4-angled or winged, invested by the calyx, embryo curved or nearly straight endosperm mealy, copious. About 800 species. Of economic importance are the pie plant (*Rheum Rhaponticum*), and rhubarb (*R. officinale*) of Thibet, the root of which contains cathartic acid and is a powerful cathartic, it also contains chrysophan $C_{21}H_{20}O_{14}$, emodin $C_{15}H_8O_2(OH)_3$, rhein $C_{18}H_8O_5(OH)_4$, and chrysophanic acid $C_{15}H_8O_2(OH)_2$. It is purgative and astringent. The canasgre (*Rumex hymenosepalus*) produces a thick root valuable for tanning leather. It is a native of the southwest. The tannin is the same

as that found in rhubarb, and rheotannic acid. The patience dock (*R. Patenzia*), pale dock (*R. altissimus*) and curled dock (*R. crispus*) are troublesome weeds, French sorrel (*R. scutellatus*) is cultivated in Europe and used as a salad. The presence of the silver plant of the west (*Eriogonum umbellatum*) is said to be indicative of gold and silver. *Muehlenbeckia platyclados* of the Samoan Islands is frequently cultivated in greenhouses. The mountain sorrel (*Oxynia digyna*) is used as a salad plant.

Genera of Polygonaceae

Sepals 6 stigmas 3

2 Rumex

Sepals 5, occasionally 4, erect in part.

Achenes triangular or lenticular

Embryo slender curved around one side of the endosperm 3 Polygonum

Broad cotyledons of embryo twisted and plaited 1 Fagopyrum

1 *Fagopyrum* (Tourn.) L. Buckwheat

Annual or perennial, somewhat fleshy, smooth, leafy herbs with erect stems; leaves petioled and alternate, hastate or deltoid flowers, small white, or greenish, paniculately-racemose, perfect; calyx 5-parted, persistent, the divisions like petals, stamen 8; ovary 1-celled, 1 ovule, style with 3 divisions, fruit an achene, 3-angled, endosperm mealy, cotyledons broad. About 6 species native to the old world.

Fagopyrum esculentum Moench. Buckwheat

Smoothish plants; leaves hastate, abruptly narrowed above the middle sheath half-cylindrical; racemes somewhat paniced, many flowered; sepals white, fragrant, with 8 honey-bearing yellow glands situated between the stamens.

Distribution. A common escape in eastern North America. Native of Eastern Europe and Western Asia.

Fagopyrum tataricum (L.) Gaertn. India-wheat

Annual, similar to the above species, leaves deltoid, hastate flowers smaller pedicel short

Distribution. In waste places from eastern Canada to New England. Native to Asia.

Poisonous properties. *Fagopyrum* contains the glucoside indican $C_{12}H_{11}NO$, found also in *Nerium* and other plants. The plant produces bloat especially if consumed before bloom.

Several years ago the writer received a complaint from a farmer stating that the feeding of buckwheat had produced a rash upon his hogs. Feeding of buckwheat and the eruptions or urticaria following are well known to veterinarians.

Dr. Millspaugh says of buckwheat

Many individuals cannot partake of pancakes made from the flour of the seeds without experiencing a severe itching especially observed about the large joints. A peculiarity of this itching is that it occurs after the removing of the clothing and when first retiring at night. The eruption incident to and following this itching takes the form of vesicles which degenerate into dry, dark colored scabs. Another symptom arising is a glutinous condition of otherwise natural feces, making expulsion quite difficult. Increased urinary discharge is also present in many cases.

2. *Rumex*. L. Dock

Coarse herbs, with small, mostly green flowers, which are crowded on generally whorled, panicled racemes, petioles partly sheathing at base, 6 sepals 3 outer herbaceous, sometimes united at base, spreading in fruit, 3 inner larger, slightly colored, enlarged after flowering and convergent on 3-angled achene veined, often bearing a grain-like tubercle on the back, stamens 6, styles 3, stigmas tufted, embryo lying along one side of the albumen slender, and slightly curved.

It has been claimed by some that the seeds of *R. Acetosella* poison horses and sheep.

Rumex obtusifolius Wood. Pale Dock

A tall perennial from 2-6 feet high, glabrous with erect stem, simple or branched above, leaves ovate or oblong, lanceolate, long, acute, pale green veins obscure racemes spike like or somewhat interrupted below spreading



Fig. 199. Two weeds of the smartweed family. A. Sourdock (*Rumex crispus*). B. Sheep sorrel (*Rumex Acetosella*). Both have been suspected. They contain a great deal of oxalate of lime. (U. S. Dept. of Agr.)



Not a specimen of *Passiflora ligularis* (L.) (see p. 471) but a specimen of *Passiflora ligularis* (L.) (see p. 471)

in fruit, pedicels nodding, shorter than the fruting calyx; valves broadly ovate with a conspicuous ovoid tubercle.

Distribution. Common throughout the northern part of the United States

Rumex crispus L. Curled Dock

A smooth perennial from 3-4 feet high, leaves with strongly wavy and curled margins, lanceolate and acute in the lower leaves bases are somewhat truncate or inclined to be ear-shaped, flowers collected in dense whorls, extended or prolonged into racemes, entirely leafless above, but below with small leaves, flower consists of 6 sepals, fruting pedicels as long as the calyx wings; wings heart shaped, erose dentate, each showing a tubercle, achene 3-angled, smooth.

Poisonous properties. The docks contain *rumin* $C_{14}H_{10}O_4$, which is a tasteless golden-yellow substance, slightly soluble in hot water. It acts as a rubefacient and discentient and is used for destroying parasites of the skin.

Rumex has found a place in Pharmacopoea and is also used in medical practice. It causes nausea, watery brown faeces, copious urination, a dry spasmodic cough, and perspiration. The *Rumex orbiculatus*, Great Water Dock, according to Dr. Johnson, is tonic, astringent, and slightly laxative.

Medical properties. It is used as a stimulant and diuretic.

3. *Polygonum* L. Smartweed

Annual or perennial herbs, occasionally woody, stem erect, climbing or floating; leaves alternate, entire, ochrace cylindrical, often fringed, flowers mostly perfect, green white, pink, or purple, calyx 4-5 parted or cleft; stamens 5-9, filaments filiform or dilated to the base, style 2 or 3 parted or cleft; achene lenticular or 3-angled, rarely 4-angled, endosperm present. About 200 species of wide distribution. The *P. tinctorum* of China furnishes the Chinese indigo. The Saghalen knotweed (*P. sachalinense*) was widely advertised as a forage plant a few years ago and is used in Japan and Manchuria as we use asparagus. The prince's feather (*P. orientale*) is cultivated for ornamental purposes. The tanweed (*P. Muhlenbergii*) smartweed (*P. Persicaria*), knotgrass (*P. aviculare*) and black bindweed (*P. Convolvulus*) are troublesome. The Pennsylvania persicaria (*P. pennsylvanicum*) is a valuable honey plant and its seed is a common impurity in clover.

Polygonum Persicaria L. Lady's Thumb

A nearly smooth and glabrous annual from 12-18 inches high, leaves lanceolate or linear, marked with a lunar blotch near the middle, acuminate ochrace somewhat bristly, ciliate spikes ovoid or oblong, erect, stamens mostly 6, style 2-3 parted, achene lenticular.

Distribution. Across the continent in moist places, naturalized from Europe.

Poisonous properties. None of the species is relished by stock, the *P. acre* and *P. Hydropiper* are very acrid and produce gastro-enteritis and erythema, like that caused by buckwheat. The following species produce similar troubles.

Polygonum acre H.B.K. Water Smartweed

A nearly smooth perennial, stems rooting at the decumbent base; leaves linear-lanceolate; ochrace strigose, fringed with long bristles, spikes erect,

panicled, flowers whitish or flesh colored, stamen 8, achenes 3-angled or 4-angled, smooth and shining.

Distribution. Common southward from Missouri to Louisiana, Texas and Mexico.



Fig. 200. Lady's Thumb (*Polygonum Perfoliatum*).
Common in moist places. (Charlotte M. King.)

Polygonum hydropiperoides Michx. Wild Water Pepper

A smooth, branching perennial, slightly or not at all acrid, 1-3 feet high, the narrow sheaths hairy, leaves narrowly lanceolate or oblong-lanceolate, spikes erect, slender, sometimes filiform, flowers small, flesh colored, or nearly white stamens 8, style 3-parted to below the middle, achene 3-angled, ovoid or oblong, smooth and shining.

Distribution. In swamps or wet soils across the continent from New Brunswick to California, Florida and Mexico.

Polygonum Hydropiper L. Smartweed. Water Pepper

Smooth, erect annual, 1-2 feet high, stem often reddish; leaves linear-lanceolate, or lanceolate, spikes, nodding, usually short or interrupted, flowers



Fig. 201 Water Pepper (*Polygonum Hydropiper*). Troublesome to sheep. (After Hochstein.)

usually greenish, stamens 4 or sometimes 6, ochreae cylindrical, fringed with short bristles, style short, 2-3 parted, achene lenticular or 3-angled, dull, granular.

Poisonous properties. This species and the doot yard knot weed (*P. aviculare*) are said to be troublesome to sheep

CENTROSPERMAE

Herbs mostly with perfect flowers, calyx present, corolla, when present, polypetalous, ovary superior; perisperm present; embryo coiled, curled or annular fruit not an achene. Generally fleshy plants, many found in saline soils. It includes the families *Chenopodiaceae*, *Amaranthaceae*, *Nyctaginaceae*, *Phytolaccaceae*, *Coryophyllaceae*, *Portulacaceae* and others. The last named contains the garden pursley (*Portulaca oleracea*), the moss pink (*P. grandiflora*), and the bitter root (*Lewisia rediviva*). The family *Aizoaceae* also of this order, contains the New Zealand spinach (*Tetragonia expansa*), and the ice plant

(*Mesembryanthemum crystallinum*) The family *Basellaceae* another of this same order contains the Madeira vine (*Bomstringaultia baselloides*) commonly cultivated. The seeds of some species are edible.

Families of *Centrospermae*

Fruit an utricle

Flowers bractless or occasionally with bracts.

Sepals green or greenish *Chenopodiaceae*

Flowers bracted.

Sepals generally with scarious bracts *Amaranthaceae*

Fruit fleshy, a berry *Phytolaccaceae*

Fruit indurated into a nut-like pericarp, base of calyx constricted *Nyctaginaceae*

Fruit a capsule, dehiscent by teeth or valves.

Sepals 5 or 4, distinct or united *Caryophyllaceae*

CHENOPODIACEAE Goosefoot Family

Annuals or perennials, frequently succulent herbs, or rarely shrubs, alternate leaves without stipules; flowers small greenish petals absent calyx free, stamens as many as the lobes of the calyx or fewer and inserted opposite them on their base, ovary 1-celled, fruit a 1-seeded, thin utricle or rarely an achene endosperm mealy or wanting, embryo coiled.

About 500 species of wide distribution, common in arid regions. Some of the economic plants of this family are sugar beet (*Beta vulgaris*), a maritime plant of Europe, and spinach (*Spinacia oleracea*) from the orient. The beet is one of the most important plants of the family, being largely cultivated in Europe as a source of cane sugar although as late as 1800, its use in that capacity was of little extent. It is also an important plant for stock food and for human food. Spinach is used extensively for greens but, in Utah, is somewhat of a weed. The Australian saltbushes are well known forage plants. Indigo is derived from *A. horizans*, a native of Tartary. The strawberry blite (*Chenopodium capitatum*) is cultivated in Europe for its leaves. The shrubby saltwort (*Suaeda frutescens*) is burned in the south of Europe for Barilla. The Russian thistle (*Salsola Kali*, var. *tenuifolia*) is used in much the same way. The Spanish wormseed (*S. Webbii*) contains an oil much like that found in *Chenopodium ambrosioides*. The tumble-weed (*Cyloloma atriplicifolium*) is common on the plains. The white sage (*Eurotia ceratoides*) is an excellent forage plant of the west.

Poisonous properties. The use of the beet leaves for fodder has sometimes caused bloat. It has been known for some time that the feeding of roots to animals causes the formation of renal calculi. These calculi consist of a combination of uric and phosphoric acid with lime. An experiment conducted by Prof. W. J. Kennedy and Mr. E. J. Robbins at the Iowa Experiment Station in cooperation with Prof. L. G. Michael indicated that sugar beets fed to rams will produce renal calculi. Prof. Michael says* in regard to the effect of mangels and sugar beets on the kidney:

Both roots seem to affect the kidney similarly.

A small calculus was found in one kidney of Ram VI. This ram was fed sugar beets. The membrane about the calculus and extending down into the urethra was pigmented, a decided black.

* Biennial Rept. Iowa State College of Agriculture and Mechanic Arts 23:142




Fig. 202 Salt Bush (*Atriplex canescens*). A common plant of saline soil in the west. (U. S. Dept. of Agr.)

In the kidney of Ram III the same kind of pigmentation occurred as in Ram VI. Ram III was fed Mangels. No calculus was present.

Miss S. Hartzell, who investigated the chemistry, reports as follows:

A post mortem examination of several valuable animals which the Experiment Station lost showed that renal calculi were present. This resulted in the metabolism experiment which was conducted by the Animal Husbandry Section in co-operation with the Chemical Section of the Experiment Station.

Thus far 42 rams have been used in the experiment, of which 11 were fed hay and corn, 9 were fed hay, corn and ensilage, 11 were fed hay, corn and mangels, 11 were fed hay, corn and sugar beets. In the case of those which were fed hay and corn, and also of those which were fed hay, corn and ensilage, the bladders and kidneys were normal while the results were the reverse in the case of those which were fed sugar beets or mangels along with the hay and corn. The sugar beets and mangels had the same effect. The kidneys were larger than normal, the gall bladders were distended, the bladders were enlarged, often very much so, and in several cases the heart was enlarged. The kidneys in all cases were of a pale color. Three animals died during the experiment. The bladder of one was highly inflamed while in the other two the bladders contained calculi which were too large to pass thru the urethra. The kidneys of two of the animals were broken

down while the third was much enlarged, and calculi were present. In one of the animals which was slaughtered the following conditions existed: bladder enlarged, ulcerated between ureters, kidney surface mottled, kidneys enlarged, cortex discolored, calyces full of calculi, gall bladder much enlarged, other organs normal.

According to Greshoff the leaf of *Kochia scoparia* contains saponin, as do the seeds of this species and of *K. arvensis*. A species of *Atriplex* native to China causes a skin disease known as "atriplicismus."

Genera of *Chenopodiaceae*

1. A spiny shrub

Fleshy herbs.

2. *Sarcobatus*.

Embryo coiled into a spiral, calyx horizontally winged

3. *Salsola*.



Fig. 203. Common Pigweed (*Chenopodium album*). Young shoots sometimes used in place of spinach. (Charlotte M. King.)

Seed with utricle; embryo coiled; calyx not horizontally winged
 1. *Chenopodium*.

Chenopodium L. Pigweed. Goosefoot. Lamb's quarter

Annual or perennial herbs usually covered with a white mealy substance, flowers inconspicuous, in sessile, small clusters, collected in spikes or panicles, perfect, calyx 5-, rarely 4-parted or lobed; stamens generally 5, styles 2, rarely 3, ovary 1-celled, becoming a 1-seeded, thin utricle, embryo coiled around the mealy endosperm.

A small genus of about 60 species of wide distribution in saline soil, around dwellings and in manured soil. Several species like the common pigweed (*C. album*), the Australian spinach (*C. quicquidum*), and the English Good King Henry (*C. Bonus-Henricus*), are used as a substitute for spinach. The quinoa (*C. Quinoa*) is an annual, native to Peru, which produces its flowers in dense, erect panicles. It is cultivated in Chili and Peru for its seeds, which are said to be very strengthening. It was the principal meal food of the Peruvians be-



Fig. 204. Good King-Henry (*Chenopodium Bonus-Henricus*). Used as a substitute for spinach. (From *The American Agriculturist*.)

from the conquest by Spain. An oil is obtained from the wormseed (*C. ambrosioides*). Several species of the genus like *Chenopodium album* and *C. hybridum* are weedy.

Chenopodium Botrys L. Jerusalem Oak

A more or less glandular, pubescent, aromatic annual, leaves with slender petioles, oblong, obtuse, sinuate, pinnatifid, flowers in leafless racemes, calyx 2-3 parted, dry in fruit, only partially enclosed.

Distribution. Naturalized from tropical America. Common in eastern North America to Oregon.

Chenopodium ambrosioides L. Mexican Tea. Wormseed

A smooth annual, leaves slightly petioled, aromatic, oblong, lanceolate, toothed or nearly entire; flowers in spikes, leafy or intermixed with leaves, calyx 2-3 parted, fruit dry, enclosed by the calyx.

Distribution. Naturalized from tropical America. Common in eastern North America to California. The fruit is official in the U. S. Pharmacopoeia.

Chenopodium ambrosioides L. var. *anthelminticum* (L.) Gray. Wormseed

An annual or perennial weed, glandular pubescent, leaves lanceolate, or ovate-lanceolate, acuminate at the apex and narrowed at the base, the lower



170

Fig. 295. Wormseed (*Chenopodium ambrosioides*). Fruit official. (From Darlington's Weeds and Useful Plants.)

lacinate pinnatifid, flowers in spikes without bracts, or the lower spikes leafy bracted.

Distribution. Naturalized from Europe in waste places, from Massachusetts to Ontario and from Wisconsin to Mexico.

Poisonous properties. Several species of the genus contain volatile oils. The *C. ambrosioides*, var. contains the volatile oil of wormseed. This oil has a peculiar, strong, offensive odor and a pungent disagreeable, but aromatic taste. It is said to contain *chenopodium*, $C_8H_{13}NO_2$. In the case of a man who took about one half an ounce of a soluble oil of wormseed, Dr Millspaugh says that the symptoms were those from a narcotic, acrid poison, affecting the brain, spinal cord and stomach. The patient was insensible, convulsed and foamed at the mouth. In another case a man who had taken a considerable quantity displayed hilarity and made futile attempts at talking like a drunken man. Death followed later. *C. mexicanum* contains *saponin*.

2. *Sarcobatus* Nees. Grease-wood

An erect, branched shrub with spiny branches; leaves alternate, linear, fleshy; flowers dioecious or monoecious, the staminate in terminal clusters without a calyx, the pistillate solitary in the axils with compressed calyx, adnate to the base of the papillose stigmas, in fruit a membranous horizontal wing; seed vertical, embryo green, coiled into a flat spiral. Species 1.



Fig. 206. Grease wood (*Sarcobatus* *Martiana*). A plant growing in alkaline soils in the Western United States. Poisonous to sheep. The sharp spines cause mechanical injury. (U. S. Dept. Agri.)

Sarcobatus Maximiliani Nees, Torr Grease-wood

A glabrous perennial with succulent foliage and spiny branches Wood hard.

Distribution. In dry, alkaline soil from western Nebraska to New Mexico, Nevada, and Montana, most abundant west of the front Rockies.

Poisonous properties. Prof. Chesnut says

A correspondent in New Mexico states that on one occasion he counted as many as one hundred sheep that had been killed by eating the leaves of this plant. It is claimed that cows are not affected by eating it at any time and that sheep can eat it quite freely in winter. Death is perhaps due more to the blighting effect than to any poisonous substance which the plant contains.

It might be noted also in this connection that the sharp spines on the plant often inflict serious injuries to persons who come in contact with it and also to animals, setting up inflammation and causing the formation of pus. It is used as a forage plant.

3. *Salsola* L. Saltwort

Bushy branched herbs, succulent when young, but rigid at maturity; leaves terete, prickly-pointed and sessile, flowers sessile and axillary, calyx 5-parted, persistent, enclosing the depressed fruit, the divisions horizontal, winged on the back, enclosing the utricle, stamens 5, ovary depressed, style 2, embryo coiled in a conical spiral. About 40 species of wide distribution, saline soils.

Salsola Koli L., var. *tenuifolia* G. T. W Meyer Russian Thistle

An herbaceous, smooth or slightly pubescent annual, diffusely branched from the base; from 1½ to 3 feet high, spherical in the mature form, leaves fleshy, alternate, succulent, linear, subterete, 1-2 inches long, pointed in the older specimens, upper leaves in the mature plant persistent, each subtending 2 leaf-like bracts and a flower, stem and branches red; apetalous flowers solitary and sessile; calyx consisting of 5 persistent lobes, enclosing the dry fruit which is usually rose colored, about 1-12 of an inch long, 5 stamens, nearly as long as the calyx; pistils simple with 2 slender styles producing a single ob-conical depressed seed, dull gray or green, without albumen, embryo spirally coiled. The plant flowers in July or August.

Distribution. Common from Minnesota to Kansas, west across the continent, Illinois and Kansas to New Jersey.

Injurious properties. The Russian thistle not only clogs the harvesters and harrow, injures horses legs so that boots have to be put on them but is equally disagreeable to come in contact with, to man. On this point, Prof. Dewey says

The sharp spines on the plants not only irritate and worry both horses and men, but often, by breaking under the skin, cause festering sores on the horses' legs, so that in many localities it has been found necessary to protect them with high boots or leggings. In handling grain or flax, in the processes of hauling and threshing, the sharp spines cause considerable irritation and consequent loss of time.

AMARANTHACEÆ. Amaranth Family

Herbs, or in some cases, shrubs, leaves simple, mostly entire flowers small, green or white with bractlets, usually in terminal spikes or heads, petals none; calyx herbaceous or membranous, 2-5 parted, segments distinct or united;



Fig. 207. Russian Thistle (*Salsola Koli*, var. *tenuifolia*). Causes mechanical injuries to man and stock (Charlotte M. King.)

stamens 1-5, mostly opposite the calyx-segments; ovary 1-celled, ovules solitary, fruit an utricle, circumscissile, irregular or indehiscent, seed generally smooth; endosperm usually copious and mealy. About 425 species in tropical countries mostly. Several like *Celosia cristata* are cultivated for ornamental purposes and several are weedy. Among the latter are the tumble-weed (*Amaranthus gracilis*), pig-weed (*A. retroflexus*), and prostrate pig-weed (*A. blitoides*). The leaves of several species are used as food.

Amaranthus (Tourn.) L. Pig-weed. Tumble-weed.

Annual, branching or erect herbs, smooth or pubescent leaves, simple, small flowers, monoecious, dioecious, or polygamous, green or purplish, generally with 3 bractlets, in spikes or axillary clusters, sepals 3-5, distinct, stamens 2-5, styles 2-3, fruit oblong, utricle. About 50 species of wide distribution, mostly of southern states.

Amaranthus retroflexus L. Pig-weed. Red Root

Roughish, slightly pubescent, annual with stout stems 2-4 feet high, leaves ovate or rhombic ovate; upper lanceolate, acute or acuminate at apex, flowers in dense spikes, bractlets about twice as long as the 5 scarious mucronate-tipped sepals, stamens 5; seed black.

Distribution. Naturalized from tropical America, found throughout the United States, especially on waste ground far northward. Also naturalized in Europe.

Amaranthus hybridus L. Slender Pig-weed

Similar to the preceding but with darker green or purple foliage, stem more slender, erect, leaves ovate or rhombic ovate, smaller than the preceding,




Fig. 208. Prostrate Pigweed (*Amaranthus blitoides*). A common weed. May cause blout. (Charlotte M. King.)

spikes linear cylindrical, forming dense terminal panicles bracts subulate, twice as long as the acute or cuspidate sepals, stamens 5, utricle but slightly wrinkled.

Distribution. Species naturalized from tropical America but rare or local in places; common southward.

Amaranthus spinosus L. Spiny Amaranth

Stout, branched stem, leaves ovate, rhombic-ovate or lanceolate, acute at both ends with a pair of rigid stipular spines, sepals mucronate-tipped 1-nerved, utricle scarcely circumscissile.

Distribution. In waste or cultivated ground as far north as Massachusetts, Illinois and common in Missouri and Southward. Naturalized from tropical America.

Poisonous properties. The spiny amaranth sometimes produces mechanical injuries. Mr. O'Gara calls attention to the injurious properties of the first species in Nebraska. He says that it doubtless causes a great deal of trouble in some parts of that state. Mr. C. C. Palmer near North Platte lost 5 head of cattle in his pasture. In all cases they were very much bloated and a post-mortem examination revealed a good deal of pig-weed in the stomachs. The animals in question had been accustomed to prairie grass pasture and broke into a field containing considerable of this pig-weed, some Russian Thistle and



Fig. 209. Pigweed (*Amaranthus retrofractus*). A cause of bloat in cattle. (From Darlington's Weeds and Useful Plants.)



Fig. 210. Spiny Amaranth (*Amaranthus spinosus*). Sometimes produces mechanical injuries. (From Darlington's Weeds and Useful Plants.)

lamb's quarter. After the death of the animals the fence was replaced and no further trouble was noticed. An experiment with animals carried on by Mr O'Gara proved negative. Many families in the vicinity of North Platte regard the weed as a bad bloater, by some considered as serious as green clover and alfalfa.

PHYTOLACCACEAE. Pokeweed Family

Generally herbs, a few tropical species, trees or shrubs, leaves alternate, entire, without stipules; flowers regular, perfect, polygamous or monoecious, calyx petal-like, of 4 or 5 sepals, or 4-5-parted, stamens 5-30 alternate with the segments of the calyx or with the sepals, of the same number or more numerous, ovary several-celled; ovules solitary

A small family of about 85 species, mostly tropical. The juice of the berries of the Umbra tree (*Phytolacca dioica*) of South America, now naturalized in Europe and other warm countries, is used to color wines. Other plants of the family, like bloodberry are frequently cultivated. The latter produces small spikes of white flowers, followed by red berries. Strong drastic substances occur in *P. littoralis* and *Amisomeria drastica*, natives of Chile. *P. abyssinica*, *Villamala peruviana* and our native pokeberry contain saponin and red coloring matter.

Phytolacca (Tourn.) L. Pokeweed

Tall, stout, perennial herbs with large petioled leaves, flowers borne in racemes, calyx of 5 petal-like sepals, stamens 5-30, ovary of 5-12 carpels united to form a ring, 5-12 celled, with a single seed in each cell; embryo around the endosperm.

Phytolacca decandra L. Pokeweed, Garget

A tall, glabrous, perennial herb, 6-9 feet tall, with strong odor, large poisonous root; leaves oblong-lanceolate, acute, or acuminate, flowers perfect, calyx white, stamens 10, shorter than the sepals, ovary green, 10-celled; berry dark purple, filled with crimson juice.

Poisonous properties. The young shoots of this plant may be boiled and eaten, the acrid property being dissipated in boiling. The leaves are eaten by the natives of the island of Guam. A tincture of the plant is used for rheumatism. The root is alterative, emetic, cathartic, and narcotic. Prof. Chesnut, in speaking of its poisonous nature, says:



FIG. 311 Pokeweed (*Phytolacca decandra*). The root of the plant is very poisonous. (C. M. King.)

Most instances of poisoning arise from an overdose when the plant has been used as a medicine, but there are also accidental cases due to eating of the root, which has been variously mistaken for that of the parsnip, artichoke and horseradish. A few fatal cases of poisoning of children have been attributed to the fruit, but whether death was really due to the seed or the pulp is somewhat uncertain. The evidence is chiefly against the

seed, for it is known to contain a poisonous substance. Pokeweed is a violent but slow-acting emetic, vomiting beginning only after about two hours. It also affects the nerves and muscles, producing retching, spasms, severe purging and sometimes convulsions. Death is frequently due to the paralysis of the respiratory organs.

Dr. Guttenberg makes a similar report in regard to effects of poisoning by pokeweed, adding that death often is a result.

The roots of pokeweed are often mistaken for other fleshy roots, such as horse-radish. The leaves, as has been said, are harmless when boiled, somewhat resembling spinach, but the root is very poisonous. The poke root was used by the Indians in medicine. Dr. Millspaugh, who values the plant not only as an emetic, but also as an efficient remedy, says

In certain forms of rheumatism, the root with lard was found to be an excellent ointment as a cure for many forms of skin disease: psoriasis, eczema, scabies, and tinea circinata, also in syphilitic ulcers.

Dr. Millspaugh says

The fresh root, gathered late in autumn or early in spring, is chopped and pounded to a pulp and weighed. Two parts by weight of alcohol are taken, and after thoroughly mixing the pulp with one-sixth part of it, the rest of the alcohol is added. After having stirred the whole, pour it into a well-stoppered bottle, and let it stand eight days in a dark, cool place. The tincture is then separated by decanting, straining, and filtering.

Thus prepared it has a light straw-color by transmitted light, at first a stinging, soon followed by a decided bitter taste, and a very slight acid reaction.

He adds

I noted in my readings several years ago that the berries had been used for pies by frugal housewives, and often since have half determined to try poke-berry pastry, distraction has, however, always overruled valor, and the much-thought-of pie is still unmade and uneaten. The young shoots, however, make an excellent substitute for asparagus, and I much prefer them, if gathered early and discriminately.

The acrid alkaloid *phytolaccin*, according to Dr. Edmond Preston, occurs in the root of this plant also *phytolaccic acid* and an amorphous yellowish brown, transparent substance, very soluble in water and alcohol. Nagi reports a toxic substance *phytolaccotoxin* $C_{24}H_{34}O_8$. The berries have been used for coloring but this is not entirely successful, because no mordant will fix the color. The juice of the berry is a delicate test for acids when lime water is added to it.

Dr. Johnson says

All parts of the plant possess acid and somewhat narcotic properties. The use of the fresh plant, or a strong decoction of the root, applied locally, may strongly irritate the skin, especially if tender or abraded. Taken internally it causes nausea, vomiting, and purging, and, in overdose, acro-narcotic poisoning. It has been employed with more or less satisfactory results in a great variety of cutaneous affections, and in rheumatism, especially when chronic or of a syphilitic origin. There is little doubt that, in view of the uncertainty which at present exists regarding it, this plant would well repay further careful experimentation.

Nagi reports that *phytolaccotoxin* resembles *picROTOXIN* and *escatotoxin*. A glucoside has also been found in common poke, saponin also occurs.

CARYOPHYLLACEÆ. Pink Family

Herbs with opposite entire leaves, frequently swollen at the nodes, flowers perfect or rarely dioecious, sepals 4 or 5, persistent separate or united with the calyx tube, petals of equal number, styles 2-5, or rarely united into 1, ovary usually 1-celled, occasionally 3-5-celled, ovules attached to a central column; seeds several or many, small coiled or curved embryo, with a mealy albumen.

A large family of about 70 genera and 1500 species, widely distributed, most abundant in the northern hemisphere. Many of the plants of this family




Fig. 211a.

Fig. 211a. Flowers of Soapwort or Bouncing Betty (*Saponaria officinalis*), Calyx, Corolla, Stamens and Pistil. (C. M. King).

are cultivated for ornamental purposes. Of these we may mention the hardy pink (*Dianthus barbatus*) and carnation (*D. Caryophyllus*). The spurrey (*Spergula arvensis*) is occasionally cultivated as a forage plant in Europe and sometimes in this country, but is a weed of grain fields in Europe. A few species like *Saponaria officinalis* and the catchfly are medicinal. A red dye is obtained from a species of *Coccus* found on *Scleranthus perennis*. The leaves of *Puronychia argentea* are used as a substitute for tea. The mitchwort (*A. crassifolia*) of Europe and some parts of the United States is poisonous to horses. The European sandwort (*Arenaria serpyllifolia*) common eastward in sandy waste places is said to cause salivation in horses. Several species of the family like *Saponaria officinalis*, *Gypsophila Struthium* of Spain *Agrostemma*, *Lychnis*, and *Herniaria* contain saponin.

Genera of Caryophyllaceae

Sepals united into a tube or cup.

Calyx ovoid or sub-cylindrical, 5 angled, not prominently nerved 5 *Saponaria*.

Calyx 5-toothed, prominently nerved.

Styles 3 2. *Silene*.

Styles 5 or 4, alternate with petals 3. *Lychnis*.

Styles 5 or 4, opposite petals, silky plants 4. *Agrostemma*.

Styles 2 1. *Gypsophila*.

Sepals distinct 6. *Stellaria*.

1. *Gypsophila* L. Gypsophyl

Glabrous and glaucous herbs, leaves narrow, flowers small, in paniculate, axillary clusters. calyx cylindrical, 5-toothed, 5-nerved without bractlets. petals 5 claws, narrow, stamens 10, styles 2. About 50 species native to Europe, 2 species introduced to North America.

Gypsophila paniculata L. Tall Gypsophyl

A glabrous or pubescent perennial, from a simple fusiform root; leaves lanceolate, narrowed at the base, flowers in paniced cymes, calyx campanulate,

segments with scarious margins; petals white or pink, slightly emarginate, larger than the calyx.

Distribution. Native to Europe and Asia. From Manitoba to Nebraska.

Poisonous properties. Used in medicine as a detergent. An allied *G. Struthium* contains *Sapotoxin* and the glucoside *saponin*. It is an acrid poison.

2. *Silene* L. Catchfly

Herbs with pink or white flowers, solitary or borne in cymes, calyx more or less inflated and five-toothed; petals 5, narrow and clawed, stamens 10, styles 3, rarely 4 or 5, ovary 1-celled or incompletely 2- to 4-celled, pod 1-celled, dehiscent by 6, apical teeth; seeds roughened.

About 250 species of wide distribution. Several like sweet William (*S. Armeria*) are cultivated for ornamental purposes. The starry campion (*S. stellata*) of our prairies and thickets might well be cultivated more than it is.

Silene latifolia (Mill.) Britten & Rendle Bladder Campion

A branched perennial, a foot or more high, with opposite glaucous ovate lanceolate leaves; flowers in loose cymose panicles, calyx bladderly, inflated, petals 2-cleft, white; seed roughened.




Fig. 212. Deptford Pink (*Dianthus Armeria*). Frequently cultivated for ornamental purpose. (Charlotte M. King.)



Fig. 213. Night-flowering Catchfly (*Silene noctiflora*) (After Fitch.)

Distribution. Native to Europe. Common in fields and along roadsides from New England to Illinois and Iowa.

Silene aoturrhina L. Sleepy Catchfly

A puberulent annual with glutinous nodes and slender stem; lower leaves spatulate or oblanceolate, petioled, upper leaves linear to subulate, flowers small in cymose panicles, calyx not inflated, but expanded by the opening pod, ovoid, petals pink, obcordate, minutely crowned, seeds small, roughened.

Distribution. Common in sandy fields, gravelly soils, and in waste places from New England and Florida to Mexico, north to British Columbia, and east to Ontario.

Silene noctiflora L. Night-flowering Catchfly

A viscid hairy annual, from 1-3 feet high, lower leaves obovate or oblanceolate, the upper sessile and lanceolate, flowers few, in a loose panicle, white or



Fig. 214. Sleepy Catchfly (*Silene aoturrhina*). A weed of sandy fields. (Charlotte M. King.)



Fig. 215. Night-flowering Catchfly (*Silene noctiflora*). Poisonous. (Charlotte M. King.)

pinkish, fragrant, opening at night, calyx tube elongated and enlarged by the ripening pod, petals 2-cleft and crowned, seeds small blackish, roughened, kidney shaped.

Distribution. Native to Europe. Common in waste places from New Brunswick to Florida, Kansas and Iowa to Manitoba.

Poisonous properties. According to Stebler and Schröter, the leaves of *Silene latifolia* are eaten by stock and it is regarded as of some value for forage purposes, but Prof. Schaffner, in his "Poisonous and Other Injurious Plants of Ohio," suggests that the sleepy catchfly may be poisonous.

3. *Lychnis* (Tourn.) L. Campion

Erect herbs, with ovoid tubular oblong or inflated calyx 5-toothed, 10-nerved, occasionally with leaf-like lobes; petals 5, or rarely 4, styles 5, rarely 4, alternate with the often appendaged petals, seeds numerous, globular or kidney-shaped pod opening by as many, or twice as many valves. A small genus of about 40 species native to the cooler regions. Several species cultivated for ornamental purposes. The scarlet lychnis (*L. chalcedonica*) is frequently cultivated in old gardens.

Lychnis Flos-cuculi L. Ragged robin

A downy, branching, pubescent annual, or viscid above, from 2-3 feet high, leaves lanceolate or linear lanceolate; flowers in loose panicles, red, bluish, or whitish; calyx glabrous, short, petals cleft into 4 lobes; capsule globose.

Poisonous properties. It contains a form of saponin called *lychnidin*.

Lychnis dioica L. Evening Lychnis

Biennial, usually dioecious, viscid, pubescent; leaves ovate-oblong or ovate-lanceolate; flowers few, loosely paniculate, white or pinkish, opening at evening; calyx tubular, becoming swollen with the ripening fruit; styles 5.

Distribution. Native to Europe, common in eastern and middle states. In the West it is not uncommon in clover fields, where it is introduced with clover seed.

4. *Agrostemma* Linn. Corn Cockle

Calyx ovoid, 10-ribbed; teeth elongated, longer than petals; stamens 10, styles 5, opposite unappendaged petals; leaves linear. Tall annual or biennial plant.

Agrostemma Githago L. Corn Cockle

A hairy annual weed, leaves linear-lanceolate, acute or long-acuminate, flowers perfect, long-peduncled, calyx lobes long, linear, surpassing the purple red petals, capsules 1-celled, large with numerous large seeds which are roughened and black.

Distribution. This plant is widely distributed from Nova Scotia to Quebec, south from New England to the southern states, and westward and northward, generally in wheat growing regions. Difficulty in screening wheat by ordinary methods has caused this weed to be generally scattered in wheat growing regions. These screenings are much used in feeding stock in some places. The farmer often sows cockle with his wheat.



Fig. 216. Meadow Lychnis (*Lychnis Floriculsi*). Contains saponin. (After Fitch.)



Fig. 217. Corn cockle *Agrostemma Githago*. a, sprays showing flowers and seed capsule, one-third natural size, b, seed, natural size b', seed, four times natural size. (U. S. Dept. Agr.)

Poisonous properties. According to Kruskal, the seeds contain *githagin* $2(C_{17}H_{29}O_{11})$. The ripe dried seeds are broken into a coarse powder and used in medicine. Dr. Millspaugh gives the proportions as follows: "Five parts by weight of alcohol are poured upon the powder, and the whole allowed to stand eight days in a well stoppered bottle in a dark cool place, shaking thoroughly twice a day." The tincture is somewhat acrid. The seeds of the cockle are frequently used to adulterate cheaper grades of flour in Europe. Dr. Millspaugh gives a case in which death followed where two 14½ oz lots of wheat flour containing respectively 30% and 45% of these seeds were fed to two calves. This amount of cockle caused severe cramps of the stomach within an hour, followed by diarrhoea and finally death. Where ducks and geese ate the seeds, death followed when sufficient was taken, and the post-mortem showed inflammation of the bowels. Prof. Pierce states that this is especially true when the seeds are crushed. A large amount of screenings are sold for chicken feed, and frequently complaints are made of poison, or at least that chickens will not eat the screenings.

In describing symptoms indicative of poisoning by corn cockle, which, Dr. Allen says, place the seeds among the cerebro-spinal irritants, he agrees essentially with Dr. Chesnut.

Dr Chesnut says:

The poisonous constituent, *saponin*, is a non-crystalline powder, very freely soluble in water, and possessing a sharp, burning taste. It has no odor, but when inhaled in the smallest quantity it produces violent sneezing. When briskly shaken with water it froths like soap. The poison is found in nearly all parts of the plant, but mainly in the kernel of the seed. Cases of poisoning have been noted among all sorts of poultry and household animals, but are rarely due to any portion of the plant as found growing in the field. The poisoning is generally produced by a poor grade of flour made from wheat containing cockle seeds. Machinery is used to remove these seeds from the wheat, but the difficulty of separating them is so great that the result is not entirely accomplished. The quantity remaining determines the grade of flour in this particular regard. It sometimes amounts to 30 or 40 per cent, but this quality is sent out only by ignorant or unscrupulous dealers or is intended for consumption by animals only. Flour containing a smaller amount has often been made into bread and eaten, sometimes with fatal results, the baking not always being sufficient to decompose the poison. The effect may be acute, or, if a small quantity of the meal is eaten regularly, it may be chronic. In the latter case it is sometimes known as a disease under the name of "githagium." The general symptoms of acute poisoning are the following: Intense irritation of the whole digestive tract, vomiting, headache, nausea, diarrhoea, hot skin, difficult locomotion, and depressed breathing. Coma is sometimes present, and may be followed by death. Chronic poisoning has not been closely studied in man, but experiments upon animals show chronic diarrhoea and gradual depression, the animal losing vigor in breathing and in muscular movements until death ensues. The action is antagonized by the use of digitalis, or of the simple extract of digitalis (*Digitalis purpurea*) a dangerous poison, which should be given only by a physician.

The more prominent symptoms as recorded by Friedberger and Fröhner are, briefly, colic, vomiting, slavering, paralysis, stupor, hyperaemia of brain and spinal cord.

Dr Chesnut also adds:

Corn cockle meal is easily detected in second and third class flour by the presence of the black, roughened scales of the seed coat. These are sure to occur if the flour has not been well doited. Its presence is otherwise detected by the peculiar odor produced when the meal is moistened and by chemical tests with iodine. Wheat containing corn cockle seeds should be rejected for planting.

It has been asserted in Europe that corn cockle is injurious in flour and bread stuffs. Dr Chesnut says:

A person eating 1200 grains of bread made from flour containing only one-half per cent of corn cockle seed would consume six grains of cockle seed, an amount which the author believes beyond a doubt to be poisonous in its effects.

The poison in corn cockle is *sapotoxin* $C_{11}H_{20}O_{14}$ and is partially decomposed while baking, but nevertheless some of it remains and the use of flour which contains corn cockle should be forbidden. It has long been suspected of being poisonous. Mr John Smith in his Domestic Botany, says:

It being difficult to separate the seeds from the grain, the value of the latter is deteriorated, and the flour is rendered unwholesome.

5. *Saponaria* Linn. Soapwort

Calyx ovoid to sub-cylindrical, 5-toothed, obscurely nerved, terete or 5-angled, smooth, stamens 10, styles 2, pod 1-celled, or sometimes 2-4-valved, and 4-toothed to apex. Coarse annual or perennial with mucilaginous juice, hence common name of soapwort because of the property of forming a lather with water.

Saponaria officinalis L. Rouncing Bet

Perennial herbs with large flowers in cymose clusters, calyx narrowly ovoid or oblong, five toothed, petals clawed or unappendaged, stamens 10, styles 2, pod 1-celled or incompletely 2 to 4-celled and 4-toothed at the apex. About

40 species in Europe, Asia, and Northern Africa. *Saponaria officinalis* is frequently cultivated in old gardens. The mucilaginous juice forms a lather with water and is valuable for taking grease spots out of wollen cloth.

Saponaria Vaccaria L. Cow herb

A glabrous annual from 1-2 feet high with opposite ovate lanceolate leaves, flowers in corymbed cymes; calyx 5-angled, enlarged and angled in fruit, petals pale red. Cow herb is another important constituent of "cockle" in wheat screenings, and like the preceding weed has been largely spread by means of wheat culture.

Distribution. Common in Europe, found in wheat fields of the east and as far west as Missouri, Kansas, the Rocky Mountain region, and Pacific Coast, and wheat regions of the northwest.

According to Sohn, it contains the substance *saponin*, $C_{62}H_{104}O_{18}$, a neutral sharp, amorphous substance, having a burning taste and producing a violent sensation. The toxic substance is partially removed by baking.



Fig. 218. Bouncing Betty (*Saponaria officinalis*). A branch with flowers. The double flowered form is sometimes cultivated for ornamental purposes. (Charlotte M. King.)

6. *Stellaria* L. Chickweed

Tufted herbs with white flowers in cymose clusters; sepals 4-5, deeply 2-cleft, sometimes none, stamens free, 10 or fewer; styles 3, rarely 4 or 5, capsule ovoid 1-celled, several to many seeded.

Stellaria media (L.) Cyrill. Common Chickweed

A nearly smooth annual or winter annual, decumbent or ascending; leaves ovate or oval, the lower on hairy petioles; flowers white in terminal leafy cymes or solitary in the axils, sepals oblong, longer than the 2-parted petals; stamens 2-10.



Fig. 219 Cow-herb (*Saponaria Vaccaria* L.). The seed of this plant is common in wheat screenings and is supposed to be poisonous. (U. S. Dept. Agr.)

Distribution. A weed in waste places, lawns and fields, very troublesome in lawns. Naturalized from Europe. Extends from New England and Canada across the continent.

Poisonous properties. The seeds of common chickweed are used as food for cage-birds and are also readily eaten by chickens, but, according to Mr Wm Carruthers, they cause disorder to the digestive system when eaten by lambs in large quantities.



Fig. 228. Chickweed (*Stellaria media*) Seeds said to be injurious. (Charlotte M. King.)

RANALES

Herbs, shrubs or trees, calyx usually of separate sepals, corolla usually present and of separate petals, ovary or ovaries superior, free from the calyx carpels 1-many, stamens mostly hypogynous and more numerous than the sepals. Contains the families *Nymphaeaceae*, *Ceratophyllaceae*, *Ranunculaceae*, *Berberidaceae*, *Menispermaceae*, *Magnoliaceae*, *Calycanthaceae*, *Anonaceae*, *Myristicaceae* and *Lauraceae*. The *Nymphaeaceae* are aquatic perennial herbs. The rhizome of water chinquapin (*Nelumbo lutea*) of the Mississippi Valley and introduced into Massachusetts by the Indians, was used for food. The sacred bean or lily (*N. nucifera*) cultivated for ornamental purposes produces an edible seed and rhizome rich in starch. The pods of wokus (*Nymphaea polysepala*) are

used as food by the Indians in the northwest. The blue flowered *Nymphaea stellata* of tropical Africa and the Egyptian lotus (*N. Lotus*) are frequently cultivated, as are the *Victoria regia* of the Amazon region and *Euryale ferox* of eastern Asia. The water lilies (*Castalia odorata* and *C. tuberosa*) are pretty water plants of North America. The family *Ceratophyllaceae* contains the water weed (*Ceratophyllum demersum*) of North America, troublesome also in Europe. The family *Myristicaceae* contains the nutmeg (*Myristica fragrans*) of which mace is the aril, the oil contains *myristicin*. Prof Cushny in an address before the Royal Society of Medicine in London, referring to Nutmeg poisoning, says that the symptoms are drowsiness, stupor, and diplopia ('seeing double'). Delirium is frequently present, and sometimes the first symptom is burning pain in the stomach, with anxiety or giddiness. The symptoms generally resemble those resulting from *Cannabis indica* (hashish). One fatal case oc-




Fig. 221 Yellow Water Lily (*Nymphaea polysepala*). (W. S. Dudgeon.)

curred in a boy who had eaten two nutmegs. From experimental work Prof. Cushny has come to the conclusion that the symptoms are to be attributed to the action of the oil of nutmeg on the central nervous system. This is depressed; but there are some signs of stimulation in the form of restlessness, slight convulsive movements, and tremor. The oil has also a marked local irritant action, whether given by the mouth or hypodermically. Several other species like *M. succedanea* and *M. fatua*, are used by the natives where these plants are indigenous.

Families of Ranales

Stamens numerous sepals distinct, petals absent or present.

Receptacle hollow enclosing the numerous pistils and achenes, leaves opposite. 5 Calycanthaceae

Receptacle not hollow; flowers generally perfect.

Fruits cohering over each other, cone-like Magnoliaceae

Fruits not cohering over each other, separate.

Anthers not opening by uplifted valves, pistils usually more than 1

Sepals 3, petals 6; shrubs or trees Anonaceae

Sepals 3-15; petals when present about as many Ranunculaceae

Anthers opening by uplifted valves except *Podophyllum*, pistil

1 Berberidaceae

Dioecious climbing vines, simple leaves Menispermaceae

Stamens 9-12 in several series, anthers opening by uplifted valves, petals absent. Lauraceae

RANUNCULACEAE. Crowfoot Family

Herbs or a few woody plants with acid juice, flowers polypetalous or apetalous, regular or irregular, calyx free, often colored like the corolla, sepals 3-15; petals 3-15 or absent, stamens numerous, pistils few or numerous, distinct; fruit a dry pod, berries or achene seed-like, embryo minute, albumen present.

A rather large, widely distributed family of plants many of which like aconite, larkspur, and marsh marigold, are poisonous. Many, such as virgin's bower (*Clematis virginiana*), *C. Jackmanni* and other species are cultivated for ornamental purposes, the *C. Jackmanni* being especially desirable. The columbines, like the European columbine (*Aquilegia vulgaris*), the Rocky Mountain columbine (*A. caerulea*), and our eastern columbine *A. canadensis*, the paeonies (*Paeonia officinalis* and *P. Montana*), and the larkspurs (*Delphinium*) are also cultivated for ornamental purposes, the most familiar of the latter being the garden annual, *Delphinium Consolida*. Several perennial species of *Delphinium* are also very showy. The seeds of stavesacre (*D. Staphisagria*), native to southern Europe and the Levant, contain an alkaloid *delphinium* $C_{21}H_{23}NO_5$ which is a powerful and acrid poison.

Nigella is said to contain an alkaloid, *nigellin*, *N. damascena* contains the alkaloid *damascenon* $C_8H_{11}NO_2$. In 1872, a German chemist found an alkaloid in *Isopyrum thalictroides*, the so-called *isopyrin* $C_{22}H_{21}NO_5$. *Thalictrum macrocarpum* contains the alkaloid *thalictrin*. *Aquilegia* is said to be free from alkaloids.

Gluconides have also been found in some of the members of this family, as adonidin $C_{20}H_{26}O_8$ in *Adonis amurensis*.

The European *Adonis aestivus* and *A. vernalis* are recorded as poisonous by Lehmann. The black cohosh (*Cimicifuga racemosa*) is also somewhat acrid.

Many of the plants of the family contain *anemonin* $C_{16}H_{14}O_4$ and some are used for medicinal purposes.

Among these are aconite (*Aconitum Napellus*), crocus, Pasque flower, pulsatilla *Anemone patens* var *Wolfgangiana* (Bess) Koch. and yellow puccoon (*Hydrastis canadensis*). The black roots of black hellebore (*Helloborus niger*) are used in medicine, as a purgative, being poisonous in overdoses. The tuberous roots of one of the crowfoots (*Ranunculus Ficaria*) resemble grains of wheat and are sometimes boiled and eaten but they have a sharp acrid taste and are known to produce blisters. The water crowfoot (*R. aquatilis* var *capillaceus*) is apparently harmless and is used as forage in England and on the Continent.



Fig. 222. European Columbine (*Aquilegia vulgaris*). Flowering branch, flower, longitudinal section of flower, pistil and stamens. (After Engelm.)

According to Greshoff *Clematis Fremonti*, *C. integrifolia*, *C. lanuginosa*, *C. orientalis*, *C. pseudo-flammula* contain HCN. He also states that saponin is of widespread occurrence in this genus and that he found it in the leaves of *C. Pitcheri*, and *C. recta* and in the leaves of *Trollius pumilus*, and *T. chinensis*.

Genera of Ranunculaceae

Flowers regular

Sepals 3-20 apetalous.

Achenes tailed 5 *Clematis*

Achenes not tailed 4 *Anemone*

Fruit a follicle, sepals yellow.. 1 *Caltha*

Flowers solitary, sepals 3. 9 *Hydrastis*

Petals and sepals present.

Petals 5, yellow or white 6 *Ranunculus*

Petals small, tubular 2-lipped. 2 *Helleborus*

Petals small, stamen-like. 8 *Actaea*

Flowers irregular.

Upper sepal spurred; petals 4.

7 Delphinium

Upper sepal hooded.

3 Aconitum

1. *Caltha* L. Marsh Marigold

Herbs with heart-shaped or kidney-form leaves, flowers yellow, white or pink, sepals large, 5-9, petal-like, petals none, stamens numerous, pistils 5-10, styles nearly wanting, pods follicles, spreading, many seeded, marsh plants of temperate and colder regions. About 8 species, 3 species native to Eastern North America, and 1 species common in the Rocky Mountains at high altitudes.



Fig. 225. Black Hellebore (*Helleborus niger*). Entire plant. The roots contain a purgative substance that is poisonous in over-doses. (From Vaseau's *Traité de Botanique*.)

Caltha palustris L. Marsh Marigold

A stout, glabrous perennial with a hollow stem from 1-2 feet high; the basal leaves on long petioles, leaves reniform, upper leaves shorter, petioled and sessile; flowers with yellow sepals.

Occurs in swamps and meadows.

Poisonous properties. The marsh marigold or cowslip is regarded as poisonous in Europe. In this country, however, it is frequently used as a pot herb. The flower buds are sometimes pickled. Coville says

By many it is considered superior to any other plant used in this way. There is no doubt that boiling dissipates the active principles found in the plant.

Stebler and Schröter say that it is poisonous in a green state, and Rusby states that when fed with hay it produces diarrhoea and stoppage of the flow of milk. According to Lloyd, it contains a small quantity of an acrid substance identical with the acrid oil of *Ranunculus*. Cattle and sheep refuse

to eat the plant. Marsh marigold is known to contain an alkaloid which is said to be identical with *nicotia* but it has not been isolated.

Dr. Millspaugh in speaking of the uses of this plant, states that it is extensively gathered in early spring and cooked for greens, making one of our most excellent pot-herbs. Rafinesque asserts that cattle browsing upon it die in consequence of an inflammation of the stomach produced by it. According to Freidberger and Fröhner it causes hæmaturia.

2. *Helleborus* L. Hellebore

Erect perennial herbs, with large, palmately divided leaves, flowers large, white, greenish or yellowish, sepals 5, petal-like, petals small, tubular, stamens numerous, carpels generally few, fruit several seeded follicles. A small genus of about 10 species, natives of Europe and Western Asia.

Helleborus viridis L. Green Hellebore

Basal leaves smooth, consisting of 7-11 oblong, acute, sharply-serrate segments; flowers large.

Distribution. In waste places from Long Island to Penn. and W. Va.

Poisonous properties. Black Hellebore is said to be a drastic purgative when used for domestic animals. The plant contains the glucosides, *helleborin* $C_6H_{10}O$, which is a highly narcotic, powerful poison, *helleborein* $C_{27}H_{44}O_{10}$, which is slightly acid and *heleboresin* $C_{18}H_{30}O_2$.

The symptoms from poisoning are Stupor followed by death with spasms. *H. foetidus* is also poisonous.

3. *Aconitum* L. Aconite

Perennial herbs with palmately lobed or divided leaves, flowers large, irregular, showy, paniculate or racemose, sepals 5, irregular, petal-like, the upper helmet-shaped or hooded, prolonged into a spur, petals 2, small concealed under helmet, spurred, 3 lower absent or very minute; pistils 3-5 forming follicles, several seeded.

About 60 species. Native of the North Temperate regions. One species, the *A. Napellus*, used in medicine, is the source of aconite. One western species is poisonous to live stock.

None of the species of this genus is weedy in Eastern North America. The three species, *A. noveboracense* Gray, *A. uncinatum* L. and *A. reclinatum* Gray, occur in Eastern North America but are very local. None of these seems to be very poisonous, but the roots of *A. uncinatum* are bitter, even in a dry state. The exotic *A. Lycotomum* and *A. Fischeri* are employed to kill wild animals.

Aconite is derived from the European *Aconitum Napellus* which produces not only poisonous stem and leaves, but also a very poisonous tuberous root which is from 2-4 inches long and sometimes an inch thick. The Indian aconite is obtained from *Aconitum ferox*, a plant growing from 3-6 feet high and bearing large dull-blue flowers, it is found in the Alpine regions of the Himalayas, and is used as an arrow poison. Among other equally poisonous species mentioned by Flückiger and Hanbury, is *A. uncinatum* growing in Eastern North America. They also state that the root of another species of *Aconitum*, *A. heterophyllum*, with large flowers of dull yellow, and purple, or blue, is poison-

ous. This root contains the chemical substance *aconin* $C_{23}H_{31}NO_9$, an intensely bitter alkaloid. The European aconite contains *aconitin* $C_{38}H_{49}NO_{12}$, *pseudoaconitin* $C_{36}H_{47}NO_{12}$, and *aconus* $H_{39}C_{39}NO_{11}$. The North American species *A. septentrionale* contains, according to Rosendal, the following alkaloids: *lappakonin*, $C_{34}H_{48}N_2O_8$, a crystallized form, *septentrionalin* $C_{34}H_{48}N_2O_8$, and *synaktonin* $C_{38}H_{51}N_2O_{12}$.

Blyth who has collected records of poisoning in Europe by aconite states that there have been two cases of murder, seven suicidal, seventy-seven more or less accidental, six were from the action of the alkaloid, ten from the root, and in two cases, children ate the flower, in one case, the leaves of the plant were cooked and eaten by mistake, in seven cases, the tincture was mistaken for brandy, sherry or liquor, in the remainder of the cases the tincture, the liniment or the extract was used. The Indian species are much used, especially *A. ferox*, which is applied to poison stock and arrows, the latter to destroy the wild animals. It is a common practice to mix a decoction of the root with water or food.

Dunstan and Anderson* summarize the alkaloids obtained from *Aconitum* as follows:—"The first, a toxic group, of which the type is ordinary aconitin, contains alkaloids which are diacyl esters of a series of poly-hydric bases containing four methoxy groups, the aconines."

"The members of this group are:—

Aconitin from *Aconitum napellus*.

Jupaconitin from *Aconitum deionorrhysum*.

Bisbaconitin from *Aconitum spicatum*.

Indaconitin from *Aconitum charmantum*."

These are all highly poisonous.

The second group is the aasin group which contains *asinin* from *A. heterophyllum* and *palmannin* from *A. palmatum*. These are non-poisonous alkaloids.

Aconitum columbianum Nutt. Western Aconite

An erect, stout perennial, 3-6 feet high, more or less pubescent above, with short, spreading or viscid hairs; divisions of the leaves broadly cuneate and toothed, lobed; flowers purple or white, in a loose terminal raceme; hood variable in breadth and length of beak.

Distribution. Grows at an altitude of 5000-10,000 feet in low grounds, near brooks and springs, from Montana, Wyoming and Colorado to the Sierras.

Poisonous properties. The chief effect of aconite results from its influence over the heart and blood vessels. It decreases the force and frequency of the cardiac pulsations. After long continued use, aconite affects the nervous system causing the loss of sensation; bodily temperature is also reduced after medicinal uses of the drug. Dr Winslow, in his *Veterinary Materia Medica and Therapeutics*, speaking of its toxicology says

The minimum fatal dose of aconite is about $\frac{1}{2}$ gr. for the horse gr. xx for medium sized dogs and gr. v. vi. for cats. The smallest fatal dose recorded in man is a teaspoonful of tincture of aconite, equivalent to about gr. xxx. of the crude drug. The minimum lethal quantity of aconitin is gr. $\frac{1}{10}$ for man, and about the same for cats. For dogs it is from gr. $\frac{1}{4}$ to gr. $\frac{1}{5}$. The writer has found that cats will die from fifteen minutes to half an hour after receiving the smaller deadly doses under the skin, but large doses produce death immediately by paralyzing the heart. Large therapeutic doses cause, in horses, restlessness, pawing the ground, shaking of the head, clamping of the jaws, increased secretion of saliva, mucus, and attempts at swallowing, probably owing to the peculiar sense of irritation

*Trans. Jour. Chem. Soc. 1905: 1650. See Blyth Poisons: Their Effects and Detection.



Fig. 224. Aconite (*Aconitum Napellus*).
The source of the aconite of commerce.
Stem, leaves and root are poisonous.
(After Faguet.)




Fig. 223. *Aconitum* (*Aconitum columbianum*) a, flowering plant, b, seed capsule — both one-third natural size. A poisonous plant of the Western U. S. (U. S. Dept. Agr.)



Fig. 224. Western aconite (*Aconitum uncinatum*). A poisonous plant in Eastern North America, from Pennsylvania to Iowa. (Ada Hayden.)

produced by the drug in the throat. Nausea and retching are observed in all animals, while vomiting occurs in dogs and cats. The pulse and respiration are weakened and generally retarded. After lethal doses these symptoms are intensified. We observe violent retching, frequent and difficult attacks of swallowing, ejection of frothy mucus from the mouth in horses copious sweating, pulse first weak and infrequent, later rapid, running and almost imperceptible, respiration slow, interrupted, and shallow, and reduction of temperature. Death is preceded by muscular twitchings, in the horse, and loss of strength so that the subject falls and is unable to rise; or in the case of cats and rabbits, the animals jump vertically into the air, topple over backwards and go into convulsions, lying helpless on their side. The labial muscles are retracted and the lips drawn back showing the teeth covered with foam. The face is anxious, the eyeballs are retracted or protruded, and the pupils more commonly dilated. Death takes place usually from asphyxia, occasionally from syncope. The post mortem appearances are simply those resulting from asphyxia.

The western aconite is bitter and retains its bitterness even on drying. It also benumbs, according to Lloyd, just as does the European aconite. The Lloyds quote Prof Power in asserting that it contains some alkaloids, one probably *aconitin*, and several other poisonous principles. According to Dr

Bartholow of Jefferson Medical College, it is a paralyzer of mobility, but does not impair the contractility of muscles or the irritability of the motor nerves. Death is caused by paralysis of respiration, the heart continuing its action some time after respiration has ceased. The *Aconitum Napellus*, affects the heart in opposite ways. Prof Chesnut says in regard to the poison of this plant

All of the parts of the west American aconite are poisonous, but the seeds and roots are the most dangerous. The active principle is not well known, but chemical and physiological experiments point to the existence of one or more alkaloids which resemble aconitin. The effect of the poison is characteristic. There is first a tingling sensation on the end of the tongue which gives rise shortly to a burning sensation and is rapidly followed by a very pronounced sense of constriction in the throat. The choking thus produced is made the more alarming by the retarding effect which the poison has upon the respiration. The tingling and prickling over the entire body is also characteristic. Besides these symptoms, there are generally severe headache, abdominal pains, confused vision, vomiting, and diarrhoea. Delirium is usually absent. Death comes from a stoppage of the respiration in from one to eight hours.

Mr R. Schimpfky, in his Important Poisonous Plants of Germany, makes this statement upon the authority of a physician. He tried the nectar of the European aconite by chewing the flower. After chewing a little while, the same was thrown away and an hour later, he felt upon the end of the tongue, a dull pain as though he had burned it. This sensation remained for three days. In Europe it is not unusual to mix the leaves of this with other salad plants. Frequently the plant is cultivated to be used in destroying insects.

Dr. Chesnut says

No specific antidote is recognized, but physicians have used atropin, or digitalis and nitrite of amyl, with good effects. The ordinary emetics and stimulants must be given. Artificial respiration should be maintained for a couple of hours, if necessary and a recumbent position must be maintained throughout the treatment.

Aconitum uncinatum L. Wild Monkshood

Plant smooth, stem slender and somewhat reclined, root thickened; leaves 3-5 lobed petioled, lobes ovate lanceolate, with coarse teeth, large blue or pale flowers with erect helmet, found in rich shady woods along streams.

Distribution Eastern North America extending into Iowa

Poisonous properties. Contains the same active principle as the other species of aconite.

4. *Anemone*

Erect perennial herbs, root leaves lobed, divided or dissected, stem leaves forming an involucre either remote or near the flower, sepals few or many, 4-20 petal-like, or in one section, petals stamen like, stamens numerous pistils numerous achene pointed or tailed, flattened, single seeded. About 80 species in temperate regions. Several as *Pulsatilla* are medicinal.

Anemone patens L. var *Wolfgangiana* (Bess.) Koch

A perennial herb, with radical leaves, appearing after the bluish or purplish flowers have blossomed, vilous with long silky hairs, flowers erect, coming from a simple stem, which is naked except for the involucre, petals wanting or abortive, stamen-like, sepals petal-like, about 1½ inches long, leaves ternately divided with the lateral divisions 2-parted, stamens numerous, pistils numerous in a head with long, hairy styles, in fruit forming feathery tails.

Distribution. Illinois and Nebraska, Wisconsin, Minnesota, Missouri, N W Territory, Rocky Mountains to Texas, British Columbia. Known as the pasque or sand flower, but very commonly and incorrectly called the crocus.

Poisonous properties. A very poisonous plant. The allied European *Anemone Pulsatilla* is also regarded as poisonous. The different parts of the plant are extremely acrid and when applied to the skin cause irritation and vesication. The acidity of the plant is due to the presence of a crystalline substance called *anemonin* $C_{10}H_8O_4$ which when heated with acids, forms *anemonic acid* $C_{10}H_8O_6$. Lloyd states

A. parts of fresh *Anemone patens* are acrid and very irritating. Dr W. H. Miller informs us that his hands have been very badly blistered, in consequence of the juice having splattered over them while pressing the plant. The vapors evolved from the fresh juice are of such an acrid nature as to have inflamed the eyes, and have closed them temporarily. For this reason, persons refuse to work with the fresh herb, and botanists have been known to severely irritate their hands simply from contact with the recent plant.

The only demand for this plant is by Homeopathic physicians. All portions of the European *Anemone patens* are very acrid, but the dried plant is merely an astringent. The plant evidently contains a volatile acid substance, which is given off when heat is applied. Our sand flower was one of the chief medical plants of the Indians of Minnesota. The plant is still used, when in a fresh condition by the Indian. Dr. Millsbaugh gives the following method for preparing it

The whole, fresh, flowering plant is chopped and pounded to a pulp and weighed. Then two parts by weight of alcohol are taken, the pulp thoroughly mixed with one-sixth



Fig. 227. European Anemone (*Anemone Pulsatilla*) A poisonous plant with acrid properties. (From Vesque's *Traité de Botanique*.)



Fig. 228. Pasque Anemone (*Anemone patens*, var. *Woffgangiana*) A plant with well known acrid properties. (Charlotte M. King.)

part of it, and the rest of the alcohol added. After thorough mixture, the whole is allowed to stand eight days in a well-stoppered bottle. The tincture thus prepared, after straining and filtering, should have a light, seal-brown color by transmitted light, an acrid astringent taste, and a decidedly acid reaction.

Dr. White in his *Dermatitis Venerea* calls attention to the irritating properties of the common wind-flower *A. quinquefolia*. The species is widely distributed in woods in Eastern North America. He states that a large wholesale dealer in medical plants regarded it as an externally corrosive poison. It is probable that other species of the genus are more or less acrid. Some of these plants like the white meadow wind flower *A. canadensis* are probably looked upon with some suspicion. Dr. Johnson in his *Manual of Medical Botany*, makes this statement in regard to the common Wind-flower

Pulsatilla is an acrid irritant which, in large doses, has often produced serious and alarming effects. In safe medicinal doses, however, its effects are by no means so well known. At various times and by numerous authors it has been highly praised as a remedy in diseases of the eye, in rheumatism, amenorrhoea, dysmenorrhoea, etc. In this country it has been employed chiefly by homoeopathic practitioners, and usually in very minute doses. Many of the results claimed for it under such circumstances are at least doubtful. Certain it is that other practitioners have not been able to confirm them.

Anemone quinquefolia L. Wind Flower

A low smooth perennial with filiform rootstock, involucre or 3-petioled, trifoliate, toothed leaves, sepals 4-7, ovate, white, pale blue or purple; carpels 15-20 oblong with a hooked beak.

Distribution. In woods from Nova Scotia to Georgia and the Rocky Mountains, also in Europe.



Fig. 229. Wind flower (*Anemone quinquefolia*). A well known plant with more or less acrid properties. (Ada Hayden.)

Poisonous properties. Leaves, roots, and stems are acrid, same properties as in the Pasque Flower

Anemone canadensis L. Meadow Anemone

A hairy perennial from 1 ft. to 18 inches high, involucre 3-leaved bearing a long peduncle or a pair with a 2-leaved involucre at the middle radical leaves long petioled, 5-7 parted or cleft, sepals white 6-9 lines long, head of fruit globose achenes flat, tipped with stout style.

Distribution. In low grounds, meadows, especially westward from Labrador to Saskatchewan, Colorado to Maryland and New England.

Poisonous properties. All parts of the plants with acrid properties

5. *Clematis*

Climbing vines or perennial herbs, more or less woody, leaves opposite, slender petioled, pinnately compound, lobed, or entire, sepals 4 or rarely more, petal-like, petals none; stamens numerous, ovaries free, the fruit an achene 1-seeded, style long, persistent, plumose, silky, or occasionally naked. About 100 species of wide distribution, abundant in temperate regions. The most common species in Northern States is the *Clematis virginiana*, in the South and Rocky Mountains, *C. ligusticifolia*, and in the South, the *C. Flornia*, and *C. Pitcheri*. The European *C. vitalba* contains *anemone*!

Clematis virginiana L. Virgin's Bower

Perennial, climbing, with leaflets mostly broadly ovate, acute, cut low; flowers axillary, clusters paniced, polygamo-dioecious, white, the style persistent, plumose. The western *C. ligusticifolia* is nearly like this species.

Distribution. From Canada to Florida, and Kansas, Nebraska, northward. *C. ligusticifolia* is common throughout the Rocky Mountains from Western Nebraska to the Pacific Coast.

Clematis Pitcheri Torr. & Gray

Perennial herb with pinnately compound leaves, high upper leaves often simple, flowers large, solitary, on long peduncles, usually nodding caryx, bell-shaped, dul. purplish sepals, with narrow and slightly margined recurved points, tails of the fruit naked or shortly villous. From Southern Indiana to Central Iowa, and Kansas and Texas.

Poisonous properties. J U and C G Lloyd, in their *Drugs and Medicines of North America*, report the medical properties of several species of the genus *Clematis* as follows

It imparts a rank taste, which, after prolonged chewing, becomes acrid and irritating, although at first it is only disagreeable.

Dr Rushy also mentioned the poisonous character of the species of *Clematis*. In Cuba, one species of the genus is used in case of tooth-ache to blister the face, and this as well as another species is used in the same way for rheumatism. Dr J C. White refers to the European *Clematis recta* as producing blisters and often ulcers, and causing the eyes to water and become inflamed.

An infusion of the plant in oil has been used to cure the itch, and violent inflammation of the skin has been produced by friction with it.



Fig. 230. Virgin's Bower (*Clematis virginiana*). Probably poisonous. (From Johnson's Medical Botany of N. A.)



Fig. 231. Clematis, Virgin's Bower (*Clematis puberula*). The flowers are dull purplish in color; the plant is more or less acrid. (Ada Hayden.)

6. *Ranunculus* (Tourn.) L. Crowfoot Buttercup

More or less acrid, annual or perennial, herbs with alternate, simple, lobed, divided, or dissected leaves, solitary or somewhat corymbose flowers, yellow, white or red, sepals 5, petals 5 with a nectariferous pit or scale at the base, stamens numerous, pistils numerous, achenes numerous in heads, generally flattened or pointed with an erect seed, tipped with the style. About 200 species in temperate and colder regions.

Ranunculus acris L. Dutch Crowfoot

A glabrous annual, 1 foot or more high, stem hollow, root leaves 3-lobed, rounded, stem leaves 3-parted, the lobes cut and toothed, upper leaves nearly entire, flowers pale yellow, petals but slightly longer than the calyx, stamens numerous, pistils numerous, in oblong and cylindrical heads.

Distribution. Wet ditches in the Northern States also in Europe.

Poisonous nature. The plant is highly acrid, blistering the mouth and skin. In Europe it is used by beggars for making sores. Other species with similar properties are *R. acris*, *R. bulbosus*, and *R. repens*.

Ranunculus septentrionalis, Pur. Creeping Crowfoot

A low, hairy, or nearly smooth, glabrous perennial, ascending, or often producing long runners, leaves with 3-stalked divisions, the terminal one broadly



Fig. 232. Curled Crowfoot (*Ranunculus sceleratus*). Common in low grounds. Contains an acrid poison. (Ada Harkness.)

wedge-shaped, 3-cleft or parted; flowers yellow, petals obovate and larger than the sepals, stamens numerous; pistils numerous, style long and attenuate; fruit an achene.

Distribution. Common in moist shady places in the northern states.

Ranunculus acris L. Tall Crowfoot

Herb, perennial, with fibrous roots, from 2-3 feet high, basal leaves tufted 3-7 divided, divisions cleft in narrow acute lobes, upper leaves short petioled, 3-parted, petals much longer than the spreading calyx, head when in fruit globose.

Distribution. Native to Europe, widely naturalized in the Northern States and Canada.

Poisonous nature. Juice acrid, poison dissipated on drying. Symptoms produced in animals are blistering, slaving, choking, vomiting, in some cases, followed by death resembling that from apoplexy.

Ranunculus abortivus L. Small Flowered Crowfoot

Smooth, branching biennial from 6 inches to 2 feet high, root leaves round, heart-shaped or kidney shaped, crenate, or lobed, stem leaves sessile

or nearly so, divided into oblong linear lobes; flowers small, sepals 5, reflexed, petals 5, yellow, head globose, carpels mucronate with a minute curved beak.

Distribution. In moist woods and meadows a troublesome weed. Newfoundland to Manitoba, Nebraska, Colorado and Florida.

Poisonous properties. The leaves of the plant have an acrid, peppery taste and cause blistering.



Fig 233. Buttercup (*Ranunculus acris*). The juice of this plant has acrid properties. (Ada Hayden.)




Fig 234. Small-flowered Crowfoot (*Ranunculus abortivus*). The leaves cause blistering. (Ada Hayden.)

According to Eschner, the oil of *Ranunculus acris*, in warm-blooded animals, as an acrid narcotic, producing, in small doses, stupor and slow respiration, in larger doses, also, paralysis of the posterior and anterior extremities, and, before death, convulsions of the whole body. The acrid action is shown by a corrosive gastritis and by hyperæmia of the kidneys, more particularly of their cortical substance. *Anemone* causes similar symptoms, but is followed by no convulsions, nor does it irritate sufficiently to corrode the organs, as the oil does.

Dr. Millsbaugh mentions especially the *R. bulbosus* as having a peculiarly powerful irritant action upon the skin, whether applied locally or internally.

Murray states that a slice of the fresh root (bulb?) placed in contact with the palmar surface of a finger brought on pain in two minutes; when taken off, the skin was found without signs of extra circulation or irritation, and the itching and heat passed away, in two hours it nevertheless returned again, and in ten hours a serious blister had formed, followed by a bad ulcer, which proved very difficult to heal.

Stebler and Schröter state that *Ranunculus acris* produces diarrhoea, abortion, and loss of flesh, and when eaten in large quantities, death ensues in a few hours. Dr Johnson in his *Manual of Medical Botany of North America*, says

The ranunculi are too acrid to render their internal use either desirable or safe. Most of them are avoided by domestic animals; one may often see *R. acris*, for example, growing luxuriantly in pastures where almost every blade of grass is cropped close. Their acrid properties have, however, led to their employment externally as rubefacients or vesicants in cases where other and perhaps better agents were not at hand, or were for any reason contra-indicated. As is well known, cases of misadventure occur in which cantharides are inadmissible on account of their effect upon the urinary organs. In some such cases ranunculus has been used with good effect. One of the fruits of this agent is its extreme violence. The fresh plant, bruised and applied to the skin, may vesiculate in an hour or hour and a half, and may possibly produce an ulcer not easy to heal. It is, therefore, far less safe as a rubefacient than mustard, and, as a rule much less desirable as a vesicant than cantharides. It has been employed to some extent in European countries as an external application in chronic rheumatism, neuralgia, etc. but never sufficiently to have obtained a place in the pharmacopoeia. In this country it is used still less, and is little more than mentioned in works on materia medica. An interesting observation regarding the possible effect of *R. acris* on pregnant cows was reported to the author by his brother Mr F M Johnson. In a herd of cows pastured for years in succession in an old field thickly beset with this weed, abortion was frequent and troublesome. As soon, however, as this pasture was broken up and the herd moved to another part of the farm in which the plant did not grow, abortion disappeared. Now although as stated above, domestic animals avoid this plant, yet when feeding where it is very abundant, they must occasionally swallow it accidentally and though there is no positive proof that the abortions were due to the plant in question, the facts as stated are interesting and significant. It is at least possible that ranunculus exerts an influence upon the reproductive organs like that which is claimed by some for pulsatilla.

Leaves, flowers, and stems of the Ranunculi have a peppery and pungent taste when eaten, reminding one of mustard. According to Lloyd, boiling water dissipates the acrid principle.

Many other species of the genus *Ranunculus* such as *R. Flammula*, and *R. arvensis* are acrid and poisonous, causing the formation of blisters.

Delphinium Tourn

Perennial or annual herbs of erect, branching habit, with racemose or paniculate showy flowers, leaves palmately lobed or divided. sepals 5, irregular, petal-like, the upper one prolonged into a spur. petals 2-4, irregular the upper one spurred, and enclosed in the spur of the calyx. stamens numerous. pistils 1-5, forming follicles in fruit, many seeded. Species about 60 in the North temperate regions.

Several species like Stavesacre (*D. Staphisagria*) are used in medicine, and many are poisonous to live stock. Several species like the field larkspur (*D. Consolida*) the rocket larkspur (*D. Ajacis*), the great flowered larkspur (*D. grandiflorum*) and blue larkspur (*D. elatum*), are ornamental.

Delphinium Penardi Hoth. Prairie Larkspur

A perennial, pubescent or hairy herb, more or less glandular above, from 3-5 feet high, leaves 3-5-parted, the divisions 2 or 3 times cleft, the lobes linear. flowers in racemes. white, slightly tinged with blue, spur horizontal straight or slightly curved upward, follicle pubescent many seeded. This is closely related to *D. azureum* of more southern distribution, with light blue flowers and downy follicles.

Distribution. The Prairie Larkspur is common on sandy soil, gravelly knolls and prairies of Illinois and Wisconsin to Manitoba, Kansas and Arkansas.

7 *Delphinium Geyeri* Greene

A hairy perennial 1 2 feet high, leaves dull green, somewhat branched; flowers in dense racemes, azure blue.

Distribution. Common on the high plains of Colorado to Montana.



Fig. 235. Carolina Larkspur (*Delphinium penardii*). Like other species of this genus it is poisonous. Common on prairies and gravelly soil. (Ada Hayden.)



Fig. 236. Purple Larkspur (*Delphinium bicolor*) is found in Montana and Westward. Poisonous. (Chas. H. U. S. Dept. Agr.)

Delphinium Menziesii DC

Glabrous below at least at the very base, pubescent above with spreading hairs, especially the inflorescence, leaves 5-parted, divisions 2 to 3-cleft, flowers large, deep-blue, in a loosely few to many-flowered simple raceme, upper petals veined with purple, spur long and slender; ovaries somewhat tomentose.

Distribution. Common from San Francisco, California, north to British Columbia, Idaho, Montana and Wyoming.

Poisonous properties. Chestnut and Wilcox say:—

Experiments have been made which show that at one stage of growth the leaves of the species (*D. Menziesii*) may be safely eaten, to a certain extent, by sheep. Dr. S. B. Nelson, professor of veterinary science in the Washington Agricultural College, in an article entitled *Feeding Wild Plants to Sheep*, published by the Bureau of Animal Industry of this Department showed that it is possible to feed as much as 24½ pounds of the fresh leaves of *D. Menziesii* to a sheep within a period of five days without causing any apparent ill effects. The stage of growth of the larkspur was not stated, but, judging from the other experiments described in the same report, it was probably in a well-advanced flowering stage.

In regard to the poisoning from this species, opinion seems to differ. Dr. Nelson states that Dr. Wilcox was in error in regard to the plant that he worked with at first. More than likely it was a *D. bicolor*, which is corroborated by the report. Under *D. bicolor*, mention is made of *D. Menziesii*. However, it is more than likely these are certain stages in the development which are more poisonous than others. Dr. S. B. Nelson from his experiments concludes as follows:

The results obtained in these eight trials with *Delphinium Menziesii*, the feeding of nine and one-half pounds of the bulbs, stems and leaves of the immature plants to sheep No. 6, the feeding of nearly twenty-five and six pounds of the plants gathered while in full bloom, to Nos. 2 and 3 respectively, and the hypodermic injections of the two extracts into Nos. 4 and 5, certainly constitute evidence sufficiently convincing to justify the conclusion that *Delphinium Menziesii* is not poisonous to sheep and they may be allowed to graze where it grows even in abundance without fear of any harm from it.

Dr. Nelson likewise carried on an experiment with *D. simplex* with similar results.

Delphinium bicolor Nutt. Purple Larkspur

A smooth or somewhat pubescent, tuberous rooted, perennial, 1-2 feet high with a cluster of finely divided leaves, the lower orbicular in outline, all deeply cleft or parted; racemes few or several-flowered; flowers dark purple, sepals and spur ½ to ¾ inch long, upper petals pale yellow, and white with blue veins, follicles smooth or minutely pubescent when young.

Distribution. Common in dry ground, Eastern Oregon, and Washington, to Utah, British Columbia, Montana, and Colorado. Grows in elevations from 4000 or 5000 feet to 10,000 feet.

Poisonous character. It is regarded as poisonous by stockmen. Experiments reported by Dr. Wilcox in the Montana Agricultural Experiment Station Bulletin show that an extract from less than one ounce of the dried leaves of this species was fatal to a yearling lamb. Chestnut and Wilcox report further experiments as follows:

The following experiments were made with a view of ascertaining whether the permanganate of potash, which it was proposed to use in many cases of plant poisoning, could be used with satisfactory results as a chemical antidote in case of this plant, and also of determining which parts of the plant were most toxic. During some seasons the purple larkspur causes extensive poisoning of sheep and calves. Cattle and horses, on the other hand, seem to eat it less frequently than the tall larkspur. In the season of 1900 conclusive evidence against this plant was obtained in only one locality. This was derived from two cases among calves in the Flathead Valley. The calves were about 5 weeks old, and at the time when the poisoning occurred, were running in a native pasture where the purple larkspur grew sparingly. The symptoms of poisoning in these two cases were similar to those already outlined from poisoning in the tall larkspur in cattle, with the exception that a slight bloating was to be observed in the case of the calves. The respiration and heart beat became exceedingly rapid as the symptoms of poisoning increased in severity. The body temperature was slightly lowered, and this was accompanied by pro-

fuse sweating. The increased perspiration may have been due in part to the violent spasms in which the animals finally died. Death occurred about four hours after the appearance of the first symptoms. No remedy was applied in these cases.

Delphinium scopulorum Gray

A glabrous or finely pubescent perennial with leafy stem 1-6 feet high from fascicled thick roots, leaves numerous orbicular, 5-7, parted, lower cuneate, and the upper consisting of narrow, cleft, and lacinate divisions racemes many flowered, sparingly pilose, flowers blue varying to white or pink on short erect pedicels spur longer than sepals, lower petals deeply notched, and upper whitish, and a little shorter than the oblong sepals; follicles about half an inch long, erect, seeds small, with a loose coat.

Delphinium occidentale Watson. Tall Larkspur

A glandular pubescent perennial from 4-6 feet high, leaves deeply 3-5 cleft, divisions broadly cuneate, somewhat 3-lobed, flowers numerous in a many-flowered sparingly-branched panicle, sepals spatulate, acuminate, dull or dark blue, very variable in size, seeds light colored, and somewhat spongy.

Distribution. At higher altitudes from Colorado to Eastern Oregon, and Nevada.

Delphinium troilifolium Gray California Cow Poison, or Poisonous Larkspur

A tall smooth perennial, 2-5 feet high, sparingly villous, hairy, leaves large, long, petioled, 5-7 lobed, lobes lacinate cleft and toothed with acuminate segments; flowers large in loose racemes, color bright blue, $1\frac{1}{4}$ inches broad, spur as long as the sepals, sepals oblong lanceolate, acuminate, sparingly villous, follicles smooth, 6-8 lines long, seeds turbinate.

Distribution. Common along the Pacific Coast from British Columbia to California.

Poisonous properties Prof Chesnut says "In Humboldt County, Cal., it is known as Cow Poison on account of its fatal effect on cattle. Its toxic character has been questioned. Perhaps it is not equally poisonous throughout all stages of its growth."

Delphinium tricorne Michx.

A simple stout perennial 6 inches-2 feet high, with a cluster of roots, leaves slender, petioled, deeply 5-7-cleft, or divided, the divisions linear or ovate, flowers in loose racemes, blue, occasionally white or purple, spur slightly bent. Follicles tipped with a short beak.

Distribution. In open rich woods or clay soil, Western Pennsylvania to Southeastern Iowa to Arkansas and Georgia.

Poisonous properties The Stavesacre (*Delphinium Staphisagria*) of Italy and Greece to Asia Minor has long been used in medicine, having been known to the ancients. Pliny mentions the use of the powdered seeds for destroying vermin on the head. It is still largely used for destroying pediculi. The eclectic physicians use it for its specific action on the reproductive organs. The disease produced by *Delphinium* may be called delphinosis.

According to Prof Hills, Stavesacre and *D. Consolida* are used in the treatment of dropsy and spasmodic asthma. The effects produced are due to



Fig. 237. California Cow Poison (*Delphinium troitiforme*). A species common along the Pacific coast, and said to be fatally poisonous to cattle. (Ada Hayden.)




Fig. 238. Tall Larkspur (*Delphinium glaucum*) of the Pacific Coast region. It is poisonous. (Chestnut, U. S. Dept. Agri.).

the alkaloid *delphinin* $C_{22}H_{35}NO_8$, very poisonous with a bitter sharp taste. Three other alkaloids have been isolated, *staphisagrin* $C_{12}H_{22}NO_5$, bitter, the poisonous *delphinoidin* $C_{42}H_{82}N_2O_7$, and *delphuin* $C_{23}H_{46}N_2O_6$, an extremely poisonous alkaloid, to which may be added the substance *caicitripin*. Dr J. C. White in his *Dermatitis Venerata* states that acute dermatitis resembling eczema may appear from the use of stavesacre seed. *Desphincurarin* $C_{25}H_{30}NO_7$, has been obtained from the root stock of several species of *Delphinium*, *D. bicolor* contains 0.27 and *D. scapularum*, 1/3 per cent.

In this country, it appears, from Mr Cheney's observation that *D. consolida* is largely sold for the European plant. A tincture of the seed is often mixed with *Lobelia inflata* and sold as a parasiticide. The Stavesacre seeds are still employed as in old times for the destruction of pediculi in human beings. For this purpose, they are converted into powder and dusted among the hair.

Results of the experiments made by Dr Crawford of the U. S. Dept. of Agr. show in 1905 in regard to larkspur poisoning. (The first batch of plants was collected April 26th, 1905)

1 c. c. injected into a guinea pig (subcutaneously), weight 730 grams. Caused no disturbance.

3 c. c. in guinea pig, no symptoms.

6 c. c. in guinea pig, killed.

6 c. c. injected into guinea pig, 285 grams, killed in 33 minutes.

4 c. c. injected into guinea pig, 352 grams, no symptoms.

Repeated

5 c. c. killed guinea pig weighing 196 grams. Died in 55 minutes.

4 c. c. injected into guinea pig, 299 grams. No symptoms.

Evidently lethal dose for this solution lay between 4 to 5 c. c.

Second Stage, Gathered May 16th, 1905

Solution corresponding to 4 c. c. of No. 1 caused no symptoms in guinea pig weighing 445 grams, while 5.3 c. c. killed one of 350 grams, but death was delayed longer than with extract of first stage.

Third Stage, Gathered in June, 1905

Solution corresponding to 4 c. c. caused no symptoms in guinea pig weighing 376 grams.

5.3 c. c. caused no symptoms in guinea pig weighing 500 grams.

6.6 c. c. caused no symptoms in guinea pig weighing 480 grams.

Evidently a lethal dose is much higher and the plant loses much of its activity in development.

This report is very conclusive in proving that the plant contains an active poison, and further in substantiating the claims of experienced observers that the plant loses much of its toxic properties as it approaches the flowering period.

Seven and one-half grams of dried purple larkspur fed to each of three rabbits on April 20th. No results.

Seven and one-half grams of fresh purple larkspur from same patch fed April 25th to each of three rabbits. Two showed slight uneasiness, and one was bloated a little. One, showing less effect than the others, had eaten but three and one-half grams.

On May 1st a like quantity from the same patch was given to the same rabbits under similar conditions. Results, two died, and the other distressed.

On June 15th, plants from the same source, being in full bloom, but the leaves and stems dry were fed to rabbits. Although very hungry, they at first refused to eat, but later ate large quantities of it without any ill effects. The experiments with tall larkspur were equally as confusing. The fact that the plants at one period of growth gave negative results was no guaranty that they would not be dangerous at another. The tall larkspur growing luxuriantly on the college campus proved to be very active, physiologically and furnished the best specimens for producing the physiological effects upon animals. In the experiments with antidotes this domesticated species was found to be very poisonous while in bloom in the middle of August.

Lloyd in *Drugs and Medicines of North America*, quotes from a letter from Wm. C. Cusick, a botanist of the West, who states that in some places the *D. decorum* F. & M. var. *nevadense* poisons cattle, but only in the early spring when they are first turned on the crop. It is thought by cattle men that the cattle pull the plants up by the roots and eat them, which really causes the

poisoning. Many die, but most recover. Prof. Chesnut, in his *Principal Poisonous Plants of the United States*, says

The percentage of fatal cases in cattle which have eaten this and other larkspurs is said to be small. A rough estimate by a cattleman places it at about 20 per cent for one species of the group, when the animals are not properly treated, and 5 per cent other wise. This is probably a low estimate, however, for in a case of poisoning from *D. Menziesii* that occurred in Montana in May, 1892 and was reported by Dr. E. V. Wilcox, nearly 400 sheep were affected, 250 of which died.

Dr. Wilcox says in regard to *D. glaucum*

The tall larkspur appears not to be eaten by sheep. All cases of poisoning from this plant observed in Montana during this year and previous years have been among cattle. As already indicated, sheep are not driven to the mountain ranges until about the middle of July, and at this season the tall larkspur is altogether too large and coarse for their consumption. It is well known, on the other hand, that cattle will feed on much coarser forage than sheep, and at the same time they are allowed to run on the high ranges in the early spring. In Montana the light green tufts of leaves of the tall larkspur first become conspicuous about the last of April or the first of May and the flowers begin to open about the middle of June. This fact is significant for the reason that light frosts of snow often occur in the larkspur belt as late as the first week in June, and, since there is then no other verdure in sight, the uncovered portion of the larkspur is in a high degree tempting to stock, all the more because it is succulent. The danger is increased by the fact that at the time of snow falls cattle seek the shelter of creeks where the larkspur is more abundant and most advanced in growth. At this period, moreover, the leaves are, as we can verify from personal observation, very bitter and they are probably then more poisonous than at any other stage of growth. The older ones, as is also the case with the blue larkspur, are not so bitter. The plant is recognized by cattlemen as dangerous to cattle from May west about the middle of June. Mr. Vard Cochran informs us that on his range in the lower basin of the Gallatin it is the main duty of our men during this period to keep his cattle away from the Indian mountainous regions where this larkspur abounds.

The symptoms are described as follows

In general, the animals affected manifest symptoms similar to those produced by overdoses of strychnine. The first signs of poisoning are usually a general stiffness and irregularity of gait. There is often a pronounced straddling of the hind legs in walking. These symptoms increase in severity until locomotion becomes difficult or impossible, and the animal finally falls to the ground. It usually falls and gets on its feet again a number of times, the muscular movements becoming more and more irregular and uncoordinated. At the same time the skin is very sensitive to touch, and the muscles of the sides and legs soon begin to quiver spasmodically. This is a very characteristic symptom, being usually exhibited for several hours. The function of the special senses is seldom impaired, the animal being apparently able to hear and see as well and as correctly as under conditions of health. Although a slight increase in the quantity of saliva is to be noticed in some cases, this symptom is never so pronounced as in cases of poisoning by death camas. During the later stages of poisoning the animal is usually attacked with violent convulsions, in one of which it finally dies. In this respect, also, the symptoms differ distinctly from those of death camas poisoning, which is usually quite without spasms. The digestive functions seem not to be affected by larkspur poisoning. The temperature is lowered slightly during the first stages, in one instance having been as low as 97°. During the later stages the pulse becomes very frequent and the breathing rapid and shallow. The cerebral symptoms are simply those of excitement, and the appetite seems not to be lost until shortly before death.

In regard to *D. bicolor*, the symptoms observed from experiments are described by the same author as follows

Summing up the results of these experiments, we find that the most prominent, easily observable symptoms were a stimulation of the respiration and a brain symptom manifested by dizziness or a rhythmical movement of the head. As it was not the object of the experiment to determine the symptoms of poisoning, these having been already secured by Dr. Wilcox, no special effort was made to determine the pulse rate, but it was to be noted that in experiment 2, which was as nearly fatal as any, the heart action was extremely rapid and weak. None of the experiments proved fatal and no results were obtained on which

a satisfactory trial of the permanganate of potassium as an antidote could be based, hence no experiments were made in that line.

Dr. B. Kennedy in the Nevada Agricultural Experiment Station, reports in regard to the symptoms of poisoning from *D. glaucum*, as follows:

This dangerous, poisonous plant has been quite frequently from Lincoln Valley to Summit Soda Springs. It occurs usually in large patches by the side of creeks in the valleys. Although sheep do not care for it, yet it is sometimes eaten by them. Sheepmen do not consider it dangerous. Cattlemen, however, are afraid of it, and consider it very poisonous. Mr. Ridinger of the Tehachapex dairy, about eight miles from Webber Lake, told us that cattle die after eating it, with the following symptoms: Trembling and shivering, succeeded by extreme weakness, which makes affected animals stagger and fall or lie down.

That Larkspur is poisonous, at least some species, appears from the experiments carried on by Dr. Geo. H. Glover and C. Dwight Marsh.

There can be no question but that the several species of larkspur growing native in the mountainous districts of Colorado are a greater source of loss to the stockmen than all other weeds combined. While the larkspur is confined to the mountainous regions, it nevertheless holds large share in the aggregate mortality throughout the state from poisonous plants; larkspur takes second place only to locoweed. We have no statistics at hand whereby we can estimate, with any degree of accuracy, the total loss, but judging from the reports of other western states and from information received from most every section of the state, it would seem that \$40,000 annually is a conservative estimate. There are four species of larkspur found growing abundantly in the middle and western portion of this state, and one found growing sparingly in the eastern plains section. Other species have been found in isolated places, but have not been especially accused of doing any harm, and their toxicity has not been proved. The four species found in the greatest abundance and named in the order of their importance, are purple larkspur, *Delphinium Nelsonii*, Greene, tall larkspur, *Delphinium elongatum*, Rydb., *D. Geyeri*, Greene, and *D. Barbeyi*, Nutt. These all have the same characteristic flowers, and are found growing in the mountains at altitudes from 5,000 to 11,000 feet. The *D. Amabile*, Nutt., has a white flower and may be seen growing adjacent to streams and in the arroyos on the plains as far east as the state line.

The *Delphinium elongatum* and *D. Nelsonii* are the only ones which contain a sufficient quantity of deleterious substance to produce poisoning. From all accounts it seems probable that the plants are most poisonous in the spring when they are fresh. When the poison has been absorbed into the system atropin is an antidote.

8. *Actaea* L. Baneberry

Perennial herbs with 2 or 3 ternately-compound leaves, sepals 4-5, petal-like, soon falling, petals 4-10, small, flat, spatulate, on slender claws, stamens numerous, free, with slender white filaments, ovary solitary, stigma sessile, fruit a berry, seeds compressed, smooth, horizontal.

Actaea rubra (Ait.) Willd. Red Baneberry

A smooth perennial 1-3 feet high with ternately divided leaves, on long, smooth petioles, leaflets ovate, sharply cut, and toothed. Calyx with 4 greenish sepals, corolla 8-10 petals, white, shorter than the stamens, stamens numerous, berries cherry red.

Distribution. Eastern North America to Hudson Bay, and the Rocky Mountains.

Poisonous properties. Prof. Chesnut states that sheep are occasionally poisoned by eating the leaves of the closely related European species. These plants are seldom, however, eaten. In its medical action *Actaea* is similar to *Cimicifuga*. It contains a resinous body which is neither acrid nor bitter, but accord-

ing to Fred Stearns, the rhizome has purgative properties. This, according to Mr Lloyd, may be somewhat overdrawn. Prof. Sayre states that it is a violent purgative, irritant and emetic. The berries are known to be somewhat poisonous.

9. *Hydrastis* Ellis. Yellow Puccoon

A low perennial herb with knotted yellow rootstock, and a single root; stem bearing 2 leaves near the summit; flowers large, greenish-white, 3 sepals, petal-like, soon falling; petals none; pistils 12 or more, 2-ovuled, fruit 1 to 2-seeded berry, crimson in color. A genus of 2 species

Hydrastis canadensis L. Golden Seal. Yellow Puccoon

Rootstock 1 and 2 inches long, knotted; berries numerous, small, fibrous root, roots and rootstocks yellow in color; leaves pubescent, palmately 3-5-lobed; calyx petal-like.

Distribution. From New York to Southern Michigan, Southern Wisconsin and Eastern Iowa to Arkansas to Northern Georgia.

Poisonous properties, also medicinal properties. The plant contains the alkaloid berberin $C_{20}H_{17}NO_4$ and hydrastin $C_{21}H_{21}NO_6$, a so-called alkaloid, but which, according to Mr. Lloyd, cannot be considered in the pure condition; also canadine $C_{20}H_{21}NO_4$ and xanthopurcin, it also contains a fixed oil of a disagreeable odor and taste, and a black resinous substance. It produces ulceration and catarrhal inflammation of the mucous surfaces. The plant acts very similarly to *Cimicifuga*. The Lloyds have given an extended account of the anatomy, structure and therapeutical properties of this plant. The alkaloid berberin $C_{20}H_{17}NO_4$ is identical with the substance found in the barberry, according to Prof. Power.



Fig 220. Golden Seal (*Hydrastis canadensis*). The rootstocks of this plant as well as the stem and leaves contain more or less acrid substances. (Charlotte M. King.)

BERBERIDACEAE. Barberry Family

Shrubs or herbs, with alternate leaves, stipulate or exstipulate flowers either solitary or in racemes, perfect, stamens as many as petals and opposite them, hypogynous, fruit a berry or capsule. A small family of about 20 genera and 105 species, widely distributed in the North Temperate region, also in Temperate South America and Asia. Some of our North American species are weedy, others woody and several are more or less poisonous. A few like the common



Fig. 240. Common Barberry (*Berberis vulgaris*). The spines often inflict mechanical injuries while the leaves and wood are more or less poisonous. (Ada Hayden)




Fig. 240a. Single flower. s, stamen, n, anther, p, pistil, u, nectar gland.

barberry (*Berberis vulgaris*), blue cohosh or pappoose root (*Caulophyllum thalictroides*), the twinleaf (*Jeffersonia diphylla*), and mandrake (*Podophyllum peltatum*) are used in medicine. The root of *Berberis aristata* and *B. vulgaris* is sometimes used as a fish poison. Probably some other members of this order are poisonous. The blue cohosh (*Caulophyllum*) contains saponin. This plant is said to be extremely bitter to the taste, but is not, however, common.

Key for genera

- | | | |
|--------------------------------|-----------|--------------------|
| Herbs with simple large leaves | . . . 1 | <i>Podophyllum</i> |
| Shrubs with unifoliate leaves. | | 2. <i>Berberis</i> |

1. *Podophyllum* L. Mandrake, May-apple

Perennial herbs with simple, smooth, erect, stem; creeping rootstocks and thick fibrous roots; stems bearing 2 leaves with large flowers, flower buds with 3 green bractlets, 6 fugacious sepals, petals 6-9; stamens twice as many

as the petals; pistil 1; stigma large, flat, sessile, fruit a large fleshy berry, 1-celled and many-seeded, each seed enclosed in a pulpy aril.

4 species, one in eastern North America and the others in India and Eastern Asia. The *P. emodi* of Asia contains the same principles as the American species and is poisonous.

Podophyllum peltatum L. Mandrake

Perennial herb, with creeping rootstocks and thick fibrous roots, flowering stems with 2-leaves, 1-flowered, the flower bud with 3 small green bractlets, which fall away early, calyx of 6 unequal sepals, corolla white, of 6-9 petals, about twice the length of the sepals, stamens 12-18, twice the number of petals, inserted below the pistils, with short stamens, anther cells opening longitudinally creeping rootstock, from 1-5 feet long, fibrous rooted.

Poisonous properties. Its medicinal virtues were well known to the Indians of North America, and an early writer, Catesby, remarked that the root was an excellent emetic. It has a bitter, acrid taste, similar to that of roots of other plants of the family. Its active properties seem to reside in the resinous substance. Prof. Power failed to find an alkaloid. The name *podophyllin* has been given to the product found in the resinous substance contained in other members of the family. This in turn contains *podophyllotoxin* $C_{31}H_{44}O_9 + 2H_2O$, and *peripodophyllin* $C_{31}H_{44}O_9 + H_2O$, producing a very bitter taste; and intensifying the action of *podophyllin*, *Berberin* $C_{20}H_{11}NO_4$, which is feebly toxic to man, and *asponin* have also been found. Cases of poisoning have occasionally



Fig. 241. Mandrake (*Podophyllum peltatum*).
powerfully purgative. (Ada Hayden).

been reported. It produces fatal prostration. It seems to exert a special influence upon the liver. Dr Rusby says

Its taste, especially when fresh, is very repugnant, and yet if eaten in quantities it would unquestionably prove fatal, as shown by the effects of over-dosage in medicine. In the Philadelphia Medical and Surgical Reporter, XIX, 308, a fatal case is recorded in which the evidence is perfectly clear that poisoning resulted from continued large doses administered by an ignorant and careless physician. The poisonous symptoms were all referable to the bowels, those of enteritis. It is also very interesting to note the peculiar effects of poisoning of the external skin by the powder and by the resin of this drug. It produces an ulcer of a very peculiar character, closely resembling one of venereal origin. Serious errors of diagnosis, leading to the gravest injustice to the reputation of the patient, have been known to occur in reference to these cases. A very serious ulcer upon the eye-bail, is among these recorded cases. The very greatly elongated rhizome of *Podophyllum*, with its very long smooth internodes, broadened nodes with their very large, low cup-shaped scars, and sparse roots underneath, is doubtless well known to all pharmacists. The plant is not only very common, but extremely abundant east of the Mississippi and is liable to be encountered almost anywhere."

In regard to its action on man, Dr. Millspaugh says

Here the same action takes place, but extends to the rectum with sufficient intensity to cause prolapsus and hemorrhoids. The first effect of the drug is an excitation of salivary and biliary secretions, followed by torpor and icterus. The symptoms of disturbance caused by the drug in doses varying from $\frac{1}{4}$ to $\frac{1}{2}$ grains of "podophyllin" and in persons working in the dust of the dried root, are substantially as follows. Inflammation of the eyes, soreness and puckering of the nose, salivation and white-coated tongue, extreme nausea, followed by vomiting severe pains in the transverse colon and abdomen, followed by an urgent call to stool thin, offensive, copious stools weak pulse, prostration, drowsiness, and cold extremities.

Among other experiments with this drug upon animals, those of Dr. Ansell seem to be the most characteristic. He found, resulting from his many applications of an alcoholic solution to the peritoneal cavity direct, that no local inflammation arose, although an intense hyperaemia occurred in the duodenum externally, and the whole of the small intestine, even going so far as to cause a breaking down of the tissues and resulting ulceration, causing discharges of glairy mucus streaked with blood, this hyperaemia ceased usually at the ileocaecal valve. Post mortem. The mucous membrane were found inflamed and covered with bloody mucus. Other observers noted that retching, salivation, and emesis, followed by purging, colic, and intense tenesmus, with low pulse and rapid exhaustion followed the administration of the drug.

Dr Schaffner says

Roots, stems, and leaves, drastic and poisonous, but the ripe fruit is not. Leaves, when eaten by cows, produce injurious milk. The ripe fruit may be eaten in small quantities.

The root of Mandrake affects the skin. Mr Lloyd, in White's work writes:

Our employees experience great trouble in working this, owing to the irritating action of the skin. We have in numerous instances had our men cease work for several days owing to its action, which causes very painful inflammation of the skin, excoriation of the eyes.

Dr Winslow says

The action is exerted mainly on the duodenum, which is intensely inflamed and even ulcerated in poisoning. *Podophyllum* directly increases the secretion of bile in small doses, while purgative quantities hasten its excretion by stimulation of the muscular coat of the gall bladder (except in the horse) and small intestine. It is probable that the intestinal secretions are somewhat augmented. The faecal movements, after medicinal doses of podophyllin, are liquid, often stained with bile, and may be accompanied by some nausea and griping.

Berberis L. Barberry

Shrubs with yellow wood, simple or compound leaves, often spiny, flowers, yellow in racemes, or rarely axillary, sepals 6-9, like petals, petals 6, imbricated in 2 series, stamens 6, fruit a berry with 1-3 seeds. About 75 species of wide distribution.

Berberis repens Lindl. Trailing Mahonia.

A smooth, trailing shrub, 1-4 ft. high, leaves petioled, pinnate, leaflets 3-7, ovate, acute, flowers several in a raceme, yellow, persistent bracts, fruit globose, bluish purple.

Distribution. From Western Nebraska to Arizona, and British Columbia, Northwest Pacific Coast from Washington to California.

Berberis Aquifolium Pursh. Oregon Grape

A low shrub 2-10 feet high, leaflets 5-9 oblong ovate, spinulose dentate above, flowers yellow in racemes and terminal clusters, fruit globose, dark in color

Distribution. Idaho to the Rocky Mountains.

Poisonous properties. Both *Berberis repens* and *B. Aquifolium* contain the alkaloids berberin, oxycanthin $C_{15}H_{13}NO_3$, and berbamin $C_{15}H_{13}NO+2H_2O$

Prof. Schaffner reports that the berries of the trailing Mahonia are injurious to birds. When eaten fresh they are emetic and cathartic.



Fig. 242. Oregon Grape (*Berberis Aquifolium*). Berries are said to be poisonous to birds. (Ada Hayden).

MENISPERMACEAE. Moonseed Family

Woody plants with alternate lobed or entire leaves, climbing without stipules, flowers small, dioecious, in panded racemes or cymose clusters, sepals 4-12 petals 6, fewer or more, stamens of the same number or fewer, fruit a 1-seeded drupe embryo long, curved endosperm scanty. About 300 species mainly in the tropics.

Moonseed (*Menispermum canadense*) is a beautiful native climber of the North with black drupes and contains menisperm. The Carolina moonseed (*Cocculus carolinus*) is common in the South. Fish poison made from *Anamaria*

pauiculata is used for destroying vermin and to poison fish. The wood of this species is very bitter to the taste and contains an alkaloid *menispermum* $C_{20}H_{34}NO_4$ and a toxic substance *picrotoxin* which is said to be a mixture of two bodies *picrotoxin* $C_{23}H_{36}O_8$ and *picratm* $C_{26}H_{42}O_7$. The Columba Root (*Jateorrhiza Columba*) is used as a medicine and contains several alkaloids, among them, *berberin* and *columbamin*. The false Columba Root (*Cascinum fenestratum*) produces a yellow dye. The alkaloid *pelonin* is obtained from *Chondrodendrum tomentosum*.




Fig. 244. Laurel Magnolia (*Magnolia virginiana*). A well known flowering shrub or small tree of the Southern states, well known as an ornamental plant.

Fig. 243. Moon-seed (*Menispermum canadense*). The fruit is blue, much the color of Concord grapes. Plants like the fig poison belong to the same order are known to be poisonous. This plant has been suspected of being poisonous. (Charlotte M. King)

MAGNOLIACEAE. Magnolia Family

Trees or shrubs, leaves alternate, leaf buds covered by membranous stipules, flowers large solitary sepals and petals hypogynous, colored alike stamens numerous, adnate anthers, carpels numerous, separate, or coherent, packed together, ripening into an aggregate fruit, seeds 1-2 in each carpel, achenes or folicles, endosperm fleshy, embryo minute. About 70 species of North America and Asia. The bark of several species used in medicine.

The family *Magnoliaceae* contains the well known magnolias planted for ornamental purposes. Among these are the great flowered magnolia (*Magnolia*

grandiflora) which has evergreen leaves and fragrant flowers and is extensively planted in the South, the sweet bay (*M. virginiana*) a shrub or small tree with fragrant flowers, the cucumber tree (*M. acuminata*) the umbrella tree (*M. tripetala*) the Yulan tree (*M. Yulan*) of China and Japan, the purple magnolia (*M. obovata*), the tulip tree (*Liriodendron Tulipifera*) frequently planted as an ornamental tree and producing the most valuable timber of the family, the star anise (*Illicium floridanum*) of the south has aromatic bark and pods. The fruit of *I. anisatum* of Japan, found growing around Buddhist temples, furnishes the poisonous *akumin*, from the same species in China is made the liquor anisette, Anise is furnished by *I. anisatum*.




Fig. 245. Crest-flowered Magnolia (*Magnolia grandiflora*). An ornamental tree of the southern United States. (After Faguet). The odor of the flowers said to be injurious to some people.

Poisonous properties. The crystalline substance *magnolin*, a glucoside, and a volatile oil occur in the large leaved Magnolia (*M. macrophylla*) of the south. The tulip tree contains the bitter principle *liriodendrin*, also an alkaloid, and a glucoside. The *Talanum macrocarpa* of Mexico contains a haemolytic substance capable of dissolving the red corpuscles of the blood. The flowers of *Michelia nitagrica* are used in perfume. The winter's bark (*Drimys Winteri*) is used in medicine.

Illicium anisatum of Japan is said to contain a poison belonging to the *pyrotosin* class, says Blyth. In 1890 five children in Japan were poisoned by eating the seeds of this plant; three died. After considerable experimentation

Dr Langaard concluded that all parts of the plant were poisonous. The poison causes excitation of the central apparatus of the medulla oblongata and clonic convulsions analogous to those produced by *pyrolotoxin*, *toxwesia* and *cicutoxin*. Small doses kill by paralyzing the respiratory center. Large doses cause heart paralysis. When animals are poisoned by small doses chloral hydrate is an efficient remedy but has no effect when large doses have been taken.

CALYCANTHACEAE

Shrubs with entire short petioled opposite leaves, without stipules, flowers fragrant, large, solitary, on leafy branches, sepals and petals numerous, stamens numerous, the inner short, sterile, pistils numerous, fruit of an ovoid pyriform receptacle, enclosing few to many smooth solitary achenes, seed erect. About 5 species of North America and Asia. Several species are cultivated for ornamental purposes.

Calycanthus L. Carolina Allspice

Flowers purple or red; sepals and petals numerous, stamens numerous, pistils numerous. A small genus of 4 species, 3 in Eastern North America, and 1 on the Pacific Coast.

Calycanthus floridus L. Strawberry Bush

A branching shrub 3-9 feet high, branches pubescent; leaves oval, soft, downy underneath, roughish above, flowers dark purple, with the odor of strawberries, sepals and petals linear-oblong.



Fig. 246. Strawberry Bush (*Calycanthus floridus*)
Commonly used as an ornamental shrub and is known
to be somewhat poisonous. (Charlotte M. King).

Distribution. In rich soil, North Carolina to Georgia, Alabama and Mississippi, cultivated in Missouri, north to Central Iowa.

Poisonous properties. This plant contains several alkaloids among them *calycanthin*. Chesnut says:

The large oily seeds of the *calycanthus*, or sweet-scented shrub are strongly reputed to be poisonous to cattle in Tennessee.

ANONACEAE. Papaw or Custard Apple Family

Trees or shrubs, generally aromatic, leaves entire, alternate, stipules absent, flowers with calyx of 3 sepals and a corolla of 6 petals in 2 rows hypogynous, anthers adnate, filaments very short, pistils many, separate or cohering in a mass, fleshy or pulpy in fruit, seeds large with a hard seed-coat, small embryo and copious endosperm. About 550 species, many in the tropics.

Annona Adans. North American Papaw

Shrubs or small trees with solitary flowers from the axils of the leaves of the preceding year, sepals ovate, petals 6, imbricated in the bud, pistils few, ripening into 1, large and oblong, pulpy, several seeded fruits, seeds horizontal, flat. A small genus of about 7 species, natives of eastern North America and of Asia.

A. triloba Dunal. American Papaw

Shrubs or small trees with thin, obovate-lanceolate leaves, petals dull purple.

Distribution. Along streams from Ontario and New York to McGregor, Iowa, Nebraska and Texas.



Fig. 247. Common Papaw (*Annona triloba*). Common in woods of the South. (Charlotte M. King)

Poisonous properties While the papaw is edible, there are some cases of poisoning on record. Care should therefore be exercised in its use. The leaves contain alkaloids. The papaw (*Polyalthia argentea*) contains *aninin*, a tasteless amorphous alkaloid.

LAURACEAE

Aromatic trees and shrubs with alternate or rarely opposite leaves without stipules, flowers small, fragrant, polygamous, dioecious or monoecious, calyx 4-6 parted, corolla absent, stamens in 3 or 4 series on the calyx, some imperfect, fruit a 1 seeded berry or drupe, endosperm none. About 1000 species, mostly tropical.

The spice bush (*Benzoia gestivale*) produces fragrant flowers and aromatic leaves. Camphor $C_{15}H_{12}O$ is obtained from *Cinnamomum Camphora* which comes from the islands of Formosa and Japan, cinnamon, a well known spice, comes from *C. zeylanicum* which is extensively cultivated in Ceylon, cassia is from *C. Cassia*. Cassia and cinnamon were well known to the ancient especially to the



Fig. 248. Cinnamon Tree (*Cinnamomum zeylanicum*) Flowering and fruiting branch. The cinnamon of commerce is the bark of this tree. (After Faguet)

Israelites who used them as incense on their altars. The oils derived from these plants are excellent antiseptics.

Cinnamomum contains a volatile oil *camphor* which is found in the roots and leaves, *eugenol* occurs in the leaves, stems and bark. Clove bark is derived from a small Brazilian tree (*Dicypellum caryophyllatum*) belonging to this family. The alligator pear or avocado (*Persea gratissima*), a native of the West Indies and tropical America, is much esteemed as a dessert fruit and in making salads. Roofs has written of its successful culture in Florida. It is said that the oil is used extensively in America in soap manufacture. Leaves of laurel (*Laurus canariensis*), native to Canary and Madeira Islands, and bay (*L. nobilis*) of southern Europe are used in culinary processes. *Acciandra Rodiei* contains *berberin*, which is identical with the *pelosin* $C_{16}H_{21}NO_3$ of *Cissampelos Pongia*. The Indian laurel contains *laurotetanol* $C_{14}H_{23}NO_3$. The California laurel (*Umbellularia californica*) is a strong local anesthetic and to be irritant and acrid. The leaves, according to the Indians, will drive flies away.

Sassafras Nees

Trees with spicy aromatic bark, small mucilaginous twigs and foliage. Flowers greenish yellow, naked in racemes, calyx 6-parted spreading, sterile, with 9 stamens, 3 inner with pair of glands, fertile flowers with rudiments of stamens. ovoid blue drupes.




FIG. 249. *Sassafras* (*Sassafras variifolium*). A well known tree of the south which furnishes the sassafras oil and bark of commerce. (W. S. Dodgson).

Sassafras varifolium (Salisb.) Ktze. *Sassafras*

Leaves oval and entire, mucilaginous, or 2-6 lobed to about the middle and often as wide as long, membranous, pinnately veined, petioled, stamens about equaling the calyx-segments, fruiting pedicles red, much thickened below the calyx.

Distribution. From E. Mass. to S. E. Iowa, Kansas and southward.

Poisonous properties. It is said to be poisonous, its bark however is used medicinally as a tonic and its wood is valuable.

RHOEADALES

Mostly herbs with regular and perfect flowers, sepals and petals usually present, polypetalous, stamens free, ovary superior, free from the calyx, compound, composed of 2 or more united carpels. It contains the families *Resedaceae*, *Cruciferae* and *Papaveraceae*. The mignonette (*Reseda odorata*) is a well known, cultivated fragrant plant of the family *Resedaceae*, which also includes the dyers weed (*R. luteola*), the latter contains the substance *luteolin*.

Families of the Order Rhoeadales

Sepals generally 2; endosperm fleshy *Papaveraceae*.

Sepals or divisions 4-8, endosperm none

Capsule 2-celled, sepals and petals 4, flowers regular, stamens, usually tetradynamous *Cruciferae*.

Capsule 1-celled, sepals and petals 4, flowers regular or irregular *Capparidaceae*

PAPAVERACEAE. Poppy Family

Annual or perennial herbs, with milky or colored juice, leaves alternate, stipules none perfect, regular, or irregular flowers, sepals 2, occasionally 3, falling when the flower expands, petals 4-12, spreading, soon falling stamens inserted under the pistils, distinct, pistil 1, many ovuled, chiefly 1-celled, fruit a capsule containing numerous oily seeds. Genera 24-26, species about 200. Widely distributed chiefly in north temperate zone.

Comparatively few of the plants of this family are weedy and quite a number are medicinal and poisonous. The common poppy (*Papaver somniferum*) is used in medicine. It is found as an escape near buildings, especially in sections where Germans have settled, undoubtedly due to the fact that they cultivate it for its beauty as an ornamental plant and use the seeds in culinary operations. The poppy is largely cultivated in China, Smyrna, Joppa, and several countries of Europe and India, for the opium. Opium yields a large number of alkaloids. The more important of these are *morphin*, and *codein*. A perfectly harmless oil equal to olive oil is obtained from the seed. The seed is also fed to birds. The red poppy (*Papaver Rhoeas*) is sometimes cultivated. A syrup is made from the petals, and also a coloring matter used in red ink. The California poppy (*Eschscholtzia californica*) is a valuable soporific, and analgesic "free from the disadvantages of opium." The Indians, according to Chesnut, use it to stupify fish. Celandine (*Cheidomum majus*) native to Europe, has been naturalized in places in the East and is occasionally somewhat weedy. The juice of this plant is yellow, while that of the common cultivated poppy is white, and that of the blood root (*Sanguinaria canadensis*) is reddish. The




Fig. 250. California Poppy (*Eschscholzia californica*) a, flower, b, fruit before, and, c, after dehiscence. The juice of this plant is a valuable soporific. (After Strasburger, Noll, Schenck and Schimper).

rhizome of the blood root is used in medicine and contains an alkaloid *sanguinarin* and a dye. The *corydalin* is found in a species of the genus *Dicentra*. The bleeding heart (*Dicentra spectabilis*), native to China, and the climbing fumitory (*Adlumia cirrhosa*) are frequently cultivated for ornamental purposes. According to Elyth, the root of the tuberous-rooted corydalis (*Corydalis tuberosa*) contains eight alkaloids, of which *corydalin* $C_{22}H_{27}NO_4$ is the most important, since, when taken in large doses it may cause epileptiform convulsions, death taking place from respiratory paralysis. The *C. lutea* contains *corydalin*. Schlotterbeck and Watkins found 5 alkaloids in the American celandine (*Styliphorum diphyllum*) among them *chelidonin* $C_{26}H_{31}NO_4 + H_2O$. The alkaloids *stylopin* $C_{18}H_{19}NO_3$, *protopin* $C_{20}H_{21}NO_3$, and *sanguinarin*, have been in part found in other plants in the family.

Genera of *Papaveraceae*

Petals 8-12, pod 1-celled 2-valved.

Petals white, rootstock short red

3 *Sanguinaria*.

Petals 4, pod 2-valved or more.

Flowers yellow

4 *Chelidonium*.

Pod 4-20 valved.

Ovary incompletely many celled

1 *Papaver*

Stigmas and placentas 4-6

2 *Argemone*.

1. *Papaver*. Poppy

Plant with milky juice, leaves lobed or dissected, alternate, flowers and buds nodding, sepals 2 or occasionally 3, petals 4-6; stamens numerous, ovules numerous stigmas united into a persistent disk; capsule globose, obovoid or oblong, seeds small, with minute depressions. About 25 species, natives mostly of

the Old World. *P. nudicaule* is found in high mountains in the Rockies and in the Alpine regions of Europe and Asia.

Papaver somniferum L. Garden Poppy

An erect glaucous herb, leaves clasping, large, oblong, wavy, lobed or toothed, flowers broad, bluish-white with purple centre, filaments somewhat dilated, capsules smooth.

Distribution. Native to Asia, but widely naturalized in Europe, and extensively cultivated in China, India, and Smyrna. Occasionally found spontaneous around gardens in North America.

Poisonous and medical properties. From the milky exudation that comes from making an incision in the unripe capsule, opium is obtained which yields not less than 5 per cent of crystallized morphin and occasionally as high as 22 per cent in Turkey opium, the usual yield being between these two extremes. The chief markets for opium are Turkey, Asia Minor, India, and Egypt, that of Smyrna being considered to be the best although good opium has been grown in the United States. This opium has a sharp, narcotic odor, and a bitter taste. Opium has been a fruitful source of a large number of alkaloids. Flückiger and Hanbury enumerate the following: "*Hydrocodarin, morphin, pseudomorphin* ($C_{17}H_{19}NO_7$) $_2 + H_2O$, *codein, thebain* $C_{19}H_{21}NO_5$, *protopin, laudanin* $C_{20}H_{23}NO_4$, *codamin, papaverin* $C_{20}H_{21}NO_4$, *rhoeadin, meconidin, cryptopin, isodonin, narcotin* $C_{23}H_{27}NO_7$, *lanthopin, norcein* $C_{23}H_{27}NO_8 + 3H_2O$, *gnoscopin*." The most important of these are *morphin* $C_{17}H_{19}NO_3 + H_2O$, a colorless or white and shining odorless substance with a bitter taste, and *codein* $C_{18}H_{21}NO_3 + H_2O$, a nearly transparent odorless substance with a faint-



Fig. 252 Garden Poppy (*Papaver somniferum*). Flower and capsule. Opium is made from the milky juice exuding from the unripe capsule. (Pursey)

ly bitter taste which occurs in amounts varying from 0.5-2 percent. *Narcotin* is found in quantities varying from 0.75-9 percent.

It may be of interest in this connection to state that the German chemists Wolfgang, Weichardt, and Stadlinger found toxins in opium. These writers expressed the opinion that the complex physiological action is due to these substances.

In regard to the properties of opium, Fluckiger and Hanbury speak as follows:

Opium possesses sedative powers which are universally known. In the words of Pereira, "It is the most important and valuable medicine of the whole *Materia Medica*;" and we may add, the source, by its judicious employment, of more happiness and, by its abuse, of more misery than any other drug employed by mankind.

There are occasionally cases of poisoning from the poppy plant. Certainly cases from overdoses of opium are frequently recorded in the annals of medical jurisprudence.

Opium may be absorbed to a slight extent by the unbroken skin, according to Winslow, and causes a mild, anodyne action. Opium diminishes the two principal activities of the digestive organs, namely secretion and motion. The action upon the alimentary tract is lessening secretion, in part a local one and partly constitutional, following the absorption of the drug. The mouth is made dry, thirst is increased and appetite impaired. Opium is absorbed rather slowly from the stomach and bowels, and stimulates the splanchnic nerve centre of the sympathetic system, which inhibits the movements of the stomach and intestines, and thus lessens peristaltic action of these organs. Opium is directly opposed to belladonna in this respect, as the latter drug paralyzes the intestinal inhibitory apparatus (splanchnic endings), and so increases peristalsis.

The most important action of opium is upon the nervous system, and its influence is more powerful upon man than upon lower animals. At first, opium exerts a stimulating influence upon the spinal cord. Ruminants are comparatively insusceptible to opium. Dr. Winslow says "Ounce doses of the drug cause, in cattle, restlessness, excitement, hoarse bellowing, dry mouth, nausea, indigestion and tympanites. Sheep are affected in much the same manner. One or two drachms of morphine have led to fatality in cattle. Fifteen to thirty grains of the alkaloid comprise a lethal dose for sheep. Swine are variously influenced, sometimes excited, sometimes dull and drowsy.

According to the same authority, its action on horses causes drowsiness, sometimes, and at other times produces no visible effect.

Four to six grains, given in the same way, cause restlessness, a rapid pulse, and moisture of the skin. The animal paws the ground and walks in a rhythmical manner about the stall. The pupils are dilated. Large doses (12 grains) are followed by increased excitement, sweating, muscular rigidity and trembling while still larger doses (four drachms of the extract of opium) cause violent trembling, convulsions, insensibility to pain and external stimulation, without coma, or (morphine, at 36 under the skin) stupor for several hours (3 hours), dilated pupils and blindness. Followed by delirium and restlessness, continuing for a longer time (7 hours) and ending in recovery. Horses have recovered from an ounce of opium, but 2 3/4 ounces of the drug, and 100 grains of morphine have proved fatal.

Dr. Winslow is here quoted upon the toxicology of opium.

The symptoms of poisoning have already been sufficiently described in previous sections. The treatment embraces irrigation of the stomach, or the use of emetics, as apomorphine hydrochlorate under the skin, and the subcutaneous injection of strychnine and atropine in phosphate in the first stages, and enemata of hot, strong, black coffee leading the animal about snapping him, or using the faradic current. Dr. Moor of New York, has apparently found in potassium permanganate the most efficient antidote for opium and morphine. Ten to fifteen grains, dissolved in eight ounces of water should be given by the mouth, to large dogs. One to two drachms of potassium permanganate may be administered to horses in two or three pints of water. Permanganate solutions oxidize and destroy morphine, and should be acidulated with a little vinegar or diluted sulphuric acid, after the ingestion of morphine salts. The antidote has been recommended to be given subcutaneously after absorption, or hypodermic injection of morphine, but this is not of the slightest use. Hypodermic injections have not infrequently caused poisoning.

Papaver Rhoeas L. Corn Poppy

An erect annual with hispid spreading hairs, lower leaves petioled, the upper, smaller sessile, pennatifid, lobes lanceolate, acute, and serrate, flowers scarlet with darker center, filaments dilated, capsule smooth with 10 or more stigmatic rays.

Distribution. In waste places along the Atlantic and Pacific coasts. It is frequently cultivated.

Poisonous properties. Poisonous like other species of this genus.

Freidberger and Fröhner give the symptoms of poisoning from this plant as colic constipation, tympanites in cattle, raging fit of fury in horses.

In India this species is a troublesome weed but the seeds are collected and a yellow acid oil obtained therefrom which is used both in medicine and as an illuminant*.

2. *Argemone* L. Prickly Poppy

Herbs with yellow juice, spiny toothed leaves and stems, flowers large, sepals 2-3, petals 4-6, stamens numerous, styles short, stigma 3-6-radate capsule prickly, oblong, opening by 3-6 valves; seeds small, numerous.

A small genus of about 8 species of the southern states, Mexico and Western North America.

Argemone mexicana L. Mexican or Prickly Poppy

A glaucous annual from 1-3 feet high, with spines or without, leaves sessile, clasping by narrow base, glaucous, runcinate-pinnatifid, spiny-toothed; flowers large, whitish or generally yellowish, calyx with 2 sepals, bristly pointed; stamens numerous, stigma sessile, seeds numerous, reticulated.

Distribution. Introduced along the Atlantic coast as far north as the middle states. Native from Florida to Texas. A most common and troublesome weed in Texas. It yields however a valuable painter's oil.

Argemone intermedia Sweet. Prickly Poppy

A spiny, leafy plant from 2-2½ feet high setose, hispid, flowers large white, sepals green, hispid, petals obovate, capsule armed with stout spines, horns with a terminal spine, seeds numerous, black sunken meshes.

Distribution. From Central Kansas, Nebraska, South Dakota, to the Rocky Mountains and Texas. Common in dry soil.

Poisonous properties. The small prickles cause somewhat painful injuries when they penetrate the skin. According to Schlotterbeck the *A. Mexicana* contains *fumarin* $C_{20}H_{19}NO_6$ and *berberin*. In Mexico used in the same way as the poppy.

Sanguinaria L. Bloodroot

Perennial with a horizontal, thick rootstock, juice red, leaves basal, palmately veined and lobed, heart-shaped or reniform, flowers white, sepals 2, soon falling, petals 8-12, arranged in several rows, stamens numerous, placentae 2, capsule oblong, dehiscent to the base, seeds smooth, crested. A single species.

* D. Hooper. Agrl. Ledger 1907:35.

Sanguinaria canadensis L. Bloodroot

Calyx, sepals 2, light green, falling as the bud opens, petals 8-12 or more, $\frac{1}{2}$ to 1 inch long, oblong-spatulate, spreading, white or slightly rose-tinted, increasing in size for two or three days after the bud opens, and then falling away; stamens about 24, in several rows, much shorter than the petals, those in the inner rows longest; anther narrow, opening longitudinally

Distribution. In rich woods, N. S. to Manitoba, Neb., Fla. and Ark.

Poisonous and Medical Properties. Lloyd in White's book on dermatitis, writes.

There are two native drugs that are very irritant to mucous surfaces, so much so that the dust is very disagreeable, and we presume that they would have a similar irritating action on the skin. Bloodroot, and *Caulophyllum thalictroides*, blue cohosh or pappoose-root.

Bloodroot has a bitter and acrid taste due to the substance *sanguinarin*. In small doses, this substance exerts a tonic influence, promoting gastro-intestinal secretion and thus aiding digestion. On its physiological action, Dr. Millspaugh says of *sanguinarin* $C_{20}H_{12}NO_4$, "This alkaloid is very acrid to the taste, and toxic, and causes violent sneezing." Millspaugh gives its physiological action as follows

Sanguinaria in toxic doses causes a train of symptoms showing it to be an irritant; it causes nausea, vomiting, sensations of burning in the mucous membranes wherever it comes in contact with them, faintness, vertigo, and insomnia. It reduces the heart's action and muscular strength, and depresses the nerve force, central and peripheral. Death has occurred from overdose, after the following sequence of symptoms: violent vomiting, followed by terrible thirst and great burning in the stomach and intestines, accompanied by soreness over the region of those organs, heaviness of the upper chest with difficult breathing; dilation of the pupils, great muscular prostration, faintness and coldness of the surface, showing that death follows from cardiac paralysis.

Rusby says

The effects of *Sanguinaria canadensis* L., or blood root are distinctly poisonous and Johnson definitely records that fatal results follow overdose. Yet the rhizome is not at



Fig. 253. Blood root (*Sanguinaria canadensis*). The colored latex contains poisonous alkaloids.

all-ible to be eaten, on account of its peculiar blood red color, which is forbiddingly suspicious, and more especially because of an exceedingly acrid taste which would render the chewing and swallowing of a poisonous quantity an act of heroism. It is exceedingly common throughout the northeastern United States, and in a number of localities within a few miles of this city. The root also contains *chelerythrin*, *homochelidonium* and *protopas*.

4. *Chelidonium* L.

Erect branching herbs, with alternate deeply pinnatifid leaves yellow juice and flowers, 2 sepals, 4 petals, stamens numerous, distinct styles, capsule linear dehiscent to the base, seeds smooth, shining, and crested.

Distribution. A genus of one species, native to Europe, but widely naturalized in North America.

Chelidonium majus L. Celandine

Flowers consisting of 2 sepals which are ovate, yellowish, soon falling, corolla 4 petals, contracted at the base, stamens numerous, shorter than the petals.

Poisonous and Medical properties. The alkaloid *chelerythrin* $C_{21}H_{17}NO_4$ is identical with the *sanguinarin* of the last plant. *Chelidomin* $C_{80}H_{116}NO_8 \cdot H_2O$, an alkaloid existing particularly in the root, is colorless and bitter. *Homochelidonium*, consisting of three basic substances is found in *Baccharis*, *Sanguinaria*, *Adonis* etc. This plant produces congestion of the lungs and liver; it is also an excessive irritant, and has a narcotic action upon the nervous system in its action resembling gamboge. On this point Dr. White says:

Mr. Cheney informs me that he has known the plant to poison the skin, if handled so as to crush the leaves or stem. To indicate this extent to which it is used in medicine, it may be stated that a collector in North Carolina offers fifteen hundred pounds of the leaves for sale.

CRUCIFERAE. Mustard Family

Herbs or rarely woody plants with acrid, watery juice alternate leaves without stipules flowers in racemose or corymbose clusters, cruciform of 4 deciduous sepals and 4 petals, placed opposite each other in pairs, spreading and forming a cross; stamens 6, 2 shorter, 1 pistil, consisting of 2 united carpels fruit a pod either much longer than broad (siliqua), or short (silicle), or in-



Fig. 254. Common Celandine (*Chelidonium majus*). Poisonous to the skin. (After Pilch).

dehiscent, separating into joints, seeds without endosperm; seed coat frequently mucilaginous, embryo large. About 1500 species of wide distribution.

The cabbage (*Brassica oleracea*), native to Europe, has long been cultivated, cauliflower, brocoli, and brussels sprouts also belong to the same species. Rape, a well known forage plant, the turnip, the Swedish turnip, and rutabaga (*B. campestris*) are native to Europe. The Chinese cabbage (*B. Pe-Tsu*) is commonly cultivated in China. Black mustard (*B. nigra*) and white mustard (*B. alba*) are extensively cultivated for their seeds, which when ground make the commercial mustard. The radish (*Raphanus sativus*), cultivated for the root, is native to Europe. Water cress (*Radicula nasturtium-aquaticum*) is much used as a salad plant in colder regions. Horse radish (*R. Armoracia*) the well known condiment is native to Europe. European pepper grass (*Lepidium sativum*) is cultivated as a salad plant, while the seeds of our pepper grasses (*Lepidium apetalum* and *L. virginicum*) are used as bird food. Sea-kale (*Crambe maritima*), native to Europe, has also been introduced into this country as a vegetable. The *Pringlea antiscorbutica* of Kerguelens Land resembles the common cabbage and is used by sailors as a vegetable when they touch that country. The Rose of Jericho (*Anastatica hierochuntica*) of North America and Syria is regarded as sacred by the natives. Many plants of this family are cultivated for ornamental purposes, among the most familiar are the candytuft (*Iberis*), stock (*Matthiola incana*), sweet alyssum (*Alyssum maritimum*), and wall-flower (*Erysimum asperum*). Dyer's woad (*Isatis tinctoria*) of China was formerly cultivated for a dye obtained from the leaves. The characteristic odor of plants of this order when crushed or when mustard seed is ground in water is due to an enzyme myrosin, discovered by Bussy in 1839. This same ferment occurs in the families *Cruciferae*, *Capparidaceae*, *Rosaceae*, *Tropaeolaceae*, *Lamnaceae*, *Papaveraceae*.

It occurs in special cells known as myrosin cells which give a marked protein reaction. The contents are finely granular, free from starch, chlorophyll, fatty matter, and aleurone grains. These cells become red with Milon's reagent, and when heated become orange red, and a violet red color when treated with copper sulphate and caustic potash. The subject is treated fully by Reynolds Green in his work on "The Soluble Ferments and Fermentation," and in Effront and Prescott's "Enzymes and their Applications."

Some of the European plants of the family like (*Erysimum crepidifolium*) cause staggering in animals. The *Cheiranthus Cheri* contains a glucoside which acts on the heart. Rape under some conditions is poisonous. Several correspondents in Breeders Gazette (Chicago) have ascribed poisoning where rape was frozen, or when the plants were wet with dew.

Genera of *Cruciferae*

Pod terete or turgid or 4-angled.

Pod obovoid, flowers yellow

5. *Camelina*

Pod linear or oblong

Cotyledons accumbent.

Pod short, flowers yellow or white..

3. *Radicula*

Cotyledons incumbent.

Pod angled or terete, flowers yellow or white

1. *Sisymbrium*

Cotyledons conduplicate, flowers yellow or white

2. *Brassica*

Pod short.

Pod many or few seeded, obcordate-triangular

Pod many seeded, obcordate-triangular

Pod few seeded, orbicular, obovate or obcordate

Pod 2-seeded, flat, notched ,

4 Capsella

7 Thlaspi

6 Lepidium

1 *Sisymbrium* (Tourn.) L.

Annual or biennial herbs with usually simple spreading pubescent, hairs occasionally forked or stellate, leaves entire, oblanceolate, divided, pinnatifid, or runcinate flowers in racemes calyx open, greenish, of 4 sepals, corolla white, yellow or yellowish, or rarely pink, small, pistils terete flattish or 4-6-sided, small entire stigma, seeds small oblong, cotyledons incumbent

A small genus of 60 species. Found in temperate regions of both hemispheres. Several species are well known troublesome weeds.

Sisymbrium officinale Scop. Common Hedge Mustard

A slender erect annual or winter annual, 1½ 2½ feet high, lower leaves divided, runcinate, pinnatifid, upper entire or hastate at base, flowers small, yellow, borne in spike-like racemes, seeds small, brown, cotyledons incumbent.




Fig. 255. Tumbling Mustard (*Sisymbrium altissimum*). Common in Canada and from Minnesota to Washington. (Dewey, U. S. Dept. Agr.)

Distribution. A very common weed along railroads, door yards, and fields from Canada south to Florida and west to Illinois, Wisconsin, Minnesota, Missouri, Dakota, Nebraska and Kansas, and Pacific North coast. Common hedge mustard is a naturalized weed from Europe. It occurs throughout Europe, Russia, Germany, Great Britain, France—except Northern Scandinavia.

Sisymbrium officinalis L. Tumbling Mustard

An erect, much branched annual from 1-4 feet high, lower leaves pinnatifid, irregular toothed or wavy margined, upper leaves smaller, thread-like, after flowering, leaves drop, leaving the stem and pods. Flowers pale yellow rather large, sepals 4, green, corolla of pale yellow petals. Pods narrowly linear divergent, seeds small, longer than broad, generally oblong in outline with rather blunt ends, radicle usually very prominent and straight, curved spirally around the cotyledons.

Distribution. This weed has spread with considerable rapidity in the Northwest. Dr. Robinson states that it was once scarcely more than a ballast weed about the large cities of the Atlantic seaboard, and records its occurrence sparingly in southern Missouri (Bush). It is common now, however from the Mississippi Valley northwest to the Pacific Coast. One of the most common weeds of Montana, Idaho, Eastern Washington, Oregon, and native to British Columbia. Found also along the Atlantic seaboard.

Poisonous properties. This plant has properties somewhat similar to those described for Mustard, therefore may produce deep ulcers which are difficult to heal.

Brassica (Tourn.) L. Mustard, Turnip, Rape

Annual or biennial branching herbs, basal leaves pinnatifid, flowers yellow racemose, pods elongated nearly terete or 4 sided, seeds spherical, 1 row in each cell; cotyledons conduplicate. About 25 species of Europe, Asia, and North Africa, introduced in North and South America, Australia. The black mustard (*B. nigra*) and charlock (*B. arvensis*) common in grain fields and waste places across the continent.

Brassica nigra Koch. Black Mustard

A tall, coarse, much branched annual, 2-5 feet high, hairy or smoothish, somewhat bristly, at least on the veins, leaves variously divided or deeply cut, and sharp toothed—large terminal lobe, the upper leaves small, simple, usually linear, flowers yellow, smaller than in charlock, pods smooth, about ½ inch long, 4-cornered, tipped with a slender beak. Seeds black or reddish brown, smaller than in charlock, cotyledons conduplicate.

Brassica arvensis (L.) Ktze. Mustard or English Charlock

Branching annual from 1-3 feet high, hispid or glabrate, lower leaves petioled with 1 large terminal lobe, and several small lateral lobes, with the divisions unequal. Upper leaves barely toothed, flowers yellow, large and very fragrant, pods 1-2 inches long, irregular in outline, appearing somewhat nodose, 2-7 seeded, or more occasionally, the upper part of pod forms the beak. Seeds round, brownish black, darker than in *B. nigra*, and more minutely pitted. When moistened, the seeds become mucilaginous.

Distribution. It is a common and troublesome weed in cultivated ground from Mass. to Oregon. The most troublesome weed in grain fields of the Northwest.

mustard, no specific toxic symptoms have been noted, but in speaking of white mustard, he states that the essential oil of mustard is a virulent, irritant poison, causing when ingested, severe burning, followed by increased heart action and if pushed to extremes, loss of sensibility, paralysis, stupor, rigors, and death. It causes immediate vesication, followed by deep ulceration, which is difficult to heal.

Dr. White in his *Dermatitis Venenata*, says

The action of the sinapium is well known. In a few minutes after its application the skin begins to feel warm, and by the end of a half-hour, if the patient bear it so long, this sensation has increased to an intolerable burning. The changes in the cutaneous tissue are, within a few minutes, a considerable degree of hyperaemia, which, after a time, increases to an intense redness, which persists for a day or two, and often leaves behind it a persistent pigmentation, at times of a dark brown color, to mark the seat of the sinapium. On this account one should never be applied upon the upper chest or other part of a woman which the dress will not always conceal. If the action be continued beyond its legitimate rubefacient effect, a period which varies greatly in persons, it may produce vesication, or even deep suppuration, effects at times very intractable under treatment. It is stated that the addition of vinegar to a mustard poultice greatly lessens its activity.

Sarepta mustard (*B. Besseriensis*) much resembles black mustard and is used in the same way. Sarson or Indian colza (*B. campestris* var. *Sarson*) is used in India in place of white mustard and oil cakes made from it serve as stock food. A brown variety (*B. Napus* var. *dichotoma*) is grown both as an oil seed and as a vegetable. Another variety, Indian mustard (*B. juncea*) is also known as of economic importance.

The mustards of India are not always easy to distinguish. They have been carefully studied by Col. Prain.

Brassica Rapa causes inflammation of the bowels, tympanites, constipation, diarrhoea, some brain irritation, and haemorrhagic enteritis. The disease is more common in Europe where rape seed is used as stock food.

3. *Radicula* (Dill.) Hill

Annual, biennial or perennial, usually glabrous herbs of pungent qualities, pinnate, entire, or pennatifid leaves; flowers yellow or white; calyx with spreading sepals; pistil with short or slender style and 2-lobed, or entire stigmas; pod usually short, varying from oblong-linear to globular-terete; seeds numerous, small in 2 rows in each cell, cotyledons accumbent. About 25 species of wide distribution.

Radicula Armoracia (L.) Robinson. Horse radish

A stout perennial with long deep roots, leaves large on thick petioles, oblong, crenate, or pinnatifid, glabrous; stem leaves lanceolate, or oblong cordate, flowers with 4 green sepals and 4 white petals, not common, pods short, globular but fruit seldom found.

Distribution. It is native to the eastern part of Europe, Turkey, Greece, and the Caspian Sea through Russia, Poland and Finland. In Germany, France, Sicily, Norway, and Great Britain, it has escaped from cultivation. Common in Northern United States.

Radicula palustris (L.) Moench, var. *hispida* (Desv.) Robinson. Marsh Cress.

An erect annual, or biennial, pubescent herb, from 1 2½ feet high, leaves pinnately cleft or parted, or occasionally the upper lacinate; the lobes toothed,



Fig. 256b. Horse radish. (C. M. King)

upper leaves nearly sessile; pedicels as long as the small flowers, generally longer than the pods; pods ovoid or oblong; styles short.

Distribution. Common in northern portions of United States, to the Gulf and west to the coast; also Canada. Native in Europe.

Medical and poisonous properties. It is certain that horse radish and other members of this genus have properties somewhat similar to those of the mustards mentioned above. Horse radish contains both *singrin* and *myrosin*. Dr. Rusby mentions that it may produce serious trouble. He says

The common horse radish, likewise, loses its irritating properties when heated or dried. These are almost identical with those of mustard, and while it would not generally be regarded as a poisonous article, yet used in excess it may become so through its powerful irritation of the urinary organs, by which it is excreted. Johnson gives a case in which this result was extreme and serious. It may therefore be borne in mind that it should not be consumed in inordinate quantity. This result, should it occur, would be found excruciatingly painful.

Dr. Johnson in his *Manual of Medical Botany of North America*, writes as follows:

The uric principles of these plants appear, clinically, to be eliminated by the kidneys, and hence, incidentally, they produce a decided diuretic effect. The urine is not only increased in quantity, but partakes also of the uric character of the plant employed. In one case that came under the author's observation, the individual, though in perfect health, so far as the genito-urinary tract was concerned, suffered extremely from vesical pain and irritation for hours after using horse-radish as a condiment. In animals it produces a violent colic.



Fig. 256c Shepherd's Purse (*Capsella Bursa-pastoris*) U. S. Dept. Agr.

4. *Capsella*, Medic. Shepherd's Purse

Annual or winter annual, erect herbs, pubescent with more or less branched hairs, flowers in racemes, small, white; basal leaves tufted; pistils with short styles; pods obcordate, triangular, compressed at right angles to the partition, valves boat-shaped, seeds numerous, small, without margins, cotyledons accumbent. A small genus of 4 species, 2 in North America.

Capsella Bursa-pastoris (L.) Medic. Shepherd's Purse

An annual or winter annual, 1½ feet high, root leaves clustered, lobed, pinnatifid, or merely toothed, stem leaves sessile, lanceolate, auricled, flowers small, white; pods triangular, truncate, or emarginate, many seeded seeds light brown, elongated with a prominent ridge, seeds mucilaginous when moistened with water; cotyledons incumbent.

Distribution. One of the most common weeds everywhere in eastern North America from Nova Scotia to Florida, west to Texas and the Pacific Coast, from Eastern Canada to Manitoba to Vancouver in B. C. Cosmopolitan, as common in Europe as in the United States. Naturalized from Europe.

Poisonous properties. It produces the same symptoms as other members of the family only somewhat less severe.

5. *Camelina*, Crantz. False Flax

Erect, annual herbs, sparingly branched; leaves entire, lanceolate, or pinnatifid flowers racemose, yellow, sepals 4, green, petals 4, yellow small pistil with stigma entire, style slender, pod obovoid or pear-shaped, flattish seeds small, numerous, cotyledons incumbent. The 5 species are native to Europe and Asia.

Camelina sativa (L.) Crantz. False Flax

An erect, glabrous annual with simple or sparingly branched stem, 1½ feet long, smooth, or slightly pubescent, hairs stellate, leaves erect, lanceolate or arrow shaped, entire or nearly so, flowers small, yellow, pedicels in fruit spreading; pod obovoid or pyriform, smooth reticulated, margined seeds light brown 1 line long, minutely pitted, cotyledons incumbent, caudicle prominent, running lengthwise. On the addition of water, the seeds become mucilaginous.

Distribution. It is common in Manitoba, south to Minnesota, Northern Iowa, and the Dakotas, where it is a well known and a troublesome weed, in flax and grain fields from Ontario to the Middle States across the continent.

Poisonous properties. The plant has a disagreeable sharp odor and causes counter-irritation.

6. *Lepidium* (Tourn) L. Pepper Grass

Erect or diffuse, annual, biennial or perennial herbs leaves entire, or pinnatifid flowers racemose, white, petals small or none, stamens 6 or fewer, pod roundish, flattened contrary to the partition, winged or wingless, seeds solitary in each cell, cotyledons incumbent or rarely accumbent. About 65 species in temperate regions. The European *L. campestre*, native to Europe cultivated for salad purposes, is occasionally spontaneous. The seeds of two of our native species are used for bird food.



Fig. 256a. Fulse Plant (*Crotalaria anagyroides*). Seeds become mucilaginous on addition of water. (After Hochstein)



Fig. 257. Small Pepper-grass (*Lepidium apetalum*). Causes anasarca. (Charlotte M. King)

Lepidium virginicum L. Large Pepper Grass

Pod circular or oval with a little notch at the upper end, seeds light brown, elongated, with a prominent ridge on one side, on the addition of water they become mucilaginous. cotyledons accumbent.

Lepidium apetalum Walld. Small Pepper Grass

Seeds light brown, elongated, with a prominent ridge on one side. Seeds become mucilaginous when moistened with water. Cotyledons incumbent.

Distribution. In nearly all parts of the United States.

Poisonous properties. Pepper grass produces counter-irritation.

7. *Thlaspi* L. Field Pennycress

Low plants with undivided root leaves, stem leaves arrow-shaped and clasping, flowers small, whitish or purplish, pod orbicular, obovate or obcordate, seeds 2-8 in each cell, cotyledons, accumbent.

Thlaspi arvense, Field pennycress, Frenchweed or Stinkweed

A smooth annual with small white flowers, pod, broadly winged, about $\frac{1}{2}$ inch in diameter, deeply notched at top. Commonly naturalized in some places, becoming more abundant in the Northwest. A common weed in grain fields. It is common in the Canadian Northwest and not infrequent in Iowa and Minnesota, abundant in the Dakotas.

Poisonous properties. Probably causes counter irritation. In the Canadian Rockies the weed is carefully avoided by stock because of its pungent properties.



Fig. 258. Field Pennycress (*Thlaspi arvense*). (After Pilch.)

CAPPARIDACEAE. Caper Family

Herbs, shrubs or, occasionally, trees, alternate leaves and cruciform flowers; sepals 4-8; petals 4 or none; stamens 6-numerous, not tetradynamous, fruit a 1-celled pod or berry with 2 parietal placentae, seeds similar to those of the Cruciferae but with the embryo coiled. An order of about 35 genera and 400 species. Generally found in warm regions, few in the United States. The plants are often acrid or pungent, the flower buds of one, the Caper (*Capparis spinosa*), are pickled. Several of the species like the Rocky Mountain bee plant are cultivated for ornamental purposes. A few of the plants are weedy.

Capparis contains the coloring matter *rutin*.

Genera of Capparidaceae

Plants clammy pubescent.

Stamens 8 or more..... 2 *Polarisia*

Plants not clammy pubescent.

Stamens 6, pod few seeded .. 3 *Cleome*

Stamens 6, pod many seeded. 1 *Cleome*

1. *Cleome* L.

Glabrous annuals, leaves trifoliate or simple; flowers in leafy bracted racemes, petals entire, with claws, stamens 6, pistil with a 1-celled ovary,

pod linear with a long stalk (stipe) many seeded, the receptacle bearing a gland beyond the stipitate ovary. A small genus of about 75 species, mainly tropical.

Cleome serrulata Pursh. Rocky Mountain Bee Plant

An annual, from 1-3 feet high, with digitate, 3-foliolate leaves and leafy, bracteate racemes; calyx 4-cleft, petals 4, cruciform, short clawed, and rose-colored, pods linear, many seeded.

Distribution. Widely distributed west of Missouri, from the Canadian Rockies to Kansas, Mexico, Arizona and Utah, eastward occasionally from Minnesota to Illinois. The plant is regarded with great favor as a bee plant.

Cleome lutea Hook. Yellow Cleome

This plant is like the preceding, but the leaves are 5-foliolate or the upper 3-foliolate, leaflets oblong or oblong lanceolate, entire, stalked or sessile, flowers yellow; pod linear, stipe longer than the pedicel.

Distribution. In dry soil from Nebraska to Washington and Arizona. The former species is particularly conspicuous west of the 100th meridian.

Poisonous and Medical properties. These plants are not generally placed with the poisonous plants, although they contain the same pungent principles that members of the Mustard family have. They are seldom eaten by stock.



Fig. 259. Rocky Mountain Bee Plant (*Cleome serrulata*). A plant with pungent properties. (Ada Hayden).



Fig. 260. Yellow Cleome (*Cleome lutea*). Common in the West. (Ada Hayden)

2. *Polanisia* Raf.

Clammy herbs with whitish or yellowish flowers, palmately compound or simple leaves, flowers produced in racemes; sepals 4, deciduous, petals with claws and notched at the apex, receptacle not elongated, bearing a gland at the base of the ovary, stamens 8-numerous, unequal, pod linear or oblong, turgid, many-seeded, seeds reticulated. About 14 species in tropical and temperate regions. Annuals, with glandular hairs, common in sandy soils or on railroad embankments.

Polanisia graveolens Raf. Clammy-weed

The near relative of the Rocky Mountain bee plant is a clammy weed with loose racemes of conspicuous flowers, petals with claws, stamens 8-32, pod linear or oblong, turgid, many-seeded.

Poisonous properties. The same may be said of this as of *Cleome*. It is a clammy, pubescent weed with very pungent properties.

3. *Cleomella* DC

Annual herbs with 3-5 foliolate leaves, calyx of 4 sepals, flowers generally in racemes; petals 4, entire, without claws, receptacle short, stamens 6, inserted on the receptacle, ovary short, long-stalked, pod linear to oblong, many-seeded. About 75 species, found chiefly in southwestern North America and Mexico.

Cleomella angustifolia Torr

A glabrous annual from 1-2 feet high, leaflets 3, linear lanceolate or linear oblong bracts simple flowers small, yellow, pod rhomboidal, raised on a slender stipe, but shorter than the pedicel, few seeded.

Distribution. From Nebraska and Kansas to Texas New Mexico and Colorado. Abundant in waste places.

SARRACENIALES

Carn vorous plants secreting a viscid liquid, radical leaves, scapose flowers, corolla choripetalous sepals generally distinct, stamens usually free ovary compound super- or Contains the families *Droseraceae*, *Sarracenaceae* and *Nepenthaceae* the genus *Sarracenia* has 6 species in eastern North America, *S. purpurea*, found as far west as Minnesota, contains the alkaloid *Sarracenia*. *Darlingtonia californica* occurs in California and Oregon *Heliamphora* is native to Guiana. The family *Nepenthaceae* with 40 species is found mostly in the India-Malayan regions, some species being frequently cultivated in green-houses. The plants of these orders are insect vorous, capable of digesting insects.

DROSERACEAE

Perennial or biennial glandular pubescent bog herbs or somewhat shrubby plants, leaves mostly from the base with tentacles, which secrete a viscid substance to catch insects circinnate in the bud flowers perfect, raremose, calyx persistent, 4-8 parted, or the sepals distinct, petals 5 free, stamens 8-20 ovary free, 1-3-celled, styles 1-5, simple 2-cleft; capsule 1-5-celled. A small order of 125 species of wide distribution. The most important genus is *Drosera*, commonly called sundew, the tentacles of which secrete a viscid fluid which catches




Fig. 268a. Various insectivorous plants: 1, *Sarracenia variowirtis*, 2, *Darlingtonia californica*, 3, *Sarracenia laciniata*, 4, *Utricularia villosa*.

insects and clings to them. The *D. rotundifolia* is commonly found in our northern bogs. The *Drosera rotundifolia* is found on the sandy hills of Portugal. The Venus fly-trap (*Dionaea muscipula*) of the Carolinas grows on the sandy barrens and feeds on insects. These plants are somewhat rare.

Poisonous properties. According to Dr. Schaffner, the common sundew is poisonous to cattle. From one species of *Drosera* two pigments have been isolated, the red having the formula $C_{11}H_8O_5$ and the yellow $C_{11}H_8O_4$. Plants of the family in Australia are said to be poisonous to sheep.

According to Greshoff the leaves of *Drosera binata* contain hydrocyanic acid. *D. rotundifolia*, and *D. intermedia*, were also found to contain a little HCN. The leaves of *Dionaea muscipula* contain the same substance.

Herbs, shrubs or trees, flowers usually polypetalous, stamens mostly perigynous or epigynous, sepals chiefly united or confluent with receptacle, carpels 1 or more, distinct or sometimes united into a compound ovary. The order includes the families *Podostemonaceae*, *Crossulaceae*, *Cephalotaceae*, *Saxifragaceae*, *Pittosporaceae*, *Hamamelidaceae*, *Platanaceae*, *Rosaceae*, *Connaraceae*, *Leguminosae* and other small families. The family *Saxifragaceae* includes the red currant (*Ribes vulgare*), black currant (*R. nigrum*), the Missouri currant,



Fig. 26b. *Dionaea muscipula* in its natural habitat, showing the arrangement and withering of the leaves and the flowers and where they are to reach the sun. (Kerner-Oliver, Dept. Agr. Univ. of Minn.)




Fig. 26c. Insect Traps. 1, *Dionaea muscipula*, 2, Section of *Dionaea*, 3, One of the Sarracenia, 4, Leaf of *Aldrovanda*, 5, Section of *Aldrovanda*, 6, Trap of *Sarracenia*, 7, Gland in the wall of trap of *Sarracenia*. (Kerner, Dept. Agr. Univ. of Minn.)




Fig. 261. Common Sundew (*Drosera rotundiflora*). Poisonous to cattle. (After Pilch).



Fig. 262. Missouri Current (*Ribes aureum*). The flowers are spicy fragrant. It is a well known shrub. (W. S. Dudgeon)

(*R. aureum*), the Crandall, a well known cultivated form the cultivated gooseberry (*R. Grossularia*), Missouri goosecherry (*R. gracile*) the cultivated hydrangeas (*Hydrangea paniculata*); the wild hydrangea (*H. arborescens*); used as a diuretic; the mock orange (*Philadelphus coronarius*), and *P. grandiflorus*, *Deutzia scabra*, the *Asiatica japonica* and the alony root (*Henckera americana*) a powerful astringent. Several members of this family are poisonous the western *Juncus americana* a pretty shrub of the Rocky Mountains with white flowers, contains an appreciable amount of HCN according to Greshoff. The same substance also occurs in the garden *Hydrangea hortensis*, *H. arborescens* an American species of the southern region. The *H. Thunbergii*, *H. Lindleyana* and *H. involucrata* all contain HCN. The HCN however, is in a transitory stage. Greshoff states that the leaves of *Philadelphus coronarius*, *P. Lemnais* and *P. microphyllus*, and the seeds of *P. grandiflorus* contain saponin as do the leaves of *Deutzia scabra*. The family *Hamamelidaceae* includes the witch hazel (*Hamamelis virginiana*) containing a bitter principle, the storax (*Liquidambar orientalis*) of the Old World which contains two resin alcohols, *sorexin* and *ester* and red gum (*L. Styraciflua*) a common tree of moist woods of the South supplying a well known commercial wood, it is also an ornamental tree the leaves of which, when bruised, are fragrant. The family *Platanaceae* contains the sycamore (*Platanus occidentalis*), a large tree which furnishes valuable wood and is also used for ornamental purposes. According to Greshoff the leaves of *Platanus acerifolia*, *P. cuneata*, *P. occidentalis* and *P. orientalis* contain HCN. The young leaves are poisonous.

Families of *Rosales*

Calyx free from the ovary, wholly superior

Simple 1-celled

Leguminosae

Ovaries 2 or more compound, stamens twice as many as the pistils

.

Crassulaceae.

Stamens inserted on the calyx, stipulate Rosaceae
 Calyx more or less coherent with the compound ovary Rosaceae (Pomeae)



Fig. 251. Red Gum (*Liquidambar styraciflua*). Furnishes a commercial wood.

CRASSULACEAE DC. Orpine Family

Succulent herbs, leaves generally sessile without stipules, flowers small, symmetrical, usually cymose, petals and sepals equal in number, from 3-20 stamens as many or twice the number, pistils distinct, fewer than the sepals, receptacle usually with small scales 1 back of each carpel, fruit a dry dehiscent follicle, usually many seeded. This is a small family of about 50 species, many being found in dry soils, rocks, etc. in North Temperate regions. A few of the species are cultivated for ornamental purposes, among these being the *Roelevia* of the Cape of Good Hope, whose flowers have narcotic properties, poisoning sometimes resulting from smelling them. Some of the *Crassulaceae* contain *crassulic* and *malic acids*. The *Cotyledon ventricosa* of South Africa is said to produce the Kenta disease of that region, although this is usually ascribed to some members of the Pulse family and is probably identical with loco disease. The leaves of wild tea (*Coltea edulis*), according to Bull. Miscellaneous Information of Kew Gardens, when chewed are said to have great sustaining powers.

Sedum (Tourn.) L. Stone Crop

Fleshy, smooth herbs, mostly perennials, with alternate leaves, flowers cymose, perfect or dioecious, calyx 4-5-lobed, petals 4-5, distinct, stamens 8-10, perigynous, pistils 4-5, distinct or united at the base, styles short, follicles many seeded. About 150 species, mostly of temperate and cooler regions of the northern hemisphere. A few are cultivated for ornamental purposes. There are several somewhat weedy species as *S. purpureum* and mossy stone crop (*S. aca*) the latter native to Europe but escaped to roadsides in the East. It has acid properties, which is also true of the live-forever, a plant of the Rocky Mountains.

Sedum purpureum Tausch. Live-forever

A stout perennial 2 feet high with fleshy oval or obtuse, toothed leaves, and flowers in compound cymes, corolla purple, with oblong lanceolate, purple petals, stamens perigynous; pistils with a short style, fruit a follicle with a short pointed style

Distribution. Native to Europe, frequently escaped from cultivation and found around dwellings and in cemeteries.

Sedum acra L. Mossy Stone Crop

A moss-like plant spreading on the ground, leaves small, alternate, ovate thick and fleshy, flowers yellow, perfect, in spike-like clusters, cymose, the central flower with 5 sepals, 5 petals, and 10 stamens, the others with 4 sepals, 4 petals and 8 stamens. follicle spreading, tipped with a slender style

Distribution. On rocks and along roadsides from New Brunswick to New York and Pennsylvania. Native to Europe



Fig. 263. Common Live-forever (*Sedum purpureum*). A plant with acid properties. (Fitch)

Fig. 264. Live Forever (*Sedum stenopetalum*). Common in the Rocky Mountains, a plant with acid properties. (W. S. Dudgeon)

Poisonous properties. *Sedum acre*, according to Dr Schaffner, produces inflammation and vesication when applied to the skin, it is used to remove the false membrane in diphtheria. Dr White, in his "Dermatitis Venenata," says with reference to the *Sedum acre* "Wood states that the whole plant abounds in an acrid, biting juice. Oesterlen says that it is sharply irritative to the skin. The National Dispensatory states that the juice is capable of blistering the skin, and that it is used upon corns and warts to soften them, and upon swollen glands as a resolvent. Mr Cheney, a wholesale dealer in vegetable drugs, informs me that the juice of the green plant is poisonous to the skin of many persons." This plant, however, is not common in the United States.

ROSACEAE. Rose Family

Herbs, shrubs or trees. leaves alternate or some opposite, stipulate frequently falling soon after the leaves appear, flowers regular, stamens generally numerous, distinct, inserted on the calyx, petals as many as the sepals or rarely wanting, pistils 1-many, generally distinct, except in Pomace where the pistil is united to the calyx, fruit various, achenes, follicles, drupes or pomes as in the apple, seeds 1-many, without albumen, embryo straight, with large cotyledons. The order contains about 90 genera and 1500 species of wide distribution in temperate and tropical regions, some boreal. But few of the plants are noxious or have noxious qualities.

The several cherries, like *Prunus serotina* and *P. virginiana* are known to cause stock poisoning, and the seeds when eaten likewise produce fatal results in man. The *P. serotina* or wild black cherry, is used in medicine, under the name of *P. virginiana*. The choke cherry (*P. virginiana*) is also used in medicine. The bark of the wild black cherry is official. It contains tannic and gallic acids, and a volatile oil resembling the volatile oil of bitter almonds. It is used as a tonic and astringent. The leaves of the laurel cherry are used for making cherry laurel water which is a sedative narcotic. From *Prunus Amygdalus*, var. *amara*, native to Asia, is obtained the amygdalin or bitter almonds, which is converted into hydrocyanic acid. This acid is deadly poison, and is obtained from a great many different plants. The leaves of the laurel cherry also contain the same substance. HCN has been found in *Prunus paniculata*, *P. pendula*, *P. Pennsylvanica*, *Pyrus Aria*, *P. pinnatifida*, *P. japonica*, *Crataegus orientalis*, *Cotoneaster integerrima*, *Autonia crenatifolia*, *Amelanchier alnifolia*, *Chamaelea japonica*. According to Greshoff the leaves of *Koeneckia angustifolia* contain hydrocyanic acid. The same writer reports this substance in the mountain mahogany (*Cercocarpus parvifolius*) of the Rocky Mountains. It must be regarded as poisonous. In laurel cherry, it is largely derived from the decomposition of *laurocerasins*. This species also contains *prunocerasins*. Several species of other orders also might suitably be mentioned here as containing substances capable of being converted into hydrocyanic acid. In this class are the toadstools (*Agaricus oreades*), bitter cassava (*Manihot nitensima*) and sorghum (*Andropogon Sorghum*).

According to Greshoff *Cornus foetida*, *Spiraea japonica* and many other plants of the family contain saponin.

The same substance is obtained from other plants of the genus *Prunus*. Oil of roses is obtained from the *Rosa damascena*, Miller, var. The mucilaginous seeds of the quince (*Pyrus Cydonia*) have been used in medicine for a long time.




Fig 265. Wild Red Cherry (*Prunus pennsylvanica*). It contains the glucoside amygdalin. (Ada Hayden).

Many plants of the family are cultivated for ornamental purposes and some are economic. In the first class is the laurel cherry (*Prunus Lauro-cerasus*), a handsome fragrant shrub of the Caucasus to northern Persia, cultivated in the southern states and very common in the Mediterranean regions of Europe. The Mayday tree of Europe (*Prunus Padus*), as well as numerous species of the genus *Spiraea*, like *Spiraea Douglasii*, *S. salicifolia*, *S. japonica*, *S. Thunbergii*, the nine-bark (*Physocarpus opulifolius*), species of the genus *Rosa*, such as the prairie rose (*Rosa setigera*), sweetbrier (*Rosa rubiginosa*), dog rose (*Rosa canina*), *R. rugosa*, *R. gallica*, and the cinnamon rose (*R. cinnamomea*) are frequently cultivated. *Kerria japonica*, *Rubus odoratus*, *Pyrus coronaria*, *P. japonica*, *P. Aucuparia*, *P. americana*, *Crataegus mollis* and *C. punctata* are also cultivated.


The family contains a large number of valuable fruits, of these we may mention the service berry (*Amelanchier canadensis* and *A. spicata*), the apple (*Pyrus Malus*), the pear (*Pyrus communis*), the quince (*P. Cydonia*), straw-

berry (*Fragaria vesca*, *F. virginiana*, var. *Illinoensis*), the *F. chiloensis*, native to Chili and the Pacific coast (the common garden strawberry is a modified form of the Chilian strawberry), and the Indian strawberry (*Duchesnea indica*). We may also mention the wild northern plum (*Prunus americana*), the Chickasaw plum (*P. angustifolia*), European garden plum (*P. domestica*), sand cherry (*P. pumila* and *P. Besseyi*); the cherries, English cherry (*P. avium*), naturalized in the southern states, especially in Virginia and Maryland, the sour cherry (*P. Cerasus*), also naturalized in the East and extensively cultivated, the wild red cherry (*P. pennsylvanica*), commonly used in the north, the Japanese plum (*P. triflora*), the apricot (*P. armeniaca*), peach (*P. persica*), almond (*P. Amygdalus*) flowering almond (*P. nana*), wild red raspberry (*Rubus idaeus* var. *aculeatusimus*), black raspberry (*R. occidentalis*), garden raspberry of Europe (*R. Idaeus*), salmon berry (*R. parviflorus*), dewberry (*R. cuneifolius*) and the wineberry of Japan (*R. phoenicolasmus*). The fruit of the Icacó plum (*Chrysobolanus Icacó*) of tropical America is edible. The wood of the wild black cherry (*P. serotina*) takes an excellent finish, and therefore is highly desirable for cabinet making and for interior finishing of houses. The wood of other species of the genus is used in the manufacture of pipes and furniture. Most of the plants of the order contain no injurious substances. *Malic acid* $C_4H_6O_5$ occurs in the fruit of the cherry (*Prunus Cerasus*), plum (*P. domestica*), the apple (*Pyrus Malus*), the strawberry (*Fragaria virginiana*), (*F. vesca*), etc. *Salicylic acid* $C_7H_6O_3$ occurs in the fruit of the strawberry, citric acid in *Rubus*, the strawberry and *Prunus domestica*. The *Quillaja Saponaria* contains *sapunin*, the bark yielding 2 per cent. Kobert distinguishes two substances *quillajic acid* $C_{10}H_{20}O_{16}$ and *sapotoxin* $C_{11}H_{22}O_{17}$.

The glucoside *amygdalin* was first obtained in 1830 by Robiquet and Boutron from the seeds of the bitter almond. Laebig and Wochler named the substance which converts the *amygdalin* into the so-called essence of bitter almonds, *emulsin*. They found that through the action of *emulsin*, sugar and *prussic acid* were formed. The name *synaptase* was given to *emulsin*.



Emulsin can also convert *salicin*, *helicin*, *phlorizin*, and *arbutin*. The change in *arbutin* is as follows:



In the cherry leaves *emulsin* occurs in the leaves and younger branches. *Emulsin* also occurs in *Penicillium glaucum* and *Aspergillus niger*.

The Kooso (*Brayera anthelmintica*) is a large dioecious ornamental tree from Abyssinia. The drug comes from the pistillate flowers which have a tea-like odor but a bitter, nauseous taste and contain *cuscutoxin* which is a muscle poison, *protocoin* and *coin* which is bitter and acrid. In medicine it is used as a taenifuge but in large doses produces vomiting and colic. *Agrimonia gryposepala*, *Guaiaco stipularia*, and *Genum urbarum* are used as astringents. The roots of the water avens (*Genum rivale*) are tonic and powerfully astringent. The soap-bark (*Quillaja Saponaria*) of Peru and Chin is used as an expectorant and is an irritant poison. According to Schneider who investigated a great many of the saponins which occur injuriously in about fifty families, they act

poisonously by dissolving the blood corpuscles. *Cholesterin* contained in the body acts as a natural antidote against them.

Several plants of the order produce members of the terpene group, rose oil, contains *rhodinol* $C_{10}H_{16}O$, supposed to be identical with geranium oil and a second terpene *roseol*, $C_{10}H_{16}O_2$. Many fruits of the order especially *Pomace*, contain *mannite* and *sorbite*. The *orbatin*, $C_{12}H_{14}O$, obtained in many plants also occurs in some plants of this order, the glucoside *hydrochinon*, C_6H_4O , occurs in the huds of pears. *Quercetria* $C_5H_8O_2$, derived from a glucoside, is found in the flowers of haw (*Crataegus*), the bark of apple trees and of *Prunus institia*. *Amygdalin* $C_{20}H_{27}NO_{11}$, occurs in the seeds of many plants of the family, especially in *Pomace* and *Prunace*, also in the bark of *Prunus Padus*, *P. serotina*, etc., and in the seeds of *Pyrus Aucuparia*.



Fig. 266 Kousso (*Brayera outalimnica*) Flowering branch. Contains a muscle poison. (After Vague). Fig. 266a. Common Wild Plum (*Prunus americana*). (C. M. King)

Genera of Rosaceae

Ovary inferior or enclosed in the calyx tube

Carpels numerous, fruit an achene

3. *Rosa*

Carpels few, fruit not an achene

Carpels cartilaginous, fruit a pome

4. *Pyrus*

Carpels bony, drupe-like

6. *Crataegus*

Ovary superior not enclosed in calyx tube.

Calyx deciduous, fruit a drupe

5. *Prunus*

Calyx persistent.

- Pistils numerous; fruit drupelets 1. *Rubus*.
 Pistils numerous, fruit an achene 2. *Fragaria*.

1. *Rubus*. Raspberries and Blackberries

Perennial herbs, shrubs or vines, vine prickly, with alternate leaves, 3-7 foliolate or simple, flowers terminal; axillary or solitary, white reddish or pink, usually perfect, calyx 5-parted, petals 5, deciduous; stamens numerous, achenes usually many, inserted on the receptacle, which is either fleshy or dry, carpels forming drupelets. About 200 species chiefly Northern.

Rubus Idaeus L. var. *aculeatissimus* (C. A. Mey) Regel & Tiling. Wild Red Raspberry

Stems biennial, upright shrubs covered with straight, stiff bristles, some hooked, and glandular hairs; leaflets 3-5, oblong, ovate, pointed, whitish, downy underneath; petals as long as the sepals, whitish; fruit light red. Spreads by suckers.

Distribution The species is native to the Appalachian Mountains as far south as the Carolinas; common at high altitudes in the Rockies. Frequently



Fig. 267. Wild Red Raspberry (*Rubus Idaeus*, var. *aculeatissimus*). The prickles of the red raspberry produce mechanical injury. (Ada Hayden).



Fig. 268. Wild black cap raspberry (*Rubus occidentalis*). This plant has thorns which are irritating. (Ada Hayden).

troublesome in fields for several years in the north and persists for a long time in gardens. A common native of the north.

Rubus occidentalis L. Black Raspberry, or Black-cap Raspberry

Stems biennial, glaucous, recurved, beset with hooked prickles, rooting at the tip, leaves pinnately 3-foliate, or rarely 5-foliate, leaflets ovate, coarsely doubly serrate, whitish underneath, flowers corymbose clusters, petals shorter than the sepals, fruit usually purplish-black, occasionally white.

Distribution. Quebec to Georgia, to Missouri, to Minn. Like the preceding species, often troublesome in fields and gardens.

Rubus villosus Ait. High Bush Blackberry

Shrubs 1-6 feet high, upright or reclining, armed with stout recurved prickles, branchlets and lower surface of leaves glandular; leaflets 3-5, ovate, pointed, terminal one stalked; flowers in corymbose clusters, petals white, fruit not separating from the juicy receptacle, blackish.

Distribution. From Nova Scotia to Georgia, Missouri, Kansas, to Minn. Troublesome like the black-cap Raspberry.

Injurious properties. No species of the genus *Rubus* is known to be poisonous, but the bristles and spines on various species frequently inflict injuries. Numerous cases of inflammation, and later pus formation, are reported from the prickles found on the common red raspberry. This is especially true of the larger prickle of the black raspberry and the dewberry. The bark of the blackberry contains *villosin*. The leaves are said to cause an irritation of the skin of berry pickers or others who walk among the bushes.



Fig. 269. Common Blackberry (*Rubus villosus*)
The common blackberry produces prickles which act injuriously in a mechanical way. (Ada Hayden).

2. *Fragaria* L. Strawberry

Perennial stemless herbs; leaves petioled, stipulate, 3-foliate, flowers white, in clusters, polygamo-dioecious; calyx persistent, with 5 bractlets, deeply 5-lobed, petals 5, stamens numerous, carpels or pistils numerous, the receptacle elongated, which become fleshy in fruit. A small genus of about 25 species containing the common cultivated strawberry (*F. chiloensis*) of the Pacific coast, our wild strawberry (*F. virginiana*), the European strawberry (*F. vesca*) and the Indian strawberry (*F. Indica* or *Duchesnea indica*).

Fragaria vesca L. European Wood Strawberry

A perennial, with ovate leaves, dentate, thin, flowers white, racemose receptacle, elongate, fruit with achenes, seeds free above the receptacle. It is very different from our common wild strawberry, in which the achenes are sunken in the flesh.

Poisonous properties. Few people would suspect that the seeds of the strawberry are injurious, but I have known people who have found that the eating of the common cultivated strawberry is injurious, and it is known that the European strawberry (*F. vesca*) produces a rash that sometimes resembles that produced by scarlatina. Dr. Millsaugh, in speaking of the European strawberry says of a lady coming under his care who had consumed the fruit of the strawberry grown in Florida:

In the afternoon of the same day the skin was hot and swollen, the patient thirsty and restless, and little sleep was gained that night. The next day the eruption began to fade, the appetite returned, and restlessness ceased. On the third day exfoliation began and was very profuse, the skin appearing quite similar to the condition existing after a severe attack of scarlatina.




Fig. 27b




Fig. 270a

Fig. 27b. Wild Strawberry (*Fragaria virginiana*). (Ada Hayden). Fig. 270a. Wild Strawberry (*Fragaria vesca*). (Willis).




Fig. 371 European Wood Strawberry (*Fragaria vesca*). This plant occurs in fields and along roadsides. Sometimes causes dermatitis. (W. S. Dudgeon).

Prof. Prentiss* reports the case of a man who, at the age of 14, had become quite ill from eating strawberries and forever afterward could not eat them without becoming ill.

3. *Rosa* L.

Erect or climbing shrubs, with prickly stems, alternate leaves adnate to the petioles, flowers showy, corymbose, or solitary, calyx urn-shaped, stamens and carpels numerous; achenes, enclosed in a berry-like calyx tube. Several species of the genus *Rosa* are more or less troublesome in fields. The *Rosa centifolia*, used for preparing rose water contains a volatile oil. A confection is made from the hips of *Rosa canina*. *R. gallica* contains a volatile oil and a yellow crystalline glucoside *quercetin*.

Rosa pratincola Greene Prairie Rose

An erect perennial shrub with densely prickly stems bearing slender bristles, narrow stipules, more or less glandular, toothed, leaflets 7-11, broadly elliptical to oblong-lanceolate, sessile or nearly so; flowers corymbose or rarely solitary, pink, sepals lanceolate, somewhat glabrous fruit smooth.

Distribution. Common on prairies of Wisconsin, Iowa to Texas, New Mexico and Montana. In Iowa, Missouri and parts of Minnesota and Arkansas it is most troublesome.

Rosa blanda Ait. Smooth Rose

An erect shrub with few straight prickles or wholly unarmed, from 1-3 feet high, leaflets 5-7, short stalked, oblong-lanceolate, cuneate, stipules dilated, naked or slightly glandular-toothed; flowers usually large, corymbose or solitary.

Distribution. From Newfoundland to Ontario and Illinois.

* Bos. Gazette 18-19.

Rosa Sayi Schw. Say's Rose

A very prickly shrub with low stem, 1-2 feet high; leaflets 3-7 broadly elliptica, or oblong, lanceolate, glandular, ciliate and resinous, stipules dilated flowers large, solitary or rarely more, outer sepals usually with 1 or 2 narrow lateral lobes.

Distribution From northern Michigan, Northwest Territory to Colorado.



Fig. 1 372. Arkansas Rose (*Rosa pratincola*) A. N. S. P. (Ada Hayden).

Rosa Woodsii Lindl. Wood's Rose

Stems usually low, 6 inches to 3 feet high with slender, straight or recurved spines and scattered prickles, or unarmed above, leaflets 5-7 obovate to oblong or lanceolate, more or less toothed, flowers corymbose or solitary, sepals naked or hispid, fruit globose.

Distribution. Prairies of Minnesota and from Montana to New Mexico and Texas.

Mechanical injuries. The prickles inflict injury to cattle, and are especially troublesome in grain fields. Before the binder came into use men were

injured by the prickles and spines of the rose which produced inflammation and caused the formation of pus.

4. *Pyrus* L.

Trees or shrubs; simple leaves, flowers in cymose clusters, white or pink, calyx urn-shaped, 5-lobed, petals 5, short-clawed, stamens usually numerous; styles mostly 5, distinct or united at the base, ovules 2 in each cavity, carpels leathery fruit a pome. About 37 species of wide distribution, chiefly in the North Temperate region. The following species of the genus are cultivated for their fruits, quince (*P. Cydonia*), pear (*P. communis*), Japan or sand pear (*P. sinensis*), apple (*P. Malus*), Old World crab apple (*P. baccata*), mountain ash (*P. americana*), (*P. nictitans* *P. sambucifolia*), European mountain ash (*P. Aucuparia*).

The fresh bark of the wild mountain ash is used in medicine; it is known to produce irritation of the alimentary mucous membranes, and a reflex nervous irritation.

Pyrus coronaria L. Wild Crab Apple

A small tree with petioled or ovate to triangular-ovate leaves, sparingly pubescent beneath, sharply serrate and often lobed, flowers rose-colored, fragrant, calyx slightly pubescent, pome fleshy, fragrant, greenish-yellow, acid. Two other species are found in eastern North America, namely *P. angustifolia*, with small leaves and few flowers, and *P. ioensis*, with firm leaves narrowed at the base, and pubescent calyx, chiefly west in the Mississippi Valley, *P. riviniana* Dougl., occurs from California to Alaska.

Distribution. Our Eastern wild crab is found from Ontario to Michigan and South Carolina; in the west it is replaced by the *P. ioensis*.

Poisonous and medicinal properties. All the species of the genus *Pyrus* contain the glucoside amygdalin, $C_{20}H_{27}NO_{11}$, which is converted by the action of the ferment into hydrocyanic acid. The bark also contains citric acid, $C_6H_8O_7$, and malic acid, $C_4H_6O_5$, both of which appear in the fruit of the apples. There may be occasionally cases of poisoning where animals are allowed to browse upon the wilting leaves of the apple.



Fig. 273. Common apple (*Pyrus Malus*). The well known cultivated apple. (W. S. Duggan)




Fig. 274. Ilex Crab (*Ilex laevis*). 1. Flowering branch. 2. Longitudinal section of flower with petals removed (natural size). 3. Fruiting branch. 4. Longitudinal section of fruit. 5. Summer branch. 1, 3, 4, 5, one half natural size. (H. M. Cheney)

Crataegus L. Hawthorn, White Thorn

Shrubs or small trees, usually spiny; leaves petioled; flowers in corymbose clusters, white or pink; calyx tube urn-shaped, lobes 5, acute, petals 5, roundish, stamens numerous or few; styles 1-5; fruit a pome containing 1-5 bony, 1-seeded stones. About 75 species, although the number is sometimes estimated as high as 125. Some species, like *C. punctata*, and *C. mollis* are ornamental.

Crataegus mollis Scheele

Shrub or small tree; shoots densely pubescent; leaves large, slender petioled, cuneate, truncate or cordate at base, usually with acute lobes, more or less densely pubescent beneath; flowers large, 1 inch across, fruit bright scarlet with a light bloom.

Distribution. Common in thickets from Eastern Canada to Iowa and Kansas, and Texas.

Poisonous properties. Large numbers of haw fruits are eaten, and several deaths due to the eating of *Crataegus* have been reported in Iowa. These were probably largely due to strangulation or indigestibility of the stony "seed." The flesh is said to be indigestible as well.



Fig. 275. Common Red Haw (*Crataegus mollis*) (C. M. King)

5. *Prunus* L. Plum and Cherry

Shrubs or trees with alternate petioled leaves and small stipules, flowers variously clustered, mostly perfect, calyx inferior free from the ovary with a bell-shaped or urn-shaped tube and 5 spreading lobes, falling after flowering, petals white or pink, spreading stamens 13-20 or more, distinct, inserted on the throat of the calyx, perigynous, pistil solitary style simple stigma capitate; ovary 1-celled 2-ovuled, fruit a drupe seed usually one embryo large, cotyledons fleshy, endosperm absent. Species about 90, of the north temperate regions, tropics, America and Asia. The sweet cherry (*Prunus avium*) sour cherry (*P. cerasus*), native plum (*P. americana*), Chicksaw plum (*P. Chickasaw*), European plum (*P. domestica*), Japan plum (*P. triflora*), the flowering almond (*P. triloba*), peach (*P. persica*), and apricot (*P. armeniaca*) are all well known in cultivation.

Prunus virginiana L. Choke Cherry

A tall shrub or small tree, bark gray, leaves thin, oval, oblong or obovate, acuminate at the apex, smooth or slightly pubescent, sharply serrate, teeth large, often doubly serrate; flowers white in rather loose racemes, terminating leafy branches, petals roundish, fruit red, turning dark or crimson. Astringent.

Distribution. Forming thickets from New Foundland to Manitoba to Texas and Georgia.



Wild Cherry (Prunella virginiana) - Common in the Blue Ridge area - Photo by Colburn

Prunus demissa Walp. Western Wild Cherry or Choke Cherry

A shrub or small tree, leaves thick and oval or obovate, acute or more or less obtuse at the apex, teeth rather short, flowers white in dense racemes, terminating leafy branches, fruit dark or purplish black, less astringent than the preceding.

Distribution. Dry soil, common in thickets and woods from Dakota to Kansas, New Mexico to California and British Columbia.

Prunus serotina Ehrh. Wild Black Cherry

Large tree with reddish brown branches, reddish wood leaves thick, oblong, or lanceolate-oblong, taper pointed, serrate, with short teeth shining above, flowers in elongated spreading or drooping racemes, petals obovate, fruit purplish black, and slightly astringent.




Fig. 276. Black cherry (*Prunus serotina*). 1. Flowering branch. 2. Longitudinal section of flower, enlarged. 3. Fruiting branch. 4. Cross section of fruit. 5. Longitudinal section of fruit. 6. Winter branchlet. 1, 3, 6, one-half natural size. 4, 5, natural size. (M. M. Cheney in Green's Forestry in Minnesota.)

Distribution. From New England to Ontario, to Florida and Texas, Kansas, Dakota and Minnesota.

Prunus pennsylvanica L. Wild Red Cherry

A small tree 20-30 feet high, light brown bark, leaves oval or lanceolate, acute or acuminate, finely and sharply serrate, glabrous, with slender petiole, flowers white in corymbose clusters, fruit small, globose, light red hue and sour

Distribution. In rocky woods. New Foundland to the Rocky Mountains to Georgia.

Poisonous properties. Many cases of poisoning have been recorded from eating the seeds of peach and bitter almonds. They contain a highly poisonous



Fig. 277 Wild Red Cherry (*Prunus pennsylvanica*). 1 Flowering branch. 2 Longitudinal section of flower. 3 Fruiting branch. 4 Longitudinal section of fruit, slightly enlarged. 5 Cross section of fruit. 6 Embryo enlarged. 7 Axil of leaf, showing stipules. 8 Winter branchlet. 1, 3, 7, 8, one-half natural size. (M. M. Cheney)

substance from which prussic acid is obtained. Mr. Chesnut says in regard to the black cherry

The fruit is rather agreeable, being but slightly bitter and astringent in taste. In some localities it is much used to flavor liquor. Poisoning is frequently caused by cattle eating the wilted leaves from branches thrown carelessly within their reach or ignorantly offered as food. Children occasionally die from eating the kernels of the seed or from swallowing the fruit whole.

Prof. Chas. D. Howard, of the New Hampshire Station, says

The poisonous property of all species of cherry leaves is due to hydrocyanic acid, popularly known as prussic acid. This compound does not exist as such in the growing leaf, but is derived from a class of substances called glucosides, of which amygdalin is the type peculiar to the cherry. This, or a closely allied body, is to be found not only in the leaves and bark, but especially in the stones of cherries, peaches and plums, and the seeds of the apple. By the action of moisture and a vegetable ferment called *emulsin*, which exists in the plant, a complex chemical reaction takes place, that begins in the leaf the moment connection with the circulatory system is cut off. The three products of this reaction are hydrocyanic acid, grape sugar and benzaldehyde, or bitter almond oil.

There is a popular opinion that the leaves of the cherry are poisonous only when cut and in the wilted condition; that cattle may safely nibble them from the growing shrub without danger of injury; and that they are quite harmless when dried. Our observations, however, prove these views to be but partially correct. As a matter of fact, distillations of samples, made within twenty minutes of cutting, show that the freshly cut leaves yield nearly as much acid as the wilted ones do when calculated on the weight of fresh material taken, and when eaten fresh, the character of the juices within the animal stomach is such as to render that organ a most favorable place for the conduct of the reaction in which prussic acid is liberated.

On the other hand, while it is true that the thoroughly dried leaves yield a comparatively small amount of acid, still they may always produce some, and as ordinarily dried in the hayfield, they may be capable of generating a considerable quantity of the poison. The desirability of carefully excluding them from all hay is therefore apparent.

One hundred grams of bitter almonds (*Prunus amygdalus communis*) in the form of pulp, yield 250 milligrams prussic acid, the same amount of kernels from cherry stones yields .70 mgr. leaves of the cherry laurel (*Prunus laurocerasus*), occurring in Europe and Mexico, 32 mgr. kernels of peach, 164 mgr., apple seeds, 35 mgr. The stones of all these species must therefore be regarded as dangerous, the fruit, in every case, so far as can be ascertained, is harmless.

The leaves of the wild black cherry are the most poisonous of the three species investigated, though all are dangerous.

Both the wilted leaves and fresh leaves are poisonous, while the dried are to be regarded with suspicion.

Vigorous, succulent leaves from young shoots, which are the ones most liable to be eaten by cattle, are far more poisonous than the leaves from a mature tree or stunted shrub.

Leaves wilted in bright sunlight to about 75 per cent original weight, or until they begin to appear slightly limp and to lose their gloss, yield the maximum amount of prussic acid.

I have seen cattle browse on the leaves of *Prunus demissa* in the Rocky Mountains. It is believed by stockmen to be poisonous. Probably the danger is not so great because the leaves are eaten direct from the tree and not wilted. The Bureau of Forestry has shown that much loss occurs which can easily be prevented by changing the trail.

It has been known for a long time that seeds of the various members of the genus *Prunus* contain poisonous properties. The bark of several of our wild cherries is also known to contain a poisonous principle, a ferment known as *emulsin*, which in the presence of water acts on the glucoside *amygdalin* $C_{20}H_{27}NO_{11}$, and produces *hydrocyanic acid*, a powerful poison, it is a clear, colorless liquid of a characteristic taste and odor, resembling that of a bitter

almond. Externally, hydrocyanic acid produces a paralytic effect. When taken internally, it acts as a sedative upon the mucous membrane. Prof. Winslow gives the following characteristics of poisoning

In poisoning, the blood becomes first a bright arterial hue, and later assumes a dark, venous color. The first condition is due to the fact that the blood does not give up its oxygen for some reason. Bruston suggests that it is because the blood is hurried so rapidly through the dilated peripheral vessels that it does not have time to yield up its oxygen. The dark color of the blood is probably owing to asphyxia and accumulation of carbonic dioxide, following the paralytic action of prussic acid upon the respiratory centre.

Prussic acid has an essentially depressing action upon the nervous system as a whole. The brain, cord and nerves become paralyzed by large doses.

The spinal cord is paralyzed at a period after coma and convulsions have appeared. The peripheral nerves and muscles are paralyzed directly by toxic doses, and not through the mediation of the central nervous apparatus.

Inhalation of the pure acid will cause death in a confined atmosphere and even inhalation of the medicinal solution will induce the physiological symptoms of the drug.

He also says of the toxicology of prussic acid

Prussic acid is one of the most powerful poisons in existence. Death may be instantaneous, or life may be prolonged for over an hour after lethal dose. More commonly the animal survives for a few minutes, and we observe the following symptoms in dogs: The animal falls, froths at the mouth, the respiration is of a gasping character and occurs at infrequent intervals. There is unconsciousness, the pupils become dilated, there are muscular tremblings, and clonic or tonic spasms. Defecation and micturition occur, and erections often ensue in the male. Respiration ceases before the cardiac pulsations.

Three stages may be distinguished in fatal poisoning. First: a very short period elapses before the symptoms appear. There are giddiness, difficult breathing, and slow pulse in this stage. Second the pupils dilate, vomiting may occur, and the animal utters loud cries. Spasmodic defecation, micturition and erections may be present with convulsions and unconsciousness. Third the last stage is characterized by collapse, spasms, general paralysis and death. The subacute form of poisoning may ensue and prove fatal, or, owing to the volatile character of the drug, complete recovery may take place within one-half or three-quarters of an hour. Occasionally dogs continue to be paralyzed for several days and get well. The minimum fatal dose recorded in man is 9/10 of a grain of pure acid, or about 50 drops of the medicinal solution. Four to five drachms of the diluted acid frequently, but not invariably, cause minute poisoning and death, in horses, within an hour. One to two drachms of the pharmacopoeial preparation usually kill dogs within ten minutes.

Prussic acid is commonly used to destroy the domestic animals. Two to four drachms of the medicinal acid are to be given to dogs and cats of the ordinary size, and certain, painless, and rapid death will occur if a fresh preparation of the drug can be obtained. The unopened, half-ounce vial, kept by druggists, is recommended. Big dogs, horses, and the other larger animals are not killed rapidly, nor sometimes at all, by great quantities of the diluted acid. Hence, shooting is a more humane and preferable mode of death for them. In the experience of the writer, one to two drachms of prussic acid saturated with potassium cyanide, failed to kill a horse, when injected directly into the jugular vein. The odor of the acid cloggers about the animal for a few hours after death, the eyes are fixed and staring, the pupils dilated, the teeth are clinched tight and covered with froth while the blood is of a very dark color. The treatment embraces emptying the stomach by large doses of promptly acting emetics, or by the stomach tube or pump syringe, either and brandy subcutaneous, and inhalations of ammonia, together with artificial respiration, and hot and cold douches upon the chest.

Hydrocyanic acid is produced by a number of other plants referred to in another connection.

Wild cherry bark (*Prunus serotina*) and leaves by distillation yield a volatile oil resembling that of bitter almond. The same is true of the *P. virginiana*. Hydrocyanic acid is formed only by the action of a ferment upon amygdalin which is present in all plants of this sub-order. Fresh leaves are generally considered harmless, but Chesnut says that cattle are frequently poisoned from

eating the wilted leaves. He also adds that the seeds of all varieties of cherries and plums, both native and introduced are subject to suspicion. The flesh of none of the species is in any way poisonous. Chesnut also says in another contribution, that no cases are on record where stock have been poisoned by eating the leaves of any species, while still on the tree. It is only after they have been cut off and are partially wilted that they are considered dangerous. The reason for this is that during the process of wilting, prussic acid is formed from non-poisonous constituents which are always present in the leaf and bark. The wilted leaves have the characteristic odor of prussic acid. The amount of amygdalin contained varies from 3.6 to 4.12 per cent and yields from 0.23-0.32 hydrocyanic acid. Recent studies indicate that an amydonitril glucoside occurs in young leaves of both *Prunus Padus* and *P. serotina*.



Fig. 278. The European May Day tree (*Prunus Padus*), poisonous. This is frequently cultivated. (Ada Hayden)

LEGUMINOSAE

Herbs, shrubs, trees, or vines with alternate, mostly compound, stipulate leaves, papilionaceous or sometimes regular flowers, calyx 3-6, or 4-5 cleft, stamens 10, rarely 5 and sometimes many, monadelphous, diadelphous or distinct, pistil, simple, free, becoming a legume in fruit or sometimes a loment, ovules 1-many, seeds generally ex-albuminous or nearly so or, in some, with copious albumen. About 6500 species of wide distribution, but most abundant in the tropics.

Economic plants

The order contains a large number of economic plants, especially food plants. The common bean, (*Phaseolus vulgaris*), was originally found in the southwestern United States, but is now cultivated in all civilized countries. Common string and golden wax beans are types of the last named. The scarlet runner, (*P. multiflorus*), generally cultivated for ornamental purposes is also used as food, although the ripe beans are unwholesome and sometimes poisonous. The three-lobed kidney bean, (*P. trilobus*), is commonly cultivated in India. The lima bean, (*P. linatus*), also native to America, supposedly Brazil, is not known in a wild state. The seed of the Adzuki bean, (*P. Mungo*, var. *glaber*), is used as food in Japan. The soy bean, (*Glycine Soja* and *G. hispida*), of which there are many varieties, is native to China and Japan and is used in large quantities by the Japanese and Chinese for food, but is little used in the United States, being here cultivated as a forage plant.

Soy beans can only be fed in moderate amounts to cattle because of their purgative properties. A loss of a considerable number of cattle occurred in England recently where soy bean cake had been used. When fed mixed no trouble was caused, but when fed alone it caused poisoning.

The cow-pea, (*Vigna Catjang*), native to China, has been cultivated for centuries by the Chinese and extensively used for food. It is also used in many other warm countries, especially the southern states, not only for human food, but also as one of the best forage plants, for which purpose it is now




Fig. 279. Soja Bean (*Glycine hispida*), used both as a food and as a forage plant (U. S. Dept. Agr.)

cultivated as far north as Minnesota; it is also a soil renovator. The *Dolichos Lablab* is used for food in the tropics. The yam bean, (*Pachyrhizus angulatus*), in some countries, forms a resource as food in case of the failure of the usual crops. The garden pea, (*Pisum sativum*), probably originated in western Asia along the foothills of the Caucasus. It has, however, long been cultivated in Europe. Some authorities believe that it may have originated from the field pea (*Pisum arvense*). The pea is extensively cultivated in Europe and Canada as a forage plant. The chick pea, (*Cicer arietinum*), is a native to Caucasus and the Caspian Sea region and has been cultivated since



Fig. 280. Garden pea (*Pisum sativum*),
a valuable food plant. (U. S. Dept. Agr.)

remote times in northern Africa and other Mediterranean countries. In the United States, it is cultivated chiefly in the arid regions both for stock and for human food. The lentil (*Lens esculenta*) has been cultivated in the Mediterranean region for centuries but its original home is not known. It is grown and used now from Central Europe south and east to India, as food for both men and stock. The peanut (*Arachis hypogaea*), probably native to Brazil, was cultivated by the ancient Peruvians but is now widely scattered in all warm countries. The nut (seed) is used as food and a fine oil is extracted from it. A plant allied to the peanut (*Voandzeia subterranea*) is used as an article of food in western and southern Africa. The pigeon pea (*Cajanus indicus*) is an important article of food in the tropics, especially in India, and is also used as a fodder plant. The sword bean (*Canavalia ensiformis*) and the Jackpea (*C. obtusifolia*), cultivated in the tropics are used as food, the skin having been



Fig. 281. Pennut (*Lonicera hyemalis*)



Fig. 282. Broad Bean (*Vicia Faba*). Cultivated both as a food and as a forage plant. (From the American Agriculturist.)

first removed. The broad bean (*Vicia Faba*) is much cultivated in Europe for both animal and human food. The seeds of *Paprika africana* are used as food by native Africans and in Abyssinia and in the Indian Archipelago are esteemed as a good substitute for coffee.

The Australian wattles (*Acacia*), of which there are many species, vary greatly in size. The bark of some of these is used for tanning purposes. The wood is valuable and takes a fine polish, *A. Gerrardi* being an example of this class. From *A. Farnesiana* is derived the oil of cassia, much used in perfumery. It is prepared by macerating the flowers in olive oil. Cassia pomade is prepared from fatty substances to which the cassia flowers have been made to impart their perfume. *C. occidentalis* is used as a substitute for coffee. Many of the legumes produce important gums. From the Algarrobe, or locust tree of Jamaica (*Hymenaea Courbaril*) is produced a gum said to be superior to shellac, the sweet pulp of the fruit is edible. The gum Kino (*Pterocarpus Marsupium*) is a native of India and yields a gum that is used both for tanning and dyeing and as an astringent. Kino contains from 40-80 percent of tannin and kino red. *P. tinctorius* produces a valuable wood, and a related species (*P. Dalbergioides*) produces a wood similar to mahogany. The Tonka bean or Tonquin (*Dipteryx odorata*) of Guinea contains the substance cumarin and is used as a snuff and as a scent in cigars. Cumarin is widely distributed in the plant kingdom, especially in such Leguminosae as *Dipteryx*, *Melilotus*, and *Myrosylon*. It occurs in species of other families, as the palm, vernal



Fig. 283. Arwort (*Coronilla varia*). A poisonous plant of Europe—adventitious in the U. S. (From Strasburger, Noth, Schenck and Schimper)

grass, madder, rue, and orchids, in such composites as *Trilisa*, in the root-stock of *Vitis serratifolia* and in *Prunus Mahaleb*. It is most easily detected when the plant begins to wilt or after it is dead. *Pseudocumarin* $C_7H_6O_2$ is like the odorous substance found in *Coronilla scorpioides*. The seeds of *Mucuna giganteum* and of *M. pruriens* are used for various purposes as watch charms, or as other ornaments.

Some species of the family which contain a good fiber are applied in weaving cloth. A species of *Crotalaria*, *C. juncea* is cultivated in India for its tough fiber, and is used for making ropes and bags. *Sesbania aculeata*, a branched annual, is also cultivated for its fiber. The stems of Shola (*Aeschynomene aspera*) native to India are used for making hats. Blue indigo dyes come from *Indigofera Anil* of the West India. The *Genista tinctoria* or Dyers' Broom



Fig. 284. Hairy Vetch (*Vicia villosa*). Cultivated as a forage plant. (U. S. Dept. Agr.)

of Europe and Asia, and naturalized in the U. S. contains a yellow coloring principle. The seeds of *Entola scandens* are used in the Samoan Islands in playing games. The *Pithecolobium dulce* contains a pulpy pod which is eaten. The plant is good fodder. Guava (*Inga vera*) is grown as a shade tree and as a substitute for coffee. It is not to be confused with the fruit producing guava (*Psidium guajava*). Important forage plants not previously mentioned are red clover (*Trifolium pratense*), native of Europe and used extensively in northern United States, alsike clover (*T. hybridum*), white clover (*T. repens*), well known as a forage plant and a good honey plant, *T. alexandrinum* the great forage crop of Egypt, known as the Berseem, Japan clover (*Lespedeza striata*) a well known forage plant of the South, French honeysuckle (*Hedysarum coronarium*), an ornamental plant, native to Spain, also used as a forage plant of Europe and Western Asia, *Desmodium intolium* used as a forage plant in the tropics, Florida beggar weed (*Desmodium illinoense*) of India, alfalfa (*Medicago sativa*) of Europe and Western Asia, the vetches (*Vicia villosa* and *V. sativa*), lupines (*Lupinus albus*) cultivated for forage purposes, besides many valuable native forage plants like the *Horackia Purshiana*, the wild pea (*Lathyrus venosus*) etc.




Fig. 285. Dyer's Broom (*Genista tinctoria*) Contains a yellow coloring matter. (After Faguet.)

Prof. N. E. Hansen through his exploration has brought into prominence the yellow-flowered alfalfas from Siberia, *Medicago ruthenica* and *M. platycarpa*, and some of the hardy alfalfas are apparently hybrids between *M. sativa* and *M. falcata*. C. V. Piper* calls attention to a number of valuable leguminous forage plants that should be cultivated in this country, such as the Lyon bean (*Strolobium Lyon*). The Kudzu (*Pueraria Thunbergiana*) a woody native of Japan, is much used as a forage plant in that country and has been cultivated in Florida. The Guar (*Cyamopsis tetragonoloba*) is an East India annual legume and is said to be very drought resistant. Tangier pea (*Lathyrus tingianus*) is a native of Northern Africa. It is excellent and said not to be poisonous like the other species. The moth bean (*Phaseolus acutifolius*) is used for food in India and is said to be a splendid forage plant. The Adzuki bean (*Phaseolus angularis*) native of southern Asia is used for food in China, Japan and India, and is said to possess valuable qualities as a hay plant. The Kuli (*Dolichos biflorus*) native to India is said to give promise in the semi-arid regions in Texas as a valuable forage plant. Under the Vetches the more recent introductions that give promise are the scarlet vetch (*Vicia fulgens*) of Northern Africa, the black purple vetch (*V. atropurpurea*) of Algeria and the woolly nod



Fig. 286. Carobtree, or St. John's Bread (*Ceratonia siliqua*). The fruit is edible and is supposed to be the "locust" of Biblical history. (After Paget.)

* Yearbook U. S. Dept. of Agr. 1908-24.

vetch (*V. dasycarpa*) of the Mediterranean region, the Carob tree or St. John's bread (*Ceratonia siliqua*) is a small tree of the Mediterranean region, the pods of which contain much mucilage of a sweet nature from which syrup was made, and is supposed to be the locust which John the Baptist lived upon in the Wilderness and is used as food, and cattle also relish it. Manna (*Alhagi maurorum*) is a dwarfed, thorny, shrubby plant which produces a kind of manna. The locust trees of the West Indies, or Courabaril (*Hymenaea Courbaril*) produces a hard timber. The pods contain bean-like seeds embedded in a white spongy mass. The Zamang (*Pathecolobum Saman*) is a large tree of Venezuela which produces thick, flat pods, containing a sweet pulp commonly used by cattle and horses for food but which are liable to cause internal disorder. The honey locust of eastern North America (*Gleditsia triacanthos*) produces a hard wood. The pod contains a gummy sweetish substance much relished by stock. The pods of the mesquite tree (*Prosopis juliflora*), native to Texas, are used by stock. The Kentucky coffee tree (*Gymnocladus dioica*) is native to the Mississippi Valley and it produces a broad tough pod which contains large, hard seeds. The pod contains a sweetish, disagreeable and nauseating material more or less poisonous. The hard wood is durable. The seed of hairy vetch (*Vicia hirsuta*) is a common impurity in grain seed. The plant is used for forage. The seeds of *Castanospermum australe* are used in New South Wales in the production of starch. The seed of the coffee astragalus (*A. boottiana*) is said to produce, when roasted, the true coffee flavor, and is much used in Sweden.

Many of the species of the order are ornamental, among which may be named the Judas tree or red bud (*Cercis Siliquastrum*), the Caraganas, shrubs with beautiful yellow flowers the broom (*Cytisus scoparius*) of Europe naturalized along the sea coast, also used in medicine the black locust (*Robinia Pseudo-acacia*), commonly planted as an ornamental tree. The laburnum (*Laburnum anagyroides*) is an ornamental tree with poisonous seeds and hard wood, used for turned work. The wisteria (*Wisteria speciosa*) a hardy species of the southern states and *W. Chinensis* of China, are cultivated as far north as central Iowa. The seeds of several species of the order are used for making necklaces, among these are the red seeds of the coral tree (*Erythrina Corallidendron*) of the West Indies, the crab's eye (*Abrus precatorius*) and the *Ormosia dasycarpa*. The Jequrity seed (*Abrus precatorius*) is used as a weight in India, according to Dr. Spafford, each seed weighing approximately 1 gram.

Medicinal Plants. Of the medicinal plants, in this family, the most important only will be mentioned here. The broom (*Cytisus scoparius*), native to Europe, is used as a diuretic and purgative. Fenugreek (*Trigonella Foenum-graecum*) was formerly used in medicine, but the powdered seeds are now used as an ingredient of curry powder and also in the preparation of stock foods, they have a characteristic odor and bitter taste. Licorice root (*Glycyrrhiza glabra* and var. *glandulifera* (Rusman), native to the warmer regions of Europe and extending eastward into Central Asia, is made into extract of licorice which is used to cover the flavor of nauseous medicines and contains the glucoside *glycyrrhizin*. Cowhage (*Mucuna pruriens*) is a lofty climbing plant with dark purple flowers of the size of the sweet pea, pods from 2-4 inches long covered with rigid, pointed, brown hairs, which if touched, enter the skin and cause

itching. The plant is used for the expulsion of intestinal worms, the hairs being mixed with honey and molasses. The young pods are used as food. The Calabar or ordeal Bean (*Physostigma venenosum*) is a climbing perennial plant resembling the scarlet runner and is native to tropical western Africa. It was formerly used by the natives to produce poisoning and is also an antidote against strychnine poisoning. The seeds contain several poisons, among them the alkaloids *physostigmine* $C_{15}H_{21}N_3O_2$, which contracts the pupil of the eye, *calabarine* a tetanizing principle, and *eseridine* $C_{15}H_{23}N_3O_2$, a purgative. The wood of Araroba (*Andira araroba*) is very bitter and is used in ointments. Balsam of Tolu (*Myroxylon toluiferum*), a native of Venezuela and New Granada, is used as an ingredient in lozenges and contains a volatile oil *tolene* $C_{10}H_{18}$.

Logwood, (*Hæmatoxylum campechianum*) is a spreading tree, native to Central America and Honduras, the bark of which is used for dyeing and staining, also for domestic purposes, such as in chronic diarrhoea, and contains *hæmatoxylum* $C_{16}H_{14}O_8$ and *hæmatein* $C_{16}H_{14}O_8$. Senna leaves are derived from the leaves of Cassia (*C. acutifolia* and *C. angustifolia*), the former being found in Nubia and the latter in southern Arabia and India. Senna leaves are used in medicine as a purgative and contain *cathartic acid*, *chrysophan* and two bitter principles *sennairol* and *sennaptein*. Several of our native species of *Cas-*



Fig. 287. European Licorice (*Glycyrrhiza glabra*)
(After Engelm.)




Fig. 244. *Acacia (Acacia arabica)*. Flowering and fruiting branch. The source of gum arabic. (After Faguet.)

like the partridge pea (*C. Chamocrista*) produce scours in sheep because of their purgative properties. The purging cassia (*C. fistula*) is a tree indigenous to India. The pulp of the pod is a mild laxative. *Cassia ornata* of the Pacific Islands is a powerful cathartic. Tamarind (*Tamarindus indica*), a large, hardy tree indigenous to tropical Africa, is now widely distributed in other tropical countries. The fruit is used in medicine as a mild laxative and also in making a drink. It contains citric, tartaric, and other organic acids. The pulp and seeds are also eaten; the latter, when boiled, make a tenacious glue. The leaves and flowers are used as mordants in dyeing. *Copaiba balsam* (*Copaifera officinalis*) is a native to South Africa. The balsam is collected by Indians and used because of its stimulating action on the mucous membrane. It contains several acids, among them *copaibic acid*, $C_{20}H_{30}O_2$. Gum Arabic is obtained from *Acacia arabica*, the finest product coming from several species of the genus *Acacia* (*A. Senegal*), a plant well known to ancients. It possesses no real medicinal value. Catechu (*Acacia Catechu*), a small tree with thorny

branches, found in tropical eastern Africa, is used in medicine as an astringent. Under another name, Cutch, it is used in medicine because of its astringent properties. The resin from Cutch is made into cakes used for dyeing and tanning. Cutch contains *catechol* $C_6H_4O_2$ and *catechutannic acid*.



Fig. 289. *Copaiba Balsam* (*Copaifera officinalis*) Used medicinally (After Faguet)

Gum tragacanth (*Astragalus gummifer*) native to western Asia, is a spiny shrub with yellow flowers and is used to give consistency to lozenges, it contains *traganthin*, $C_{12}H_{16}O_8$ and *arabin*.

Poisonous and medicinal plants Most of the substances occurring in the poisonous plants are mentioned under the species described. Many of the *Leguminosae* contain alkaloids, few of these are, however, found in *Mimosaceae* although alkaloids have been found in *Acacia tenerrima*, *Albizzia lucida*, and *Pithecolobium Soman* which are *Mimosae*. Of the *Papilionaceae*, the *Sophoreae*, *Podalyriaceae* and *Genisteae* frequently contain alkaloids. *Cytisin*, found in seed of the *Ulex europaeus*, *sophornin* in *Sophora speciosa*, and *baptizoxin* are identical with *cytisin*. *Mustrin* $C_{11}H_{11}N_2O$, an alkaloid, resembling *lupanin* occurs in the root of *Sophora angustifolia*. *Anagyrin* $C_{15}H_{23}NO_2$ and *cytisin*, both occur in the seeds of *Anagyris foetida*, *Baptisia* and many other genera.




Fig. 240. Tamarind (*Tamarindus indica*). Fruit, flower and leaves. Fruit used in making a refreshing drink, seeds furnish a glue. (After Faguet.)

Retamin $C_{14}H_{24}NO_{10}$ is obtained from the young branches of *Genista sphaerocarpa*. The seed of *Trigonella Foeniculum-astrucum* contains *trigonellin* $C_8H_7NO_4$, the same alkaloid is said to also occur in the pea, hemp and oats. *Physostigmin* $C_{14}H_{21}N_2O_6$, or *eserin* occurs in the ripe seed of *Physostigma venenosum*. The so-called *calabrin* is a secondary product. *Robins* *Acon* is said to contain *nucovin*. *Pancos* $C_{27}H_{39}N_3O_5$ is found in the fruit of *Pentaclethra macrophylla*.

Quite a number of the Leguminosae also contain glucosides. One of the earliest discoveries made in connection with glucosides was of *glycyrrhizin* found in some species of *Astragalus*, *Ibura pectoratus*, and the root stock of *Polypodium vulgare* and other plants. The root of our wild licorice also contains a glucoside to the extent of 8.53 per cent. *Glycyrrizin*, found in licorice, has the formula $C_{42}H_{64}O_{16}$.

Ononis spinosa contains a glucoside *ononin* and a second glucoside *ononin* $C_{28}H_{44}O_{12}$. *Lupinin* $C_{24}H_{34}O_{10}$ is a glucoside found in the seedlings of *Lupinus intens*, which through hydrolysis forms *lupigin*, $C_{14}H_{22}O_4$. *Gastrolobin* is found in the leaves and young branches of *Gastrolobium b. obam*, *bap*.




Fig. 291 Purging Cassia (*Cassia Fistula*) Flowering and fruiting branches. The pulp of the pod is a mild laxative. (After Faguet.)



Fig. 292 Cow Pea (*Vigna Catjang*) A well known forage plant of the South. To the right a legume, to the left a flower and a part of a branch in the lower left hand corner (W. S. Dudgeon.)

ham, $C_{28}H_{52}O_{14}$, occurs in *Baptisia*, and *lephrosin*, a poisonous substance in *Tephrosia toxicaria*. A very toxic unnamed glucoside, $C_{28}H_{52}O_{16}$, occurs in the bark and root of *Derris elliptica*, *Mundulea suberosa*, and *Lonchocarpus violaceus*. Power, however, failed to find this glucoside in *Derris wagnara*. *Turboin*, $C_{24}H_{42}O_{10}$, occurs in *Tephrosia toxicaria*. *Gallotannic acid*, $C_{16}H_{16}O_8 + 2H_2O$ so called, occurs in the pod of *Cassalpinia coriaria*. The bark of certain species of *Acacia* contains 30 per cent of tannic acid. The wood of *A. Catechu* is colored red by *catechin*, a crystallizable substance.

Lindley in the earlier edition of his Vegetable Kingdom, states, that the plants of this family are on the whole wholesome and nutritious, and later declares that the family must be considered poisonous. The species used as food must be considered an exception.

In Australia there are several members of this family that are poisonous. Dr Gray in an American Agriculturist of Oct., 1878, says

What a pity that our cattle are not better acquainted with the corrected rule. In Europe and in the Atlantic States, no harm is known to come to cattle from want of proper discrimination. But when European flocks were taken to Australia and to pastures and forage almost wholly new, thousands of sheep perished in the Swan River Valley Colony in consequence of cropping the leaves of some leguminous plants to which they were attracted. What made the matter worse for the botanists, was that the very plants which did the mischief had been recommended by one of them (M. Presl, a German) as the best thing the Agricultural Society could cultivate, as artificial food for stock. But another botanist, Drummond, a savvy Scotchman, made some experiments, that proved that the people were right in charging the damage to these very species of *Gastrolobium* which the German botanist, on general principles, expected to be innocent and useful.

The Australian *Gastrolobium* are all more or less poisonous. Baron Müller having long ago reported *Gastrolobium grandiflorum* as poisonous. The *G. edyemum* known as the York road poison has a toxic base *cyanic cyanic acid* $C_2H_{10}O_4$ which decomposes and forms *gastrolobic acid* $C_7H_{10}O_4$. The following species are recorded as poisonous by Maiden in Australia: *G. trianthemum*, *G. polystachyum*, *G. grandiflorum*, poisonous to sheep and goats, the seeds being especially toxic but not to pigeons. The diseased animals have difficulty in breathing, then they stagger and die, death occurring in from 3-6 hours. The poison enters the circulation, stops the action of the lungs and heart. The raw flesh is said to poison rats, and the blood, dogs. The boiled or roasted flesh is, however, eaten by the natives and is not poisonous. The blossoms are very poisonous. The *Mirbelia racemosa* is also poisonous to sheep, cattle, and goats. Two species of *Goodia* according to Maiden are poisonous, the *G. lotifolia* and *G. medicaginea*. These plants produce what locally goes by the name "back scours." The animals become weak, emaciated, and die. The *Gastrolobium* and *Erotarium* are stock killers in Australia though used as forage in South Australia. Maiden reports that the bean tree (*Castanospermum australe*) is poisonous to stock especially the beans, when cooked, however they are eaten by the Aborigines. The box poison (*Oxylobium paniculatum*) is said to be a very poisonous plant to stock. The *Gompholobium uncinatum* is very injurious to sheep in New South Wales. The *Suaresona Greyana* and *S. coronataetolia* are poisonous. Sheep that eat them are called indigo-eaters. Both species act much like the loco weeds of the United States, "sheep go wrong in the head," horses also act strangely. "The eyes stand out of their heads." This disease is identical with the "Nenta" disease of South Africa and the "Pea eating" disease of Australia. The South African disease is produced by *Lessertia*. The symptoms

from *Astragalus mollissimus*, *Gompholobium*, *Sophora secundiflora*, *Cytisus procylliferus*, are all cerebral. Mac Owen regards them all as belonging to the same category and that *Lathyrus* caused by *Lathyrus sativus* is allied to them. That the well known forage plants *Lotus corniculatus*, and *L. australis*, of excellent repute, are often injurious to stock, but perhaps only from causing indigestion, as stated by Maiden, is worthy of note in this connection. Moussu and Desain report the deaths of a flock of 54 sheep due to poison resulting from eating another plant of this order, *Galega officinalis*, 80 others in the same flock were badly affected. Ecchymoses were found in the walls of the alimentary tract and in the fatal cases a large amount of serum had collected in the pleural cavity. In subsequent experiments, it was found that 3 kg. of *G. officinalis* was sufficient to poison a sheep, the plant, however, seems not to be poisonous to rabbits. It is of interest to note that the genera *Robinia*, *Indigofera*, *Wisteria*, *Caragana*, *Colutea*, *Swainsona*, *Galega*, *Lesqueria* *Astragalus*, and *Sesbania* all of which contain poisonous species, many of which are exceedingly toxic, belong to the tribe *Gageae* of this family. *Glycyrrhiza* of the same tribe is not poisonous nor are all of the species in genera like *Astragalus*, *Caragana* etc.

The *Erythrophium guineense* contains an alkaloid *erythrophium* which acts like digitalin and picrotoxin. The Indigo of Australia (*Indigofera australis*) is regarded as poisonous in Australia. The *Lathyrus sativus*, *L. cicera*, and *L. clymenum* are poisonous but the active principles have not been isolated. Of other poisonous genera *Robinia*, *Baptisia*, *Gymnocladus* *Thermopsis* may be mentioned, but the treatment will be given more at length in another connection.

According to Dunstan and Henry, *Lotus arabicus*, when moistened with water and crushed, produces prussic acid. The glucoside *lotusin* is converted by the enzyme *lotase* into prussic acid. We may mention here that many of the spiny *Acacias* (*A. pallens*) of Australia may be injurious in a mechanical way.

The poisonous substance of Jequirity (*Abrus precatorius*) is a toxalbumin called *abrin* (found also in *Cassia huspidata* of Mexico) which is easily decomposed by heat. Behring has produced an antitoxin against the *abrin* or *A. precatorius*. The beans when cooked are eaten in Egypt.

A poisonous resin has been found in *Wisteria chinensis*, and a glucoside *wisterin*. *Colutea arborescens*, a well known southern European ornamental plant is poisonous, the leaves being so strongly purgative that they are frequently substituted for the genuine Senna. European authors list it among the poisonous plants.

The *Tephrosia purpurea* of Australia is poisonous to stock, and is used to stupefy fish. A large number of other plants of the order are used as fish poisons. Mention may be made of *Derris*, *Abrus* and *Clethra*, others are mentioned in Part I. Some like *Azadirachta* and *Pithecolobium* are used as arrow poisons. The seeds of the jequirity plant (*Abrus precatorius*) are much used in India for the purpose of poisoning especially in criminal cases of cattle poisoning less than 2 grams of the powdered seed causing death in 48 hours. The usual method of the "Chamar" or "Skinner" caste is to prepare small spikes, first soaking the seeds in water, then pounding them, and drying them in the sun, they are then sharpened upon a stone attached to a handle, and driven under the skin and left there. Daggers are rendered poisonous by being dipped into the powdered seed.

According to Greshoff the leaves and also the seeds of *Cassia marylandica* contain saponin, leaves of *Prosopis juliflora*, *Galega officinalis*, *Psoralea macrostachya* as well as the seeds contain saponin. *P. tenuiflora* is regarded as poisonous and is avoided by cattle.

Corolla not papilionaceous or only slightly so, endosperm copious.

Flowers perfect, leaves abruptly pinnate. 1 *Cassia*.

Flowers polygamous or dioecious 2 *Gymnocladus*.

Corolla papilionaceous without or with endosperm, stamens usually 10, usually diadelphous or monadelphous.

Stamens 10, distinct.

Leaves palmately 3-foliolate.

Pod inflated 4 *Baptisia*.

Pod flat 3 *Thermopsis*.

Leaves pinnate 5 *Sophora*.

Stamens monadelphous, diadelphous, or rarely distinct

Anthers of 2 forms, stamens monadelphous

Leaves simple.

Pod inflated 6 *Crotalaria*.

Pod flat.

Leaves 1-3-foliolate 8 *Cytisus*.

Leaves usually 7-11 foliolate 7 *Lupinus*.

Anthers all alike.

Leaves generally 3-foliolate.

Flowers in racemes, pods coriaceous 9 *Melilotus*.

Flowers in heads, pods membranous 10 *Trifolium*.

Flowers in spikes or heads, pods curved 10 *Medicago*.

Leaves pinnately foliolate.

Leaves not tendril bearing, plants not climbing

Herbs with glandular dots 12 *Psoralea*.

Herbs without glandular dots; pods flat

Leaves odd pinnate.

Herbs. Flowers large 13 *Tephrosia*.

Trees or shrubs 12 *Robinia*.

Leaves evenly pinnate 15 *Sesbania*.

Pod turgid inflated.

Leaflets not toothed, or only at the apex

Keel tipped with an erect point 17 *Oxytropis*.

Keel not tipped with an erect point 16 *Astragalus*.

Leaflets toothed all round 20 *Cicer*.

Herbaceous plants, leaves with tendrils or climbing.

Leaves with tendrils, style bearded at the apex 18 *Vicia*.

Style bearded down one side 19 *Lathyrus*.

Herbaceous climbers, not tendril bearing 21 *Phaseolus*.

1. *Cassia* L.

Herbs, shrubs, or in tropical regions, trees, with abruptly pinnate leaves, calyx of 5 sepals united at the base, petals 5, somewhat unequal, spreading, imbricated, and clawed, stamens usually 10, or 5, often unequal, and some imperfect, anthers all alike, or the lower larger, opening by 2 pores at the apex,

ovules numerous. About 220 species, mostly in warm and temperate regions. A species well known in medicine is Senna (*C. acutifolia* and *C. angustifolia*) with leaves which are laxative.

Cassia Chamæcrista L. Partridge Pea

An annual, spreading, 1 foot long; leaves with a sessile gland on the petiole, leaflets of 10-15 pairs, flowers large, showy, petals yellow, with a purple spot at the base, anthers 10, elongated, and unequal, 4 yellow, the others purple.

Distribution. In dry or sandy soil from Maine to South Dakota, Texas to Florida.



Fig. 291. Pea Partridge (*Cassia Chamæcrista*) c. Pod.
a. Petiole. b. Stamens. (C. M. King)

Poisonous properties. This plant is common in hay and when the seeds are consumed in large quantities, has a cathartic action. Cases of mild poisoning to sheep have been reported to the writer. *C. marilandica*, a plant with curved pods that are somewhat hairy at first, possesses similar properties. *C. hirsutula* contains abrin.

2. *Gymnocladus*, Lam.

Trees with large, bipinnate leaves, and showy, white, dioecious or irregular, polygamous flowers, calyx elongated-tubular below, 5-cleft, the lobes narrow, nearly equal, petals 5 (rarely 4), oblong or oval, stamens 10, distinct, short,



Fig. 294. Wild Senna (*Cassia Marylandica*). 1. Flower. 2. Pod. A plant growing in the Eastern Atlantic States as far south as North Carolina. Laxative like the Common Partridge pea (Selby Ohio Agri. Exp. Stat.)

inserted on the petals; ovary rudimentary, or none in the staminate flowers, sessile and many-ovuled in the pistillate, pod oblong, thick, large, and coriaceous.

Gymnocladus dioica (L.) Koch. Kentucky Coffee-tree

A large tree with rough bark, leaves large and ample, 2-3 feet long, 7-15 leaflets, ovate or acute, glabrous or pubescent on the veins beneath, racemes many-flowered flowers slender-pedicelled; seeds hard, $\frac{1}{2}$ inch across, imbedded in a sweet, but disagreeable, and somewhat mucilaginous, material.

Distribution. From Western New York to Pennsylvania, Eastern Nebraska, and Arkansas.

Poisonous properties. Cases of poisoning are not uncommon. The alkaloid *cylasin* $C_7H_{14}N_2O$, a crystalline, rather bitter, and caustic substance which causes dilation of the pupil, is reported to have been found according to Chesnut, in the leaves and soft pulp of the fruit of the coffee bean. The pulp has long been used, when mixed with milk, to poison flies. In speaking of the symptoms and treatment, Prof Chesnut says:

Few accidental cases of poisoning arise, but the pulp, in one instance, caused severe illness in a woman who ate a small quantity mistaking it for that of the honey locust (*Gleditsia triacanthos*) which is frequently eaten by children. The symptoms were not fully noted at the time, but are described from memory as conspicuously narcotic. The effect began within five minutes and lasted several hours. The treatment should probably be the same as that for laburnum, viz., emetics, stimulants, injections of coffee, and an alternately hot and cold douche to the head and chest.




Fig. 295. Kentucky Coffee Tree (*Gymnoclad dioica*). 1. Leafy branch from sterile tree. 2. Portion of branch with long winged seed. 3. Long winged seed. 4. Longitudinal section of winged seed. 5. Longitudinal section of flower. 6. Longitudinal section of flower. 7. Portion of branch bearing a long winged seed. 8. Longitudinal section of flower. 9. Longitudinal section of flower. 10. Longitudinal section of flower. 11. Longitudinal section of flower. 12. Longitudinal section of flower. 13. Longitudinal section of flower. 14. Longitudinal section of flower. 15. Longitudinal section of flower. 16. Longitudinal section of flower. 17. Longitudinal section of flower. 18. Longitudinal section of flower. 19. Longitudinal section of flower. 20. Longitudinal section of flower. 21. Longitudinal section of flower. 22. Longitudinal section of flower. 23. Longitudinal section of flower. 24. Longitudinal section of flower. 25. Longitudinal section of flower. 26. Longitudinal section of flower. 27. Longitudinal section of flower. 28. Longitudinal section of flower. 29. Longitudinal section of flower. 30. Longitudinal section of flower. 31. Longitudinal section of flower. 32. Longitudinal section of flower. 33. Longitudinal section of flower. 34. Longitudinal section of flower. 35. Longitudinal section of flower. 36. Longitudinal section of flower. 37. Longitudinal section of flower. 38. Longitudinal section of flower. 39. Longitudinal section of flower. 40. Longitudinal section of flower. 41. Longitudinal section of flower. 42. Longitudinal section of flower. 43. Longitudinal section of flower. 44. Longitudinal section of flower. 45. Longitudinal section of flower. 46. Longitudinal section of flower. 47. Longitudinal section of flower. 48. Longitudinal section of flower. 49. Longitudinal section of flower. 50. Longitudinal section of flower. 51. Longitudinal section of flower. 52. Longitudinal section of flower. 53. Longitudinal section of flower. 54. Longitudinal section of flower. 55. Longitudinal section of flower. 56. Longitudinal section of flower. 57. Longitudinal section of flower. 58. Longitudinal section of flower. 59. Longitudinal section of flower. 60. Longitudinal section of flower. 61. Longitudinal section of flower. 62. Longitudinal section of flower. 63. Longitudinal section of flower. 64. Longitudinal section of flower. 65. Longitudinal section of flower. 66. Longitudinal section of flower. 67. Longitudinal section of flower. 68. Longitudinal section of flower. 69. Longitudinal section of flower. 70. Longitudinal section of flower. 71. Longitudinal section of flower. 72. Longitudinal section of flower. 73. Longitudinal section of flower. 74. Longitudinal section of flower. 75. Longitudinal section of flower. 76. Longitudinal section of flower. 77. Longitudinal section of flower. 78. Longitudinal section of flower. 79. Longitudinal section of flower. 80. Longitudinal section of flower. 81. Longitudinal section of flower. 82. Longitudinal section of flower. 83. Longitudinal section of flower. 84. Longitudinal section of flower. 85. Longitudinal section of flower. 86. Longitudinal section of flower. 87. Longitudinal section of flower. 88. Longitudinal section of flower. 89. Longitudinal section of flower. 90. Longitudinal section of flower. 91. Longitudinal section of flower. 92. Longitudinal section of flower. 93. Longitudinal section of flower. 94. Longitudinal section of flower. 95. Longitudinal section of flower. 96. Longitudinal section of flower. 97. Longitudinal section of flower. 98. Longitudinal section of flower. 99. Longitudinal section of flower. 100. Longitudinal section of flower.

3. *Thermopsis* R. Br.

Perennial with finely appressed pubescence. 2-3 feet high. leaves rhombic-ovate leaves and lobaceous stipules. flowers large yellow or purple borne in racemes. calyx bell shaped or short tubulate, with equal and separate lobes or the upper united. standard nearly orbicular as long as the wing and the keel, stamens 10, separate and incurved. pistils sessile or short stalked, frequently flat, linear, oblong or curved. ovules numerous. A small genus of about 15 species of North America and Asia.

Thermopsis mollis (Muhl.) M. & A. Curtis. Alleghany. The virginian

Perennial with finely appressed pubescence. 2-3 feet high. leaves rhombic-ovate lanceolate 1-3 inches long entire and nearly sessile. stipules ovate or lanceolate,

racemes chiefly terminal, flowers yellow, pod short-stalked and narrow and somewhat curved.

Distribution. In the mountains of Virginia, North Carolina and Tennessee.

Thermopsis rhombifolia Richards. Prairie *Thermopsis*.

An erect perennial from 1 2½ feet high, appressed, silky pubescent, stem angular, leaves with broad conspicuous stipules, leaflets obovate, at length nearly glabrous, bracts oval, flowers yellow, in a rather short raceme of few flowers pod linear and curved, spreading, several seeded.

Distribution. In sandy soil and foot hills of the mountains from Manitoba to South Dakota, Nebraska and Kansas, west to the Rocky Mts., and in Utah, Wyoming and Montana.

Poisonous nature. This plant is very common in the foot hills, and is supposed to produce poisoning of stock. It is often consumed by sheep. It is said that the seeds of the plant are poisonous and the Canadian Department of Agriculture reports several cases of poisoning to children where the seeds were eaten. *T. montana* is a species occurring from western Nebraska and Kansas to the Pacific Coast. Species of *Thermopsis* are said to contain *cytisin*.

4. *Baptisia* Vent

Perennial herbs with palmately 3-foliate, or rarely simple leaves, basal sheathing scales, flowers large, in racemes; calyx 4-5-toothed, corolla with a



Fig. 296. Yellow-flowered Bitter weed (*Thermopsis montana*). Plant is exceedingly bitter. (From U. S. Dept. Agr.)

large standard, but not longer than the wings, stamens 10, distinct, pods stalked, roundish, oblong, inflated, and many-seeded, seeds often spreading and rattling. About 16 species in Eastern North America.

Baptisia australis (L.) R. Br. Blue False-Indigo

Tall, smooth, stout perennial 4-5 feet high; leaves short-petioled, leaflets oblong or oblanceolate, obtuse, stipules conspicuous, persistent, racemes terminal, loosely flowered, 1-2 feet long, erect, flowers blue.

Distribution. From Western Penn. to Arkansas and Kansas to Ga.

Baptisia tinctoria (L.) R. Br. Yellow or Indigo Broom

A smooth, slender perennial herb 2-3 feet high, leaves nearly sessile, leaflets obovate or oblanceolate, sessile or nearly so, racemes few-flowered, flowers yellow.

Distribution. In dry soil from Maine to La., west to Minn.

Poisonous properties. *Baptisia* which is probably the same as *cytaria* occurs in *Baptisia tinctoria*. The glucoside *baptisin* $C_{16}H_{23}O_{14}$ occurs in some species of the genus.

Dr. Milspaugh states that disturbances produced by taking considerable quantities of the tincture are

Vertigo, dull, heavy headache with weakness and weariness of body, and tendency to delirium, soreness and lameness of the eyeballs, with hot, flamed face, tongue coated white, yellow or yellowish-brown, loss of appetite, nausea, and burning in the stomach,



Fig. 297 Wild Indigo (*Baptisia bracteata*). Said to be poisonous. (Ada Hayden.)

dull pain in the region of the liver, especially at the site of the gall-bladder; face yellow, with burning cheeks, constant pain and aching in the abdomen, followed by marked distention, and soreness on pressure.

According to Dr. Hughes, *Baptisia* excites true primary pyrexia in the human subject. This pyrexia is very much like that of the early stages of typhoid.

Baptisia leucantha T & G. Large White Wild Indigo

A smooth, erect perennial herb, petioled leaves, leaflets obtuse, rounded, or sometimes slightly emarginate, stipules deciduous, racemes lateral; flowers white or cream color. Prairies and alluvial soils, Ont. to Minn., to Fla. and La.

Poisonous properties. According to Hyams it is a violent emetic and cathartic when taken in large doses and in small doses a mild laxative.

Baptisia bracteata (Muhl.) Ell. Large-bracted Wild Indigo

Perhaps more common in sandy soil in the west than *B. leucantha*. It is also shorter and flowers earlier in the season. Prairies, Mich. to Minn., La. Tex.

Poisonous properties. Dr. Schaffner states that the blue wild indigo and the yellow wild indigo are emetic, and that the latter species is regarded as



Fig. 298. Wild Indigo (*Baptisia leucantha*). This plant is a native of prairies and alluvial soils. Said to be poisonous. (Ada Hayden.)

poisonous. The taste of the plant is rather disagreeable, and it is not generally eaten by stock.

5. *Sophora* L.

Shrubby or herbaceous perennials, leaves odd pinnate with numerous leaflets, flowers white, yellow or violet in terminal racemes or panicles, calyx bell-shaped, with short teeth, standard rounded or obovate, wings obliquely oblong, keel oblong, nearly straight; stamens all distinct or nearly so, pistil short-stalked, style incurved, pod stalked and terete, constricted between the seeds. About 25 species, of warm and tropical regions.

Sophora sericea Nutt Silky Sophora

A low perennial herb from 6-12 inches high, more or less silky canescent, stipules subulate, deciduous, leaflets about 21, elliptical or cuneate oval, flowers in short terminal racemes, nearly sessile, corolla white, pod dry leathery, finely pubescent and few seeded.

Distribution. The prairies of Nebraska and Colorado to Texas and Arizona.

Poisonous properties. Mr Chesnut says of this plant

The silky sophora, of the Southern Great Plains region, has been somewhat vaguely reported as one of the plants that "loco" horses in that region. The seeds contain a very poisonous alkaloid.



Fig. 299. Silky Sophora (*Sophora sericea*). A plant of the plains and thought to be poisonous. (Charlotte M. King.)

Sophora secundiflora DC Coral Bean

A stout shrub or small tree, with deep green leaves of about 9 elliptical, oblong, obtuse, coriaceous leaflets, terminal racemes of showy violet, fragrant flowers, and large, woody pods 3-5 inches long, containing 3-4 round red beans as large as small marbles.

Distribution. Common from the Gulf Coast to the Pecos and less abundant in mountain canons to New Mexico. It is mostly shrubby, but becomes a tree 30 feet high and forms groves in the vicinity of Matagorda Bay.

Poisonous properties. Mr. Cheanut says

The beautiful bright red berries of the Frijolillo or coral bean of southern and western Texas contain a powerful poisonous alkaloid. The plant is said to have poisoned stock in Texas and in northern Mexico.

It contains *sophorin*, an amorphous alkaloid, which, according to Czapek, is probably identical with *cytisin*. The beans are somewhat used by the Indians to produce intoxication.

6. *Crotalaria* (Dill) L. Rattle-box

Herbs or occasionally somewhat woody plants, with simple or 3-7 foliolate leaves, yellow flowers borne in racemes, calyx 5-cleft, somewhat 2-lipped, standard large, heartshaped, wings oblong or obovate, keel curved, stamens monadelphous, with anthers of 2 forms, pod inflated like the pea, but shorter and many-seeded. About 250 species found chiefly in the tropics.

Crotalaria sagittalis L. Rattle-box

Annual from 3 inches to a foot high, with a small straight root; stem branched, vilious, terete or wing margined; leaves oval or oblong-lanceolate,



Fig. 300. Rattlebox (*Crotalaria sagittalis*). a, whole plant, b, cross section of seed pod—both one-third natural size. The cause of crotalism in horses. (U. S. Dept. Agr.)

from 1/2-1/3 of an inch wide, edge of the leaf entire or somewhat wavy and hairy stipules united and decurrent on the stem, inversely arrow shaped, peduncles few-flowered, flowers yellow, about 1/4 of an inch in diameter, calyx 5-cleft, standard of the flower large, heart-shaped, keel scythe-shaped, stamens monadelphous, anthers of 2 sizes, 5 smaller and roundish pod large, inflated, bears a close resemblance to the garden pea, greenish at first, becoming blackish, seeds from 1/10-1/12 of an inch in diameter, flattish, kidney-shaped, which when mature break away from the point of attachment and rattle in the pod, hence the name "rattle-box."

Distribution This plant is common in sandy soil from Maine to Minnesota, South Dakota, Iowa, Nebraska, and Northern Texas. The plant is extremely common on the sandbars of the Missouri river, where it may be collected by the wagon load.

Poisonous nature The earliest mention of the poisonous nature of the weed was made by Drs. Stalker and Hessey. Dr. Stalker who performed some experiments with the plant gives the following symptoms:

The disease had been known in this region for three or four years, but had not until the present summer (1884) prevailed to such an extent as to attract general public attention. But now the loss in horse stock on some farms was not to be counted by hundreds, but by thousands of dollars. The disease proved to be one that had not hitherto come within the range of my experience nor had I any intimation of anything exactly identical with it. I spent several days among the farmers on the lower side of the Missouri river taking careful notes of the symptoms, and gathering the history of the progress of the disease. On some farms I found almost all the horses affected, and on others but a few individuals. Deaths were an almost daily occurrence, and the farmer who owned a large stock of horses did not know today whether he would have teams for his farm work a week later. The disease in most cases is very slow in its progress, but proving almost uniformly fatal after a number of weeks or months. There is a general decline of bodily vigor throughout this period, and the only abnormal symptom in many cases is that of marked emaciation and consequent weakness. Horses that have been kept at pasture through the summer without work, and where the grass grew in greatest abundance were so thin in flesh that they walked with the greatest difficulty. A closer examination of many of these patients revealed nothing more than the condition resulting from starvation. This was not uniformly the case. In a number of instances there was marked coma or stupor, the animal often falling asleep while eating. In some instances the animal would remain standing for a whole week, sleeping much of the time with head resting against some object. In a few instances the animal lost consciousness, and broke through fences and other obstructions. A number of the diseased animals were placed at my disposal and assisted by Dr. Farchild and Dr. Miles. I made post mortem examinations of five subjects with the most perfect informality as to the lesions presented. In every instance there were marked hæmorrhagic effusions into the fourth ventricle, the liver and spleen were enormously dense, the walls of the intestines were almost destitute of blood, and the stomach enormously distended with undigested food. The stomach with its contents in some cases was weighed as much as seventy pounds. These post mortem conditions, together with the clinical symptoms led me to believe the animals were obtaining some poisonous principle with their food. The symptoms in some cases bore such a resemblance to those produced by eating *Asaragolus multiflorus* or "locoweed" of the Western plains, as to direct my investigations to that family of plants. A careful examination of the meadow and pasture areas was not rewarded by the discovery of a single "loco plant."

It took but little investigation, however to find a closely related plant growing in great abundance, both in the meadows and pastures. This was the *Crotalaria sagittalis*, or rattle-box. This is also known as the wild pea, and is accounted by many farmers as the best of forage plants. Knowing the bad reputation of some of its near relatives, I determined to make some experimental tests with the plant. I employed a boy to collect about thirty pounds of the green plants, which I brought with me on my return to the college. I procured a strong young horse, affected with incurable catarrh, and attempted to induce him to eat the plant. This he persistently refused to do, though I sharpened



Fig. 300a. White Lupine (*Lupinus albus*). A forage plant introduced from the Mediterranean region. Seeds contain a bitter alkaloid. U. S. Dept. Agr

his appetite by a protracted fast. It is a matter of common observation that animals eat it with the greatest relish in localities where it grows. Failing to induce the animal to take the plant voluntarily I prepared a strong infusion, and by means of the stomach pump gave the preparation obtained from about ten pounds of the plant. In twenty minutes stupor began to ensue, the eyes were closed, the head was rested against the side of the box, the breathing became stertorous, and all the symptoms developed that were to be seen in the patients previously examined. At the end of six hours the stupor began to disappear, the eye began to regain its brightness and in another hour the horse began to eat. The following day, when he had apparently recovered from its effects, he was given half the quantity of the drug as on the previous day. In this instance the symptoms were developed much more rapidly the animal became unconscious in a short time and died in an hour and a half. The post mortem revealed the same condition of the brain as in the cases examined in the Western part of the State. I now resolved to make a second experiment, in which the animal should receive a small quantity for a number of days in succession. Having procured another subject for experimentation, and a bushel of mature fruit or pods of the plant, I commenced on Sept. 5th, to give daily the infusion obtained from about one quart of the pods. On the fifth day of the experiment the characteristic stupor came on. The animal rested its head against the box and slept while standing. The symptoms grew more marked till the thirteenth day of the experiment, when the animal died. The post mortem showed the same as in the other cases. These experiments leave no doubt in my mind that the trouble along the Missouri river is occasioned by the animals' feeding on this little plant. It is from eight inches to a foot in height, with branching stems bearing yellow flowers in July and developing large pods resembling the pea, but containing a number of black, hard seeds. It grows on sandy bottom land, and is very abundant in the meadows and pastures in portions of the Missouri bottom. It is seldom seen among the tame meadow grass in any considerable amount. It thrives best among the wild grasses. Animals, doubtless, eat it much more than formerly, when the wild pasturage was better than at present. Cattle sometimes, though not often, suffer in the same way as horses.

The disease is also known as the Missouri Bottom disease. Hundreds of horses in the Missouri Bottom in Western Iowa and Eastern Nebraska die from eating this weed, it being most common in unbroken fields. Horses should be kept from all suspected fields. Only cultivated grasses and forage plants should be grown. Some doubt has been expressed that this plant is the cause of the trouble. The writer a number of years ago, in collaboration with Dr. Miller, investigated an outbreak near Council Bluffs. This disease occurred only in the bottoms, where the weed was common, and a large number of horses die from it annually. No other injurious plants could be found except some ergot on wild rye. A decoction of the weed found here was fed by Dr. McNeill to a horse but no injurious symptoms followed. A decoction of the seeds was fed to a guinea pig without any serious symptoms. Dr. F. B. Power however found a small amount of an alkaloid in the seeds which caused slight illness in a kitten. From all of these experiments we may conclude that rattie box is injurious under some conditions.

7. *Lupinus* (Tourn.) L. Lupine

Herbs or rarely shrubs with generally palmately compound leaves, stipules adherent to the base of the petiole; flowers showy, in long, dense racemes, calyx deeply toothed and 2-lipped, corolla with an orbicular or ovate standard with margins reflexed, wings oblong, or obovate, lightly cohering, and enclosing the keel, which is incurved or beaked, stamens monadelphous, anthers of 2 forms; pistil with an incurved style and sessile ovary, pod flattened, somewhat constricted. About 100 species of temperate regions, or a few in warm regions. The North American species are chiefly west of the 100th meridian. The *Lupinus perennis* occurs in sandy soil from New England to Minnesota and Louisiana, *L. albus*, *L. luteus* and *L. angustifolia* are cultivated for forage in Europe, the seeds being used as a substitute for coffee.

Lupinus argenteus Pursh. Hairy Lupine

A much branched perennial, slightly shrubby, from 2-3 feet high, silky pubescent hairs appressed, leaves with small stipules, petioles equalling or longer than the leaves, leaflets sessile, narrowed at the base, flowers in rather dense, terminal racemes, purple, pod silky, pubescent, generally 3-5 seeded. A very variable species.

Distribution. Prairies of South Dakota to Western Nebraska to New Mexico, Utah and from Arizona to Montana. Abundant in the foothills.

Lupinus perennis L. Wild Lupine

Perennial, somewhat hairy, erect stems, 1-2 feet high, leaves compound, 7-11 oblanceolate leaflets, flowers showy, purple-blue, in a long raceme, pods broad, very hairy, 5-6-seeded.

Distribution. Sandy soil from New England to Minnesota, Missouri, and the Gulf region.

Lupinus plattensis Watson. Nebraska Lupine

Somewhat like the preceding, with appressed silky-villous hairs, and a glaucous hue, leaflets spatulate, flowers in loose and short peduncled racemes, petals pale blue.






Fig. 30. Wild Lupine (*Lupinus Flattensis*). Causes lupinosis. Charlotte M. King

Lupinus leucophyllus Dougl.

Leafy, densely silky-tomentose perennial, compound leaves, 7-10 oblong-lanceolate leaflets, flowers in sessile racemes, densely flowered, petals blue or pink.

Distribution. Rocky Mountains, Colorado to Washington, and Northern California.

Lupinus holosericeus Nutt

A perennial, shrubby plant with silvery-canescens leaves, 12-20 inches high, compound leaves, 5-9 lanceolate leaflets, flowers in whorls or scattered calyx bracteolate, the upper slightly 2-cleft, petals bright blue.

Distribution. Oregon to California.

Lupinus species. There are many other species of *Lupinus* in the Rocky Mountains and along the Pacific Coast. A great many of these have been looked upon with suspicion.

Poisonous properties. European white lupine, *Lupinus albus*, *L. luteus*, and others contain the glucoside *lupinin* $C_{24}H_{34}O_{10}$, a crystalline substance with a bitter taste and a frothy odor, *lupindin* $C_{11}H_{14}N_2$, a pale yellow, heavy, oil with a pungent, bitter taste, *lupinum* $C_{10}H_{14}NO$, also bitter with an apple-like odor. *Lupinus angustifolius* contains *lupinin* $C_{12}H_{24}N_2O_4$, an intensely powerful alkaline substance. The substance *arginin* $C_6H_{14}N_4O_2$, found in the etiolated cotyledons of the lupine and the Soy bean, is a proteid. Prof. Chesnut says in regard to the *Lupinus leucophyllus* Dougl.

The above species is very abundant in Montana, where it is said to have caused the death of a very large number of sheep. There is some question whether the animals were killed by a poisonous constituent of the plant or merely by bloat. The seeds of all the lupines are probably deleterious in the raw state. In Europe, however, the seeds of *Lupinus albus*, after the bitter taste has been removed by steeping and boiling, are eaten by human beings as well as by cattle.

The so-called *nitrogen* obtained by European chemists from some of the lupines can be extracted by weakly alkaline water and is to be regarded as an active poisonous principle. Some European investigators, however, think that the alkaloids are not the cause of the poison. To the above poisonous species we may add *L. linifolius*, and *L. hirsutus*. The disease caused by these has long been known in Europe and has received the name of lupinosis. It is common where lupines are used for forage purposes. According to Freyberger and Fr. Jurek from one-half to three-fourths of the animals perish. According to Arne and Schneidermuhl the disease can be produced experimentally with *cupinotoxin* in sheep, horses, goats, and pigs. This substance occurs chiefly in seeds and pods. Dry heat does not destroy it but steam under pressure does. There is a probability that the poison is produced by metabolism.

Chesnut and Wuxox, in their paper on Stock-poisoning Plants of Montana, make the following statement in regard to the Lupine poisoning of Montana:

So far as we have been able to observe, lupines are not very extensively eaten by sheep during the spring and summer. This statement is at least true for normal conditions, where sheep are acquainted with the range and are not being trailed or driven. Horses and cattle take kindly to lupines and eat them in large quantities during their immature stages. When sheep are being trailed through strange country, or when they have just been unloaded from cars, and are in a hungry condition, they eat lupines ravenously in any stage of growth. The lupines are not considered valuable as forage plants for sheep until after early fall frosts, or until other forage plants have become dry and uninviting.



fodder. In late fall, and especially after early snowstorms, the lupines constitute one of the chief forage plants on some of the mountain ranges. It should be remembered that the leaves of lupines remain green and the plants offer slightly succulent forage after other plants have become dry.

The first case of poisoning from lupines which was brought to our attention occurred in August, 1896. A band of sheep, while being moved from one range to another was driven rapidly, and was constantly in a very hungry condition, when it was allowed to feed in a field of lupine for a short time. Within two hours after beginning to eat the lupine a number of sheep manifested violent symptoms of poisoning, and a few died within one hour after the appearance of the first signs of poisoning. Of the 200 sheep in the band 140 had died before the following morning. The season of 1896 was rather late and at the time when the poisoning occurred the lupine pods were fully formed, but the seeds were not quite ripe. In this case the sheep were driven away from the lupine as soon as the first symptoms of poisoning had been noticed and some of the sheep had eaten only small quantities of the plant. About 150 out of the 200 were affected, and as only 50 of these ultimately recovered it will be seen that the death rate was very high.

The owner of these sheep during the same season cut a quantity of lupine hay during the second half of July. In the winter of 1897 a band of 110 bucks belonging to the same sheep raiser were kept in a covered corral and were fed on cultivated hay. On one afternoon during the winter these bucks were given a liberal quantity of the lupine hay. About three hours after feeding this hay a noisy disturbance was noticed among the sheep. Upon investigation the owner found the sheep in a frenzied condition, and during the night about 90 of them died. No more lupine hay was fed and no more trouble was experienced.

They state further that the lupine poisoning occurred in various parts of the state. In 1898 about 2,000 having been poisoned. 1,150 sheep died out of a single band of 2,500 sheep. They also state that one sheep raiser in Deerlodge Valley lost 700 sheep from the poisoning of lupine. They report another case which occurred on June 28, 1900, near Livingston, in two bands of sheep, each numbering 3,000, which were being trailed westward from Livingston. The sheep were liberally salted before being started on the trail, the first day they traveled about 5 miles, and camped on opposite sides of a small stream. After watering, one band was driven across the creek and camped on a bench about 30 feet higher than the stream. On the following morning, the sheep which had been driven across the stream manifested symptoms of poisoning. Ultimately 1,900 died. This poisoning is referred to as lupinosis, a disease of which in Europe both acute and chronic forms are recognized, but in the United States the chronic form only has been recognized. The marked symptoms of poisoning are acute cerebral congestion, and great mental excitement. The sheep rush about in different directions, often running against the herder or other persons. The first stage of frenzy is followed by a second stage in which there is pronounced irregularity of movements and violent spasms, and falling fits. In many cases death occurs in from 1 to 1 1/2 hours. The pulse during the attack is strong and regular. Lower animals are attacked by convulsions, and these convulsions resemble those caused by strychnine poisoning. The excretion of the kidneys is increased, and sometimes it is bloody. The post mortem conditions are described by Chestnut and Wilcox as follows:

Post mortem examinations of the sheep poisoned by lupines revealed conditions very similar to those found in the acute cases of locu disease, already described above, with the exception that in loco disease the kidneys were not affected. The lungs were slightly congested, but this condition was not so pronounced as in cases of strychnine poisoning. The cerebral membranes were in all cases congested. In the more violent cases small blood vessels had been ruptured in various parts of the body, which may have been due either to increase of blood pressure or to the struggles of the animal.

In regard to the treatment, Dr. Wilcox recommends as follows:

No remedies have been used in cases of mock poisoning from American species of lupine. From our general experience with potassium permanganate it seems reasonable

to suppose that this substance would probably destroy the lupine alkaloids in the stomach if administered promptly after the first signs of poisoning. In the main, however, reliance should be placed upon prevention. With regard to the use of lupine hay, our experience and observations indicate that this is always dangerous for sheep if cut at a time when the seeds are retained in the hay. Since the limit of the period during which lupines are not poisonous can not be determined for the present with any certainty it seems advisable to abandon entirely the use of lupine hay for sheep, except after a preliminary test in feeding large quantities of the hay to one or two sheep. If it should prove to be non-poisonous, it may then, of course, be fed with safety.

The poisonous principle in all plants which have been fully investigated varies in quantity according to the stage of growth of the plant, and is located more abundantly in one part of the plant than in another. These facts seem to be strikingly true of lupine, since, as already indicated, the plants are sometimes eaten in large quantities with impunity, while at other times the plants cause extensive losses, especially among sheep. The evidence thus far collected regarding this matter indicates that the seeds are the most poisonous part of the plant.

Mr. O'Gara of Nebraska, in speaking of the Lupine says this:

There are three species of Lupines in the western part of the sand-bill region and throughout the foot-hills, which are worthy of attention. So far as can be learned, cattle and horses either do not eat them or are not harmed by them, but sheep men say that they are extremely poisonous to sheep when eaten after the pods have formed and have begun to ripen. Many sheep owners are very careful to avoid patches of Lupine in driving their sheep from one range to another, and never trust the flock to a green herder who is unacquainted with the range.

The three species common to the regions mentioned are the Nebraska Lupine (*Lupinus Platanus* S. Wats.), the silvery Lupine (*Lupinus argentus* Pursh.) and the Low Lupine (*Lupinus pusillus* Pursh.). The last named is a small hairy plant four to eight inches high, much branched near the root, bearing commonly five leaflets at the end of the leaf-stalk. The densely clustered blue flowers are borne on a stalk four to eight inches long. The pod is finely-hairy and is three-fourths to one inch long.

Dr. Nelson conducted some experiments in poisoning from three species of *Lupinus* *L. ornatus*, *L. sericeus*, *L. leucophyllus*, from which the following results were obtained with reference to the effect of feeding quantities of these plants. In regard to the first of these species, negative results were obtained in part, to sheep fed as early as May 30th, June 8th, July 14th, July 31st, and August 2nd, partly in the year 1898, and partly in 1901, these experiments being made in 1898 and 1901. In 1904 experiments were made with two sheep fed between November 16th, and December 22nd, receiving 274 pounds of this hay. This hay was eaten fairly well, and the sheep were given no other food except the lupine, and had constant access to water.

Some loss of flesh occurred in both sheep and one of them became affected, December 29th, with an attack of stomatitis with quite well developed ulcers in the mouth. He practically recovered by January 1st. No other untoward symptoms were manifested during the course of the experiment.

In regard to *Lupinus sericeus*, the results were in part negative, but June 28th, 1899, a sheep was fed 2 pounds of lupine that was in full bloom and partial fruit, gathered a few days previous. On the morning of the 29th, the sheep was drowsy, and kept a recumbent position. In the afternoon the comatose condition was more marked. He walked with an unsteady gait and pressed his head against the fence when he happened to reach it, showing a partial paralysis. The animal died on June 30th, slightly bloated.

The ventricles of the heart were partially and the auricles completely filled with a black coagulated blood. The lungs were congested, the stomach filled with partially digested food, otherwise apparently normal.

The experiments with *Lupinus leucophyllus* were negative.

8. *Cytisus* L.

Shrub with trifoliate or unifoliate leaves, showy flowers, chiefly in terminal racemes, calyx 2-lipped, with short teeth, standard ovate or orbicular, keels straight or curved, anthers large and small ovary with many-ovuled, and incurved style pod flat, oblong or linear. About 40 species, natives of Europe, Western Asia, and Northern Africa.

Cytisus scoparius (L.) Link. Broom

A stiff, nearly glabrous shrub, elongated, straight, angled branches, lower leaves 3-foliate, obovate, leaflets which are mucronate-tipped, upper leaves sessile, often unifoliate, flowers bright yellow, in elongated leafy racemes.

Distribution. Along the seacoast of Nova Scotia to Virginia and very common along the Pacific Coast.

Poisonous properties. The Scotch Broom (*Cytisus scoparius*) common on the Pacific and Atlantic coast but naturalized from Europe, contains the alkaloid *cytisin* $C_{11}H_{14}N_2O$ and is poisonous. Blyth records 400 cases of poisoning from this. The symptoms in stock are shivering, vomiting, staggering and general paralysis. *Cytisin*, occurring in many of the *Genisteae*, was found in 1818, in *Laburnum anagyroides* and since then, has been found in many other species of the genus *Cytisus* and in *Ulex europaeus*, *Sophora* sp. *Thermopsis* sp., *Baptisia tinctoria*, *Anagyris foetida*, *Lotus inaequalis*, *Colutea cruenta* and *Euchresta Horsfieldii*. Some species of the genera *Genista* and *Cytisus* do not contain *cytisin*. *Cytisus scoparius* also contains a volatile alkaloid *spartein* $C_{18}H_{25}N_9$, a single drop of which, according to Blyth, killed a rabbit that showed symptoms similar to those of nicotine poisoning.

9. *Melilotus* Tourn.

Annual or biennial herbs with trifoliate leaves, small white or yellow flowers in racemes with the odor of cumaria, teeth of the calyx short and nearly equal, shorter than the pod, corolla deciduous with obovate or oblong




Fig. 362. Flowers of Sweet Clover (*Melilotus alba*). 1, Standard above. 2, Showing wings and keel. 3, Showing stamens and pistil in keel.

standard, obtuse keel, free from the stamen tube, stamens diadelphous, pod wrinkled, straight, ovoid or globose—1-2-seeded. A small genus of 20 species, native to Europe, Africa and Asia. 3 species naturalized, found in North America, 2 of them quite weedy.

Melilotus alba Der Sweet Clover

An erect annual or biennial from 2-4 feet high, rather distant compound leaves, leaflets obovate, oblong, obtuse, serrate, narrowed at the base, truncate, emarginate or rounded at the apex, flowers with white petals, small fragrant pod ovoid, reticulated and smooth.

Distribution Abundant in waste places in the eastern and Atlantic states, also in the southern states and throughout the Mississippi valley, the Rocky Mountain region and the Pacific coast. Sweet clover is one of the most common weeds in pastures, and along roadsides.

Melilotus officinalis (L.) Lam. Wild Yellow Sweet Clover

An upright, yellow flowered herb from 1-4 feet high, leaflets oblong, or ova the apex more or less obtuse corolla yellow pod with irregularly reticulated veins

Distribution Common in waste places in the irrigated districts of the west, becoming more or less common in the Mississippi Valley and along the Atlantic Coast

Melilotus indica (L.) All Sweet Clover

An upright annual like the preceding, but with much smaller yellow flowers

Distribution Native to Europe, introduced in balast along the Atlantic coast and abundant on the Pacific coast

Poisonous properties The sweet clovers contain the substance *cumarin* $C_9H_8O_2$, which is found in the Tonka bean, sweet vernal grass, vanilla grass, etc. In Europe the sweet clover is suspected of being poisonous. This plant is used as a forage plant in the South, and Mr. Cohagen of Iowa, has had excellent results in feeding this plant to stock. Its protein content is equal to that of alfalfa. It is probable that some forms are entirely inert. Some years ago, the writer conducted an experiment in feeding considerable quantities of sweet clover, but without any injurious symptoms resulting. A tincture prepared by mixing the fresh flowers with alcohol has a vanilla like odor, and a bitter taste. Dr. Milspaugh states that in large doses, *cumarin* causes nausea, vomiting, vertigo, great depression of the heart's action, and cold extremities. Dr. Schaffner states that both of the sweet clovers are objectionable in wheat because of the foul odor the seed imparts to the flowers. According to Friedberger and Frohner sweet clover causes paralysis of the muscles. Dr. MacOwen states that in New South Wales, the *M. indica* is said to cause paralysis of horses.

10. *Medicago* L. Medick, Alfalfa

Herbs with pinnately 3-foliolate leaves, leaflets dentate toothed flowers small, yellow or violet in axillary racemes or heads, calyx teeth short nearly equal standard obovate or oblong, stamens diadelphous, ovary 1-ovuled, pod curved or spirally twisted, indehiscent 1-few seeded. About 50 species native to Europe and Asia. Bur clover (*Medicago lupulina*) and hop clover or black

medick (*M. lupulina*) are used as forage plants on the Pacific coast but eastward are regarded as troublesome weeds, the former injurious to wool.

Medicago sativa L. Alfalfa

An upright, smooth perennial, leaflets obovate, oblong, toothed, obtuse emarginate or mucronate, stipules entire; flowers in a short raceme violet, pod spirally twisted. A valuable forage plant.

Distribution. Common in the irrigated districts of the West, also frequent eastward, but common southward, spontaneous from New England to Minnesota, Kansas northward and westward, native to Europe and Asia.

Poisonous properties. A large amount of the green fodder is said to produce tympanites, but alfalfa is, however, one of the best of forage plants.




Fig. 303. Flowers of Red Clover 1 a. Leaf c. Standard 2 & 3. Wings, 4, keel 5, wings, 6, keel

11. *Trifolium* (Tourn.) L. Clover

Herbs, leaves mostly 3-foliate, palmately or pinnately, stipules united with the petioles, leaflets usually toothed, flowers in dense heads or spikes calyx persistent lobes 5, nearly equal, corolla withering or persistent, claws alternate to the stamen tube, stamens diadelphous or the tenth one separated for a part of its length, pods small and membranous, indurulent or dehiscent 1-6 seeded.

A large genus of about 250 species mostly in the northern hemisphere. Many are valuable forage plants, among these are red clover (*T. pratense*), alsike clover (*T. hybridum*) and white clover (*T. repens*). Several are weedy as yellow hop clover (*T. agrarium*), low hop clover (*T. procumbens*), and stone clover (*T. arvense*).

The alsike clover (*T. hybridum*) and red clover (*T. pratense*) occasionally produce bloat.

Dr. Jacob Moses and A. M. Harcourt have recently described a disease sometimes caused by alsike clover.¹

¹ Bull. Tenn. Agr. Exp. St. 18:28 (1905)



Fig. 304. Alfalfa (*Medicago sativa*) a, b, seed pod \times , seed. An excellent forage plant, sometimes causes bloat. (U. S. Dept. Agr.)

"The cause of this trouble among horses and mules is not positively understood. Whether the toxic effect is due to the plant itself which possibly undergoes some change within the digestive tract and subsequently liberates a poison, or whether it is due to the presence of a mould in connection with alsike clover is still undetermined. The mould has been strongly suspected. To determine this point will require further investigation. It is known, however, that the principle lesions are produced on the skin and mucous membranes.

"The symptoms of this disease vary to some extent, depending upon the location of the lesions and the length of time the animal remains on the alsike pasture, after the symptoms begin to develop. The cases which came under observation in Marshall County showed marked similarity of symptoms, involving principally the skin, the mucous membranes of the mouth, and the eyes.

"The prevailing symptoms of the disease are as follows

On the skin are inflamed areas, appearing at first as more or less rounded vesicular swellings, varying from one-half inch to five or six inches, or more, in diameter. The hair over the affected areas stands erect, and has a dull appearance, indicating loss of vitality. Later the skin becomes hard and puffed out, as the result of the formation of pus underneath. Finally the deadened skin is cast off, leaving a deep, raw, angry-looking ulcer, which eventually heals, with the formation of a conspicuous scar, covered with more or less white hair. These changes in the skin may occur on any part of the animal,

but especially on the limbs, body and croup. The eye symptoms consist of a marked conjunctivitis, with swelling of the eyelids, sensitiveness to light, and a watery discharge from one or both eyes. The mucous membranes of the mouth become inflamed (stomatitis), ulcers form, and the animal slobbers and refuses to eat. The advanced cases are frequently accompanied by emaciation. The tongue is usually affected, and the inflammation may extend throughout the entire digestive tract. The functions of the liver may be disturbed, and a yellowish (jaundice) coloration of the tissues follows. In such cases symptoms of colic are not uncommon, and the respirator, tract may become involved and pneumonia develop. Some observers in other countries have noticed marked nervous symptoms, such as excitement, convulsive movements, staggering gait, and paralysis of the throat, with inability to swallow, the paralysis at times becoming generalized, the animal getting down and being unable to rise. In the cases observed in this state, the nervous symptoms, except the general depression, were not very noticeable.

"The outcome of the disease depends upon the location and extent of the lesions upon the horse or mule affected. If they are situated on the exterior the animal will readily recover as soon as removed from the alkali pasture. If the vital organs are involved, such as the brain, lungs and liver, the disease may readily produce death. Among those cases occurring in this State, not a single fatality has been heard of at the Station. But even though the death rate is small where the ordinary precautions are taken, the disease



Fig. 305. Red Clover (*Trifolium pratense*). Occasionally the cause of blout.
(U. S. Dept. Agr.)

has considerable economic importance, since it leaves the animal more or less disfigured by the formation of scars, which materially depreciate his market value.

"The treatment is comparatively simple. As soon as the disease is recognized the animal should be removed from the alsike clover pasture and the wounds subjected to ordinary antiseptic treatment, such as frequent washing with 5 per cent solutions of carbolic acid or creolin, and the application to the ulcers on the skin of drying powders, consisting of boric and tannic acids in equal amounts."

The so-called clover sickness is supposed to be caused by the clover rust which has been described elsewhere. No doubt some of the trouble arising from feeding clover hay is caused by moulds found on the hay. Dr W. D. Gilchrist says that he has observed several cases of the kind in this state. The animals showed extreme restlessness followed by coma, bloody discharge from faeces followed by diarrhoea, weakness and debility. Change in fodder caused the trouble to cease.

I have recently received a similar complaint from Dr C. J. Scott of Knoxville, Iowa, three animals having succumbed.

Trifolium incarnatum L. Crimson Clover

A soft pubescent, slightly branched, annual, leaves long petioled; broad stipules; leaflets nearly sessile, obovate or obcordate cuneate at the base,



Fig. 306. Crimson Clover (*Trifolium incarnatum*). Sometimes produces phytohemagglutinin, which may cause death. (U. S. Dept. Agr.)

denticulate, flowers in elongated, oblong or ovoid, heads, sessile, calyx hairy, lobes plumose pointed, corolla crimson.

Distribution. Used as a cover crop and a forage plant in the south and east. Found on ballast from Maine to Pennsylvania. Native to Europe.

Injurious properties. According to Prof. Coville it produces phytobezoars and occasionally causes death in animals.

Trifolium repens L. White Clover

A smooth perennial with slender creeping and spreading stems, leaflets inversely heart-shaped or notched, obscurely toothed; stipules narrow peduncles very long, flowers in small loose heads reflexed when old, calyx shorter than the white corolla; pods 4-seeded.

Distribution. In fields and waste places throughout eastern North America, the Northwest and the Rocky Mountains.

Poisonous properties. Said to cause tympanites in cattle and slobbering in horses.

Psoralea L. *Psoralea*

Perennial herbs, usually sprinkled with glandular dots, leaves generally 3-5 foliolate, flowers spiked or racemed, white or mostly bluish-purple, calyx 5-cleft, persistent stamens diadelphous or occasionally monadelphous, pods about as long as the calyx 1-seeded.



Fig. 307. White Clover (*Trifolium repens*). Occasionally the cause of tympanites in cat. c. (Lamson-Scribner, U. S. Dept. Agr.)

About 100 species of wide distribution, many native to the plains, the roots of some being tuberous and farinaceous. The Indians used the tuberous roots of the *P. esculenta*, known as Pomme Blanche, or Pomme de Prairie, of the voyageurs for food. The roots of *P. hypogaea* and *P. cuspidata* were also used.

Psoralea argophylla Pursh. Silver-leaf Psoralea

Densely silvery, pubescent with appressed hairs, stem zig-zag, divergently branching, from 1-3 feet high; leaflets elliptical lanceolate, flowers spicate, interrupted, blue, pod oval, membranaceous, plant seldom seeding.

Distribution. From Wisconsin to Kansas and New Mexico to the Northwest territory

Poisonous properties. This plant was suspected of being the cause of a severe case of poisoning in Story County, Ia., two years ago. This is the first time the writer has known plants of this genus to cause poisoning, but he has had some correspondence with the parties concerned and thinks there can be no doubt that the poisoning was caused by the plant in question. A child was seriously poisoned by eating the seeds of this plant, but she finally recovered. It was thought this poisoning might have been caused by *Astragalus caryocarpus* but the plant sent me was the above.

13. *Tephrosia* Pers. Hoary Pea

Herbs or somewhat shrubby plants, odd-pinnate compound leaves, flowers in racemes or short clusters, red or white; stipules small; calyx 5-cleft; petals 5, standard roundish, usually silky outside, turned back, about as long as the coherent wings and keel. About 120 species, native of warm and tropical regions, a few are found in the United States.

Tephrosia virginiana Pers. Goat's Rue. Catgut

Perennial with villous or silky and whitish hairs, stem erect and ample, 1-2 feet high, leaflets 17-25, linear-oblong, terminal oblanceolate, narrowed to cuneate at the base, emarginate at the apex; flowers yellowish purple in long, dense racemes.

Distribution. In dry and sandy soil from Maine to Louisiana, west to Minnesota and Eastern Iowa, to Mexico.

Poisonous properties. This species, along with others, was formerly used to poison fish. The Mexican species, *T. toxicaria*, gets its name from its supposed toxic properties, and in South America, one species is commonly employed by the natives to poison the fishing streams.

The root is also poisonous to frogs and guinea pigs.

Tephrosia toxicaria contains the glucoside *tephrosin*. Several active substances have been obtained by Hanriot* from one species, *Tephrosia Vogellii*. Three substances were isolated, one, *tephrosol* $C_{10}H_{16}O$ is toxic especially to fish, a second toxic substance is *tephrosin* $C_{31}H_{48}O_{10}$.

14. *Robinia* L. Locust Tree

Trees or shrubs, stipules often prickly or spiny, leaves compound, odd-pinnate, the oblong leaflets with short stipules; flowers showy in axillary racemes.

* Compt. Rend. 1907, 454-451. Journal Chemical Society Abs. 92 386, 1907.




Fig. 308. Black Locust (*Robinia Pseudo-acacia*). 1. Flowering branch. 2. Flower. 3. Tube of stamens. 4. Longitudinal section of pistil. 5. Diagram of flower. 6. Legumes. 7. Pod open, showing seed. 8. Seed. 1, 2, 4, 7 $\frac{1}{2}$ natural size. (M. M. Cheney in Green's Forestry of Minnesota.)

mes, calyx short 5-toothed, and slightly 2-lobed, standard large, about as long as the wings and equal, stamens diadelphous, pod flat, several seeded and margined. A small genus of 6 species, native of North America and Mexico. Several species are cultivated for ornamental purposes, like the *Robinia viscosa*, which is native from Virginia to North Carolina and Georgia, and the *Robinia hispida*, native to the mountains of Virginia and Georgia. The *R. neo-mexicana*, with purple flowers, native to S Colorado and New Mexico, is frequently cultivated.

Robinia Pseudo-acacia L. Locust Tree, False Acacia or Black Locust

A large tree with rough bark, spiny stipules, 9-19 stalked leaflets obtuse, emarginate, or mucronate, flowers in loose drooping racemes, white, fragrant; pods smooth, standard yellowish at the base

Distribution. Widely planted as an ornamental tree. It produces valuable timber which is extensively used for posts. This species is, however badly infested with the borer.

Robinia viscosa Vent. Clammy Locust

A small tree with rough bark, stipules short, occasionally spiny twigs and petioles glandular, leaves 11-25, stalked leaflets obtuse and mucronate, nearly smooth, racemes dense, flowers in rather dense racemes, pinkish, not fragrant, pedicels glandular, hispid, pod hispid.

Distribution. Southwestern Virginia to Georgia, occasionally escaped from cultivation northward.

Poisonous properties. The bark and leaves of this species contain a powerful poison which has proved fatal to persons eating them. Children have been poisoned by eating the roots. It is true, however, that the flowers of the plant are often eaten with impunity and that bees collect from them large quantities of nectar. Dr. Rusby states that the occasional poisonous properties of honey are due to its origin in these flowers, though there are good theoretical reasons for doubting this. The bark of young twigs is sometimes pounded to a pulp, and used to make a tincture which is used in medicine as a tonic and cathartic, while the medicinal use of the flowers is mildly narcotic. It contains the substance robinin $C_{22}H_{40}O_{10}$ an aromatic glucoside which resembles the glucoside quercetin and is found chiefly in flowers, also the substance obigenin $C_{15}H_{16}O_6 + H_2O$. The seeds are also poisonous, and Dr. Milspaugh quotes Dr. Shaw as follows, in regard to the symptoms produced by poisoning from eating the seeds "Inability to hold the head upright, nausea and attempts to vomit, with a tendency to syncope, when in an upright position, voice, respiration and heart's action feeble, as from exhaustion, a painful paralytic condition of the extremities, which become shrunk on the fifth day. All the symptoms seemed like those produced by a long-continued diarrhoea, though in this case purging was not present." Dr. Johnson states that the symptoms of poisoning are those of Belladonna poisoning, a fact also noted by Dr. Waldron in the case of a horse that had eaten the bark. Friedberger and Frohner state that the animals have colic, tympanites and paralysis.

Dr. Rusby comments upon the poisonous character of the common black locust as follows:

Of this Dr. Johnson records that by eating the roots children are poisoned with symptoms like those of Belladonna poisoning, and that the bark and leaves are emetic. Prof. F. W. Power has experimented upon himself with the stem bark of this tree, proving the very actions effects which it produces and he has examined its composition with the result of showing that the poisonous constituent is an obnoxious substance, thus confirming the general character of that family, the Leguminosae. The most positive and prominent case recorded in regard to this article is that of Dr. Z. P. Emery. In the latter part of March, 1887, the two boys, inmates of the Brooklyn Orphan Asylum, were poisoned at one time by eating a bark which was being stripped in the vicinity for the making of fence posts. None of the cases terminated fatally. The prominent symptoms, stated in the order of their occurrence, were the vomiting of a rosy mucus, flushing of the face, dilated pupils, dryness of the throat, feeble pulse, extremities cool, face pale, vomiting of blood, cold extremities, heart feeble and intermittent, face deathly pale and ashy. The symptoms as I have named them are seen to be progressive. A rash similar to that of Belladonna poisoning was also present, but very feebly. In the beginning there was a high fever. Treatment consisted of sinapisms over the stomach, subcarbonate of bismuth, camphor and brandy.

A farmer in Dallas County, this state, informs me that sometimes the leaves are macerated in water and used to kill flies.

A case of poisoning to a horse was recorded in *Breeders' Gazette* in 1909. In this case the horse had eaten some of the bark of a tree. The symptoms were similar to those recorded by Dr. Waldron.

15. *Sesbania*. Scop. Seshan

Tall, smooth, branching herbs or shrubs with pinnate leaves and yellow flowers in axillary or compound racemes, calyx bell shaped, obliquely truncate, 5-toothed, standard short, orbicular, wings oblong, keel blunt, stamens diadelphous, style short, incurved at the apex, legume oblong, stalked, compressed, the endocarp membranaceous, at length separating from the coriaceous epicarp and enclosing 2 seeds. A small genus of 15 species of warm or temperate regions.

Sesbania platycarpa Pers.

A tall, smooth, branching annual vine, leaflets 10-35 pairs, mucronate, pale beneath, racemes shorter than the leaves; corolla yellowish purple spotted, with membranaceous saked pods.

Distribution. From the Carolinas to Florida, Missouri, and Texas.

Poisonous nature. In 1897, Dr. A. P. Anderson sent this to the writer with a letter from some stockmen from South Carolina, who stated that it was suspected of poisoning his cattle. Mr. Chesnut records a similar statement as follows:

In 1897, the United States Department of Agriculture received from South Carolina the seeds of this plant, which were found in the stomachs of cows.




Fig. 309. *Sesbania* (*Sesbania platycarpa*). This plant is common in the southern states, found along roadsides and woods, known to be poisonous. (Charlotte M. King.)



Fig. 370. Loco Weed (*Astragalus mollissimus*).
Charlotte M. King

16. *Astragalus* L. Milk Vetch

Chiefly perennial herbs with odd pinnate or occasionally simple leaves, with stipules; flowers spicate or racemose, calyx 5-toothed, tubular corolla clawed, usually long and narrow, standard erect, wings oblong, keel nearly as long as the wings; stamens diadelphous; pod several to many-seeded, dehiscent or indehiscent, 1-2 celled. A large genus of about 1000 species of wide distribution in the United States, largely western. Several species are used in medicine. One plant is widely known as the loco weed. One species native to gravelly knolls or dry places of the Mississippi Valley is known as the ground plum or hog pea (*A. carvocarpus*). This species is said to be edible. It should, however, be used with caution.

Astragalus mollissimus Torr. Loco Weed

A stout, short-stemmed perennial with membranous stipules. leaflets 19-27, ovate oblong, flowers in dense spikes, violet purple, rather large, pod oblong, glabrous somewhat compressed, sessile, furrowed at both sutures, at length curved.

Distribution. Found on plains of Nebraska, Kansas, Colorado to Texas and New Mexico.



Fig. 313. Ground Plum (*Astragalus caryocarpus*). It is not known whether this species is poisonous or not, although some of the related species are known to be poisonous and belong to the loco weeds. It is said that the pods of this species are edible. (Ada Hayden.)




Fig. 314a. Map showing distribution of Loco Weed or Purple Loco Weed (*Astragalus mollurimus*). After Marsh, U. S. Dept. Agr.

The term loco weed is applied to a great many plants. In addition to the species mentioned above *A. oocarpus*, *A. Crotolaria* and *A. lentiginosus* must be added, species of *Oxytropis* are considered in another connection. The *Sophora sericea* of this family, *Malvastrum coccineum* of the Mallow family and corydalis (*Corydalis aurea* var. *occidentale*) have at times been classed as loco weeds. Thus far none of these plants except those belonging to the pulse family have been found to cause loco poisoning.

Astragalus Bigelovii Gray Rattle-box

A subcaulescent, soft, silky, villous perennial; long, scape-like peduncles; flowers in dense spikes, violet pod oval-oblong, densely woolly, sulcate

Distribution. From S. W. Colorado to Texas and Mexico.

Astragalus Pattersoni A. Gray

Perennial, robust, 1-2 feet high, with appressed pubescence or sometimes smooth; leaflets oblong, rather thick, peduncles racemously many-flowered; corolla white, the keel sometimes purplish at the tip, pods smooth, sessile or stipitate; abruptly contracted within the calyx.

Distribution. Southwestern Colorado to Utah.

Astragalus Hornii A. Gray

A glabrous or minutely pubescent perennial, slender stems; leaflets about 21, narrowly oblong, peduncles longer than the leaves, flowers in a dense head, or short spike, numerous, small, calyx teeth subulate; pods ovate, from a broad base, straight, villous pubescent.

Distribution. From Southern Utah to California.

Poisonous properties. It is regarded as a very troublesome weed. Colorado has passed a law for its extermination, the state having paid out nearly \$200,000 in bounties between 1881 and 1885 to check its ravages. Much has been written on the subject of the poisonous properties of the woolly loco weed and other members of the genus. Brewer and Watson, in their Botany of California, state that the last species described is said to be poisonous to sheep, and Prof. Chesnut states that stock are affected by this loco weed in the southern part of California. It has certainly been regarded as a poisonous plant for a long time, and numerous investigations have been carried on in regard to its poisonous properties by Dr. Stalker, Prof. Sayre and others, Dr. Stalker coming to the conclusion that the loco poisoning might be brought about through the action of intestinal worms.

Prof. Sayre reported the death of a jack rabbit with symptoms similar to those recorded for horses and cattle. He was, however, unable to find a poisonous principle. Miss C. M. Watson reported a small amount of an alkaloid from the stemless loco weed, *Oxytropis Lamberti*. Dr. Mary Gage Day in Dr. Vaughn's laboratory found that when a half grown kitten was fed with milk containing a decoction of the root, stem and leaf, emaciation, convulsive excitement, and, finally death occurred, when an adult cat received 60-70 cc of a more concentrated solution death occurred on the thirteenth day and subcutaneous injection of a concentrated decoction in frogs and chickens caused death in 1-2 hours from heart paralysis. Dr. J. Ott experimented with *Astragalus mollissimus* and found that it decreased irritability of the motor nerves

affected the sensory ganglia of the nervous system, preventing them from readily receiving impressions and killed by arresting the heart action.

Dr. B. T. Galoway states that the loss from "loco" poisoning in Colorado alone has reached the sum of one million dollars per annum. It might be said in this connection that Prof. Power and Mr. Gambier investigated the subject, but were unable to locate definitely the kind of alkaloid. These gentlemen state

One kilogramme of the *Astragalus* herb gave 0.2 of a gramme of the alkaloid, equivalent to 0.036 per cent. Nothing further was determined concerning its nature, as it does not appear to be especially active. An extract from one kilogramme of the seed of the *Crotalaria* gave 11 grammes of an alkaloid, 0.036 per cent of the weight of the seed. It had a bitter taste, and seems to be more potent than that obtained from *Astragalus*.

From these investigations these authors conclude that both the *Astragalus* and the *Crotalaria* contain very small amounts of toxic alkaloids, to which the symptoms of poisoning may be reasonably attributed. Prof. Sayre, who has not gone into the details in the paper referred to, however reiterates what he has stated in several previous ones, that it is a question whether so small amount of a kaoid could produce such grave physiological disturbances.

The symptoms of poisoning are very well given in a paper by Prof. Cheanut.

Horses, cattle and sheep are affected by loco, but the principal damage is done to horses. The effect is not acute, but in its slow progress stimulates diseases caused by bacteria, worms, or other parasites of such as are caused in man by the continued use of alcohol, tobacco, or morphine. Two stages are recognized. The first, which may last several months, is a period of hallucination or mania accompanied by defective eyesight, during which the animal may perform all sorts of antics. After acquiring a taste for the plant it refuses every other kind of food, and the second stage is entered in. This is a lingering period of emaciation, characterized by sunken eyeballs, sunken hair and feeble movements. The animal dies as if from starvation in periods ranging from a few months to one or two years.

Dr. Carl Rued. isolated an acid (*loco acid*) from it to which he attributed the poisonous qualities of the plant.

Astragalus is said to cause considerable trouble in Nebraska. "In regard to the treatment," says Mr. O'Gara,

There is little to be said. All medicines that have been tried seem to have been of doubtful effect. More can be done by keeping animals away from loco than in any other way. As long as there is a plentiful supply of grass, there is little to be feared, but when pastures and ranges run low stock should be closely watched. At the very first appearance of trouble affected animals should be removed to some place where they cannot gain access to loco. Good nourishing food should be given. If the disease has obtained a strong hold on the animal there is little hope of recovery under the best of treatment hence the need of early treatment and prompt removal from the source of the trouble.

Dr. Mayo suggests the following treatment

	Ounces
"Sulphate of iron, pulverized	1
Gentian root pulverized	4
Ammonium chloride pulverized	1
Potassium nitrate, pulverized	1

Mix thoroughly and give from a heaping teaspoonful to a tablespoonful, according to the size of the animal, in the food three times daily."

Dr. Dwight C. Marsh and Albert C. Crawford under the direction of Dr. R. H. True who has charge of the Poisonous Plant Investigation of the U. S. Department of Agriculture have arrived at conclusions wholly at variance with previous investigation. They conclude that *Oxytropis Lambertii* poisons horses, sheep, and cattle and that *Astragalus mollissimus* poisons horses, but does not

poison cattle because they rarely eat it. Dr. Marsh states that the symptoms described by stockmen were corroborated, these are

The lowered head, rough coat, slow, staggering gait, movements showing lack of muscular coordination, sometimes more or less paralytic symptoms, a generally diseased nervous system, and in the later stages of the disease extreme emaciation.

The principal pathological changes are pronounced anemia of the whole system diseased stomach walls, and in acute cases a congested condition of the walls of the stomach, while in chronic cases there are frequently ulcers. Generally speaking, locoed cattle have ulcers in the fourth stomach. There is an excess of fluids in the various cavities of the body. This is especially noticeable in the epidural space of the spinal canal. Here the effusion is more or less organized, presenting the appearance of a gelatinous mass, which is especially abundant in the lumbar region and about the exits of the spinal nerves. In most locoed females the ovaries are found in a diseased condition.

Dr. Crawford from his laboratory work concludes

The symptoms described in stock on the range can be reproduced in rabbits by feeding extracts of certain loco plants. Those especially referred to here under the term "loco plants" are *Astragalus mollissimus* and *Astragalus Lambertii* (*Glycyrrhus Lambertii*).

The production of chronic symptoms in rabbits is a crucial test of the pharmacological activity of these plants.

It is the inorganic constituents, especially barium, which are responsible for this poisonous action at least in the plants collected at Hugo, Colo., but perhaps, in the future, loco plants from other portions of the country may be found to have other poisonous principles.

There is a close analogy between the clinical symptoms and pathological findings in barium poisoning and those resulting from feeding extracts of certain of these plants. Small doses of barium salts may be administered to rabbits without apparent effect, but suddenly acute symptoms set in analogous to those reported on the ranges.

The administration of sulphates, especially epsom salts, to form insoluble barium sulphate would be the chemical antidote which would logically be inferred from the laboratory work, but of necessity these sulphates would have to be frequently administered, and their value, after histological changes in the organs have occurred, remains to be settled. But the treatment of acute cases of barium poisoning in man is not always successful, even when sulphates combined with symptomatic treatment are employed. The conditions under which sulphates fail to precipitate barium must be considered.

Loco plants grown on certain soils are inactive pharmacologically and contain no barium. In drying certain loco plants the barium apparently is rendered insoluble, so that it is not extracted by water, but can be extracted by digestion with the digestive ferments. To be poisonous the barium must be in such a form that it can be absorbed by the gastro-intestinal tract.

There are plants with barium salts which are not injurious.

Dr. Marsh found that it is easy to kill the woolly loco weed (*A. mollissimus*) in fenced pastures because it occurs in small patches. The stemless loco weed is of wider distribution and, when in fenced pastures, can be killed but not so easily when it occurs on the ranges. He recommends treatment as follows:

In regard to the second phase of remedial work, it was found that locoed cattle can in most cases be cured by a course of treatment with strychnin, while locoed horses can generally be cured by a course of treatment with Fowler's solution. The animals under treatment must not be allowed to eat the loco weed and should be given not only nutritious food but so far as possible, food with laxative properties. To this end, magnesium sulphate was administered to correct the constipation which is almost universal among locoed animals. It should be noted, too, that magnesium sulphate may serve to some extent as an antidote to the poison.

Dr. C. Dwight Marsh¹ in a recent publication in speaking of this plant as well as the White Loco Weed and the experimental work of the Department of Agriculture, states that they are the weeds which produce the disease from Montana to northern New Mexico, Arizona and in western Texas, but there are many locoed animals where these two species do not grow. In California,

¹The Loco-Weed disease. Farmers' Bull., U. S. Dept. Agr. 380 1909.

Arizona and New Mexico there occur other leguminous plants which are known as loco weeds. Some of these are poisonous and the symptoms of loosed animals and the pathological findings are similar to those produced by the plants found in the eastern Rockies, especially in Colorado, where the purple loco is abundant. Dr Crawford finds barium in them which he thinks may be connected with the poisonous effects of the plants. In California the *A. diffractus*, *A. arizonicus*, *A. Thurberi* and *A. Bigelovii* are called "rattleweeds" and have been suspected. They are being studied by Dr Crawford of the Bureau of Plant Industry.

In his most recent paper on Loco Poisoning* Dr. Marsh affirms Dr Crawford's opinion that the purple loco weed (*Astragalus malinus*) is more poisonous than the white species (*Oxytropis Lamberti*). Dr Marsh adds this caution, that since such animals are extremely sensitive to strychnin, it has been found necessary to give it in small doses. He says

The daily doses should not ordinarily exceed three-twentieths or four-twentieths of a grain, or 0.009 to 0.012 of a gram. Large animals may take as much as one-half grain, but this is a maximum dose and often will be found too much. It was also found that sodium cacodylate when given to cattle in hypodermic injections of 5 grains, or 0.4 grams, daily commonly gave beneficial results. The best results, however were obtained from the use of strychnin and Fowler's solution as already outlined.

It may be added, in regard to the question of immunity, that loco poisoning comes on in a slow and cumulative manner so that there is no possibility of animals becoming immune.

It does not seem that the above is final in regard to all of the loco weeds. It is hardly likely that Profs. Power, Sayre, Gambier, and others have been entirely wrong in regard to their conclusions. When we find that related plants have strongly toxic properties we may expect to find the same properties also in some of these plants.

17 *Oxytropis* DC. Stemless Loco Weed

Perennial herbs or sometimes shrubby, generally acaulescent, with numerous tufts of short stems covered with scaly stipules. Flowers in racemes or spikes; calyx teeth nearly equal, petals clawed, standard, erect, keel erect, stamens appendaged, diadelphous pod more or less 2-celled or 1-celled sessile or stalked. About 125 species of the North Temperate regions. Most of our species are western. Several of our species are known to be poisonous to live stock. The following species of this genus are classed with the loco weeds: *O. Lamberti*, *O. deflexa* and *O. multiflora*. According to Greshoff the young leaf of *Oxytropis lapponica* has an extremely bitter taste, there is also an indication of saponin. On analysis the leaf was found to contain hydrocyanic acid. Greshoff also found hydrocyanic acid in the seeds of *O. sulphurea*.

Oxytropis Lamberti Pursh. Purple or Stemless Loco Weed

Nearly acaulescent perennial herbs or shrubby plants with tufts of very numerous short stems coming from a hard and thick rootstock containing many scaly stipules, stems and leaves are covered with silky and fine appressed hairs, or smooth, leaves pinnate, leaflets linear, flowers racemose or spicate, rather large and elongated, purple, violet, or sometimes white, stamens diadelphous, keel tipped with a sharp projecting point.

* Farmers Bull. U. S. Dept. Agr. 380:16.

Distribution. Western Minnesota, Western Iowa, and Missouri to Texas, and New Mexico, north to British Columbia, and northwest territory

Poisonous properties. The stemless loco weed is one of the most characteristic loco weeds of the West. The symptoms of poisoning are similar to those produced by the woolly loco weed described at length elsewhere. The poisonous substance has not been isolated. An alkaloid, however, has been reported by Prof. Prescott Chesnut and Wilcox, in speaking of the history of the loco poisoning in Montana, say that in Colorado the plant which is most commonly known as loco weed is *Astragalus mollissimus*. In Montana, on the other hand, the plants most generally called loco weeds by the stockmen are species of *Aragalus* (*Oxytropis*). "The species which is most concerned in causing the loco disease in Montana is the *Aragalus spicatus* and is closely related to *Oxytropis Lambertii*." Stockmen are of the opinion that a condition somewhat similar to loco poisoning may be brought about by eating undue quantities of alkali soil.



Fig. 313 Stemless or Purple Loco Weed. a, Plant. b, Seed pod. c, Seed. U. S. Dept. Agr.

It should be stated also that the larvae of sheep bot flies, which are frequently found in the frontal sinuses of the head, can not possibly be considered the cause of the nervous symptoms characteristic of the loco disease, for the reason that these larvae are not found in greater abundance in locoed than in healthy sheep. For the same reason the presence of the common tape worm (*Taenia serrata*) in the small intestines and bile duct of sheep can not be considered as the cause of the locoed condition. These worms are almost

universally present in the intestines of sheep, and under ordinary conditions do not cause any recognizable disturbances. The disease of sheep known as "gid" is not to be mistaken for the loco disease and, furthermore, is not prevalent in this country. No indications were found during the post-mortem examinations that the walls of the stomach were affected to any appreciable extent by the action of loco weeds, although these plants were invariably found in the stomach contents of such sheep. In the majority of cases no apparent changes have been produced in the spleen, liver, or kidneys. In some instances a slight congestion of the intestines was noticed. The cerebral membranes were in all cases somewhat congested. This condition is probably one of the immediate physical causes of the mental excitement exhibited by locoed animals. Post-mortem examinations of locoed horses disclosed the same conditions as those found in the sheep.

They made a number of experiments with a young Belgian hare and other rabbits, using the water extract of the leaves, and it was shown that this was not an acute poison if from 10 to 15 cubic centimeters of the liquid were administered. An acute case of loco disease was observed in an old ewe with a lamb at her side. She had eaten considerable quantities of the white loco weed (*Aragalus spicatus*-*Oxytropis spicata*). A slight locomotor ataxia was manifested. The eyelids twitched rapidly and there was a slight champing of the jaws. Each attack lasted from 1 to 2 minutes, and the intervals between the attacks were about 5 minutes.

The lips and eyelids twitched violently and the jaws were moved upon one another with such force that the sound could be heard for a distance of 200 yards.

Similar symptoms were observed in the lamb, which died in the afternoon.

Locoed sheep are exceedingly difficult to herd.

It is the universal experience of sheep raisers that locoed sheep are exceedingly difficult to herd. The sheep may, without a moment's warning, stray away from the band, each one in a different direction, and it is easy to understand how nearly impossible it is to prevent such a band of sheep from becoming separated. Besides giving the herder much trouble in directing the course of the band on the range locoed sheep also refuse to enter the corral at night, and under any and all circumstances may suddenly manifest perplexing stubbornness.

These writers did not observe many locoed cattle, but the symptoms are essentially the same as in sheep and horses. In regard to post-mortem conditions, they say:

Numerous autopsies made on locoed sheep and horses revealed conditions which, though fairly uniform, did not constitute a well-defined series. We made a large number of post-mortem examinations upon bodies of locoed sheep which had been killed and bled immediately before examination. In these cases there was no lesion or marked change in the alimentary tract. A slight congestion of the membranes of the brain was to be observed in all cases. The lungs and heart were apparently not affected. The voluntary muscles were of a paler color than under normal conditions, and the fat tissue was considerably reduced in quantity.

As to remedies, the following suggestions have been made:

Locoed sheep should be removed from the band and fattened for market on alfalfa or other forage plants, as above explained.

The immediate isolation of locoed sheep is advisable in order to prevent the habit from spreading in the band.

It seems desirable to give sheep a regular and abundant supply of salt in order to prevent the development of any perversion of the appetite.

Locoed horses are used to the best advantage as draft animals, but they must be maintained in good condition and prevented from eating loco weeds.

Dr. Marsh recommends to cut the roots below the crown of buds. A man with a spade can destroy a large number of plants in a day. The seeds however retain their vitality for some years, hence the field will have to be gone over again. That this method will effectively destroy the plants has been demonstrated by the U. S. Dept. of Agr. at Hugo, Colorado. The larvae of a moth (*Mourina amorphella*) feeds on the purple loco weed and this insect, Dr.




Fig. 313b. Map showing distribution of stemless loco weed (*Oxytropis Lambertii*). The plant reaches western Iowa on the Iowa bluffs where it is abundant. The disease however has never been reported so far as I know from this section of the state. Map after Marsh U. S. Dept. Agr.

Marsh thinks, will help to keep the weed in check if the insect will not lose its efficiency in the course of a few years.

18. *Vicia* (Tourn.) L. Vetch or Tare

Herbs, generally of trailing or climbing habit, with pinnate tendrils bearing leaves. Flowers generally racemose, calyx 5-cleft or 5-toothed, divisions nearly equal; corolla with the standard clawed and the wings adherent to the keel; stamens diadelphous or monadelphous, pod flat, 2-valved with several seeds, seeds globular, embryo with thick cotyledons. About 120 species, widely distributed. Some species used for forage, especially in Europe. The hairy vetch (*V. villosa*) has been widely distributed in the west because of the drought resisting qualities. Our most common native species is the American vetch (*V. americana*) which might well be introduced as a forage plant.

Vicia sativa L. Common Vetch

A smooth or slightly pubescent annual from 1 2½ feet high with simple stem. Leaflets 5-7 pairs, obovate-oblong to linear, notched or mucronate at the tip, the 1 or 2 nearly sessile flowers are borne in the axils of the leaves, flowers bluish purple, calyx teeth about as long as the tube, pod linear, several seeded, seeds black.

Distribution. From eastern Canada to Northwest Territory, New England to the Carolinas, west to Missouri and northward, generally in the wheat growing sections of the northern and western states. This is another weed commonly found in wheat screenings, abundant in the northwest.

Poisonous properties. In Europe it is the cause of tympanites. Dr. Schaffner, in *The Ohio Naturalist*, states that caution must be observed in feeding this plant to pigs. It is not injurious to cows. The seeds of this Vetch are often



Fig. 313b. Common Vetch. (*Vicia sativa*)



Fig. 314. Common Vetch (*Lathyrus silvestris*). The seed of this plant poisonous to stock. Charlotte M. King

found in screenings and fed in large quantities to cattle. As far as the writer knows, there are no cases of poisoning recorded from eating the screenings of this seed. The substance *vicin* $C_8H_{16}N_2O_4$ has been found in the seeds of this species. *Convician* $C_{10}H_{16}N_2O_4 + H_2O$ also occurs in this species and in *V. Faba*. Citric acid $C_6H_8O_7 + H_2O$ is found in *V. sativa*.

19. *Lathyrus* (Tourn.) L. Vetchling. Everlasting Pea

Mostly perennial, herbaceous vines although there are a few erect herbs, generally smooth, with pinnate, usually tendrill-bearing leaves, flowers in racemes or solitary, calyx oblique or gibbous at the base, upper teeth sometimes shorter than the lower, corolla larger than that of *L. sativa*, wings adhering to keel, style dilated and rather flat above, hairy along side next to free stamen, stamens 10 (9 and 1, or monadelphous below), ovules numerous, pod flat, sometimes terete, 2-valved, continuous between the seeds, dehiscent.

About 100 species are distributed throughout North America and a few others are found in South America and the mountains of tropical Africa. One species, *L. sylvestris*, is considered poisonous in its native home in the Carpathian Mountains. It contains certain alkaloids which by the process of cultivation have become eliminated so that in many localities at the present time it is considered a good forage plant and is relished by horses. In the western United States, the prairie vetchlings *L. arvensis* and *L. polymorphus*, and the marsh vetchling *L. pratensis* are considered valuable forage plants, the latter forming a very important part of the hay and soiling materially to its feeding value. *L. venosus* and *L. ochroleucus* occurring in similar localities are much less valuable. A form of intoxication, known as Lathyrism, is said to be caused by different species of *Lathyrus*.

In Dr Wilson's "American Text Book of Therapeutics," Victor C. Vaughan translates the following account of Lathyrism from Hubert's work "Intoxicationen."

By Lathyrism we mean an intoxication that was known to the contemporaries of Hippocrates, and which was caused by the seeds of at least three species of vetch, *Lathyrus pratensis*, the red vetch, *Lathyrus sativus*, the German vetch, and *Lathyrus Cymmarum*, the Spanish vetch. In Spain, France, Italy and in certain parts of Africa and India there have repeatedly appeared, from the eating of the seed of the vetch, epidemics of a disease that especially affects males and which induces a transverse myelitis with motor and sensory paraplegia. The paralytic symptoms gradually disappear but there remains spastic paresis with heightened tendon-reflexes attributed by Prevost to secondary degeneration of the internal columns, while Strömper considers the case a typical spinal paralysis. However the symptoms may wholly disappear and recovery be apparently complete. Men and animals, children & horses, are affected in the same manner. Duvernoy described the disease in 1765. Four saw it follow the eating of vetch-bread in 1785. Desmarches observed it in France in 1819, and Pettenkofer in the Mâconnais mountains in 1847. Reports of the disease were made by Irving in India, in 1861 and 1862, and by Bourge in Africa in 1882. In 1883, Marie published in *Le Progrès Médical* a review of the literature of the subject and more recently Schewarch has done the same.

Hogs are killed quickly by the vetch. Horses suffer from paralysis of the recurrent laryngeal nerve, necessitating tracheotomy. More chronic poisoning causes paralysis of the posterior extremities, and death. Méricourt believes the disease here-here is due to a similar intoxication but this is denied by Marie and others. In horses there is atrophy of the muscles of the larynx, especially of the cricothyroideus, pericard and cæcæ, also of the thyroarytenoidæus. The left recurrent laryngeal nerve is much wasted. Microscopic examination shows the muscle greatly atrophied, without striation, and undergoing fatty degeneration. In the central nervous system one finds atrophy of the ganglion-cells in the vagus center and of the multipolar ganglion-cells in the anterior horns of the cord.

Attempts to isolate the poison have not succeeded. Teilleux found an acid that induced typical effects upon rabbits. Bourlier found an active alkaloid in the alcohol-ether extract of the seeds, and poisoned birds with it. Astier isolated a volatile alkaloid by the Stas method, and he thus explains the fact that long-continued heating at a high temperature renders the seeds inert.

20. *Cicer* L. Chick pea

Calyx tube oblique or gibbous posteriorly, lobes nearly equal or the two upper somewhat shorter, conniving, standard ovate or nearly orbicular, narrowed into a broad claw; wings obliquely obovate, free, keel somewhat broader, incurved, dilated; anthers uniform; ovary sessile 2-8 ovuled, style filiform, incurved or bent, beardless, stigmas terminal, legume sessile, ovoid or oblong, turgid, 2-valved, seeds sub-globose or irregularly obovoid; funiculus scarcely dilated; hilum small, cotyledons thick, radicle short, slightly incurved or nearly straight.

Cicer arietinum L. Chick pea

Annual, herbs, or perennial often glandular-pubescent, leaves pinnate, petiole terminating in a small tuft of spinescent hairs or in an odd leaflet, scallots dentate or incised without stipels, stipules foliaceous oblique, often dentate or incised, flowers white, blue or violet, solitary pedunculate, or few pediceled; bracts small, bractlets 0. About 14 species, especially in the eastern Mediterranean and in Central Asia; extending westward.

Distribution. Cultivated in the Rocky Mountains and in the Southwest. Also extensively in Southern Europe and tropical Asia. Considered an excellent food plant.



Fig. 315. Chick pea (*Cicer arietinum*).
(After Faguet).

Poisonous properties. Friedberger and Frohner in Veterinary Pathology give the symptoms from Cicer poisoning.

In horses roaring and difficult breathing, owing to paralysis of the laryngeal muscles; paralysis, weakness in the loins, suffocation. Post-mortem reveals nothing of moment. Therapeutics: Change of fodder, tracheotomy.

21 *Phaseolus* L. Bean

Usually vines with pinnately 3-foliate leaves, stipules and racemose flowers; calyx 5-toothed or 5-cleft, the upper teeth more or less united, standard orbicular recurved, spreading keel spirally coiled enclosing the stamens and style, stamens diadelphous, 9 and 1, style bearded. pod linear 2-valved, several seeded seeds with large embryo. About 170 species mostly of tropical regions, 12 species native to southern states. The common bean *Phaseolus vulgaris*, native to tropical America is widely cultivated. The scarlet runner (*P. multiflorus*) is widely cultivated as an ornamental plant and is said to be poisonous. The *P. Mango* is cultivated in the tropics.

Phaseolus lunatus L. Lima Bean

A twining plant with racemose flowers shorter than the leaf, pod broad and curved, acuminat shaped seeds few, large and flat some dwarf, some with long twining stems.

Distribution. Widely cultivated, native to South America.

Poisonous properties. Several cases of poisoning from the use of lima beans are reported. L. Guignard, according to an abstract in the Experiment Station Record, has determined this to be due to hydrocyanic acid. He says "A number of forms that have been described as distinct species are by the author believed to be varieties or cultural forms of *P. lunatus*. Those principally studied were the white and colored Java beans, Burma or white Indian beans, Sieva beans, Cape beans, which are extensively cultivated in Madagascar, and Lima beans. These different varieties are widely cultivated and extensively used as food, although a number of fatalities have been attributed to their use. Descriptions of the different varieties and detailed reports of the chemical studies are given.

Practically all varieties of *P. lunatus*, whether wild or cultivated were found to contain the principle which when acted upon by an enzyme yields hydrocyanic acid. The proportion of hydrocyanic acid varied from almost unappreciable amounts in some of the more improved forms, like the Lima bean, to as much as 60 to 320 mg. per gm. dry weight in certain varieties of Java beans. It was found impossible by cooking to remove all the cyanogenetic compound in Java beans. Prolonged boiling extracts the greater part, but it is merely withdrawn and not destroyed, and if the water is absorbed it presents the same danger as the beans themselves. Since either in the alimentary tract or in the blood are sufficient ferments to act upon the dissolved glucoside, resulting in the liberation of hydrocyanic acid.

GERANIALES

Herbs, shrubs or trees, petals usually present and generally polypetalous; sepals mostly distinct, stamens few, rarely more than twice as many as the sepals, opposite them when as many, compound ovary superior. Contains the

families *Geraniaceae*, *Oxalidaceae*, *Tropaeolaceae*, *Linaceae*, *Erythroxylaceae*, *Zygophyllaceae*, *Rutaceae*, *Simarubaceae*, *Burseraceae*, *Meliaceae*, *Malpighiaceae*, *Polygalaceae* and *Euphorbiaceae*. The family *Tropaeolaceae* contains the nasturtiums (*Tropaeolum majus* and *T. minus*) frequent in cultivation, the fruits of the species being used for pickles. *T. Lobbianum* is a showy greenhouse plant. The *T. tuberosum* of Peru produces a tubercous root used for food in Bolivia cooking dispelling the unpleasant flavor.

Dr. Hatasted states that some persons have suffered from an inflammation on the hand caused by handling the garden nasturtium.

T. majus contains *glucotropaeolum* similar to the essential oil of mustard. The family *Burseraceae* contains *Commiphora abyssinica* furnishing myrrh, and Aburango (*Bursera simaruba*) the most characteristic tree of Porto Rico and one which furnishes a resin known in commerce as "chihon". The family *Meliaceae* includes mahogany (*Swietenia Mahagoni*) a valuable timber tree of the Antilles, myrtle (*Myrsine Ascardi*), widely cultivated in the South as an ornamental plant the fruit of which contains mangroera and is said to be poisonous, the West Indian cedar (*Cedrela odorata*) which furnishes a valuable wood used for furniture, cigar boxes, shingles, etc., and *Tectaria emetica* which yields an oil and is low. The carapa made from the seeds of *Carapa procera* is toxic for insects. The family *Zygophyllaceae* furnishes iguay-vitae (*Gnaphalium officinale*) a heavy wood used in machinery and casting work.

The ratbane broken back or mendis (*Charitella toxicaria*) of the family *Chenopodiaceae* is much used in Sierra Leone country of Africa for poisoning. It is placed in water to poison chickens and live stock. According to Dr. Renner "No one in this colony it would appear, dies from natural causes." Dr. Renner found the cause of this mysterious trouble to be due to poisoning from ratbane poison. In one case a laborer was poisoned from having eaten some fish on which the ground fruit of *Charitella toxicaria* had been strewn for the purpose of killing rats. This shrub and an allied species are common in Upper Guinea and Serenambila. Drs. Frederick B. Power and Frank Tutin who made chemical and physiological examination of the fruit of *Charitella toxicaria* found that the fruit contained neither an alkaloid nor a cyagenetic glycoside although a glucoside of this character is said to occur in South African *Cymosa*. The ratbane contains a resinous substance which is extremely poisonous, but a toxin could not be isolated. The syrup prepared from the resin when given to a dog caused tetanus and epileptiform convulsions soon followed by death. Drs. Powers and Tutin found that the fruit of this plant contains two active principles, one of which causes cerebral depression or narcosis and that the poison which causes convulsions is cumulative in its effect.

To the family *Erythroxylaceae* belongs coca (*Erythroxylon Coca*) which contains a number of alkaloids as follows: *cocain* $C_{17}H_{21}NO_4$, *cinnamoylcocain* $C_{21}H_{25}NO_4$, *truxillin* (a) $(C_{17}H_{21}NO_4)_2$, *truxillin* (b) $(C_{17}H_{21}NO_4)_2$, *benzoylcocain* $C_{19}H_{23}NO_4$, *tropacocain* $C_{17}H_{19}NO_2$, *hygrin* $C_8H_{13}NO$, *corypharin* $C_{17}H_{23}NO_4$. The injurious effects of cocain are well known. Dr. Winslow

Solutions of cocain (4-10 per cent) applied to mucous membranes, produce perfect local anæsthesia by paralyzing the sensory nerve endings. Cocain exerts a local anæsthetic action upon the gastric mucous membrane, and in this way lessens the appetite and sometimes stops vomiting. Intestinal peristalsis is increased by moderate doses, but is decreased and destroyed by the paralytic action of large doses. The action of cocain upon the heart and



Fig. 316. Coca-tree (*Erythroxylon coca*). Furnishes the coca of commerce. (After Faguet)

vessels is not very marked, except in poisoning. The alkaloid is, however, a slight cardiac stimulant in moderate doses, increasing the pulse-rate and tension. The action upon the heart is caused by depression of the cardio-inhibitory centres, and sometimes as well by depression of the cardiac inhibitory ganglia. Vascular tension is increased because of stimulation of the medullary vasomotor centres, smooth muscle of the walls, and because of the increased action of the heart. On the other hand, both minute and large doses may diminish the pulse rate.

Cocain is a respiratory stimulant in medicinal doses, but a paralyzant in toxic amounts. The respiratory centres are first stimulated and the breathing is made deeper and quicker. Depression and paralysis of the respiratory centres follow, cyanosis supervenes, and the respirations are shallow and irregular. Death occurs from asphyxia. In man an amount of cocaine exceeding gr. $\frac{1}{4}$ should not be employed under the skin, or upon mucous membranes, and death has occurred in susceptible patients from even small doses. The most powerful action follows the use of cocaine in very vascular parts, as about the face. One half a grain of cocaine given subcutaneously to a girl eleven years old, was followed by a fatal result in 40 seconds, and the writer has seen violent convulsions produced by the instillation of a few drops of a 2 per cent. solution into the eye of a man. On the other hand, spontaneous recovery has obtained in the human subject after the ingestion of 22 grs. of the alkaloid. In the horse, the toxic dose of cocaine causes restlessness and excitement, dilated pupils and salivation, culminating within an hour in a state of acute mania and intense excitement. These symptoms are followed by gradual recovery after a space of a few hours. Three grains of cocaine given under the skin, will sometimes induce nervous excitement in susceptible horses. The treatment of dangerous forms of cocaine poisoning, with respiratory and heart failure, consists in the use of rapidly acting stimulants,—as nitroglycerin upon the tongue, and strychnin, atropin and brandy subcutaneously.

Families of Geraniales

Flowers regular or nearly so, petals present usually as many as the sepals
flowers perfect; leaves not punctate

Herbs.

Capsule splitting into 5 carpels, leaves 3-foliate or dissected.

Geraniaceae.

Capsule 2-5 celled not splitting into carpels.

Stamens 2-3 times as many as the petals.

Leaf 3-foliate

Oxalidaceae.

Stamens as many as the petals

Linaceae.

Trees or shrubs with compound leaves, leaves often punctate.

Leaves punctate

Rutaceae.

Leaves not punctate

Simarubaceae

Flowers irregular, petals 3, stamens diadelphous or monadelphous Polygalaceae.

Flowers regular generally apetalous, monoecious, carpels mostly 3, generally herbs with milky juice Euphorbiaceae



Fig. 317. *Nasturtium* (*Tropaeolum majus*) Flowering branch. (After Faguet)

GERANIACEAE

Herbs with alternate or opposite leaves. Flowers perfect, regular, 5-nerved, hypogynous, stamens as many or twice as many or more than the petals, ovary 1, usually 5-lobed, ovules 1-2 in each cavity, fruit capsular. About 450 species of wide distribution. Native to the tropics and temperate regions of both hemispheres. Many plants of this order are frequently cultivated, among these are the South African pelargonium, commonly called the geranium, which contains *geraniol* $C_{15}H_{18}O$. The sharp points of the fruit of some are injurious.




Fig. 318. Cultivated Geranium (*Polemonium anisale*). 1, 4, Flower, 2, 3, d, Stamen, a, Stigma.

frequently entering the flesh and in some cases this mechanical injury has produced death in sheep, just as in the case of *Stipa*. The *Geranium Robertianum* has a disagreeable bitter taste.

Erodium L'Her. Storksbill

Herbs with opposite or alternate stipulate leaves, flowers nearly regular, axillary or umbelate, sepals 5, imbricated, petals 5, hypogynous, the upper slightly smaller, glands of the disk 5, alternate with the petals, stamens 10, anther bearing 5, and as many sterile filaments, ovary deeply 5-lobed and 5-celled, beaked by the united styles, 5 in number, lobes of the capsule 1-seeded, the style when mature breaks away elastically and is coiled spirally, tube of carpels hairy on the inside, seeds not reticulated. The 65 species found in temperate and warm regions. Some species have become widely distributed because the seeds cling to the fleece of animals. Some species are troublesome in western United States.

Erodium cicutarium L'Her. Allsilaria or Storksbill

A hairy, tufted annual with low spreading stems. plant viscid or sticky. leaves pinnate or once to twice pinnatifid, flowers in umbel-like clusters, purple or pink, fruit hairy on the inside and spirally twisted when ripe. The *E. moschatum* is a stouter plant which occurs occasionally eastward.

Distribution. This plant is common upon the Pacific Coast especially California, occurring in grain fields and waste places. It is also abundant in dry soils in the Salt Lake basin and from Colorado to Texas, occasionally found in the eastern states and Manitoba. Native to the Old World. The weed is commonly scattered by animals. It is injurious to wool.

Injurious properties. Species of *Erodium*, like those of *Stupa*, have in some cases sharp pointed calluses which bury themselves in the flesh and inflict injuries to animals. Our common species is but slightly troublesome in this way. The carpels of the *Erodium* get into the fleece of sheep and thus the wool is rendered somewhat less salable.

E. moschatum is injurious.



Fig. 319. Hemlock Stork's Bill (*Erodium cicutarium*). This widely distributed plant sometimes causes mechanical injuries to animals. (Charlotte M. King)



Fig. 320. Musk Erodium (*Erodium moschatum*). (After Peck)

OXALIDACEAE

Generally herbs, frequently with bulbs, acid juice, leaves palmate, with obcordate leaflets. flowers regular, 5-merous, stamens 10-15, ovary 5-celled, carpels with few or many ovules, loculicidal. A small order of 250 species chiefly tropical.

Oxalis L. Sorrel, Oxalis

Annual or perennial herbs with sour juice, often bulbous with alternate, digitately compound leaves of 3 leaflets, flowers in umbel-like clusters, solitary or several flowered, regular, often dimorphic or trimorphic, sepals 5, petals 5,

stamens 10, pistil 1; ovary 5-celled; ovules several in each cell; 5 separate styles; pod 5-celled, opening loculicidally, seeds 2 or more in each cell, the outer coat dehiscant, embryo large, endosperm present. The 250 species chiefly found in the tropics. The *Oxalis violacea* with violet corolla is a common plant in woods and prairies. The *O. corniculata*, a yellow flowered species, occurs from Pennsylvania to Illinois. The fresh juice of this is said to be an antidote against poisoning from the seeds of Jimson weed. Several South American species like *O. flava* and *O. Origines* are cultivated indoors. The *O. tetraphylla* and *O. lasandra* with their crimson flowers are also handsome for indoor cultivation. The *O. crenata* of Peru is cultivated for its tuberous roots.

Oxalis violacea L. Violet Wood Sorrel

Perennial with brownish bulb and ciliate scales, leaves smooth; leaflets obcordate, the midrib sometimes sparingly hairy; flowers in cymose clusters; sepals 5; petals 5, violet purple, capsule ovoid, seeds flattened, rugose-tuberculate.

Distribution. New England to Florida and New Mexico.

Poisonous properties. Dr Schaffner notes a case of poisoning as follows:

"A case is recorded of a boy being thrown into violent convulsions by eating a considerable quantity of the leaves."

Prof Hyarna states that children have been known to die from constantly eating the raw herbs of *O. grandis*.

LINACEAE. Flax Family

Herbs, rarely shrubs, stipules small or none; flowers regular and symmetrical, hypogynous; sepals 5, rarely 4, imbricated and persistent; petals 5, or rarely 4, convolute, stamens 5, monadelphous at the base, alternate with the petals; pistil 1 2-5-celled, styles 2-5, from capsular, seeds 1-2 in each cavity; cotyledons large, flat, without endosperm or with a small amount. A small order of 4 genera and 90 species, mostly in the genus *Linum*.

Linum (Tourn.) L. Flax

Herbs, sometimes with a woody base with tough fibrous bark, leaves sessile, stipules wanting or a pair of glands, flowers in cymes, racemes, or panicles; sepals 5, petals 5, soon falling, stamens 5 pistil 1, ovary 4-5-celled or becoming divided by false partitions, making 10 cells seeds shining with a mucilaginous coat large cotyledons. Several species commonly cultivated for ornamental purposes. The blue-flowered *L. perenne* of the Rocky Mountains, and the red garden flax (*L. grandiflorum*) a hardy annual from North Africa, are cultivated.

Linum usitatissimum L. Flax. Linseed

Annual, stem corymbosely branched at the tip, acuminate sepals; flowers broad, petals large, blue. Widely cultivated in the North and frequently spontaneous.

Poisonous properties and uses. The blue-flowered annual (*L. usitatissimum*) has been cultivated for centuries. The fiber has been found among the remains of the Swiss Lake dwellers. The ancient Egyptians as well as the Greeks and Romans also used the fiber for the manufacture of cloth. It is extensively cultivated in various European countries, fine fiber being produced in Belfast, Ire-

land, Brussels, Belgium, in Russia, and the Nile region. The seeds are also used extensively for making linseed oil. The chief regions where it is cultivated in North America are the Dakotas, Minnesota, Nebraska and Manitoba. Russia also cultivates the plant for the oil. The seed contains *linoleic acid* $C_{18}H_{32}O_2$, and is rich in oil. The compressed refuse is manufactured into oil cake, which is used for cattle food. The flax oil found in the seed of the plant is about one-third of its weight. Commercially, between 20 per cent and 30 per cent are obtained. When fresh it is without color and has little taste. The commercial oil is yellow and has a repulsive taste. On exposure to the air after having been heated with oxide of lead, it dries up to a transparent varnish consisting chiefly of *linoxyn* $C_{18}H_{14}O_{11}$. In medicine the flax seed is used in the form of a poultice, which is made of the pulverized seed. When oil cake or oil meal is fed in concentrated form it produced digestive trouble to hogs, frequently resulting in death. Dr. Schaffner states that it causes death to cattle, probably due to the prussic acid evolved from the plant when wilting. This substance has been reported.

Friedberger and Frohner state that it causes violent colic, inflation, diarrhoea, staggering, palpitation, death with convulsions, autopsy shows gastro-enteritis and signs of asphyxiation.

Linum rigidum Pursh. Large-flowered Yellow Flax

An herbaceous glaucous or slightly puberulent annual with rigid angled branches from 1-2 feet high. leaves narrow, erect, usually with stipular glands, flowers large, yellow; sepals acute or awn-pointed, glandular, serrulate, petals cuneate-obovate longer than the sepals, styles separate only at the summit, capsule 5-valved and avoid.

Distribution. Loess soil of western Iowa to Missouri, Texas, Mexico to Arizona and Manitoba.

Poisonous nature According to Chesnut the plant is reported as poisonous to sheep in the Pecos Valley, Texas.

RUTACEAE Rue Family

Trees, shrubs, or herbs with simple, compound, alternate or opposite leaves, glandular, with punctate dots without stipules, flowers mostly in cymose clusters, polygamio-dioecious hypogynous or perigynous, sepals 4-5, petals 4 or 5, stamens of the same number or twice as many distinct, inserted on the receptacle; pistils 2-5, of stamot or one compound 2-5 carpels raised in an annular disk, embryo large, curved or straight endosperm fleshy or none.

About 875 species, mostly in tropical regions of South Africa and Australia. Few representatives in North America. Two species of prickly ash (*Zanthoxylum americanum* Mill and *Z. Cuneo-Herckens* L.) and our hop-tree (*Ptelea trifoliata*) are common in the United States. The fruit of the hop-tree is used in Russia as a substitute for hops. A bitter alkaloidal principle occurs in *Xanthoxylum*. The gas plant (*Dictamnus albus*) a viscid glandular plant with strong aromatic scent is commonly cultivated. The common rue (*Ruta graveolens*), a native to Europe, is sometimes cultivated in country gardens. It has a strong disagreeable odor, and is so acrid that it will even blister the hands. It contains an acrid narcotic poison. The cork tree (*Phellodendron amurense*) from the Amur region, is occasionally cultivated. The most important genus of the

order is *Citrus*. The orange (*Citrus Aurantium*) is extensively cultivated in California and Florida. *C. Aurantium* var. *vulgaris* is the bitter orange which has run wild in Florida and other parts of the world. It is used in the manufacture of candied orange peel. The citron (*C. Medica*) produces the oil of citron, the thick peel being used to make the citron of commerce. The emon (*C. Limonum*), wild in northern India, introduced into Europe by the Crusaders, is now well known in cultivation in California. The lime (*C. Limetta*) is cultivated in all tropical countries, and with the lemon is used to make lime juice. It is a refreshing drink and on sea voyages is used as an antiscorbutic. The lemon and lime are forms of *C. Medica*. The mandarine or tangerine (*C. Aurantium*) having a small flattened fruit with a thin rind and rich fruit, is grown in California and China. It is harder than the orange, but probably a form of it. The shaddock or grape fruit (*C. decumana*) with large and somewhat bitter fruit, is native to Polynesia, and in recent years has become much better known in the United States. The kumquat (*C. japonica*), native to Japan and China, produces a small and pleasantly flavored fruit. The Aegle sepiaria (*C. trifoliata*) hardly as far north as Washington, is a spiny shrub producing a many seeded, yellow, austere fruit. Hybrids of the species and *C. Aurantium* with better and larger fruit, have been produced by Webber. The so-called orange or Naranja (*C. Bigaradia*) of Porto Rico and Florida is used for stock and plantings on moist lands because it resists the foot rot which affects other varieties. The Beal fruit (*Aegle Marmelos*), native of India, with fruit about the size of an orange, produces a delicious fragrant material used in medicine. Jaborandi (*Pilocarpus pennatifolius*) native to Brazil contains the alkaloid *pilocarpin* $C_{11}H_{11}N_2O_2$ and is a powerful diaphoretic. The administration of more than 5 grs. of the alkaloid is dangerous to horses when given subcutaneously. Atropin is an antidote. The alkaloid *jaborin* $C_{27}H_{42}N_4O_4$ resembles *atropin*, also the alkaloid *pilocarpidin*. The bark of angustura (*Cusparia febrifuga*) native to Venezuela contains *cusparin* $C_{70}H_{10}N_2O_8$ three other alkaloids and the bitter principle *angustarin*. The *C. toxicaria* of Brazil is poison



Fig. 321. Orange (*Citrus Aurantium*). 1. Flowering branch. 2. Longitudinal section of flower. 3. Longitudinal section of fruit. 4. Seed. (After Wendell)

ous. *Limonia amara* contains a toxic glucoside. Citric acid is found in fruits of lemons, lime and other members of the genus *Citrus*. The glucoside *hesperidin* $C_{60}H_{60}O_{27} \cdot 4H_2O$ occurs in ripe and unripe fruits of *Citrus*, the resinous principle *naringin* $C_{28}H_{34}O_{12}$ in *C. decumana*. The essential oil of lemons is one of the terpenes $C_{10}H_{16}$, the oil of bergamot similar to the preceding is from Bergamot *limetta* $C_{10}H_{14}O_2$ is the bitter principle of *Citrus Medica* = *C. Limetta*. The leaves of Buchu (*Barosma crenulata*) act as a mild diuretic. It contains a volatile oil of which 30 per cent is *disophenol*, also a crystalline glucoside (*asosmin*). The *Commiphora abyssinica* contains a volatile oil consisting of *cuminal* and *eugenol*. *Aurantioic acid* occurs in several species of *Citrus*.

SIMARUBACEAE. Ailanthus Family

Trees or shrubs with bitter bark. leaves pinnate, alternate, without punctate dots, stipules minute or none, flowers in axillary panicles or racemose clusters, regular dioecious or polygamous. calyx 3-5 lobed, petals 3-5, stamens of the same number as the petals or twice as many, pistils 2-5 and 1-5 celled disk elongated or annular.

A small family of 125 species of warm or tropical regions. The most widely known member of the family in the United States is the tree-of-Heaven or Chinese sumac (*Ailanthus glandulosa*). The quassia (*Q. amara*, of Guiana) is used in fevers and as a substitute for hops to impart bitter flavor to beer. It contains *quassin* $C_{32}H_{48}O_{10}$ a bitter principle. The bark of other plants of the order is bitter like the *simaruba* bark. The cedron (*Simaba Cedron*) of Central America is used in the tropics for snake bites. The bitter fruit of *Simaba valdivia* contain a glucoside $C_{18}H_{26}O_{10}$.

Ailanthus Desf.

Large trees, leaves compound, odd pinnate, flowers in panicles, greenish white; calyx short, 5-cleft. 5 spreading petals, disk 10-lobed, 10 stamens of the staminate flowers inserted at the base of the disk, ovary of the pistillate flowers deeply 2-5 cleft, 1-celled, stamens 2-3, winged fruits 2-5. Three species native of China and Eastern Asia.

Ailanthus glandulosa Desf. Tree-of-Heaven

A tall tree with ample leaves, smooth or slightly pubescent, 13-41 stalked leaflets ovate or ovate lanceolate flowers, greenish pedicelled, the staminate ones badly scented.

Distribution. Commonly escaped from cultivation, along roadsides from Southern Ontario to Kansas, Southeast Iowa, hardy as far north as Central Iowa.

Poisonous properties. The bark is known to be poisonous. Dr. White, in his *Dermatitis Venenata* states that he read an account in some medical journal of the suspected poisoning by this tree during its flowering season, and the statement was made that a case of marked dermatitis of the face, had been attributed to the emanations of a tree of this species, growing very near the sleeping chamber of the patient. He records a case where a lady was poisoned by contact with it. Dr. Halsted states that when the flowers are handled they produce an irritation of the skin.

In the Medical and Surgical Reporter of Philadelphia for 1872, this statement is quoted by Dr. Kusby in regard to the poisoning coming from the roots of this plant.

A case in which four persons were apparently poisoned by this root. They were members of one family and were successively, that is, at intervals of a few days, attacked, with no other possible cause than their drinking water which they took from the well of a neighbor. They all drank water exclusively, except the husband, who was the last to be taken. Others who drank of this water occasionally suffered similarly but to a slight extent. All immediately began to recover as soon as the drinking of this water was stopped. The symptoms, which had been slight for many weeks, appeared in a violent form in November at which time an *Albathus* tree growing in the vicinity of the well must have shed its leaves, and to a great extent its fruit also, if a pessilace tree which fact was not stated. On examination the soil at about the well was found to be thickly permeated with the roots of this tree, and these were also supposed to extend into the water, though an investigation regarding this was apparently not made. Inasmuch as the symptoms had existed in a mild form before the fall of the leaves, it is fair to assume that the roots had contributed toward the result while the violent outbreak in November would seem to indicate a sudden increase in the cause due to the accumulation of the leaves in the well. The symptoms were jaundice, a dingy aspect of the face and eyes, countenance fixed and anxious, pulse frequent and soft, yellowish fur on tongue except on the tip and edges, tenderness over the liver and right costal region, a permanent pain over the stomach with paroxysmal vomiting, pain in the back, difficult micturition and obstinate constipation. The symptoms were thus apparently to a great extent those of chronic gastritis.

Dr. Schaffner says that oaks will not eat grass near the young roots. *Quercus* occurs in the leaves. They also contain the bitter principle *quartin*.

POLYGALACEAE. Milkwort Family

Herbs or rarely shrubs, stipules none. Flowers perfect, sepals 5, petals 3 or 5, free, stamens 4-8, monadelphous, or chadelphous, anthers 1 celled, open-



Fig. 322. Tree of Heaven (*Ailanthus glandulosa*).
The bark is supposed to cause dermatitis. (Ada Hayden)

ing at the top by a pore, ovary 2-celled, ovules 2, fruit a 2-seeded pod. A small order of about 700 species, found chiefly in the tropical and temperate regions. Some species of the order produce a strong fiber

Polygala (Tourn.) L. Milkwort

Herbs or shrubs; simple entire, dotted leaves without stipules, flowers perfect, irregular, occasionally cleistogamous; calyx of 5 sepals, the 2 lateral known as wings, large, colored, the other small, greenish, petals 3, free, connected with each other and the stamen tube, stamens 6 or 8, filaments united below or in 2 sets, pistil 1, ovary 2-celled, ovules 1 in each cell. fruit mainly capsular, seeds with a caruncle, anatropous, embryo large little endosperm. About 250 species, of wide distribution, chiefly of warm regions. A genus of little economic importance

Polygala Senega L. Seneca Snakeroot

Plants clustered, several from a woody and knotty rootstock, simple 6-12 inches high. leaves lanceolate or oblong-lanceolate with rough margins, dense spike, long peduncle, flowers white, none cleistogamous; wings round-obovate; crest short; seeds hairy

Distribution. In rocky woods or clay soil. New Brunswick to Minnesota, Central Iowa to the Rockies in Canada.

Poisonous properties. The dried root is gathered when the leaves are dead,



Fig. 323. Seneca Snakeroot (*Polygala Senega*). Dried root is made into a powder which is very acrid. (From Johnson's Med. Bot. of N. A.)

and made into a powder and a tincture prepared. This tincture has a peculiar acidity. Dr. Millspaugh says

After tasting the tincture or chewing the rootlets, a very peculiar sensation of acidity and enlargement is felt at the root of the tongue, which, once recognized, will always mentally associate itself with this plant. The root contains polygalic acid $C_{22}H_{34}O_{16}$. It is a white, odorless, acid, amorphous powder.

This acid forms a frothing, saponaceous solution in boiling water, and breaks up into *sapogenin* and amorphous sugar, to which the name *senegin* has been given, which by some has been regarded as identical with *saponin*. According to the author quoted above, in doses of 10 minims of the tincture to a scruple of the powdered root, it produces

Anxieties, with dullness of the head and vertigo; aching and weakness of the eyes, with lachrymation, pressure in the ball, flickerings, dazzling vision, and contracted pupils, sneezing, pyalism, inflammation of the fauces and oesophagus, with thirst with anorexia, nausea, mucous vomiting, burning in the stomach, cutting cold roughness and irritation of the larynx, with regurgitation of blood to the chest accompanied by constriction, aching, soreness, and oppression, general debility, restless sleep and profuse diaphoresis.

Senegin resembles other *saponins*. Recent investigations indicate that the plant also contains *quillagic acid* $C_{18}H_{30}O_{14}$, *sapotoxin* and two *senega saponins*. The *saponin* of *Polygala virginiana* has the formula $C_{42}H_{72}O_{17}$. Other species of *Polygalaceae* like *P. venenosa* contain *saponin*.

EUPHORBACEAE. Spurge Family

Herbs, shrubs or trees usually with a milky acid juice, opposite alternate or verticillate leaves, monochous or dioecious flowers, much reduced, subtended by bracts resembling a calyx or corolla; ovary usually 3-celled; ovules 2 in each cell pendulous; stigmas as many or twice as many as the cells, styles generally 3, fruit a capsule, separating elastically into a 2-valved capsule, fleshy or oily endosperm, seeds with flat cotyledons.

A large family of 4000 species, chiefly tropical, many of which possess noxious qualities. Some species of the genus *Manihot* found in tropical Amer-



Fig. 324. Manchineel Tree (*Hippomane Mancinella*) Furnishes an arrow poison. (From Vauquelin's *Traité de Botanique*).

ica are poisonous. The fresh juice of bitter cassava administered to dogs and cats causes death in twenty minutes. The starch from this is used for sizing. Cassava (*Manihot utilisima*) is extensively cultivated in tropical America and to some extent also in Florida. The sweet cassava roots are used as food for cattle and man. Tapioca is the starch which settling from the water used to wash cassava meal, is afterward dried. An intoxicating drink is made from cassava bread. Rubber plants of the order are the *Hevea*, *Microdus* and *Manihot*. The manchineel tree (*Hippomane Mancinella*), the celebrated poison-tree of tropical America furnishes an arrow poison. The fruit, though tempting, contains an acrid poison, which causes blisters to form. The poisonous properties are said to rival those of the deadly Upas tree (*Anthonia toxicaria*).

The following is an extract from "West India Sketches"

The branches contain a milky juice which will certainly blister the skin, and it has been a common trick among the negroes to apply it to their hands in order to excite the compassion of those who might mistake it for the effects of beating.

Kingsley, in his charming "At Last," writes of it

We learnt to distinguish the poisonous manchineel, and were thankful in serious earnest that we had happily plucked none the night before, when we were snatching at every new leaf, for its milky juice by mere dropping on the skin burns like the poisoned tunic of Nessus, and will, even, when the head is injured by it, cause blindness and death.

Dr. White in his *Dermatitis Venenata*, speaks of the use of the plant in the West Indies as follows:

This large family of *Euphorbiaceae* contains some of the most poisonous plants. One of the most virulent is the manchineel, a small tree, bearing fruit resembling an apple, which grows in Southern Florida.

Loudon states that it abounds in a white milk which is highly poisonous, and so very caustic that a single drop placed upon the skin instantly causes the sensation of a hot iron, and in a short space of time raises a blister. It is a common belief that to sleep under it causes death. While woods on the seacoast of Martinique have been burned in order to clear the country of such a dangerous pest. The fruit is highly poisonous.

Mr. Combs in his paper on Cuban Medical plants, states that its poisonous effects may be overcome by the use of *Tecoma leucoxylon* or *Jatropha gossypifolia*. The uncooked rhizome of *Maranta arundinacea* is sometimes used for the same purpose. The latex of the sandbox tree *Hura crepitans* is also very poisonous and when applied to the skin, causes eruptive pustules resembling those of erysipelas. It also produces injuries to the eyes. It contains a sharp acrid poison. When taken internally it produces vomiting and diarrhoea. The seeds are used as emetic.

The Nigeria species of Mahogany (*Ricinodendron africanus*) also produces a valuable wood.

The tallow tree (*Saprum sebiferum*) is cultivated in tropical countries for wax found on the fruit, which is made into candles. The candlenut tree (*Aleurites triloba*) is cultivated on the islands of the Pacific Ocean for oil found in the seeds, which is made into candles, soap, etc. The seeds of *A. moluccana* are roasted and eaten. The Kalo Nut (*A. Fordii*) according to Holmes* is poisonous having produced toxic symptoms in five children. It is the source of tung or Chinese wood oil. The seeds of pinhoen oil (*Jatropha Carcas*) are eaten. They are nutty and have a pleasant flavor, but when eaten in excess, produce serious trouble and death often results. The drastic principle of Croton

* Pharm. Jour. 4:25, 231, 241. Brit. Yearbook of Pharm. 1906:240.



Fig. 325. Sandbar Tree (*Ilex coccinea*). 1 Flowering and fruiting branch; 2 Part of a large branch. Latex causes dermatitis. (From Vesque's *Traité de Botanique*.)

oil (*Croton Tiglium*) has not been definitely determined, but it has been refined to a dark brown oil known as *crotonol*. The plant contains *tiglic acid* $C_8H_{16}O_2$, *Crotonic acid* $C_8H_{14}O_2$ and *Croton oil* are derived from the *Croton Tiglium*, that is cultivated in southern India. This plant is a drastic purgative, capable when given in excessive doses, of causing death. The resin produces vesication. The seeds, according to Blyth, are very poisonous. The fixed oil has the formula $C_{18}H_{34}O_2$.

The bark of *Cascarilla* (*Croton Eluteria*), native to the Bahamas, is used as a tonic and contains *cascarillin* $C_8H_{16}O_2$. The milky juice of *Euphorbia resinsfera* of Morocco is used as a purgative and is so intensely acrid that people in collecting it are compelled to tie a cloth over their nostrils and mouths. It contains the substance *euphorbin* $C_{17}H_{34}O_2$, which has a burning taste. The milky juice of *agallocha* (*Excoecaria Agallocha*) of tropical Asia is very acrid and burns the skin. It is said that if the juice drops into the eye as sometimes happens to the woodcutter, blindness may be caused. *Excoecaria glandulosa* contains *excoecarin* $C_8H_{16}O_2$. The *Homalanthus Leschenaultianus* is said to be poisonous. The fruit of *Hypena poison* (*Toxicodendron capense*) of South Africa is very poisonous, and is used to destroy beasts of prey. Gum elastic or Para rubber is derived from the South American *Hevea brasiliensis*. Other plants of the family yield caoutchouc, which contains hydrocarbons that are readily soluble in chloroform. The alkaloid *drummin* occurs in *Euphorbia Drummondii*. Several species of *Euphorbia* like *Poinsettia* (*Euphorbia splendens*) and *E. heterophylla* are cultivated for ornamental purposes. From the



Fig. 326. *Croton* (*Euphorbia* *Tiglium*). Flowering and fruiting branch. The source of croton oil. (After Panquet.)

glands and hairs covering the fruit of kamala (*Mollotus philippinensis*) a dye is made. The fruit is also used as a vermifuge, it contains rothlerin $C_{22}H_{30}O_8$ and isorothlerin. Many species of the genus are regarded as poisonous. Maiden states that the *E. Drummondii* is poisonous to stock in New South Wales. It is known as the milk plant and is especially troublesome to sheep. It causes the head to swell to an enormous size so that the animal cannot support its head. Suppuration frequently follows. *E. alata* is also poisonous to sheep in the same country. *E. crenophylla* is another suspect in that country. *E. heptagona* is an arrow poison. Some species of this genus are used as fish poisons. Emanations of *E. characias* at one time were supposed to cause malarial fever which, however, was an erroneous assumption. Lehmann, a German writer on poisonous plants lists the following species as poisonous: *E. Lathyrus*, *E. Heisteria*, *E. prostrata*, *E. lula*, *E. Cyprinus*, *E. palustris*, *E. Pepus*, *E. exigua*. The *E. antiquorum* of the East Indies, *E. canariensis* of the Canary Islands, and *E. Reinhardtii* of the Transvaal contain a milky acid poisonous juice*. The resin from *Euphorbia* produces sneezing, irritation of face and skin, vomiting and diarrhoea and when used in large doses, death. Where the drug is manufactured, workmen must protect themselves, but, even then, headache, dizziness and weakness follow. To poisoning from members of the genus *Euphorbia*, Friedberger and Frohner ascribe such symptoms as constipation, severe and bloody diarrhoea, feeble pulse and tympanites.

* Bull. Misc. Inf. Roy. Gard. Kew, 1908, 154.



Fig. 326. Caper spurge (*Euphorbia Lathyris*). a, upper half of plant, one-third natural size; b, seed capsule, natural size. Listed by Lehmann as poisonous. (Chesnut, U. S. Dept. Agr.)

According to Greshoff the leaves of *Andrachne cordifolia* and other members of the family contain hydrocyanic acid.

Key for genera of Euphorbiaceae

- | | |
|--|-------------------------|
| Flowers without a calyx enclosed in a cup-shaped involucre | 5. <i>Euphorbia</i> . |
| Flowers with a calyx, involucre absent | |
| Flowers apetalous in panicles, stamens 10 | |
| Calyx corolla-like, plant with stinging hairs. | 4. <i>Jatropha</i> . |
| Flowers in terminal racemes or spikes covered with scurfy or stellate hairs, glandular | |
| Flowers spiked or glomerate, ovary usually 3-celled | 1. <i>Croton</i> . |
| Flowers in axillary spikes or paniculate, stamens 8 or more | |
| Fertile flowers in the axils of leafy bracts, stamens usually 8. | 7. <i>Acalypha</i> . |
| Flowers in interrupted axillary spikes, stamens 8-20 | 6. <i>Mercurialis</i> . |
| Flowers paniculate, stamens very numerous, filaments branched | 3. <i>Ricinus</i> . |
| Flowers apetalous in racemes or spikes, stamens 2 or 3 style simple | |
| Flowers racemose, hirsute or pubescent | 2. <i>Trapa</i> . |
| Flowers spicate, glabrous | 8. <i>Strlingia</i> . |

1. *Croton* L.

Stellate, pubescent herbs or shrubs, leaves generally alternate, occasionally with glands at the base of the blade; flowers apicate or racemose, the staminate above, calyx 4-6 parted, petals usually present, small or rudimentary, alternating with the glands, stamens 5 or more, pistillate flowers with calyx 5-10 parted, petals usually wanting, ovary mostly 3-celled, with a single ovule in each cell.

Croton capitatus Michx. Hogwort

An annual, dense, soft woolly herb, somewhat glandular from 1-2 feet high, occasionally branched, leaves entire, lanceolate oblong, with long petioles, sterile flowers with 5-parted calyx and as many glands alternating with the obovate lanceolate petals which are fimbriate, fertile flowers several, capitate or crowned calyx 7-12 parted, 5 petals wanting, styles twice or thrice forked; seeds gray, smooth.

Distribution. A common weedy plant from Missouri to Texas, from New Jersey to Georgia, Iowa and eastern Kansas.



Fig 327 Hogwort (*Croton capitatus*). Suspected of being poisonous. (After A. M. Ferguson, Rep. Mo. Bot. Garden)

Croton texensis Muell

A branching annual from 1-2 feet high, covered with close stellate pubescence; leaves narrowly oblong-lanceolate to linear, dioecious, calyx lobes 5, unequal, petals none, staminate spikes short, stamens 10 or more, style 2 or 3 times dichotomously 2-parted, capsule stellate, tomentose and roughened, seeds ovoid or oval.

Distribution. From South Dakota to Colorado, Texas, Mexico, Missouri and Alabama.

Poisonous properties. Several species of the genus *Croton* are used in medicine. The *Croton Tiglium* contains an oil which given internally is a powerful cathartic, but when applied externally, is a rubefacient. Loss of the hair follicles and of hair may occur. When gently rubbed into the skin it produces, after a short time, a considerable degree of itching, redness, and burning, and within a few hours small red papules may develop. If more of the oil is applied the papules are more abundant and are often surrounded with a bright red halo. They often become pustular and scars fill the pustules.



Fig. 338. Texas Croton (*Croton texensis*). Sometimes causes irritation of the skin. (After Mrs. M. H. D. Irish, Rep. Mo. Bot. Garden.)

Croton oil contains several fatty acids, such as stearic, palmitic, myristic and lauric acids. The volatile part of the acids contains an acid called *nigelic* $C_9H_8O_2$, which is the same as *angelic acid*. The drastic principle of Croton oil has not been definitely determined, according to Flückiger and Hanbury. *Crotonol* $C_{18}H_{34}O_4$ is a non-purgative body causing irritation of the skin. According to Winslow, in his *Veterinary Materia Medica and Therapeutics*, "10 drops of croton oil will kill a dog unless vomiting occurs. 30 drops prove fatal to a horse, intravenously. The treatment of poisoning includes the use of emetics or stomach tube, demulcents and opium." None of our native species is mentioned as poisonous by Dr. Schaffner or Prof. Chesnut, but a few years ago I had a query through the *Wallace Farmer* in Des Moines, from a correspondent in Western Nebraska who suspected that the Texas croton was poisonous. The writer has eaten a few seeds of our southern *Croton capitatus* with slight uneasiness. On the other hand, a few seeds of the Texas croton produced powerful irritation which lasted for an hour, and then disappeared. It is listed by Bessey and O'Garra as possibly poisonous in Western Nebraska. Prof. Chesnut states in his paper on Plants used by the Indians in Mendocino county, California, that the bruised leaves of *Croton setigera* are used to stupefy fish. The common name, fish soap-root, indicates its use.

The bark of the cascarrilla (*Croton Eluteria*), native of the Bahama Islands is used as a tonic.

Tragia L. *Tragia*

Monocious herbs or shrubs, usually armed with stinging hairs, leaves alternate; flowers in racemes with bractlets, apetalous, sterile flowers with a 3-5 cleft calyx, fertile flowers with a 3-8-parted calyx, divisions entire or pinnate-



Fig. 129. Spurge Nettle (*Tragia sp.*). This spurge is common in some places in the South and has hairs that are irritating like those of the common nettle. (Charlotte M. King.)

fid, styles 3, capsule 3-lobed, separating into three 2-valved carpels. A small genus of 50 species.

Urtica urens L. Common Nettle or Tragia

A dull green, pilose plant with pilose or hirsute hairs, erect, branched stems; leaves obovate, or ovate-linear; short, petioled, pistillate flowers, several at the base of the racemes, with a 5-6 lobed calyx, capsule short-pedicelled.

Distribution. From Virginia to Florida and Texas.

Urtica nebotaeifolia Cav. Tragia or Nettle

A somewhat hispid, erect, or slightly twining plant, bearing stinging hairs, leaves ovate, or triangular-lanceolate, base cordate or truncate; short petioled, racemes many-flowered, pistillate flowers with a 5-lobed calyx, seeds chestnut brown.

Distribution. From Kansas to New Mexico.

Poisonous properties. The hairs have the same stinging property as those of the common nettle.

3. *Ricinus* (Tourn.) L. Castor Oil Bean

A tall stout herb or tree in tropics, glabrous and glaucous, large, alternate, peltate leaves. flowers in large, paniced clusters, the fertile above the staminate below; calyx 5-parted, stamens numerous, styles 3, united at the base, each 2 parted, red; capsule subglobose, or oval, separating into 3. 2-valved carpels, cotyledons large; endosperm fleshy and oily. A single species naturalized in warm countries, probably native to Asia.

Ricinus communis L. Castor Oil Plant

A tall, smooth, branching herb with palmately-lobed leaves. seeds oblong, shining, variegated with white.

Distribution. Widely cultivated as an ornamental plant, and an escape from cultivation from New Jersey to Texas.

Poisonous properties. The seeds furnish the well known castor oil, which is a mild and safe purgative. It contains *ricinolein*, or *ricinoleic acid glycerid*, $C_{18}H_{34}(C_{18}H_{34}O_2)_3$, an acrid principle, also *palmitin*, *stearin*, and *myristin*. The purgative principle found in it is unknown. Castor oil is not poisonous, but the pulp contains an acrid, albuminous substance, *ricin* $C_{18}H_{34}N_2O_6$. Dr. W. now in speaking of the poisonous character, says, the seeds "contain 50 percent of oil, and an acrid, poisonous substance. Three seeds have caused death in man, and they are ten times more purgative than the oil." A few seeds eaten entire by a child might produce serious symptoms. According to Chesnut, the seed eaten accidentally by horses has caused death. They are used also to poison sheep, according to the same authority. The oil cake is said not to be poisonous to poultry and cattle. A case is known of a young lady whose eyes became inflamed when in contact with a mere trace of the material in the laboratory. The toxin is very poisonous, but animals may be rendered immune, and the seeds then fed to them. Behring has produced an anti-toxic serum against the *ricin* or toxin of the castor oil bean.



Fig. 230. Castor Oil Plant (*Ricinus communis*). Furnishes the well known castor oil of commerce. (After Faguet.)

The symptoms of poisoning are vomiting, gastric pain, bloody diarrhoea and dullness of vision.

Sillmark¹ states that the toxalbumin of castor oil bean, when injected into the circulation is more poisonous than strychnin, prussic acid, or arsenic.

Quite recently Dr. W. N. Bingham² reported on several cases of poisoning in Cuba from eating the seeds of the castor oil plant. Some persons showed peculiar susceptibility. In one case poisoning occurred from eating a single seed while in another a great many were eaten. In both cases the seeds caused nausea, vomiting and purging of a violent nature.

Toxic substances similar to ricin occur in the following plants, *Abies procumbens* (labrum), *Jatropha curcas* (curren), *Croton tiglium* (croton), *Robinia pseudo-acacia* (robin), *Brayera anthelmintica* (castorin). According to Ceni and Beata a toxin also occurs in *Liriodendron* seedlings, *Aspergillus flavus* and *A. fumigatus*.

¹ Dorpat. Arch. 2 (1895).

² Am. Jour. Med. Sci. 128: 319-321.

4. *Jatropha* L. Spurge Nettle or Bull Nettle

Monoecious, or rarely dioecious, perennial herbs, with bristly hairs, entire or lobed leaves, flowers in cymes, calyx colored like petals in sterile flower, mostly salver-shaped, and 5-lobed, enclosing 10-30 stamens, pistillate flowers in the lower forks of the cymes, capsule ovoid or subglobose, separating into 2-valved carpels. A small genus of 4 or 5 species.

Jatropha stimulosa Michx

A branching perennial plant with a stout root, 6-12 inches high, and stinging hairs, leaves round, heart-shaped, 3-5 lobed or variously cleft, calyx of the staminate flower salver-form, white or pinkish; stamens 10, filaments almost separate, seeds oblong-ovoid, smooth and mottled.

Distribution. In dry sandy soil from Virginia to Texas.

Poisonous properties. Mr John Smith says that a plant growing at Kew was placed on his wrist, and produced in a few minutes, serious symptoms extending to the upper part of his body; the lips became swollen, and the whole of a livid red, fainting coming on in ten minutes. The writer was told of numerous instances of poisoning in Texas where it is much dreaded.

Jatropha urens, known as the Brazilian stinging net, is considered to be one of the most poisonous plants known. The Cuban physic nut (*Jatropha Curcas*) is used as a purgative.



Fig. 331 Spurge Nettle. Low Pammel.



Fig. 332. Spurge nettle (*Jatropha stimulosa*). The *Jatropha* has stinging hairs that produce injuries similar to those produced by nettle but much more powerful. (After Hochstein)

5. *Euphorbia* L. Spurge

Monoecious shrubs or herbs with alternate or opposite, verticillate leaves. flowers involucrate, involucre resembling a calyx or corolla, bearing a large thick gland in the sinuses, staminate flowers consist of a single jointed stamen on a filament-like pedicel, pistillate flower solitary at the bottom of the involucre consisting of a 3-lobed and 3-celled ovary, capsule at maturity breaking into 3-lobed 1-seeded carpels, seeds frequently caruncled, smooth, variously pitted. About 700 species, chiefly in warmer regions. A few are weedy, some poisonous and some planted for ornamental purposes. The milky juice of the Brazilian *E. heterodoxa* produces a ferment which acts much like papain.

Euphorbia Prentis Guss. Large Spotted Spurge

An ascending, erect annual from 1-2 feet high, opposite oblique leaves, which are ovate, oblong or oblong-linear, falcate, serrate, usually with a red spot or red margins; stipules triangular, flowers collected in a loose terminal cyme; appendages entire, white or red, pod smooth, angled; seeds small, blackish, ovate, obtusely angled, wrinkled, and tubercled.

Distribution. Common in eastern North America west to the Rocky Mountains.

Euphorbia maculata L. Spotted Spurge

A prostrate spreading, hairy annual, leaves oblong-linear, pubescent or smooth, oblique at base, serrate above, small brownish spots on leaves stipules lanceolate, fimbriate; flowers monoecious, included in a 4-5-lobed involucre, glands of the involucre minute, peduncles as long as the petioles, in dense clusters, pods minutely pubescent, seeds sharply 4-angled, having 4 shallow grooves, whitish.




Fig. 333. Spotted Spurge (*Euphorbia maculata*) Common roadside plant. Probably poisonous. (C. M. King)

Distribution. Common along roadsides, walks, etc., from New England to the Rocky Mountains and the Gulf States.

Euphorbia marginata Pursh. Snow on the Mountain

An erect, stout annual from 2-3 feet high, stem hairy or somewhat smooth, leaves sessile, scattered, ovate or oblong, entire, deciduous stipules, uppermost leaves opposite or whorled with conspicuous white petal-like margins, involucre bell-shaped in umbels, glands of the 5-lobed involucre with broad and white appendages, seeds ovoid, globose, terete, dark ash colored, reticulate.

Distribution. Frequently cultivated in gardens from whence it has escaped. Found in Ohio, Illinois and Indiana. Native from western Minnesota, Iowa to Colorado, and Texas.

Euphorbia corollata L. Milkweed or Flowering Spurge

Perennial with a long, stout rootstock, glabrous or sparingly hairy, leaves ovate, lanceolate, or linear obtuse, short-petioled, or sessile, inflorescence in umbel-like clusters, involucre long peduncled with white conspicuous appendages, seeds thick, ovoid, slightly pitted, ash-colored.

Distribution. In rocky or sandy soil, Mass to New York, New Jersey, Florida, Wisconsin, Minnesota, and Texas.




Fig. 234. Common Flowering Spurge (*Euphorbia corollata*). A plant with milky juice which has irritating properties common in sandy fields. It has white bracts resembling flowers. (Charlotte M. King.)

Euphorbia Lathyris L. Myrtle Spurge

A glabrous annual or biennial, simple below, branched above, from 2-3 feet high, leaves thick, linear or oblong, scattered, the upper lanceolate



Fig. 333. Snow on the mountain (*Euphorbia marginata*) a, whole plant, one-third natural size b, seed capsule, natural size. Cultivated in old garden. (Chesnut, U. S. Dept. Agr.)

or linear-lanceolate, inflorescence umbel-like, bearing 4 crescent-shaped glands, prolonged into horns, seeds oblong-ovoid, terete, usually wrinkled.

Distribution In waste places. New Jersey to North Carolina, Iowa, and California. Native to Europe.

Euphorbia Ipecacuanha L. Wild Ipecac

Perennial from 5-10 inches high, long perpendicular root, entire, smooth leaves, varying from obovate or oblong to narrowly linear, nearly sessile, involucre long peduncled and 5 transversely elliptical or oblong green glands, seeds smooth, ovate, white, pitted, and obscurely 4-sided.

Distribution In sandy soil from Connecticut to Indiana and Florida



Fig. 336. Wild Ipecac (*Euphorbia Ipecacuanhoe*.) Plant that possesses irritating properties and is also a purgative. (Millspaugh and Charlotte M. King.)

Euphorbia heterophylla L. Cruel Plant

An erect, smooth annual from 1-3 feet high, leaves alternate, petioled, linear lanceolate to orbicular, undulate, entire or toothed, the upper leaves usually fiddle-shaped, with a red base, involucre in terminal clusters, 5-lobed, with a single or a few almost sessile glands, seeds nearly round, transversely wrinkled and tubercled.

Distribution From Illinois and Missouri to Nebraska.

Euphorbia Cyparissias L. Cypress Spurge

A bright green perennial from 6-12 inches high with running rootstocks; stems clustered, occurring in patches; stem leaves linear, entire, densely crowded, those of the flower heart-shaped and entire, flowers in umbellate clusters, umbel many rayed, glands crescent-shaped, pods granular, seeds oblong and smooth.

Distribution. Native to Europe, but widely scattered in eastern North America. First introduced as a cultivated plant in North America.

Poisonous properties. All of the species are more or less irritating and in drying give off very disagreeable odors. Many of the species of the genus are used by quacks to remove warts and freckles; the juice produces an erysipelatous-like inflammation, and in one case mentioned by Dr. White, the whole abdominal wall became gangrenous.

The milky juice of the plant causes itching and inflammation. The general effect is very much like that of poisoning from the poison ivy. In Texas, according to Chesnut, the juice of *E. marginata* is used to brand cattle. The honey



Fig. 337. Yellow Flowering or Cypress Spurge (*Euphorbia cyparissias*). A branch with large bracts and small flowers. (Stranburger, Koll, Schenck and Schlimper.)

obtained by bees from the plant is poisonous and is rendered unfit for use. The acrid properties of this species were described some years ago by Dr Schneck. The juice of *E. corollata*, according to Dr Haisted and many other observers, is acrid, and, on the authority of Dr Bigelow, formerly was used for blistering purposes. The bruised root will vesiculate the skin. According to Dr J. C. White, the dust of this species produces painful swelling and vesicles upon men who handle the plant. It is used as an emetic, and is troublesome to those who collect it.

The *Euphorbia pulchra* is used as a sedative in spasmodic conditions of the respiratory apparatus. It produces dermatitis. Dr White, in his *Dermatitis Venenata*, has this to say of the species of the genus:

More than one hundred species of *Euphorbia*, or spurge, grow in the United States, either indigenous or immigrants from Europe. Of every species London says the juice is so acrid as to corrode and ulcerate the body wherever applied, and of *E. resinifera*, from which the official *euphorbium* is obtained, Pliny and Dioscorides, according to the Dispensatory, describe the method of collecting juice, so as to prevent irritation of the hands and face. This substance is used as a plaster to prolong suppuration.

Van Hasselt states that the juice of several species is used by quacks to remove warts, freckles, as depilatory, etc., and that the application of the juice, powder, and extract produces not only erysipelatous, pustular, and phlegmonous inflammation, but even gangrene. In one case mentioned the whole abdominal wall became the seat of gangrene.

Of our native species, Bigelow says that the juice of several was used in his day to destroy warts, and Gray describes them all as containing an acrid, poisonous juice. The most active of them are *E. corollata*, *E. iperacuanhae*, and *E. Lathyris*. The first of these, commonly called snake-milk, according to Bigelow, has been used for blistering purposes, and the Dispensatory states that the bruised root will vesiculate the skin.

Mr. Cheney informs me that the juice of *E. iperacuanhae* is quite troublesome to many who collect and handle it; and Basin states that the dust of *E. Lathyris*, growing both in Europe and in this country, causes redness, painful swelling, and vesicles upon the workmen employed in handling it.

With reference to the poisonous nature of the juice of the several species, nothing very definite is known. *Euphorbia* $C_{29}H_{48}O_7$ has been found in *Euphorbia Iperacuanhae*. This euphorbon acts as an irritant to the mucous mem-



Fig. 118. Large Spotted Spurge (*Euphorbia Preslii*). Supposed to cause "slobbers" in horses. (Charlotte M. King)

branes throughout the alimentary tract. The caper spurge (*Euphorbia Lathyris*) is poisonous, and the following physiological actions are described by Dr. Millspaugh:

Bright, staring, wide-open eyes, dilated pupils; death-like pallor of the countenance; retching and vomiting, violent purgation, stools frequent, copious, and in some cases bloody, irregular pulse, whole body cold and rigid, followed by heat and perspiration. M. M. E. Sudour and A. Caraven-Cachin state that emesis always precedes purgation, and that the seeds have an irritating action upon the mucous membrane of the intestinal canal, principally in the larger intestines. They divide the effects into three stages, *o*, the cold stage, including vomit-

ing and diarrhoea. *b*, the stage of excitation, including nervousness, vertigo, and delirium; *c*, the state of reaction, including heat and copious sweat.

With reference to the physiological action of the common spurge (*Euphorbia Prespit*), the following statement is made by Dr. True

Headache with frontal fullness and heat heat about the eyes; languor and drowsiness, oppression of the stomach, and constipation. The juice applied to the eyes causes severe irritation, with smarting and burning, lachrymation, and momentary blindness this we have experienced twice while gathering the plant. It is supposed that this species causes the affection in horses called "slobbers."

6. *Mercurialis* L. Mercury

Annual or perennial herbs, with opposite pinnately veined leaves; flowers dioecious or monoecious in interrupted axillary spikes, apetalous, calyx small, green, 3-parted; capsule 3-lobed.

Mercurialis annua L. Annual Mercury

A leafy stemmed, erect, annual herb, leaves lanceolate or ovate lanceolate, crenate serrate, carpels hinged, indigenous to Europe, found in waste places from Nova Scotia to Ohio, and South Carolina. The *M. perennis* differs from *M. annua* in having a creeping perennial root, and hairy leaves.

Poisonous properties. Both species are acrid and poisonous.

7. *Acalypha* L. Three seeded Mercury

Herbs or shrubs, leaves alternate, petioled, flowers stipulate in spikes or spike-like racemes or solitary, calyx of staminate flowers 4-parted, calyx of



Fig. 139. Annual Mercury (*Mercurialis annua*). Staminate and pistillate branches. An acrid, poisonous plant. (After Paget.)

the fertile flower 3-5 parted, subtended by a foliaceous bract, petals wanting in both staminate and pistillate flowers, stamens 8-16 united at their bases, capsule consisting of 3 globular 2-valved carpels, each 1-seeded. About 250 species chiefly tropical, 3 species in the central and eastern states. *A. gracilens* has smaller leaves than *A. virginica*. *A. ostryaefolia* has echinate fruit and occurs from New Jersey to Texas and Kansas.

Acalypha virginica L. Three-seeded Mercury

A smoothish or hairy annual from 1-2 feet high often turning purple, especially in the autumn, leaves ovate or oblong ovate, sparingly serrate, long petioled, sterile spike few-flowered pistillate flowers 1-3 at the base of staminate peduncle capsule 3-lobed subglobose seeds ovoid, reddish striate.

Distribution. From Nova Scotia to Florida, Texas, Kansas and Minnesota.

Poisonous properties. This has been sent to me several times as supposedly poisonous. It is distasteful to cattle and they refuse to eat it in the pasture.

8. *Stillingia* L. Queen's Root

Smooth upright herbs or shrubs, leaves alternate or rarely opposite, frequently with 2 glands at the base, flowers in spikes, apetalous, calyx 2-3 cleft or parted staminate flowers, several together in the axils of the bractlets, stamens 2 or 3 pistillate flowers solitary in the axils of the lower bractlets, capsule 3-celled and 3-seeded. About 15 species of tropical America and the Pacific Islands.

Stillingia sylvatica L. Queen's Delight

A bright green herb 1-3 feet high, leaves nearly sessile lanceolate or elliptical, 2 glandular base, flowers lemon-colored subtended by small bracts with saucer-shaped glands, calyx cup-shaped, capsule depressed, seeds ovoid, light gray, minutely pitted and a flat base.

Distribution. From Virginia to Florida, Texas and Kansas in light sandy soil.

Poisonous properties. This plant is commonly used in medicine. It is said to be an efficient alterative. It contains an acrid resin *sylvestral* and an acrid fixed oil.

SAPINDALES

Trees, shrubs or herbs, petals usually present and separate, sepals usually distinct stamens rarely more than twice as many as the sepals or fewer opposite or alternate, ovary superior, compound ovule pendulous. Contains many tropical plants some with milky juice. In the family *Burseraceae* is the common box (*Bursera sempervirens*) which is used as a hedge plant and furnishes the best wood for wood engraving. The plant is an acrid poison. It is sometimes substituted for hops in the manufacture of beer and thus becomes the occasion of serious accident. The edible crowberry (*Empetrum nigrum*) belongs to the family *Empetraceae* and occurs far northward in America and Europe. The bladder-nut (*Staphylea trifolia*) of Atlantic North America belongs to the family *Staphyleaceae*. The family *Sapindaceae* contains the balloon vine (*Cardio-*



Fig. 340. Three-seeded Mercury (*Acalypha virginica*). Supposed to be poisonous to cattle. (Ada Hayden.)

spermum Hahcacadum), a well known ornamental climber, but a weed in the South. A refreshing drink is made from the seed of guarana (*Paulinia Cupana*) of South America, it contains *caffein*, *saponin* and an acid green fixed oil. The fruit of *Sapindus Saponaria* contains a great deal of *saponin* and is used as a substitute for soap. The soapberry tree (*Sapindus marginatus*) is used as a shade tree in the South where it is a native. A sheilac is derived from the Indian *Scheuchera trijuga* and marcassa oil is obtained from the seeds of the same plant. The Indians of Brazil use the honey collected by wasps from the flowers of *Serjania lethalis* to poison their arrows. It is also used as a fish poison and contains a narcotic principle which causes death. Another fish poison is furnished by the black seeds of *S. curassavica* of Brazil. The natives use the same substance for criminal purposes on man. The nectar obtained from the flowers is also poisonous. Lehmann lists as poisonous *S. nodosa* which is used by the natives of Brazil as an arrow poison. The fruit of *S. trifolatus* of India contains *saponin*. The same substance occurs in other



Fig. 341 Common Box (*Buxus sempervirens*). The plant is acridly poisonous. (After Faguet.)

species, notably in the seeds of the Brazilian *Magonia*. Narcotic principles occur in the following genera *Serjania*, *Nephelium*, *Magonia* and *Harpulia*. The fruit of the litchi (*Nephelium Lit-chi*), a native of China and the Philippines and cultivated in the tropics, is something like a plum and is eaten fresh or dried. The *Blighia sapida* of West Africa is cultivated for its edible arillus; the *Koeleruteria paniculata* of China is cultivated as an ornamental plant.

The family *Cornaceae* contains the genus *Cornaria*. The leaves and bark of the *C. myrsifolia* of southern Europe contain much tannin which is used in dyeing. The *C. ruscifolia* of New Zealand contains a black dye. The fresh leaves are used in making an intoxicating drink. *C. myrsifolia* and *C. thymifolia* of Mexico contain a toxic principle known as *coriamyrtin* which resembles *picrotoxin*. Many species of the genus are poisonous. *Coriaria sarmentosa*, *C. arborea*, and the tree-root (*C. Tutu*) of New Zealand are poisonous. Easterfield & Ashion* have isolated a crystallin glucoside called *tutu* $C_{11}H_{20}O_7$ which appears to be closely allied to *coriamyrtin* $C_{13}H_{22}O_7$. Tutu plants are highly toxic to animals that have not become immune by first becoming accustomed to small quantities. Blyth says

For the native cattle in the Tutu districts apparently consume moderate amounts of the shrubs with impunity, whereas other cattle become seriously ill. Both *coriamyrtin* and *tutu* belong pharmacologically to the *picrotoxin* group of substances. *Tutu* is somewhat less toxic than *coriamyrtin*. There is first depression, followed by agitation, the pulse is slowed, the respirations increased in frequency and finally tonic convulsions occur. 1.25 mgm. killed a kitten weighing 1 kilogramme in 48 minutes. 3 mgm. induced in a cat, 2 kilograms in weight, a convulsive seizure, and the animal did not recover for 24 hours.

Other important families of this order will be described farther on.

Families of Sapindales

Flowers regular

Ovary 1-celled, fruit a drupe Anacardiaceae.

Ovary 2 or more celled.

Leaves simple.

Seed with an aril Celastraceae.

Seed without an aril Aquifoliaceae.

Leaves simple, palmately veined or compound.

Leaves opposite Aceraceae.

Flowers irregular

Leaves palmately compound, fruit a leathery capsule

Trees or shrubs Hippocastanaceae.

Succulent herbs, capsule elastically dehiscent Balsaminaceae

ANACARDIACEAE. Cashew Family

Trees or shrubs with acid properties, milky or resinous juice, alternate or opposite leaves, flowers small, frequently polygamous, regular, calyx 3-7-cleft, petals of the same number stamens as many or twice as many as the petals, inserted at the base of the disk, ovary 1 or sometimes 4 or 5-celled, and 1 ovule in each cavity, styles 1-3, fruit generally a small drupe, endosperm scanty; cotyledons large.

There are about 500 species in temperate and tropical regions. The cashew (*Anacardium occidentale*) is much cultivated in the tropics. According to Dr. Cook the fleshy receptacles of the fruit are used in the West Indies in

* Jour. Chem. Soc. Trans. 1901.

preparing conserves and have an acid flavor which is very palatable, the pericarp, however, contains an irritant substance *cardol* $C_{21}H_{30}O_2$, which is black, acrid and vesicating, and is used to protect books and furniture from insects, cashew oil, equal to the finest almond oil and superior to olive oil, is also a product of the plant. The juice of the shell of the nut produces poisoning similar to that of poison ivy. The kernels of the cashew may be eaten raw or roasted like chestnuts, but the fumes coming from the roasting nuts are very caustic. The pistachia nut (*Pistacia vera*) produces a fruit about the size of a plum, which contains a seed much prized for eating. The mango (*Mangifera indica*) a native of India, is now cultivated in most warm countries for its fine edible fruit. The bark of many species, like the smooth sumach (*Rhus glabra*) of the North, and the European *R. Coriaria*, contains a valuable tanning material. The smoke tree (*Rhus Cotinus*) and the stag horn sumach (*Rhus typhina*) are frequently cultivated for ornamental purposes. The fruits of *Spondias dulcis*, *S. purpurea* and *S. lutea* are edible, the last of these is called the hog plum, being so named because the hogs are fond of it. The juice of another member of the family (*Comocladia*) causes an eruption similar to that from poison ivy.

Tannic and pyrogallic acids are derived from the Chinese Indian *Rhus semi-alata*. China turpentine (*Pistacia Terebinthus*) well known to the ancients, produces red galls that are used for tanning morocco leather. The mastic (*Pistacia Lentiscus*) native to the Mediterranean region, was formerly used for making varnishes.

The *Quercubria Lorenzii* and *Q. Balanar* of Argentine and Paraguay produce a very hard red wood which contains a great deal of tannin and gallic acid. The fruit of *Dracontomelon mangiferum* of the Sunda islands is used much like lemons. The ink tree of India (*Semecarpus Anacardium*) and *Hongaria ferruginea* contain *cardol*.

The pepper tree (*Schinus Molle*) cultivated in California, is a native of Peru. The saw dust of sneezewood (*Placoxylon utile*) produces sneezing.

Rhus L. Sumach

Trees or shrubs with alternate, simple trifoliate or odd-pinnate leaves, small polygamous flowers in panicles, calyx deeply 5-parted, petals 5, spreading, stamens 5 inserted below the flattened disk, fruit small 1-seeded. About 20 species in the temperate regions common in southern Africa. Some species are poisonous. The Japanese *Rhus vernicifera* and *R. succedanea* are cultivated in Japan for the lacquer which is taken from incisions made in the trees. Dr. White states that some of the embossed Japanese papers which are used in houses have caused severe inflammation, and according to Dr. H. N. Allen, natives as well as Europeans in the East are often affected with "varnish poisoning."

Rhus Toxicodendron L. Poison Ivy Three Leaved Ivy

A climbing or trailing shrub, sometimes erect, with 3 leaflets, plant erect or climbing by means of its aerial rootlets, flowers inconspicuous, polygamous in loose and slender axillary panicles, fruit globular, glabrous, whitish and waxy, frequently remaining on the plant until late spring.

Distribution. This plant is distributed from Nova Scotia to Wisconsin, Utah, Arkansas and Florida.

Rhus Vernix L. Poisonous Sumac or Dogwood

A shrub or small tree with pinnately compound leaves, leaflets 7-13, obovate-oblong entire, smooth, or somewhat pubescent, flowers polygamous in loose slender axillary panicles, drupe white, globose, oblong.

Distribution. Found in the swamps from New England to Ontario to Minnesota, Missouri, to Louisiana and Florida.



Fig. 342. Poison Ivy (*Rhus Toxicodendron*) a, spray showing axillary panicles and leaves, b, fruit—both one-fourth natural size (Chesnut, U. S. Dept. Agr.)

Rhus diversiloba Torr. and Gray California Poison Ivy

Nearly glabrous, erect or climbing shrub, leaflets 3 or rarely 5, obtuse or deeply pinnately lobed, flowers in loose axillary panicles, drupes subglobose.

Distribution. Common on the Pacific Coast from California to Washington.

Poisonous properties. All three species are poisonous to many persons, some persons being much more sensitive to irritation from the plants than others. Dr. J. C. White describes the effect of poison oak and poison ivy. He had collected freely of the plant for many years without any disturbance. Specimens were picked on September 28th, Oct. 6th, and Oct. 10th. He felt



Fig. 343. Poison Sumac (*Rhus Vernix*), showing leaves, fruit and seed pods, one-fourth natural size. (Cesnut, U. S. Dept. Agr.)

a sensation of irritation about the eyes and throat from the specimens of poison oak collected on the first named date, while working with the plants under an Argand gas burner, but nothing further was noticed. No unpleasant symptoms were observed from the poison oak (*R. venenata*—*R. Vernix*) collected Oct. 6th. From that collected Oct. 10, he experienced symptoms similar to those observed Sept. 28th. On Oct. 12th a single vesicle with a peculiar thick cover appeared the next day another and larger appeared on the wrist, two others came on the fourth day, others continued to appear up to Nov. 3rd, after which date the effervescences gradually subsided and were no longer perceptible. In another case described by Dr. White, the head was greatly swollen and features greatly disturbed. The skin of the face and neck was deeply oedematous and largely covered with vesicles of all sizes "many of which were seated on an erythematous base, others being still in their papular stage

of development." There were also large excoriations from which fluid was exuding freely, which on drying formed small crusts. The hands were also covered. "The subjective symptoms were great retching and burning of the parts affected, with the feeling of local discomfort, consequent upon so great swelling of the features. The eyes were nearly closed. There was a slight general febrile action." Dr White also reports the death of a child from a severe case of poisoning from poison ivy. The child though healthy was not robust. A recent case was reported from Packwood, Iowa, where a fourteen-year-old girl died after terrible suffering from the effects of coming in contact with the ivy, her face alone showing the eruption from the poison.

Hundreds of persons are poisoned every year from the three species. Dr White says:




Fig. 344 Poison oak (*Rhus diversiloba*), showing leaves, flowers, and fruit, one-third natural size. (Chesnut, U. S. Dept. Agr.)

Taking the simple vesicle, with scarcely any erythema surrounding it or any very perceptible infiltration of the underlying tissues, as the type of the eruption, whether occurring singly or in groups, we may have in a small percentage an abortive attempt at vesiculation, and an arrest of the development at the papular stage—a failure, that is, of the force exudation to force apart the layers of epithelial cells or a considerable infiltration into the papillary layer may elevate a cluster of the vesicles noticeably above the general surface or they may be surrounded by a well-defined erythema or congestion of the tissue immediately surrounding them in consequence mainly of the scratching and itching, which are the only subjective symptoms present.

In the severe cases, we have greater areas of simple erythema, a multiplication of the number of vesicles—either single or massed in close contiguity and covering large surfaces or by fusion forming blebs—a greater infiltration into the underlying corium with proportionate dimension of the capillaries and external redness, and a free exudation of serum into

the cutis. The overfilling of the vesicles causes a rupture of some of their epidermal coverings, and the discharge of their fluid contents upon the surface, forming moist, excoriated surfaces, covered in part with crusta.

With reference to a sequelae question of duration, there is a diversity of opinion. There is a popular belief that within a year after the first attack there will be a repetition of the original manifestations upon the skin which may be repeated for several seasons. Dermatologists think that a variety of cutaneous affections are developed in consequence of the action of the Poison Ivy. Dr. White considers that there are good grounds for this belief and in referring to his exhaustive researches on the subject, states that he was unable to find a single instance on record of the poisonous *Rhus* on the lower animals. After placing a notice in the "Spirit of the Times," a physician wrote him that once or twice while hunting where ivy abounded, his dog's eyes had been closed by swelling which he attributed to the action, but he had never observed any eruptions. The poisoning has been attributed to *toxocodendro* $C_{27}H_{43}O_n + 4H_2O$.

Remedies. The most popular remedy is to wash with sugar of lead (acetate of lead). Prof. Chesnut says

In practice it is not desirable to use strong alcohol which is apt to be too irritating to a sensitive surface but a weaker grade of from 30 to 75 per cent should be preferred. To this the powdered sugar of lead is to be added until no more will easily dissolve. The milky fluid should then be well rubbed into the affected skin, and the operation repeated several times during the course of a few days. The itching is at once relieved and the further spread of the eruption is checked. The remedy has been tried in a large number of cases and has always proved successful. It must be remembered, however, that the lead solution is itself very poisonous if taken internally.

Much has been said in regard to the relative poisonous character of these three plants. It has been generally claimed that the poison sumac is the most poisonous, and after it comes, first the poison ivy and then the poison oak. These conclusions were arrived at from the occasional experience of individuals who were poisoned by handling one species when supposedly immune to others. Experience teaches, however, that immunity is somewhat variable in the same individual, and therefore these general statements can not be accepted without more careful experimental evidence.

Annie Oakes Huntington in her recent book on Poison Ivy and Swallow Sumach says regarding the treatment

Soap, water, and a scrubbing-brush seem altogether too simple a method of treatment to advise for the painful eruption brought on by handling these two poisonous plants. Yet if we begin with this old-fashioned country remedy and study the various methods of treatment from one generation to another, we return at last, through the most recent scientific investigations, to our original starting-point. The only effective measures are preventive ones; the only remedy is a wash which mechanically removes the poisonous oil from the skin. In this lies the sum and substance of the entire method of treatment.

She made an experiment in which it was shown that only preparations spread the poison and that constant washing with soap and water removes the poisonous or which causes the trouble. The toxic principle is soluble in alcohol and this may consequently be used. A weak solution, 50 or 75% is advisable but the treatment must be renewed. One part of hyposulphite of soda to 3 parts of water is another good solution recommended by her.

Syme in Dr. Remsen's laboratory, has come to the conclusion that the active principle of Poison Ivy is a glucoside and not an unknown volatile oil as stated by Pfaff. The glucoside as determined by Syme, is a compound of *rhamnose*, *gallic acid* and *ficetin*. It can be precipitated by a lead acetate. Syme tested the toxic action of the various fractions upon himself and was able to determine the chemical nature.

In a recent paper by Dr. Ford there seem to be some evidences for concluding that immunity may be obtained. That such immunity exists may be taken from the clinical symptoms that different persons are sensitive to even small amounts of the poison and in other cases persons who have been poisoned become accustomed to it. Synge in his experiments upon himself found that after four or five months he was no longer susceptible to the poison.

The experiments performed by Dr. Ford are of interest. The experimental material was obtained in the alcoholic fluid extract of the native plant prepared by Parke, Davis & Co.

It had already been shown by Pfaff that the internal administration of his non-volatile oil produced definite lesions in rabbits, the animals dying of an acute nephritis at the end of 16 to 18 days. Occasionally the rabbits died in acute convulsions without any microscopic brain lesions. The subcutaneous administration of the fluid extract of *Rhus Toxicodendron* produces the same effect upon rabbits as those described by Pfaff. Rarely the rabbits die in convulsions within 24 to 48 hours, but the majority of inoculated animals succumb in from 8 to 13 days. In addition to the nephritis an extensive necrosis and slough is found at the point where the poison is introduced beneath the skin. Following the inoculation we have a fairly long latent period during which the weight of the animals remains stationary. After seven or eight days in a typical case, the animal loses weight rapidly, the necrosis and slough develop, and the animal dies of the nephritis after the lapse of about two weeks. At times the skin lesions are less marked, the damage to the kidney being the important change. An intraperitoneal inoculation seems able to produce these kidney changes more rapidly than does the subcutaneous method. In addition to rabbits we have found that guinea pigs are susceptible to the drug, the lesions being produced with a water certainty and regularly. With these animals the necrosis and slough at the point of inoculation are more extensive, while the animals die of the kidney changes in about the same time. The fatal dose of the poison can be estimated for both animals with tolerable accuracy. For guinea pigs of 250 gram weight 0.25 c.c. of the alcoholic extract always represents a fatal dose, and a guinea pig of 150 gram weight practically never survives a dosage of 1 c.c.

The fatal dose for rabbits of 910 grams is 1 c.c. Certain rabbits of this weight die from smaller doses, but not regularly and larger animals show greater resistance. Animals of 1,800 to 2,000 gram weight occasionally survive 2 to 3 c.c. doses but not more than this amount.

Experiments were made to determine whether animals which had withstood some doses of the poison were susceptible to the amounts of poison capable of killing untreated animals. The experiments were made with four guinea pigs, varying in weight from 450 to 900 grams, and on rabbits varying in weight from 800 to 2,400 grams. The initial doses were small when the animals regained weight larger doses were given. It was found that the serum of immunized animals contained substances neutralizing the poisonous principle when both were injected into susceptible animals. Goats have been immunized and it is probable that immunity may be obtained in other animals.

How to Treat the Poison Ivy Patch. Various methods of treating the poison ivy have been tried. The iron sulphate, 100 pounds to a barrel of water is only partially successful. Covering with tatted paper creosoted below is effective according to Dr. G. H. Stone. This writer has also shown that sodium arsenate (commercial) at the rate of 2 pounds to 10 gallons of water poured around the roots is effective.

Rhus glabra L. Smooth Sumach

A low shrub from 2 to 12 feet high, stems with large pith, brownish bark compound leaves 11 to 31, leaflets pointed serrate whitish beneath green above flowers borne in a large panicle greenish, polygamous, calyx small 5-parted petals 5, stamens 5, inserted under the edge of a flattened disk, pistil with a 1-celled and 1-ovuled ovary stylets 3, terminal, fruit red, small 1-seeded.

Distribution. Common in Northern United States to Florida, Mississippi, Arizona, Utah, British Columbia to Nova Scotia. A troublesome weed at times in rocky soils.

Poisonous properties. There is a popular impression that the red drupes of this species are poisonous, but I know of no record where the eating of this fruit has produced poisoning. The fruit is decidedly acid.

AGNUSFOLIACEAE. Holly Family

Trees or shrubs with small, simple, mostly petioled, alternate leaves; flowers in axillary clusters, chiefly polygamo-dioecious, calyx minute, free petals 4-8 or more, stamens free, as many as the petals, pistil 1, ovary superior, 4-8 celled, ovules 1-2 in each cavity, fruit a small berry-like drupe, seeds with small embryo, endosperm present.

A small order of 160 species. The American holly (*Ilex opaca*) from Maine to Florida, Missouri and Texas, is much used for Christmas decoration. The European holly (*Ilex aquifolium*) is used for a similar purpose. The berries of this species, though eaten by birds, are said to be poisonous. Whether the seeds of the American species are poisonous is not known although it contains the principle *shon*. The *Ilex Cassine*, which occurs from southern Virginia to Florida and Louisiana along the coast, was used by the Indians during their religious ceremonies to make what they called their "black drink" an emetic intended to clear the head and stomach. It contains *caffein*. The yerba or maté (*Ilex paraguayensis*), native to southern Brazil and Argentine Republic is used like the Chinese tea and is stimulating. Properly the term "maté" is applied to the drinking cup made from a small gourd. Yerba is an important article of commerce in South American countries. The leaves contain the same active principle, *caffein*, that is found in the tea. Two other species *I. theae* and *Symplocos lanceolata* also furnish the maté. The wood of the larger trees of the genus, like *Ilex opaca* and *Ilex Aquifolium* is white and is used by cabinet makers.

Celastraceae. Staff-tree Family

Shrubs, or trees, with simple leaves, stipules small or absent, flowers regular, usually perfect, calyx 4-5-lobed, petals 4-5, stamens inserted on a flat or lobed disk, pistils with 3 or 5-celled ovary, ovules 2 in each cavity, fruit 2 to 5-celled, fleshy, seeds with an aril, embryo large, and fleshy endosperm. About 350 species of wide distribution.

The burning bush or wahoo (*Euonymus atropurpureus*) is a well known native, frequently cultivated and is a most desirable shrub. The *Coffea edulis* of Arabia is extensively cultivated and is used as coffee by the Arabs. The leaves are also chewed by the natives, having a stimulating effect similar to that of cocaine. It contains the alkaloids *caffin* and *celastrol*. The *Elaeodendron australe* of New South Wales is used for cabinet work. The genus *Pachistima* is represented in the Rocky Mountains by *Pachistima Myrsinites*, and in the Alleghany Mountains by *Pachistima Canbyi*; both of which are pretty shrubs.

Celastrus L. Staff-tree. Bitter-sweet

Mostly climbing shrubs; leaves thin, flowers racemose or paniculate, polygamo-dioecious, calyx 5-lobed, petals 5, crenulate, inserted under the disk,



Fig. 343. Washoe (*Enonymus alropurpureus*). 345a. Bitter sweet (*Celastrus scandens*)

pod globose, orange color, 2-4-celled dehiscent into as many valves, seeds enclosed in a scarlet aril, endosperm fleshy. About 30 species.

Celastrus scandens L. Shrubby or Climbing Bittersweet

Chiefly climbing shrubs with alternate leaves; flowers small, polygamo-dioecious, calyx 5-lobed; petals 5; stamens 5, inserted under the 5-lobed disk; capsule globose, orange-colored, 3-celled and 3-valved; seeds 1-2 in each cell, enclosed in a pulpy aril. About 30 species. The *C. articulatus* and several other species are commonly cultivated and are hardy. Several species are natives of the Cape of Good Hope.

Distribution From Quebec to Manitoba in Canada, and from Kansas to Indian Territory, New Mexico and the Carolinas.

Poisonous properties. The aril of "Bitter-sweet" has a sweetish, somewhat disagreeable taste. The leaves of the plant are said to be poisonous to horses. The plant *Euonymus* contains the amorphous bitter, odorless substance, *eunonymus*, which acts as a powerful heart poison. The washoe acts as a drastic purgative. The symptoms are those of deathly nausea, vertigo, prostration and cold sweat.

ACERACEAE. Maple Family

Trees or shrubs with opposite, simple or compound leaves, flowers polygamous or dioecious in cymose or racemose clusters, calyx 5-parted, petals of the same number or none; stamens 4-12, inserted on a fleshy disk, ovary 2-lobed and 2-celled, styles 2, fruit a samara, exalbuminous cotyledons thin folded. There are 3 genera and about 100 species most of them in the genus *Acer*, the maples being widely distributed in temperate regions. The maple, (*Acer*) is commonly used for the manufacture of furniture and for inferior finishings, floorings, etc. The most highly prized are the hard maples (*Acer nigrum* and *A. saccharum*). Sugar maple is also derived from these species. Curly maple is only a form of wood of these and of the *A. macrophyllum* of the Pacific Coast, which is also much prized for cabinet work. The maples, including the box elder (*Negundo aceroides* or *A. Negundo*), are also used for shade trees. The silver maple (*A. saccharinum*) is widely distributed in the United States. The red maple (*A. rubrum*) is less commonly used. The bark of *A. rubrum* was used by the Indians as a remedy for sore eyes.

HIPPOCASTANACEAE Buckeye Family

Trees or shrubs, leaves opposite, petioled, digitately 3-9-foliate, flowers in terminal panicles, irregular and polygamous, calyx 5-lobed or 5-cleft, petals 4-5, unequal, clawed, disk entire; stamens 5-8, ovary sessile, 3-celled; ovules 2 in each cavity, style slender, capsule leathery, smooth or spiny, 1-3-celled, seeds large shining; cotyledons very thick. Only 2 genera and 15 species in America and Asia.




Fig. 346. Sugar maple (*Acer saccharum*). 1. Branch bearing staminate flowers. 2. Branch bearing pistillate flowers. 3. Fruiting branch. 4. Staminate flower, enlarged. 5. Longitudinal section of staminate flower, enlarged. 6. Pistillate flower, enlarged. 7. Longitudinal section of pistillate flower, enlarged. 8. Longitudinal section of fruit, enlarged. 9. Longitudinal section of seed, enlarged. 10. Embryo, enlarged. 11. Winter branchlet. 1, 2, 3, 8, 11 one-half natural size. (H. M. Cheney in Green's Forestry of Minn.)

Aesculus L. Buckeye

Trees or shrubs with opposite petioled digitately 3-9-foliate leaves, flowers in panicles, irregular, polygamous, calyx 5-lobed, lobes unequal, petals 4-5, unequal, clawed, stamens 5-8, filaments long, often unequal, pistil with 3-celled

ovary and two ovules in each cell, capsule leathery, seeds large with shining coat, cotyledons thick and fleshy. A small genus of 15 species native of America and Asia. The horse chestnut (*Aesculus Hippocastanum*), escaped from cultivation is planted for ornamental purposes, as are others of the genus, like the species described below and *A. parviflora*, a small shrub. By washing and boiling the starch in the seed may be utilized, and this is done in France with the horse chestnut. The wood is light and brittle. The wood of the Ohio buckeye is used for making violins.

Aesculus glabra Willd. Ohio Buckeye

Trees with long-petioled leaves rough and fetid bark, flowers pale yellow, in large panicles, polygamo-monoecious, calyx bell-shaped, stamens exserted, curved, petals unequal. fruit slightly prickly when young smooth when old.

Distribution. Western Pennsylvania to Central Iowa, Kansas and Indian Territory.

Aesculus Pavia L. Red Buckeye

Shrubs with 5-7 digitate, nearly smooth, leaflets, acute or short acuminate, pubescent when young, becoming smooth, flowers in loose peduncles, calyx tubular, bright red, petals bright red.

Distribution. In fertile valleys from Virginia to Florida. Arkansas and southern Missouri.

Aesculus Californica Nutt. California Buckeye

Usually a shrub from 10-15 feet high, or occasionally a tree from 25-40 feet high, 3 feet in diameter, leaflets 4-7, usually 5, smooth, oblong lanceolate, acute, petiolate, flowers in a close panicle, calyx 2 lobed, petals somewhat unequal, white or pale rose, $\frac{1}{2}$ inch long, stamens 5-7, ovary densely pubescent, fruit usually 1-seeded.

Distribution. In California.

Poisonous properties. The leaves and fruit of the above species are regarded as poisonous. Many farmers claim that this is true only at certain seasons of the year. The seed produces sneezing and enters into the manufacture of snuff. The California species, according to Chestnut, causes abortion in cows. Dr. Rusby states that in southern states the seeds are crushed and thrown into water to stupefy fish just as the bark and roots of the relatives are in the tropics. Fatal cases of poisoning of children are reported from Texas. Suspicion has been attached to the common horse chestnut. The European chestnut is said to be useful in affording food for live stock, especially sheep and goats. This species contains *arsenulin* $C_{15}H_{16}O_8 + H_2O$, a glucoside found in the bark of many trees of the order *Sapindaceae* and the glucoside *aesculetin* $C_9H_8O_4$ and *parin* $C_{17}H_{34}O_{20}$. the testa of the seed contains *quercetin* $C_{30}H_{48}O_{12}$, *argyrescin* an acid, amorphous glucoside, *aphrodisiacin*, also an acid, amorphous pink pile and *saponin* $C_{12}H_{24}O_8$, a glucoside which is also found in the roots of *Polygala Senega*, and other plants. Dr. Milspaugh states that the horse chestnut causes inflammation of the mucous membranes of the respiratory and digestive tracts, and especially of the rectum, constant burning in the stomach and epigastrium, followed by nausea, retching, and violent vomiting with great tenderness and colic throughout the ab-




Fig. 147. Horn Chestnut (*Aesculus Hippocastanum*). Flowering branch. Entire flower. Stamen. Pistil. Entire seed and longitudinal section. Wood used in making valves. (After Faguet.)

down, are markedly present. The buche is an irritant of the cerebro-spinal system, the more prominent symptoms being confusion of mind, vertigo, stupefaction and coma.

BALSAMINACEAE. Balsam Family

Succulent herbs. leaves alternate thin, petioled. flowers axillary showy, irregular, sepals 3, the two lateral small green, the posterior large and petal-like, spurred; petals 3 or 5, some 2 cleft; stamens 5; ovary oblong 3 celled, style short or none; stigma 3-toothed or 5-lobed; ovules several in each cell, fruit a capsule in *Impatiens* coiled elastically, expelling the seeds, seeds



Fig. 348. Ohio Buckeye (*Aesculus glabra*). Causes irritation of the cerebral spinal system. (Lies Paumot.)



Fig. 349. Red Buckeye (*Aesculus parva*). a flowering branch, b, seed both two ninths natural size. (Chestnut. U. S. Dept. Agr.)

ridged, embryo straight. About 200 species mostly of tropical Asia. One genus with 2 species is native to eastern North America. The Balsam (*Impatiens Balsamina*) is frequently cultivated. The sap of some species contains a dye.

Impatiens L. Jewel Weed

Succulent herbs with simple, thin, petioled leaves, sepals 3, the 2 alternate small, green, the posterior one largest, and forming a spurred sac, petals 5, or 3, with 2 of them 2-cleft into dissimilar lobes, stamens 5, short, ovary oblong, 5-celled, fruit an oblong or linear capsule, deluscent elastically into a coned valve, scattering the seeds, endosperm none, embryo nearly straight, cotyledons flat.

Impatiens biflora Walt. Spotted Touch-me-not

A glabrous annual from 2-4 feet high, leaves ovate or elliptical, pale and glaucous beneath, flowers orange-color, thickly spotted with brown, peduncles 2-4 flowered.

Distribution. In moist grounds from Eastern Canada to Florida, to Kansas and northward to Oregon and Alaska.

Poisonous properties. Dr. Schaffner states that the plant is emetic and suspected of being poisonous to stock.

RHAMNALES

Shrubs, vines or small trees, leaves generally alternate, flowers small, regular, sepals generally more or less united, petals distinct or wanting, stamens as many as the calyx lobes and alternate with them, opposite the petals when present, ovary superior, compound, ovules erect. They contain two families, the *Rhamnaceae* and *Vitaceae*. The genus *Litis* embraces 40 species found in warm and temperate regions. The European grape (*Litis vinifera*) native from Eastern Europe to Central Asia is now widely cultivated in California, Spain, Germany, the Cape region and elsewhere. The Warden and Concord grape (*V. labrusca*) of eastern North America are also widely cultivated. Other species are, the small grape (*V. acinacalis*), the wild blue grape (*V. bicolor*), native from New York to Wisconsin, the southern fox grape (*V. rotundifolia*) with a silky flavor cultivated in the South, the cultivated northern fox grape (*V. riparia*) with very fragrant flowers. Improved forms are the Janeville and Clinton. The fruit of the mustang grape (*V. californica*) of Texas is very acid. The Virginia creeper (*Pteris quinquefolia*) is a well known ornamental climber. The Boston or Japan Ivy (*P. tricuspidata*) native of Japan is a handsome climber scarcely hardy north. The *P. heterophylla* another ornamental from China and Japan is a hardy plant with small blue berries. It does not cling. The *Ficus incensura* of Japan contains *torreodendro*, and is poisonous.

RHAMNACEAE. Buckthorn Family

Shrubs, often climbing, or trees, often thorny, with astringent or bitter qualities, leaves chiefly alternate, stipules small, deciduous, flowers in cymes or panicles, small, regular, calyx perfect or polygamous, 4-5-toothed, petals 4-5, inserted on the calyx or none, stamens 4-5, inserted on a disk which lines the calyx tube, which is often united with the single 2-5-celled ovary, ovules 1 in each cell. Fruit often mucilaginous and drupaceous. A small family of 550 species, of temperate and warm regions.

The supple jack (*Berchemia scandens*) is a pretty climber of southern woods. The buckthorn (*Rhamnus cathartica*) is frequently cultivated for hedges in the north. The juice of the unripe drupe was formerly used for staining maps and the ripe drupe is the sap green of painters. The sap has strong purgative properties. This and *R. Frangula* are local irritants. This plant contains *rhamnetin*, $C_{14}H_{12}O_7$, the *rhamnin* of earlier authors, a gluconide found in the berry. Dyes are obtained from *R. infectoria* and a dye for silks from *R. lahurca* and *R. tinctoria*. The *R. Frangula* contains *frangulin* and is an ornamental shrub, the charcoal from which is used for making gun powder. Probably all are more or less injurious. Cascara sagrada is obtained from *R.*

Purshiana, a shrub or small tree native to the Pacific coast which contains *purshianin* and is an excellent laxative. The *Paliurus australis* contains oil of wintergreen. The jujube (*Zizyphus sativa*) is used as food in South Europe and Western Asia. The Chinese *Z. Jujuba* is extensively cultivated in India and China. The fruit of *Z. Lotus* is made into a kind of bread, used by the natives of West Africa, but the ripe fruit is said to be injurious. The root of the New Jersey tea (*Ceanothus americanus*) is used as an astringent and expectorant and is also said to have been used in place of tea during the Revolutionary war. According to Greshoff the seeds of *Ceanothus americanus*, and *Ceanothus ovatus* contain a small amount of saponin. It is therefore not strange that some members of the family are poisonous. The genera *Zizyphus*, *Tapura* and *Gouania* furnish fish poisons. *Saponin* occurs in *Gouania tomentosa* of Mexico. *Colubrina fermentis* is used as a substitute for kapa. The wood of *Colletia spinosa* of South America contains a bitter principle.

The brownish-black berries of coyotillo (*Kartensia Humboldtiana*) of Texas are said to be very poisonous and Dr. A. Mitchell of San Antonio writes of that the plant is poisonous to goats.

MALVALES

Herbs, shrubs or trees. leaves simple, generally alternate, flowers regular, usually perfect, sepals separate or more or less united, corolla polypetalous or rarely wanting, stamens usually numerous, ovary superior compound, parietal axile. The more important families of this order are *Tiliaceae*, *Malvaceae*, *Bombaceae* and *Sterculiaceae*. The first family contains the basswood (*Tilia americana*), a well known timber and ornamental tree of North America, used for making boxes, lumber, excelsior, etc. *T. cordata* is also frequently cultivated under the name of linden tree. The inner bark of this produces an elastic fiber. Several species of the genus *Greenea* are used as fish poisons. Jute is derived from *Corchorus capsularis* of the East Indies and is a valuable fiber. The bromweave (*C. stenosus*) of the West Indies furnishes a substitute for tea. The fruit of *Apeiba* is edible. Corkwood (*Ochroma Lagopus*) of the family *Bombaceae* is used by the fishermen of Trinidad on their nets in place of cork and is one of the lightest of all woods. The monkey bread tree (*Adansonia digitata*) produces valuable fiber in its bark. The big cotton or kapok tree (*Eriodendron auriculatum*) furnishes a soft fiber used in upholstery. The seed known as "kapok" seed is used in the Indies as fire and in making oil cake and according to Reinders and Kuhn, is an adulterant of insect. The *Durio zibethicus* of the Malayan region produces a large edible fruit. The family *Sterculiaceae* of tropical countries includes the cacao (*Theobroma Cacao*) which produces the well known cacao beans. Cacao is a nutrient food and contains the alkaloid *theobromin*, $C_7H_8N_4O_2$, a caffeine alkaloid. The cola nut (*Cola acuminata*) of West Africa is a stimulant. The negroes of Brazil used large quantities because of its stimulating properties. It is also used for similar purposes by the negroes of the south, and the "Cola habit" is increasing among the negroes of that region. It is probable that the cola is frequently adulterated with injurious ingredients. It is a muscle stimulant used by the Alpine chambers of Europe. Fresh cola nuts do not contain caffeine, but a glucoside *kolanin* which is converted into *kolanin* ($C_{14}H_{18}(OH)_8$). An



Fig. 350. Jute Plant (*Corchorus capsularis*). A valuable fiber plant from the East Indies. (From American Agriculture, 4)

oil is manufactured from *Sterculia foetida*. The *Abramo augusta* yields a fine fiber.

To the family *Elaeocarpaceae* of the same order belongs the *Echinocarpus Sigus*, a poisonous plant which contains hydrocyanic acid. Several members of this family are economic plants. The maqui fruit is obtained from *Arctostaphylos Macqui*, native to Chili. The seeds of *Sloanea dentata* are eaten like chestnuts. The seeds of *Muntingia* are edible.

MALVACEAE. Mallow Family

Herbs, shrubs, or, in tropical countries, trees, with mucilaginous properties, tough fibrous bark and stems; leaves alternate and small, stipules small, deciduous, flowers regular and generally perfect; sepals 5, usually more or less united, petals 5, hypogynous, stamens numerous, monadelphous, several-celled; pistils several, styles united, projecting beyond the stamens above, ovary several-celled, seeds nearly exalbuminous, embryo curved. A family of about 800 species of wide distribution and of great economic importance. The cotton plant is the most important member of the family and comprises several species, the most important in the United States being *Gossypium herbaceum*, which




Fig. 35 Cacao Tree (*Theobroma Cacao*). 1 Branch with flowers and fruit. Flower cut through longitudinally. 3 Seed. 4 Fruit with a portion of pericarp removed. The beans are well known articles of commerce. (After Wessillo)

is cultivated for its fiber. The cotton fibers are plant hairs coming from the seed. Cotton seed is used to manufacture an oil serving in place of olive oil, and cotton seed meal, the latter a valuable stock food. The refuse material is used as a fertilizer. The highly explosive gun cotton is made by soaking cotton in sulphuric and nitric acids. Other cottons are Sea Island cotton (*G. barbadense*) and tree cotton (*G. arboreum*). The bark of cotton root has very active principles. Marsh mallow (*Althaea officinalis*) used in medicine as a stimulant and in confectionery contains *asparagin* $C_4H_4N_2O_4H_2O$ and *bassorin* $C_{14}H_{20}O_{10}$.

Okra or gombo (*Hibiscus esculentus*) is extensively cultivated in the south and in Europe for the young mucilaginous pods which are edible. The *H. ficulneus* is used in a similar way. The fleshy red calyx of the Jamaica sorrel (*H. Sabdariffa*) is used in making jellies and sauces, the fiber makes good cordage material.

The wood of the Cuba bass (*Hibiscus elatus*) native to the West Indies, produces a timber of greenish color used in cabinet work. The lace-like inner bark is used for wrapping and is known as lace bark. From the fiber of *H. tiliaceus* a strong paper can be made very cheaply. This is also used by the natives of the Pacific Islands in making ropes.

Fibers are obtained from *Hibiscus tiliaceus*, *H. cannabinus*, *Arena lobata*, *Abutilon indicum*, *Sida retusa* and *Napaea laetis*. A beautiful wood is derived from the magar (*Thespesia grandiflora*) of Porto Rico, the color being red when fresh black when dry. The hollyhock (*Althaea rosea*) is well known.

cultivation, as are some species of the genus *Abutilon* and the poppy mallow (*Calthraoe involucrata* and *C. triangularis*)

Chorana of eastern South America furnishes a soft fiber. The seeds of *Pachira macrocarpa* indigenous to Brazil contain a valuable oil resembling that found in cacao, kapok oil is obtained from the seeds of *Eriodendron anfractuosum*. Musk seed is obtained from *Hibiscus abelmoschus* of the East Indies and is used in perfumery. The *Malva moschata* of Europe produces a similar odorous product. The cheeseweed or dwarf mallow (*M. rotundifolia*) is a troublesome weed in gardens, waste places and barnyards. *Saponin* is found in the roots of *Sida jamaicensis* and *Hibiscus Sabdariffa*. The *Sida paniculata* is used as an anthelmintic. The ripe capsules of Queensland hemp (*Sida rhombifolia*) causes the death of fowls that feed on it.



Fig. 352. Dwarf Mallow
(*Malva rotundifolia*) (After Flich.)

Key for Malvaceae

Flowers involucrate.

Flowers small, seed solitary and not covered with cotton

2 *Malvastrum*.

Flowers large, seed covered with cotton

1 *Gossypium*.

Flowers not involucrate

3 *Abutilon*.

Gossypium L. Cotton

Herbs, shrubs, or rarely trees from 2-10 feet high. Leaves alternate, palmately veined, and lobed, stipulate, involucre of 3 heart-shaped leaf-like bracts, flowers large, regular, white or whitish, sepals 5, petals 5, stamens numerous, anthers borne along the outside of the tube of the filaments, ovaries 3-5-celled as many as the cells of the pod, seeds numerous bearing cotton. The description of one species only is given below, the Sea Island cotton (*G. barbadense*), cultivated in Florida and along the Gulf Coast. Tree Cotton *G. arboreum* is cultivated in the tropics. The fiber is long, silky and an inch or more in length. But little is produced, its use being restricted, it is said, to making thread for turbans for the priestly class.

Gossypium herbaceum L. Common Upland Cotton

An annual, 3-6 feet high, leaves with 5 short and roundish lobes. Flowers large, pale yellow, turning rose color. seeds covered with cotton. Prof. L. H. Dewey thinks that our upland cotton should be referred to *G. hirsutum* (*G. herbaceum*), which is a native of Mexico.* The crop in India, according to Dr. H. J. Webber is derived chiefly from *G. herbaceum* and in Egypt, the crop is obtained chiefly from *G. barbadense*. In warm climates, cotton is a perennial. Cotton was cultivated long before the Christian era. It is one of the most important crops of the world. Dr. Webber states that in 1792 the crop was 60,880 bales; in 1820 6,000 bales; in 1840, the product increased to 4,443,311 bales, reaching 8,547,468 bales in 1862, and in 1904, 13,693,279 bales. In a single century from 1804 to 1904, the crop increased from 130,000 bales, valued at \$13,000,014, to 13,693,279 bales valued at \$557,147,306. In the early history of cotton cultivation, the seeds were not valued at all. Growers were troubled to know how to get rid of them. But in 1904 the seeds alone were valued at \$90,258,227.86, making the total value of that year's crop, unmanufactured \$647,405,534.51.




Fig. 353. Cotton (*Gossypium herbaceum*) 1, a plant with flowers and leaves, 2, a cotton boll. 3, seed. (Strasburger, Noll, Schenck and Schumper.)

Distribution. Commonly cultivated in the Southern States.

Poisonous properties. The root is commonly used in the south and tropical countries to produce abortion. Dr. Johnson says:

Cotton acts as an abortifacient. Its action upon the uterus is similar to that of ergot and it is used instead of the latter in cases of uterine inertia during parturition, and in amenorrhoea, dysmenorrhoea, and scanty menstruation. Whether its action upon the system at large be similar to that of ergot is unknown but worthy of investigation.

* As to the botanical status of the different species of the genus *Gossypium* the following papers should be consulted: L. H. Dewey *Cyclopaedia of Agr.* 2: 281. Advance article on cotton by Webber in an earlier part of the same work (4: 348). The Nutzpflanzen Togus Notizblatt Konigl. Bot. Garten Berlin. App. 2. No. 2, p. 49. J. F. Cook, Origin of the Hindi Cotton Cir. Bur. Pl. Ind. 42. (This last paper gives some of the literature. Wall. St. George, The Wild and Cultivated Cotton Plants of the World, London 1907.

The fact that the bark of cotton root should be possessed of such active properties is an interesting and suggestive one, for it affords the only instance of specific virtues attributed to a plant of the order *Malvaceae*. It will be noted that it is only the bark of the root that possesses these properties, all of the other parts of the plant partaking of the bland character of the order—the seeds especially, which not infrequently possess the most active principles of plants, in this instance yielding a bland oil which is largely used as an adulterant of olive oil. Now is an order of plants of such strongly marked characteristics as the *Malvaceae*, the discovery of striking properties in any individual should stimulate investigation of other related individuals, for it may reasonably be assumed that investigation which starts with a rational clue will be more fruitful in results than if conducted entirely in a haphazard manner. Cotton seed is often injurious to animals.

In Friedberger and Frohner's *Veterinary Pathology*, Capt Hayes writes as follows:

Only young animals are usually affected. They display disorder of digestive and urinary apparatus, tympanites, diarrhoea (afterwards bloody), haemorrhage from the urinary organs, albuminaria, strangury and paralysis of the bladder. Duration is variable. Autopsy reveals in the acute form haemorrhagic gastro-enteritis, opaque swelling of the liver and kidneys, oedema of the lungs, dark red urine, collection of liquid in the cavities of the body. There is no enlargement of the spleen.

The liver is opaque and swollen. In chronic cases only, one finds general emaciation and dropsical phenomena. In dairy cows it induces garget and mammitis. Cotton seed oil cake contains a poisonous principle ricin.

The investigations Prof. M. B. Hardin of South Carolina made in 1892 indicated the presence of meta and pyrophosphoric acid in cotton seed meal. He suggested that these acids are poisonous. Dr. Crawford of the U. S. Dept. of Agrl. has recently investigated the subject and concludes that the pyrophosphoric acid is toxic. The seed from upland cotton is more toxic than the Sea Island. The toxic effect may be increased by heating, when the temperature rises high, this is due to the conversion of the orthophosphoric acid into the pyro form. The investigations of Dr. Crawford are very important.

Malvastrum Gray

Herbs with entire cordate or divided leaves, flowers solitary or racemose, short pedicelled, involucrete, or none, calyx 3-cleft, petals 5, notched at the end or entire, styles 5 or more; stigmas capitate; carpels indehiscent or imperfectly 2-valved, seed kidney shaped. About 75 species, natives of America and South Africa.

Malvastrum coccineum (Pursh) Gray. False Mallow

A low hoary perennial herb, with dense silvery stellate pubescence. Lower leaves pedately 3-5 parted, flowers small, red, in dense, short racemes, usually without bractlets. calyx lobes shorter than the pink red petals. carpels 10 or more, indehiscent, rugose, and usually 1 seeded.

Distribution. From South Dakota to Texas, New Mexico to British Columbia.

Poisonous properties. Has been suspected of being poisonous, but there is no evidence to support this view. Profs. Chesnut and Wilcox regard the plant as non-poisonous. Some ranchers consider that it may be the cause of loco poisoning.

Abutilon (Tourn.) Mill. Indian Mallow

Herbaceous or shrubby plants, or in the tropics, trees with soft, pubescent leaves and stems. involucrel bracts none. calyx 5-cleft. petals 5. styles 5 or

more, carpels 2-valved, 2-9 seeded, seeds reniform. About 90 species in tropical or warm temperate regions. Many of the species are under cultivation.

Abutilon Theophrasti Medic. Velvet-leaf

A stout, strongly-scented annual, 2-4 feet high, with tough, fibrous stems, leaves roundish, heart-shaped, tapering to a point, velvety; peduncles shorter than the petioles, flowers yellow, carpels 12-15, pubescent, opening at the apex, each valve beaked by a slender awn.

Distribution. Common in the northern states. Naturalized from India.

Poisonous properties. It is reported as poisonous, the strong odor is so very objectionable that it is not likely that much of the plant will be consumed by stock.

PARIETALES

Herbs, shrubs, or trees, flowers generally complete, perfect and regular or in some cases irregular, sepals distinct or united, imbricated or convolute, petals nearly always present and distinct; stamens mostly numerous ovary compound superior or inferior in some, placentae mostly parietal. The order includes a number of important families. *Caricaceae* contains the pawpaw (*Carica Papaya*) the edible fruit of which is from 6-10 in. long of yellow color and contains the alkaloid *carpene* $C_{11}H_{21}NO_2$, acting similarly to digitalis, and a glucoside *caricin*, it also contains the enzyme *papain*, which resembles *trypsin* in its proteolytic action, converting animal proteins into proteoses forming peptones. This does not occur in vegetables. These changes do not go on in acid and alkaline substances, but act best in a neutral medium. The most active changes occur best when the medium is from 33 to 40 degrees C. A similar ferment occurs in the leaves of *Carica quercifolia* of Argentina. *Caryocaraceae* includes gamboge (*Garcinia Hanburyi*) a cathartic the mammy apple (*Mammea americana*) cultivated in the West Indies, the *Calophyllum Calaba* which furnishes the Calaba balsam, the mangosteen (*Garcinia Mangostana*) of the Moluccas, widely cultivated in the tropics. The *Dipterocarpaceae* includes the genus *Dipterocarpus*, which furnishes resin, and *shorea*, furnishes Chalia resin. "Piney resin" is obtained from *Vateria* and is used as a substitute for dammar. The *Tamaricaceae* contains the ornamental *Tamarix gallica*, an excellent honey plant the *Tamarix mannifera* from which a manna-like sugar is derived the *Pouqueria splendens* of Mexico frequently cultivated and producing the ocotilla wax, and the *Myrsine germanica*, the twigs of which are used as a substitute for hops. *Cistaceae* includes the rock rose (*Helianthemum canadense*), the pinweed (*Lechea minor*) and the *Cistus polymorphus* furnishing laudanum.

The *Begoniaceae* includes the commonly cultivated genus of *Begonia* with 530 species. The *Bixaceae* includes *Bixa Orellana* which furnishes the Annatto used to color butter, silk, etc. In the *Camellaceae*, is *Canella alba*, from which is obtained the Canella bark of commerce, often called wild cinnamon and which is used as a condiment in the West Indies and Florida where it is a native.

The family *Flacourtiaceae* of the tropics includes a number of plants like *Gynocardia odorata* of India, *Hydnocarpus renemato* and the *Kayalaris africana* of Ceylon, the seeds of which contain hydroxyanic acid. The cocon oil is obtained from *Myrsine* and is used in perfumery. The *Casuarina esculenta* of Asia and Australia is a purgative. The family *Turneraceae* contains a few medicinal

plants. A variety of *Turnera diffusa* furnishes a laxative. The *Datisca cannabina* of Southern Europe belongs to the family *Datisacaceae* and furnishes the yellow coloring matter *datiscin* used to dye silk, some members of the family *Dilleniaceae* are cultivated for their beautiful flowers. The *Marcgraviaceae* are occasionally cultivated and the *Marcgravia umbellata* is used in medicine the family also contains the "Bitter-sick tree" *Datisca glomerata* which is used, according to Chesnut, by the Indians of California to poison trout.

Families of Parietales

Calyx gamiosepalous.

Throat of the calyx with a fringed crown

Passifloraceae

Throat of calyx without a crown

Loasaceae

Calyx with sepals, sepals generally distinct and persistent

Flowers irregular

Violaceae

Flowers regular

Trees or shrubs, leaves alternate

Theaceae.

Herbs or shrubs, leaves opposite or whorled

Hypericaceae.

THEACEAE Tea Family

Trees or shrubs with alternate simple leaves without stipules; flowers large, showy, regular hypogynous, mostly axillary, sepals 5, or rarely more, often with 2 bracts petals 5 or rarely more stamens more or less monadelphous; anthers 2-celled; ovary 2-5-celled ovules 2 or more in each cell, fruit a woody capsule embryo large, endosperm scant. A small order of 160 species, mostly native of warm regions. The tea plant (*Camellia Thea*) is native to Assam, and extensively cultivated in Japan, China, and India. The *Thea sinensis* with the varieties *viridis* and *holeris* furnish tea. Successful attempts at cultivation have also been carried on in South Carolina. The black and green teas come from the same species. Tea is an important article of commerce in all civilized countries, Russia, England and the United States using large quantities.

The active principle found in tea is *caffein* or *thein* $C_8H_{10}N_4O_2$, a feebly alkaline, bitter alkaloid which is a cerebral and cardiac stimulant. Thea also contains the alkaloid *caffeidin* $C_7H_{12}N_2O$, *theophyllin* $C_7H_8N_4O_2 + H_2O$, *theobromin* $C_7H_8N_4O_2$, which is a bitter alkaloid having a physiological action similar to that of *thein*. The principle alkaloid *thein*, or *caffein*, as it is known, is a feebly basic, proximate substance, obtained from the tea plant, from the dried seeds of coffee, and from some other plants. This alkaloid has no particular action upon the digestive tract, unless it is used in large quantities, when it may cause gastro intestinal irritation. *Caffein* increases the blood pressure causing the heart to beat more forcibly and rapidly. It is a certain and direct stimulant. It produces wakefulness and restlessness and stimulates the reasoning and imaginative faculties in man. In the lower animals, according to Dr Winslow, it often causes "the most intense cerebral excitement and mania when given in large doses,—produces restlessness, increased reflex excitability and convulsions in the lower animals." Dr Winslow states further that *caffein* is a spinal and muscle poison to the frog, and tetanic convulsions occur in the batrachian similar to those produced by *strychnin*, but there is also muscular rigidity. The symptoms of poisoning in dogs, cats, and mammals generally, are restlessness, occasionally vomiting in dogs, rapid breathing, primary reduction followed by rise in temperature, clonic or tonic convulsions, muscular weakness, and

general paresis. Tea contains some saponin, but it is found especially in *Thea Sasanqua* of China and Japan and a saponin-like substance called *assamin* occurs in *T. assamica*, the flowers of the former species are used to flavor tea. The flowers and leaves of *T. Kuhn* are used as an insecticide.

The *Camellia* (*C. Hongkongensis*) is a well known evergreen shrub cultivated in green houses. Two east North American genera of the order are *Sinaria* and *Gordonia*.

The leaves of many plants are used as substitutes for tea, among them are Gromwell (*Lithospermum officinale*), Willow herb (*Epilobium angustifolium*), Willow (*Salix*), Ash (*Fraxinus* sp.), European Mountain Ash (*Pyrus Aucuparia*) Mulberry (*Morus alba* and *nigra*), Coffee (*Coffea arabica*), Camellia (*Camellia hongkongensis*), Cherry (*Prunus spinosa*), Rose (*Rosa canina*) Strawberry (*Fragaria vesca* and *virginiana*), Meadow Sweet (*Thalictrum flavum*), Wistaria (*W. chinensis*), Hydrangea (*H. Hortensia*), Boxelder (*Aegundo aceroides*), Oak (*Quercus*), Alchua (*A. quivota*), Blueberry (*Vaccinium Myrtillus*). In Russia the leaves of *Vaccinium* and *Arctostaphylos* are made into tea. In North America, in addition to plants previously named the leaves of Labrador tea (*Ledum latifolium*), New Jersey tea (*Ceanothus americanus*) Oswego tea (*Monarda didyma*) and Mexican tea (*Chenopodium ambrosioides*), are frequently used.

In South America the following are tea substitutes: *Lippia pseudospathulata*, *Stachytarpheta jamaicensis*, *Psoralea glandulosa*, *Myrtus* (qui), *Symplocos Alstonii*, *Capraria biflora*, *Angracum fragrans*, and *Frutichium guaphaloides*.

In China *Sageretia theezans* is used as a tea substitute, and in Australia various species of *Myrtaceae* are used for the same purpose.

HYPERICACEAE. St. John's-wort Family

Herbs or shrubs or occasionally small trees with opposite entire punctate, dotted leaves, without stipules. flowers solitary or cymose, paniculate, perfect and regular, sepals and petals 4 or 5, sepals persistent. stamens numerous, hypogynous in 3 or 5 sets. ovary 1-7-celled, and as many styles. pod 1-celled with 2-5 parietal placentae. seeds numerous. small endosperm absent. About 275 species, some cultivated for ornamental purposes. Kalm St. John's wort (*H. Kalmianum*) of the lake region, shrubby St. John's-wort (*H. prolificum*) and the great St. John's wort (*H. Ascyron*) are desirable species for ornamental planting. The *H. perforatum* and other species yield a yellow dye. The *Vismia viridiflora* of Guiana yields a resin called American gamboge.

Hypericum (Tourn.) L. St. John's wort

Herbs or shrubs with opposite punctate leaves, flowers borne in cymose clusters, sepals 5, slightly unequal petals 5, yellow. stamens numerous, distinct, or somewhat united in sets, fruit a capsule 1 to 5-celled. seeds numerous. About 200 species, of wide distribution.

Hypericum perforatum L. Common St. John's-wort

A perennial, much branched herb, with numerous sterile shoots at the base. leaves sessile, oblong or linear, black, punctate dots, flowers borne in cymose clusters, yellow sepals shorter than the yellow and black-dotted petals. stamens many in 3-5 clusters.

Distribution. Common in the eastern states, rare west of the Mississippi and in the southern states.

Hypericum punctatum Lam. Spotted St. John's-Wort

Herbaceous, perennial, 1 or more feet high, copiously marked with black, pellucid dots, leaves sessile, oblong or ovate-lanceolate, cymes terminal, many-flowered, flowers crowded, petals pale yellow, large, longer than the oblong sepals, styles mostly not longer than the pods.

Distribution. In moist soil, Maine and Ontario to Minn., Florida, Kansas and Texas.



Fig. 354. St. John's-wort (*Hypericum perforatum*). Flowering branch, fruit, section of fruit, part of leaf. Contains an oil and an acrid resin. (From Vauquelin's *Traité de Botanique*.)

Hypericum Ascyron L. Great St. John's-wort

Large stems, from 2-5 feet high, branched, 2-4-angled, leaves oblong, partly clasping petals narrowly obovate, cymes terminal, few-flowered, flowers large, bright yellow, stamens in 5 sets, styles 5, united below, capsules ovoid 3-celled, seeds small, numerous.

Distribution. From Vermont, Canada and Manitoba to Kansas, Illinois, eastward, also found in Europe and Asia.

Poisonous properties. It is believed that, where the plant is common, it is poisonous. It is said to cause eruptions on cows' udders and on the feet of white-haired animals. A writer in *Breeders' Gazette* reports the former species as poisonous. Prof. Chesnut says:

This species and the spotted St. John's-wort (*H. maculatum*, *H. punctatum*), were brought into the Department by Dr. G. W. Bready, from Norwood, Md., who stated that five horses were poisoned in May, 1898, by eating meadow hay which contained nearly 50 per cent of these plants. One horse died from the effects of the poison, and two were killed to prevent their further suffering.

The oil of *Hypericum* is obtained from the European St. John's-wort (*H. perforatum*) and is apparently found in the black dots of the petals and fruits. The resin, however, found in the plant, is acrid and slightly bitter. The physiological action of the St. John's-wort is mental depression and exhaustion; vertigo and confusion of the head; dilation of the pupils, and increased heart action.



FIG. 135 Great St. John's-wort (*Hypericum decurrens*). (Charlotte M. King.)

VIOLACEAE. Violet Family

Usually herbs or rarely shrubs or trees, caulescent or acaulescent, with alternate, simple, entire or lobed leaves with stipules, flowers mostly irregular sepals 5; corolla of 5 petals, 1-spurred, hypogynous; stamens 5, short, filaments broad and flat, often cohering with each other around the pistil; ovary simple, 1-celled, with 2 parietal placentae, fruit a capsule; seeds anatropous. About 300 species, of wide distribution. The best known is the pansy (*Viola tricolor*) running into numerous varieties, native to the Old World. It is sometimes used in skin diseases. The fragrant violet (*V. odorata*) is also frequently cultivated. Some of our native species are very handsome. Among these are the bird-foot violet (*V. pedata*), common in gravelly soils from Maine to Florida, west to Minnesota and Iowa and the *V. pedatifida* from Illinois to Kansas and Minnesota. Our most common eastern yellow violet is *V. pubescens*, the *V. Nuttallii*, also yellow, occurs from central Kansas westward. The white Canadian violet (*V. canadensis*) is common in the north and in the mountains. *V. striata* is common from the New England states to Minnesota and Missouri. The white-flowered violet (*V. blanda*) occurs in swamps. Brazilian species is



Fig. 86b. Pansy Violet (*Viola tricolor*). Possesses emetic qualities. (After Fitch.)



Fig. 86c. 14. Yellow violet (*Viola pubescens*). b. Spurred petals. c-c, Pansy c-taxis. d, Pistil. a, Antherapur.

derived from *Hybanthus sphenoloba*. The sweet pulp of the fruit of the *Leonis glycyrrhiza* of Peru is eaten by the natives, the fruit being about the size of a peach. We shall describe only a single species characteristic of the order—the common pansy—which is a common weed in the South.

Viola L.

Acaulescent or leafy stemmed herbs, annual or perennial, flowers solitary or rarely 2, smaller than the cultivated pansy, sepals 5, petals 5, unequal, the lower spreading at the base, stamens 5, anthers erect, united in many of the species early blossoms are conspicuous, the later being cleistogamous.

Viola tricolor L. Heart's-ease

Plant usually smooth, stem angled, branched, leaves round above or the lower oval, often heart-shaped, petals variable in color or variegated, yellow, whitish, violet-blue, and purple.

Distribution. Common in dry or sandy soil from New England to Kansas, especially southward. Also common on the Pacific Coast.

Poisonous properties. The substance *violins*, an acid bitter principle, has been extracted from the above species, and also is found in some of our native varieties. It is a pale yellow, bitter powder. The substance *violaquercetin* $C_{42}H_{42}O_{24}$ is a coloring matter which is found in the pansy. The emetic effect of violets is well known and is supposed to be due to the presence of *violins*, Dr. Milspaugh states.

The most characteristic symptom of its action is an offensive odor of the urine, like that of the cat. The pains caused by this drug are of a stitching character, while its action seems spent almost entirely upon the skin, and the male sexual organs. On the skin it causes burning, stinging, and itching, followed by breaking down of the tissues into either squamous spots, or any grade of incrustated eruptions, the eruption pours out a thin yellow fluid.

Dr. Schaffner states that the sweet violet is somewhat poisonous, the underground parts being emetic and cathartic.

Dr Rusby says

Many violets are noted for their spasmic properties, yielding a glucoside called *violin* long confused with the *emetin* yielded by ipecac. They may be classed among the emetico-cathartics, and a large quantity might easily be productive of serious results to a child.

According to Spatzier, the seeds of violet contain *myrosin* and a glucoside.

PASSIFLORACEAE. Passion Flower Family

Shrubs or herbs climbing by axillary tendrils, leaves alternate, simple, generally 3-lobed, flowers perfect, regular, axillary, calyx tube persistent petals usually 5, inserted on the throat of the calyx tube, which is fringed with a crown of a double or triple row of long, slender fringe, stamens 5, monadelphous, enclosing the style of the ovary, pistil 1, ovary with 3-5 parietal placentas, styles 1-5, fruit a berry or capsule, usually many seeded.

A small family of about 300 species of warm and tropical regions. The common blue passion flower (*Passiflora caerulea*) of South America is frequently cultivated. The *P. edulis*, native of the West Indies, about the size of a hen's egg, is eaten, the grandilla (*P. quadrangularis*) producing a fruit 6 inches long, often weighs 3 pounds. The common maypop of the South (*P. incarnata*) with a fruit about the size of a hen's egg, is eaten. This species and the *P. lutea* are sometimes weedy. The Tacsonia yields *hydrocyanic* acid.

LOASACEAE. Mentzelia Family

Herbs with rough, often stinging hairs, leaves without stipules. flower regular perfect, whitish, yellow or reddish, calyx tube adherent to the ovary, lobes 4-5, petals 4-5, inserted on the calyx, stamens numerous pistil 1 usually 1 celled, with 2 or 3 parietal placentae, fruit a capsule, 1 celled with the persistent lobes of the calyx, endosperm scant.

About 200 species, nearly all native to North America. The *Barbarea*, however, being found in Africa. Species of the genus *Blumenbarhia* (*B. grandiflora*) produce pretty flowers and are cultivated in greenhouses. The *Mentzelia graveolens*, from Mexico and Texas, is also cultivated. The leaves of *Mentzelia ornata* and other species produce hooked hairs which are often annoying to man.

Mentzelia (Plummer) L. *Mentzelia*

Herbs with erect stems, alternate leaves with barbed hairs. flowers usually showy, terminal, solitary or clustered, calyx tube cylindrical or club-shaped lobes persistent petals 5-10, regular, spreading, falling, usually turning black in drying, stamens numerous, inserted on the throat of the calyx tube, styles 3, more or less united capsule dehiscent at the summit, many seeded seeds flat. About 50 species. American, chiefly west of the Mississippi river.

Mentzelia ornata Torr and Gray. Showy *Mentzelia*

A rough herb from 1-2 feet high, leaves oblong-lanceolate, deeply toothed or pinnatifid, flowers solitary, much larger than the lanceolate calyx lobes petals 10, yellowish-white, 2-3 inches long capsule 1½-2 inches long, seeds numerous, with narrow margin.

Distribution. Northwestern Iowa to Dakotas to Central Kansas and Texas.

Injurious properties. The hooked hairs of the plant cause the leaves to stick to sheep, clothing, etc. Prof Goodale of Cambridge Mass., is quoted by

Dr. White, in his *Dermatitis Venenata* is saying: "Mentzelia has grown in our garden, and has always been amazingly irritating to us all. Some species are even said to have stinging hairs." The writer has had considerable experience in collecting these species in the West, and he has learned from experience that the barbed hairs are quite irritating. Dr. Halsted states that *M. oligosperma* has the same properties. This plant is frequently cultivated.

MENTZELIA

Fleshy plants, usually spiny with jointed stems, leaves small, flowers mostly solitary regular, calyx tube adnate to the ovary with a many lobed limb, stamens numerous, inserted on the throat of the calyx, filaments filiform ovary 1-celled, ovules numerous, fruit a berry. Contains the important family *Cactaceae*.



FIG. 357 Showy *Mentzelia* (*Mentzelia albicaulis*). The hispid hairs of this plant produce mechanical irritation. (Charlotte M. King.)

CACTACEAE Cactus Family

Fleshy plants, leafless or with small leaves, stems flattened, columnar or globular, generally abundantly spiny, flowers solitary, sessile, perfect calyx tube adnate to the ovary, limb many lobed, petals numerous, imbricated in several rows, mostly distinct, ovary 1-celled; ovules numerous, anatropous, borne on several parietal placentae, fruit a 1-celled berry or a dry fruit, endosperm wanting or copious.

About 1000 species, chiefly in the regions west of the Missouri river. Many species occur in the arid regions of the southwest. One of the most interesting is the giant cactus (*Cereus giganteus*) of Arizona, with fluted columns 50 or 60 feet high. The common night blooming cereus (*C. grandiflorus*), with white flowers opening at night, is well known in cultivation. The most commonly cultivated species is *C. speciosissimus*, with crimson red flowers that open during the day. The old man cactus (*C. senilis*) is cultivated because of its long white hanging hairs. Species of the genus *Echinocactus*, with stem of many ribs, are often cultivated, the most common being *E. texensis* of southern Texas and Arizona and the *E. Ottoma* of Brazil. A very good quality of leather has been produced from *E. Wislizeni* and *Cereus giganteus*. Species of *Mammillaria* with tufted stems covered with nipple shaped tubercles, are often found in cultivation. The *Euphybia truncatum* from Brazil, with flattened, leaf-like stems and flowers 2-3 inches long, and the *Phyllocactus*, native to South America and Mexico, are also cultivated. The large genus *Opuntia*, of over 150 species, is entirely American. The *O. vulgaris*, Mill, naturalized in southern Europe, extends from southern New England west to 1 south. *O. Rafinesquii* extends from Michigan west and *O. polyacantha* to Wisconsin westward. Several Mexican species are cultivated. The Indian pear or prickly pear (*O. Ficus-Indica*) of the West Indies and South America, produces an edible fruit. Extensively naturalized in North America, South Europe and Asia and as hedge plants. It grows on the lava slopes of Mount Vesuvius, converting the lava into soil. The cochineal plant (*Nopelia coccinellifera*) a native of Mexico, is cultivated as the host of the cochineal insect, from which a warlet carmine dye is obtained. *O. Tuna* (*O. Dillenii*) and *Pereskia* also act as hosts of the same insect. The fleshy stem of some *Opuntias*, after the spines are removed, are used as work food. The berries of some species like *Opuntia Larrea* which belongs to the *O. Ficus Indica* group (*O. Stupida*antha and *Echinocereus stramineus* etc.), are eaten. A recent paper by Hare and Griffith described many details of their uses. The Mexicans call them tunas. An alcoholic drink is also made from the fruit. Some species are used as soil binders in Texas and elsewhere.

Injurious properties. The barbed trichomes penetrate the flesh and are difficult to remove. Death of animals has been caused by "hair balls" phyto-bezoars from them being formed in the stomach. Dr. William Trelease, who had occasion to examine the "hair balls" produced in some Mexican animal, gives the following account of this phyto-bezoar.

"The hair balls were a little over 3½ inches in diameter and weighed 7½ ounces. One ball was probably 4 inches in diameter." It was stated by the physician who sent them to Dr. Trelease that 16 such balls had been taken from the stomach of a bull at the Hacienda de Cruzes, it appears that the chief food of cattle at this time of the year consisted of *Opuntias*, and that the particular animal in question being allowed to roam at large, sought such food as could be found. These phyto-bezoars were brown in color, "and in appearance somewhat suggest felt or rubbed sole leather, and on examination prove to be composed, aside from the small nucleus at the center, of the barbed hairs with which the pulvini of the *Platopuntias* are armed. To the barbs with which these hairs are covered is due their power of felting together, and there is every indication that, starting about some small nucleus of vegetable fiber, they have been compacted into the dense, felty texture by the visceral movements of the

anima, to which, causing friction against one another, their perfectly round form is attributable." As is well known, the *Opuntias* produce spines and two kinds of trichomes. In the *Cylindropuntias*, each spine is invested by a deciduous sheath, "which is downwardly barbed, so that a person or animal brushing carelessly against a plant is certain to remove some of the barbed sheaths." In the *Platopuntias*, to which the ordinary flat-stemmed prickly pears belong, the spines, when present, are destitute of such a sheath. The protection to the plant is afforded simply because of their rigidity and pungency. The spines have their origin in pulvini, and in this particular genus of cacti are coated with delicate flexible hairs, divided into partitions. These hairs are lightly attached to the epidermis of the plant, so that when the pulvino is touched they are almost certain to be removed in considerable numbers. The points of the stiffer hairs penetrate the skin, the barbs with which they are closely beset preventing their ready withdrawal. Dr. Trelease, in summing up the injurious effect of cacti, says:

It is a frequent practice in Texas to cut the branches of cacti which are fed to stock into half-inch lengths. In this way every one of the obliquely set (singer) spines of *Opuntia Engelmanni* (and of some other species which are so used) is almost certain to be cut off so that the danger from the spines is removed. This treatment, however, does not destroy the barbed hairs of the pulvini, of which the bezoars under consideration are composed. It is also the practice, in some places, to roast the fragments as a means of completely removing the spines and barbed hairs, but this is objected to by some feeders, because the roasting has been ascribed to add to the laxative properties of the cactus. Where some such treatment has not




Fig. 35B. Prickly Pear (*Opuntia Engelmanni*), from the barbed trichomes of which phytobezoars are sometimes formed. (U. S. Dept. Agr.)

been resorted to, injury to the animals not infrequently results and in the bulletin referred to, Dr. Vasey gives a number of instances in which cattle have died from an accumulation of spines in the mouth and stomach, an effect somewhat comparable with that caused by the awns of *Hordenia* when cattle feed upon these.

The eating of the fruit of some species of *Opuntia* produces diarrhoea.

Under the name of Pellate (*Anhalonium* sp.), the Indians of the Rio Grande Valley of Mexico have for ages used the tops of this plant which they commonly call "mesca, button" or "mescal bean." The use has extended to Indians in Oklahoma and Indian Territory and, it is said, to the Tama Indians of Iowa. The Kiowa Indians use 14-15 grams (4-5 buttons) to produce the peculiar sensations. The so-called mescal beans are 1 1/2 inches long and about 1/4 inch in diameter, brittle when dry but soft when moistened. They have a bitter, disagreeable taste. Prentiss and Morgan were the first to call attention to the character of the drug dried from *Anhalonium*. During intoxication, the pupils become dilated, there is muscular relaxation, the pulse is somewhat slower, there is loss of sense of time, partial anaesthesia, weakened heart action in some nausea and vomiting and wakefulness. In man the influence has been described as causing an incessant flow of visions of infinite beauty grandness, and variety of color and form. Intoxication closely resembles that produced by *Cannabis indica*. Dr. Lewis found that an aqueous extract given to lower animals produced convulsions causing death by respiratory failure.

The *A. Lewisii* contains the alkaloid *anhalonium* $C_{11}H_{17}NO$ *mesconin* $C_{11}H_{17}NO_2$ and *anhalonidin* $C_{11}H_{17}NO_2$. The *A. fissuratum* contains *pellatin*. Although this substance according to some authors, is inactive, it has been used as a calumative on insane patients and in many cases causes sleep to come on. The cactus found in some species is a cardiac stimulant. It appears also that in addition to the above species the same or a few substances occur in *A. prismaticum*, *A. tomentosum* and *A. Jordanianum*.

Anhalonium is closely related to the genus *Cactus*. It bears a dense penicillate tuft of long soft hairs which persist above the apical region of the plant as matted wool.

According to several recent investigators especially Hawley and Daley, alkaloids seem to be widely present in the family *Cactaceae*. *Pectenin* is found in a species of *Cereus*, *purpureus* $C_{11}H_{17}NO$ occurs in *Pitocereus Sargentianus*, the alkaloid *pellatin* $C_{11}H_{17}NO_2$ is found in species of *Anhalonium* and *Lophophorum* occurs in *Anhalonium Lewisii* and allied species. The alkaloidal substances appear to the extent of 11 per cent in dried material. *A. Lewisii* is a cardiac and respiratory stimulant. *Saponin* also is found in several species of the family among them in *Cereus quinquatus*. Quite a number of other species of the family are used in medicine. The night-blooming cereus (*Cereus grandiflorus*) contains a glucoside which acts much like *Digitalis*. The *Opuntia Karwinskiana* contains an astringent principle. Several species such as *Rhipsalis* and *Opuntia* have anthelmintic properties.

MYRTIFLORAE.

Mostly shrubs or trees. Leaves simple. Flowers incomplete, calyx inferior, 4-5 lobed or entire, corolla usually wanting, stamens twice as many as the calyx lobes or fewer, ovary 1-celled; ovule 1.

Species of the genus *Cuphea* of the family *Lythraceae* are cultivated in the South as border plants. the cape myrtle (*Lagerstroemia indica*) of the East

Indica, of the same family, is a handsome ornamental shrub, the seeds of which contain a narcotic principle; Henna (*Lawsonia inermis*) yields a yellow dye which is used in Egypt and Arabia as a cosmetic for the hands.

Tansu occurs in the root of *Lythrum Salicaria*. *Cuphea viscosa* of Mexico contains a substance similar in action to *Digitalis*. The fresh leaves of *Ammanium baccifera* of India contain a vesicating substance.

The family *Lecythidaceae* contains the well known Brazil or Para nut (*Bertholletia excelsa*, and *B. nobilis*), the Sapucaya-nut from the snake-pot tree (*Lecythus allaria*) and other species.

Narcotic and poisonous principles occur in the fruit of *Planchonia lida*, native to the Molucca Islands, and in the seeds of some species of *Lecythus*. The roots and fruits of Chinese and Japanese species of *Borringtonia* are used as fish poisons. The family *Punicaceae* contains the pomegranate (*Punica granatum*) from the orient, cultivated in green houses in the North and out of doors in California and the South for its acid fruit, which is about the size of a small apple. The bark is used as a vermifuge and is an active irritant, its medicinal properties being due to a number of alkaloids present, of which four



Fig. 359. Clove Tree (*Eugenia caryophyllata*) native of the Molucca Isles. (After Faguet.)



Fig. 360. Brazil Nut (*Bertholletia excelsa*) (W. S. Dudgeon.)

have been isolated. The Mangrove (*Rhizophora Mangle*) belongs to the family *Rhizophoraceae*. Its wood is used in boat building. The Indian Almond (*Terminalia Catappa*) through its bark and leaves furnishes a black dye. The family

Myrtaceae contains the guava (*Psidium Guajava*), a most delicious fruit from which is made a jelly that is considered a great delicacy, the bay-berry (*Amomum caryophyllata*), from the dried leaves of which, when placed in water, an essential oil may be distilled which is used in the medication of rum, rose apple (*Jambosa malaccensis*) which yields an edible fruit and furnishes a materia suitable for baskets, hoops, sugar casks, etc., Jambuse berries (*Jambosa vulgaris*) cultivated in the tropics, the spices, piment or allspice, (*Pimenta officinalis*) and cloves, (*Eugenia caryophyllata*) from the Moluccas, common articles of commerce. Cloves contain an essential oil which is an excellent antiseptic and contains *caryophyllin*, $C_{15}H_{18}O$, and *eugenin*, $C_{15}H_{12}O_2$. Cajaput, (*Melaleuca leucodendron*) is an excellent stimulant containing *cajaputol* $C_{15}H_{18}H_2O$



Fig. 361. Buffalo Berry (*Shepherdia argentea*). Fruit used for jellies, etc. (From American Agriculturist.)

The oil of *Myrcia* is obtained from the leaves of *Pimenta acris* of the West Indies and contains *eugenol* and *chavicol*. The volatile oil of Cheken (*Eugenia Chequen*) contains *cineol*, *pinene*, a volatile alkaloid, and a glucoside. The leaves of Myrtle (*Myrtus communis*) yield a substance used in perfumery. The Eucalypts, natives to Australia, are important timber plants, producing valuable wood for interior finishing, furniture, parts of vehicles, etc. A few species are extensively cultivated in California, of which *Eucalyptus globulus* is one of the most frequently seen. The oil of *eucalyptus* and *eucalyptol* are used in medicine, and have valuable antiseptic properties. The *E. rostrata* produces Kino. The family *Combretaceae* contains many plants rich in tannin like the bark of *Terminalia Catappa* of Asia. The fruit of Myrabalans (*T. Chebula*) is rich in tannin. The *T. fagifolia* of Brazil has cathartic properties. Some of the plants of this family are used in arrow poisons. Several plants of the family *Melastomaceae* produce edible fruits, some of them like *Tocora guianensis* yield a black dye. The only indigenous genus in the Northern States is *Rhexia*. The family *Trapaceae* contains the edible Chinese water-nut or water chestnut (*Trapa natans*) naturalized in New England and New York. The family *Haloragchidaceae* contains several water plants of common occurrence like the water milfoil (*Myriophyllum spicatum*) the mare's tail (*Sagittaria vulgaris*) and the mermaid-weed (*Proserpinaca palustris*)

Shrubs.

KEY TO FAMILIES OF MYRTIFLOREAE

Leaves green, seeds pendulous	Thymelaeaceae
Leaves silvery, scurfy, seed erect	Elaeagnaceae
Herbs or rarely shrubs, calyx tube almost wholly adnate	Onagraceae

ELAEGNACEAE Oleaster Family.

Shrubs or small trees, leaves silvery, scurfy; flowers perfect or dioecious, calyx regular, simple, colored, calyx tube becoming pulpy and berry-like in fruit, strictly enclosing the achene, seed erect, ascending. A small order of 20



Fig. 367 Sea Buckthorn
(*Hippophae rhamnoides*). Pro-
duces a berry (After Fitch.)

species and 3 genera. The buffalo berry is well known. The *Shepherdia argentea* is a thorny shrub from 5-18 feet high, from Western Iowa and westward, the acid fruit of which was much used in early days for jams, jellies and pies. The Russian oleaster or wild olive (*Elaeagnus angustifolia*), a well known ornamental shrub, adapted especially to the North West, is hardy and handsome. It has spiny branches which bear fragrant flowers. The wood is durable and makes an excellent post. The Goumi (*E. multiflora*) of Japan, produces edible fruit. The *E. hortensis* formerly included two species, the *E. angustifolia* and *E. orientalis*. Prof Hansen has introduced the edible form of the shrub into the Northwest. The Arabs dry the berries and make a kind of cake. The *Hippophae rhamnoides*, an ornamental plant from Europe, is known under the name of sea buckthorn. It is commonly used for fish sauce in Russia. The plant is hardy in South Dakota. Prof Hansen says that the berries of the tree contain a narcotic poison which is eliminated by boiling.



Fig. 363. Goumi (*Elaeagnus multiflora*)
Cultivated for its edible fruit. (From American Agriculturist.)

THYMELAEACEAE. Mezereum Family.

Shrubs or small trees with acrid, tough, fibrous bark, simple opposite entire leaves, flowers in spikes or umbels, regular, calyx petal-like, tube, urn-shaped, petals present or absent, stamens twice as many as the lobes of the calyx, and borne on it, ovary free, 1-celled and 1-ovuled, fruit a berry like drupe, embryo straight, endosperm scanty or none.

About 400 species of wide distribution, most largely represented in Australia and South Africa. The leatherwood or moosewood (*Dirca palustris*) with a tough fibrous bark, is used by the Indians for thongs. The Mezereum (*Daphne Mezereum*) with fragrant flowers and bright red berries, of Europe naturalized from Europe in New England, and the handsome *D. Genkwa*, with rosepink flowers, are cultivated. The berries and leaves of the Mezereum cause blistering, it is an acrid poison. Bark paper (*D. cannabina*) native of the Himalayas to Japan, produces a tough bark which is made into paper. Lace-bark (*Lagerflora*) of Jamaica, with bark that separates into layers, was formerly used for veils, bonnets, etc. The bark of *Wikstroemia viridiflora*, of the Polynesian Islands, is used for making fishing nets, ropes, etc. The bark of *Punifera* of Brazil, causes vesication like that produced by the *Dirca palustris*. Several exotic plants of the family are poisonous like the *Pimelia trichostachya* of Australia. The fruit and leaves of *Gordia coccinea* are emetic.

Daphne. L. Laurel

Shrubs, with alternate leaves, and small purple, pink, or white flowers in fascicles heads or racemes. perianth tubular with 4 spreading lobes; stamens 8, attached to the calyx tube, filaments very short, disk none, ovary sessile, stigma large, calyx deciduous or persistent. About 40 species, native of Europe and Asia.

Daphne Mezereum L. Spurge Laurel. Lady Laurel.

A small shrub with young twigs somewhat pubescent. leaves thin, oblong-lanceolate or oblanceolate, petioled. flowers in sessile fascicles, very fragrant; perianth tube pubescent, rose-purple. drupe red.

Distribution. Escaped from cultivation from Quebec to New York native to Europe and Asia. frequently cultivated as an ornamental plant.

Poisonous properties. Some of the European species, like *Daphne Genkwa* contain acrid poisons. The plant produces blisters. The bark is used internally and in the form of an ointment. According to Loudon, in France the bark is applied to the skin for the purposes of a "perpetual blister." The bark when fresh or when soaked in water reddens the skin, when applied to it and at length, occasions vesicles followed by ulcers. Oesterlen remarked that all parts of the plant produced on contact, irritation and inflammation. Schimpfky mentions this among the twenty-six important poisonous plants of Europe and states that the bark and berries are most poisonous, and that the pleasant odor of the flowers produces headache, for which reason, therefore, they should not be placed in a living room. Linnaeus seems to have recorded cases of poisoning from this plant. *Daphne* contains the glucoside *daphnin* $C_{20}H_{24}O_{10}$, bitter and astringent, an acrid resin *mezereum*, *daphnetin* $(C_8H_8O_4)H_2O$, also with an astringent taste, *coccognin* $C_{20}H_{27}O_8$, and the glucoside *aesculin* $C_{21}H_{30}O_{11} + H_2O$. Friedberger and Fröhner state that animals poisoned by the *Daphne* have stomatitis, salivary colic and a feeble pulse. In Europe the fruit is

sometimes used as a substitute for pepper, in some cases with fatal results. Blyth says "There are a few cases of poisoning on record, and they have been mostly from the berries. Thus, Lanne has recorded an instance in which a little girl died after eating twelve berries. The symptoms observed in the recorded cases have been burning in the mouth, gastro-enteritis, vomiting (giddiness, narcosis, and convulsions, ending in death. The lethal dose for a horse is about 30 grms. of powdered bark, for a dog, the oesophagus being tied, 12 grms., but smaller doses of the fresh leaves may be deadly."



Fig. 364. *Mezereum* (*Daphne mezereum*). Ac. acid. (after Pich.)

Dirca, L.

A small shrub with tough, fibrous bark, short petioled leaves. flowers yellowish, in peduncled fascicles of 2-4 scaly bracts at the nodes or twigs of the preceding season. stamens 8, borne on the calyx, the alternate ones longer. filaments very slender, perianth bell-shaped, or funnel-form. disk obsolete. vary nearly sessile. corolla red, oval, oblong. 2 species known. 1 in Eastern North America, and 1 in California.

Dirca palustris, L. Leather-wood

A shrub with yellowish green twigs. leaves obtuse, bracts 3 or 4 oval, with brown hairs. deciduous. style longer than the stamens.

Distribution. In woods and thickets, Eastern Canada to Minnesota. Central Iowa to Missouri and Florida.

Poisonous properties. The bark is acrid, like that of the *Daphne*. all parts of the plant having a nauseous, acrid taste. The principle, however, is unknown. The fresh bark applied to the skin causes redness and vesication, the sores thus produced being quite difficult to heal.

ONAGRACEAE. Evening Primrose Family

Herbs or rarely shrubs, with alternate or opposite leaves, generally without stipules, or stipules glandular. calyx adnate to the 2-4-celled ovary. petals 2-4. stamens as many as the petals or twice as many, ovules numerous. About 300

species. A few of the plants are medicinal. The great willow herb (*Epilobium angustifolium*) is occasionally used in medicine. The hairs of the seeds of some species are used in the Arctic as lamp wicks. Many species of the family are used for ornamental purposes, especially some of the western species of the genus, *Oenothera*, the *Clarkia elegans* of the gardens and the greenhouse *Fuchsia*. The genus *Oenothera* with many species, some southwestern and some western, contains very pretty plants.



Fig. 365. Leather wood (*Dirca palustris*). This plant is well known in northern woods, especially on the banks of streams; occasionally found on high land. The bark is very tough and regarded as poisonous. (Charlotte M. King.)

Gaura, L. *Gaura*.

Annual, biennial, or perennial herbs with alternate sessile leaves, flowers white, pink or red in spikes or racemes; calyx tube narrow, prolonged beyond the ovary, the limb usually 4-lobed, reflexed, petals clawed, unequal; stamens usually 8, with a small scale before the filament, frequently declined, ovary 4-celled; styles declined, fruit hard and nut-like, 3 to 4-ribbed and angled. About 18 species.

Gaura biennis, L. *Gaura*.

An erect, soft, hairy or downy annual or biennial, leaves lanceolate or oblong-lanceolate, denticulate, flowers in slender spikes, white, turning pink; fruit oval or oblong, acute at each end, 4-ribbed.

Distribution. From Ontario to Georgia, Arkansas and Nebraska and Minnesota.



Fig. 366 Willow herb (*Epilobium angustifolium*, Sw.) *Gaura angustifolia* or *Gaura angustifolia* (L.) Hitchc. (After Fitch.)

Gaura portiflora, Dougl.

A hairy, branching, soft pubescent annual from 2-3 feet high. leaves lanceolate or ovate lanceolate, acute or acuminate, sessile, repand, denticulate, covered with long soft hairs, the pinkish flowers about $\frac{1}{2}$ inch long, borne in long flexuose spikes 2-3 feet long. fruit contracted at the base, obtusely 4-angled, glabrous.

Distribution. Common in dry soil from South Dakota to Missouri, Louisiana, the Rocky Mountain region and New Mexico and Mexico. A common weed along irrigation ditches.

Gaura coccinea, Pursh. Scarlet Gaura.

An erect or ascending, much branched smooth or canescent herb, leaves lanceolate, linear-oblong, repand or entire, flowers red, turning scarlet. fruit canescent, terete below, and narrowed above.

Distribution. From Western Minnesota, Iowa, Nebraska and Texas to Utah, Arizona, and Mexico.

Poisonous properties. The *Gauras*, or at least one species, the *Gaura coccinea* have been suspected of being poisonous to live stock in the West. This is an excellent honey plant.

UMBELLALES

Herbs, shrubs or trees, flowers nearly always with petals, divisions of the calyx and petals usually 5, stamens 4 or 5, ovary compound inferior, adnate to the calyx, epigynous ovule 1 in each cavity.

FAMILIES OF UMBELLALES

Fruit a drupe or berry.

Flowers umbellate; stamens 5....

Araliaceae

Flowers not umbellate; stamens 4....

Cornaceae

Fruit dry splitting into 2 mericarps....

Umbelliferae

ARALIACEAE Gingseng Family.

Herbs, shrubs or rarely trees, leaves alternate or whorled, flowers in umbels, heads or panicles, calyx tube adherent to the ovary, usually 5 petals inserted on the calyx, stamens as many as the petals, inserted on the disk ovary 1 or more celled, 1 ovule in each cell, fruit a several-celled drupe.

About 50 genera and 450 species, of wide distribution. Genera common to eastern North America, China and Japan. Some of the species are occasionally cultivated for ornamental purposes. One of the best known of these is the Hercules Club (*Fatsia horrida*), native from Florida west to Missouri and Texas, and the common European ivy (*Hedera Helix*) well known in cultivation.

Few of the *Araliaceae* have injurious properties, however, the prickly spines of *Fatsia horrida* of the Pacific Coast, are quite irritating.

Several species of the genus *Aralia* and *Panax* are used in medicine. The



Fig. 367. Ginseng. (After Vaguet.)

most important of these is the Ginseng, (*Panax quinquefolium*), which is native from eastern Canada to Alabama and in woods from Kentucky to Iowa, Missouri, Nebraska and Minnesota. This species is now widely cultivated, large quantities of Ginseng being exported to China, where the roots are in great demand. The Chinese Ginseng is *P. ginseng*. Several other species are used in medicine, among them the Spikenard (*Aralia racemosa*), and Wild Sarsaparilla (*Aralia nudicaulis*). These plants are not official, but they are quite commonly used. They have aromatic and stimulating properties. *Barringtonia*, $C_{18}H_{30}O_7(OH)_1$, is found in *Barringtonia*, a Japanese *Aralia*, *Panax*, and other genera. *Aralin* occurs in the roots of *Palsia horrida*. The terpene *aralen*, $C_{15}H_{24}$, occurs in *Aralia nudicaulis*. Some members of this order are occasionally weedy, especially the *Sarsaparilla*. Rice paper is made from *Tetrapanax papyrifera* native to Formosa. It is a small tree about 10 feet high. The tree is cut into to obtain the pith, which is divided into thin slices and the paper cut with a sharp knife.

UMBELLIFERAE Carrot Family

Herbs with alternate compound or sometimes simple leaves, petioles often dilated at the base, rarely with stipules, flowers 5, small, in compound or simple umbels or heads, frequently polygamous, calyx tube adnate to the ovary, limb obsolete or 5-toothed, petals 5, inserted on the margin of calyx, stamens 5, inserted on the disk, pistils with 2 styles, fruit dry, composed of 2 carpels, generally spreading from each other at maturity.

About 1600 species of wide distribution in tropical and temperate regions. A number of the plants of the family are of economic importance, among them the Carrot (*Daucus Carota*), native to Europe, cultivated before the Christian Era. The thickened roots of the carrot are important as food for man and



Fig. 368. Turnip-rooted Celery or Celeriac (*Apium graveolens*). (W. S. Dudgeon.)

domestic animals. Parsley (*Carum Petroselinum*—*Petroselinum hortense*), native to the Mediterranean regions of Europe and Asia Minor, is used for garnishing. Celery (*Apium graveolens*) is indigenous to Great Britain and other European countries, and is found growing in low lands. There are two types, the turnip rooted, cooked and eaten as a salad, and the blanched leaf stalks. Celery was known to the ancient Greeks and Romans. It is said that in a wild state the plant is somewhat poisonous. Some people are known to be sensitive to the cultivated plant. Dill (*Anethum graveolens*) containing dill oil and caraway (*Carum Carvi*), were known to the ancients. The essential oil of caraway is obtained from the seed, which is used in Europe to flavor bread and meats and contains *carvot* $C_{10}H_{16}O$. Dill (*Pucedanum graveolens*) is commonly used in flavoring pickles and salads. Cumin seeds from *Cuminum zaitvum* resemble those of caraway in odor and taste.

Many members of the order have medicinal properties. Among the more important of these are the Indian pennywort (*Centella asiatica*), and poison hemlock (*Conium maculatum*) which contains the alkaloid *coniin* which is deadly poisonous. The caraway "seeds," fennel "seeds" (*Foeniculum vulgare*), the latter indigenous to the Caspian Sea regions, and yielding *anethol*, also contain *fenchone*, *trigonellin* and *choin*.

The anise seeds (*Pimpinella Anisum*), containing the oil of anise, are used in confectionery. *Asafoetida* (*Ferula Nothera*) native to Thibet and western Asia, obtained from the milky juice of this plant, is used in medicine, and by the Persians as a condiment. The bulton snake root (*Eryngium yuccaefolium*), the cow parsnip (*Hieracium lanatum*), and the water hemlock (*Cicuta maculata*), are common plants in the northern states. The leaves and roots of the fennel (*Foeniculum vulgare*), used in southern Europe as table vegetables, and in Germany to flavor bread and cakes, contain *phellandrene* $C_{10}H_{18}$ and *choycol*. Lovage (*Levisticum officinale*) is found in salt marshes along the Atlantic coast from Labrador to Connecticut and in Europe. The *Arracacia xanthorrhiza* of Peru is much used in the Andes region. The same species, known in Venezuela as *Arracacha* and introduced into Porto Rico, is said by Fairchild to be one of the most important of food plants to the poor. The roots are large and fusiform. The roots of sea holly (*Eryngium maritimum*) when candied, boiled or roasted resemble chestnuts in taste. Gum Galbanum is derived from *Ferula galbanifera* and was used by the ancients for incense and perfumery. It contains *cadimene* and *g-pinene*. This is referred to in Exodus 25:10. Coriander (*Coriandrum sativum*) contains *coriandrol* $C_{15}H_{24}O$. Musk root or Sambil (*Fernia Sambil*) contains *umbelliferone* $C_{11}H_{16}O_3$ and *angelic acid* $C_8H_8O_2$ used as an antispasmodic. The plant occurs in Asia. Sweet Cicely (*Osmorhiza longistylis*) yields an oil similar to anise and contains *anethol*. In the *Umbelliferae* the substance *puceadone* $C_{15}H_{24}O_6$ occurs in the roots of *Imperatoria Ostruthium*, and *Pucedanum officinale*. *othamontin* $C_{20}H_{32}O_7$ is found in *Pucedanum Orcaschium lascarpin*, $C_{21}H_{34}O_8$ is from roots of *Laserpitium latifolium*. *pimpinellin* is obtained from the roots of *Pimpinella Saxifraga*. *Oenanthe crocata* contains *oenanthotoxin* somewhat similar to *cicutoxin*. *thymol* is found in the fruit of a great many of the *Umbelliferae*, *cumin oil* is secreted from the fruits of *Cuminum* *Cuminum* and other plants of this order, and contains *cymene*. Anise seed or anise fruit contains anise oil which resembles that found in star anise, the *Oenanthe Phellandrium*.



Fig. 369. Coriander (*Coriandrum sativum*) Flowering stem. (After Vague)

contains *phellandrene* a native lovage, *Ligusticum canadense* is used to flavor tobacco.

This family contains a large number of plants with active principles, some of which are entirely harmless, but others must be considered among the deadly poisons. The water drop-wort (*Oenanthe crocata*), with its parsnip-like roots, and the *O. Phellandrium*, poisonous European plants, are *Umbelliferae*. Friedberger and Fröhner state that the former causes stomatitis and paralysis. Blyth

states that the chemistry of the plant has not yet been worked out, but that all parts are poisonous, the root especially deadly. Lehmann states, however, that the first species is not as poisonous as was formerly supposed, sheep and hogs eat it, although it is poisonous to horses. In the latter it is said to produce paralysis of the hind legs. *Berula erecta* of Europe and North America is poisonous, especially the root. It is a smooth aquatic perennial, with compound simple pinnate leaves leaflets linear oblong, serrate to cut-toothed, flowers white, and fruit globose.

In Australia according to Maiden the *Apium leptophyllum* when grown in damp soils is poisonous. The wild parsnip of that country is one of the most poisonous plants of Australia, no antidote to it being known. The *Chaerophyllum temulum* of Europe causes colic and stupor. The parsley is not ordinarily considered poisonous but is said to be injurious to birds. The gum resin *aspidiotum* found on the stem of *Dorema Ammoniacum* is acrid. The resin results from the sting of an insect. The genus *Ferula* from which Asafoetida is derived causes hæmaturia and bleeding at the nose.




Fig. 378. Creeping Water parsnip (*Berula erecta*). Very poisonous. (After Fitch.)

Genera of Umbelliferae

Flowers yellow.....	7	Pastinaca.
Flowers white or greenish.		
Fruit bristly, winged.....	9	Daucus.
Fruit, not bristly, winged.		
Fruit winged, dorsally flattened		
Flowers greenish	5	Angelica
Flowers white.		
Leaves pinnate or ternate, clustered, tuberous roots	6	Oxypolis.
Leaves ternately compound, root not tuberous	8	Heracleum
Fruit wingless flattened dorsally or laterally	4	Aethusa
Fruit ovoid or oval.		
Flowers white		
Biennial plant.....	1	Conium
Perennial, roots usually fascicled.		

Oil tubes solitary.....	2 Cicuta
Oil tubes 1-3.	3 Sium

Conium L. Hemlock

Smooth biennial herbs with spotted stems and pinnately compound leaves, flowers small, white, in compound umbels, calyx teeth obsolete, petals small, obcordate or entire, fruit glabrous, somewhat flattened laterally, carpels wavy-ribbed, oil tubes none, two species, one in Europe and Asia, the other African readily poisonous. Plant well known to the ancients.

Conium maculatum, L.

An erect, branching, smooth herb, with spotted stem and pinnately compound leaves. Flowers small, white, in compound umbels, calyx teeth obsolete, petals white, fruit smooth, ovate, flattened, with prominent wavy ribs. Oil tubes absent.

Distribution. In waste places, Canada to Indiana, California, Utah and Mexico. Native to Europe.

Poisonous properties. The plant is very poisonous. It was used by the ancients to poison criminals condemned to death, and it is said that Socrates was poisoned by it. The plant is avoided by stock because of its strong odor, but the dried plants are not so poisonous. The alkaloid *conium* $C_8H_{17}N$ is derived from it. *Conium* is volatile in vapor of alcohol or water, and somewhat volatile at ordinary temperatures. It has an alkaline reaction and burning taste and causes dilation of the pupil. Two other principal alkaloids occur namely *coniine* $C_8H_{15}N$ said to be 18 times more poisonous than *conium*, *conhydrin* $C_8H_{17}NO$, *pseudoronydrin* $C_8H_{17}NO$ and *methylaconin* $C_8H_{17}N$ occur in small amounts. The percent of *conium* in fresh leaves is 0.095. The ripe seed contains 0.7 percent.

Mr. Chesnut says. Recent cases of poisoning have arisen accidentally from eating the seed for that of anise, the leaves for parsley or the roots for parsnips, also from blowing whistles made from the hollow stems. It has recently been shown that some of the anise seed in both foreign and domestic markets is contaminated with hemlock seeds, but it is not known whether serious consequences have resulted therefrom.

Symptoms. The symptoms in man are due to a general and gradual weakening of muscular power. The power of sight is often lost, but the mind usually remains clear until death ensues, as it soon does from the gradual paralysis of the lungs. The poisoning differs from that of the Water Hemlock (*Cicuta maculata*) in the absence of convulsions. Many domestic animals have been killed by eating the plant, the prominent symptoms described for cows being the loss of appetite, salivation, bloating, much bodily pain, loss of muscular power and rapid, feeble pulse.

This plant, though called hemlock, should not be confused with the hemlock tree, which belongs to the family Coniferae. It paralyzes the ends of the motor nerves, then trunks and lastly the motor center itself. Respiration is quickened and pupils contracted. The fatal dose according to Blyth is 23 grams.

2. *Cicuta L.* Water-Hemlock

Tall, smooth, erect perennial herbs with pinnate or pinnately compound leaves and serrate leaflets, umbels terminal. Flowers white, calyx teeth acute,

fruit ovate or oblong with solitary conspicuous oil tubes, corky ribs, the lateral ones strong, marsh berber. Eight species, of north temperate regions. The European *C. viscosa* is deadly poisonous. Hundreds of people have been poisoned in Europe. It acts much like our native cowbane, the symptoms being violent gastro-enteritis, dizziness, trembling, suggestive of hydrophobia, prostration, paralysis and convulsions.

Cicuta maculata L. Cowbane

A smooth marsh perennial from 2-5 feet tall, and with fasciated fusiform roots, leaves pinnately compound 2 or 3 times pinnate, long petioled, the coarsely serrate leaflets lanceolate to oblong lanceolate, stalks of the umbellifers numerous and unequal, flowers white, fruit broadly ovate to oval, small, about $1\frac{1}{2}$ inches long.

Distribution Grows in marshes and low grounds in the Dakotas, Nebraska, the Rocky Mountain region of Colorado, Wyoming and Montana to the Uintahs, east of New Brunswick and Florida.

Poisonous properties The European *C. viscosa* contains *coniin* $C_8H_{17}N$ found also in *Conium maculatum*, and the bitter principle *cicutarin*, an amorphous, resinous substance with a disagreeable taste. The poison resides in the root, stem, and leaves, but more particularly in the root. It seems to occur in an oily aromatic fluid.



Fig. 371 Water Hemlock (*Cicuta maculata*) showing section of spindle shaped roots and lower stem, the leaves, flowers, and fruit. One half natural size also fruit and cross section of seed, enlarged five times. A very poisonous plant. (U. S. Dept. Agr.)

The resinous cicutoxin, according to Boehm, is an uncrystallizable bitter body. The fatal dose, according to Chesnut, is 50 milligrams for each kilogram of body weight when administered through the mouth and 7 milligrams when injected hypodermically.




Fig. 372 European Water Hemlock (*Cicuta verna*). A poisonous plant containing cicutoxin. (From Vasey's *Traité de Botanique*)

Mr Chesnut says

The true chemical nature is not definitely known, but it is probable that it contains the alkaloid *cicuta*, and the bitter principle *cicutarin*, the latter of which is characteristic of the European water hemlock (*Cicuta verna*). Both are powerful poisons, but the latter is the more violent and produces most of the symptoms characteristic of the plant. The American water-hemlock is one of the most poisonous plants native to the United States. Its victims include both men and animals. The underground parts are the most poisonous, and are especially dangerous, because they are often washed or frozen out of the soil and thus exposed to view.

There are quite a number of cases of human poisoning on record in Wisconsin, Iowa, and Minnesota. Stock is also poisoned. People who are poisoned generally mistake the roots for parsnips. In Iowa it is often called wild parsnip. The roots of this plant are fasciated and never conical as in the true parsnip. During one season five children were poisoned in the state of Iowa, three dying, from eating the roots. Several cases of stock poisoning have occurred in Iowa and are referred to by the writer. The following interesting experience is related by Mr J. A. Minter, who says

I have just had a strange experience with my cattle having lost a four year old cow and a yearling calf. I think that they were poisoned on some kind of weed root found in the slough. I locate it on a spot where a hay stack stood about two years ago. It had been removed except the spoiled hay in the bottom. Last fall being dry, I pitched it up, dried and burned the old hay, sowed rye and timothy seed, ran the disc harrow over it several times and noticed that we turned out lots of roots like small sweet potatoes, except that they were all connected at the top. I thought they were the root of a weed that grows a stalk similar to a seeded parsnip, have a straggly top similar to an elder berry when in bloom. The stalk when mature is hollow. Now I am not certain that I am right about the top as it had been mowed before I discovered the tubers. I never thought of them doing any harm, not thought we had torn them out so they would die and do me no harm, but as the cattle 17 in number were

brought up Sunday evening they appeared to be all right until they came into the barn yard, when a cow fell down and seemed to have a spasm. It only lasted a few minutes when she got up walked about 100 feet and fell again, got up and walked about thirty rods, fell again and died in about thirty minutes. The yearling was all right until turned into the lot. In about 20 minutes she was taken in the same way except a little more severely, rose 2 or 3 times and died in about 15 minutes. I was satisfied that they were poisoned, but the cause worried me for a while then I remembered the tubers I saw in the slough. I went next morning before turning the cattle out and found that the cow and yearling had eaten some of the roots. I gathered up nearly one-half bushel of the tubers, turned out the cattle and have had no trouble since. On opening the cows I found considerable of the tubers in the stomach, and the inside of the stomach was very black.

The plant above ground likewise affects horses and evidently the poison may reside in the leaves for considerable length of time even after they are dried. The following experience of a correspondent in Ruthven, Iowa, calls attention to the danger of using hay that contains cowbane.

I enclose herewith a small paper box which contains some seed of which I sent you a specimen last summer. This species of hemlock as you call it, I picked out of a manger & a situation which look suddenly sick this morning. Sickness lasted but a short spell. Do not know whether this had anything to do with this sickness but am terribly persuaded against it. Another instance a few days ago of a colt taking violent sick all at once apparently no cause, there being considerable of this weed in the hay and I had two cows lose their calves a short time ago. The cows had access to this kind of hay. This quite frequently occurs hereabouts. On a neighboring farm where this weed abounds, they lost nearly all their calves two years ago. Apparently no cause but of course there is a cause somewhere. I am satisfied some stock will eat the leaves of this weed.

Dr. Erwin F. Smith, in referring to the poisonous nature of this weed, speaks of a case as follows:

During the warm days which melted the snow and brought back the birds and gave indication of spring time, some children of a neighborhood on the outskirts of the city gave vent to their feelings by digging and eating some waterhemp which grew upon some low ground bordering a brook. Two of these boys were taken violently sick and one of them eight years old, died, within an hour after he had eaten the root.

Dr. Smith states that upon an examination of the stomach and the root from which he ate, it was proven beyond a doubt that *Cicuta maculata* was the cause of death.

Professor A. A. Crozier calls attention to the poisonous nature of cowbane in this state and refers to a case occurring in northern Iowa as follows:

Hon. Eugene Secor, of Forest City, this state a member of the Board of Trustees of the Iowa Agricultural College, brought me today a fleshy root of a plant of the Water hemlock (*Cicuta maculata*, L.). The circumstances which brought it to his notice were as follows: A neighbor of his by the name of Mr. Brown, a farmer of about fifty years of age while dragging some potato ground upon bottom and about two weeks ago discovered one of the fleshy roots of this plant, and supposing it to be an artichoke ate it and gave a portion of it to his two sons. He soon began to feel queer or "funny" as he expressed it, and went to the house where he was taken with a spasm, followed by two or three others, when he became unconscious and within half an hour, before a physician could be summoned from the village, two miles distant, he was dead. The children had probably eaten less of the root and being given an emetic recovered. The plant is very common in the state and the roots are so pleasant to the taste as to make it particularly dangerous. I may add that I ate a piece of the root the size of a filbert with little or no unpleasant effect."

One season the writer had a record of five cases of poisoning in this state. From a press bulletin issued by the writer the following facts were given to the public:

Ira, aged ten, and Ross, aged eight years, children of T. Y. Johnson, died last night from eating the roots of a poisonous water plant that grows in front of their house on the Keg Creek flat east of the Ridgeway lumber yard. A third child, John, the seven year old son of Mrs. Amanda Kingery, also ate the root but it seems was not made so sick as the others.

Shortly before six o'clock the children came into the house showing Mrs. Johnson what they had been eating. Not knowing what it was she had them spit it out and throw away

what they had in their hands. They went out to play again, but in about twenty minutes the two Johnson boys fell to the ground as if in a fit, soon passing into convulsions. The Kingery lad was able to walk to his home, but was soon taken with spasms. Dr. Hester was called and by a vomiting process in a few hours had the child out of danger.

Dr. Lyon was summoned to the Johnson home, getting there at 6:30. The little fellows were already in terrible convulsions and nothing could be done for them, one dying at 6:45 and the other at 7:15.

Dr. Millspaugh, in his *American Medicinal Plants*, Fascicle 4, No. 67, has recorded the following observations concerning the physiological action of the *Cicuta maculata*:

"Many cases of poisoning from the root of this species have been reported, all showing, by the symptoms, that *Cicuta* produces great hyperæmia of the brain and spinal cord. The following case reported by letter to Dr. Bigelow* by Mr. E. Hazeltine (1818) gives us



Fig. 371. Purple Stemmed Poison Hemlock (*Cicuta Douglasii*) of the Pacific Coast. It is very poisonous. (Chestnut, U. S. Deer Age)

*Bigelow, Amer. Med. Bot. Vol. III, p. 121.

the symptoms noted by other observers in various other cases. A boy had eaten of certain tuberous roots, gathered in a recently plowed field, supposing them to be artichokes but which were identified as the roots of *Cicuta maculata*. His first symptom was a pain in the bowels, urging him to an ineffectual attempt at stool after which he vomited about a tea-cupful of what appeared to be the recently masticated root, and immediately fell back into convulsions which lasted off and on continuously till his death. The doctor found him in a profuse sweat and convulsive agitations, consisting of tremors, violent contractions and distortions, with alternate and imperfect relaxations of the whole muscular system, astonishing mobility of the eyeballs and eyelids, with widely dilated pupils, stridor denture, trismus, frothing at the mouth and nose, mixed with blood and occasionally violent and genuine epilepsy.

The convulsive agitations were so powerful and incessant that the doctor could not examine the pulse with sufficient constancy to ascertain its character.

At the post-mortem no inflammation was observed, the stomach was fully distended with flatus and contained "about three gills of muciform and greenish fluid, such as had flowed from the mouth, the mass assumed a dark green color on standing."

Chestnut in his paper on Some Poisonous Plants of Northern Stock Ranges says that Dr. Wilcox and himself observed 105 cases of water hemlock poisoning among sheep of which 50 were fatal, and 36 among cattle of which 30 were fatal. The loss was \$4,000, only a fraction of what occurs in Oregon. The *C. vagans* and *C. Douglasii* are poisonous, the latter along the coast.

Mention may be made in this connection of a series of most valuable papers on "The Medicinal Plants of North America" by Dr. T. H. M. Merck's Reports.* In one paper he discusses the anatomy as well as the poisons found in this very poisonous plant. The effect of the poison is similar to that of *Cicuta viridis* and is due to a resinous substance *cicutoxin* and to the volatile alkaloid *cicutin*, which has been obtained from the fruits.

Cicuta vagans Greene. Oregon Water-Hemlock

A smooth perennial with glaucous stem and vertical rootstock divided into horizontal chambers. plant 2-3 feet high, with compound leaves. flowers white.

Distribution. From Idaho to British Columbia and west to northern California.

Poisonous properties. Same properties as the preceding. Professor Hedrick estimates that 100 head of cattle are killed by it every year in Oregon. A piece about the size of a marble of the winter rootstock is believed to be fatal to man. Professor Hedrick says:

It is hard to estimate the number of cattle killed yearly in Oregon by eating *Cicuta*. One hundred would be a low estimate in my judgment. Animals eat the underground portion of *Cicuta* in getting the tops which form about the first green herbage in early spring. As they browse the foliage, the roots, being on a partly subterranean, and growing in a soft soil, are pulled up and eaten. A piece the size of a walnut or a found by experiment, is sufficient to kill a cow. It is probable that the poisonous constituent is found only in the underground stem and the roots.

While the victims of the plant are chiefly cattle and they are not exclusively so. The poisonous parts are often mistaken for Parsnips, Artichokes and Horseradish, and thus human victims are not infrequent. A number of cases of poisoning from *Cicuta* are annually reported in the United States. A writer in a local paper a few months ago, reported the case of two cattlemen in Southern Oregon, who after eating Wild Parsnip, presumably *Cicuta*, died in a few hours. Falk reports, that in Europe in thirty-one cases of poisoning from *Cicuta*, 45 per cent died.

The observations made by Prof. French and reported by Prof. U. P. Hed-

*Medicinal Plants of North America, 24. *Cicuta maculata*. Merck's Rep. Vol. III 35-38 11-12. Feb. 1909.

rick formerly of the Oregon Experiment Station have shown experimentally that this species is poisonous. A bulb was cut in small pieces, mixed with a carrot and fed to a two year old grade heifer. The animal was fed at 8:00 a. m. and at 9:30 it was dead. A post-mortem examination showed that pieces



FIG. 374. Oregon water hemlock (*Cicuta occidentalis*). a, plant with leaves, one with natural size; b and b', rootstock and horizontal roots, showing section, half size; c, terminal umbels, one with natural size; d, flowering spray. (U. S. Dept. Ag.)

of the root occurred in the rumen, and in the reticulum or second stomach. Only a very small portion of the root had been eaten by the animal. Two grams were found in the stomach. The lungs were highly congested.

The following day a one year old calf was fed with the poisonous roots. The temperature of the animal was normal, $102\frac{1}{2}^{\circ}$. Two bulbs the size of an egg were cut and mixed with carrots. This material was eaten under protest. The roots were fed at 9:15 a. m. and at 9:40 the temperature was 103 , at 10:00, $103\frac{1}{2}^{\circ}$ at 10:25, 104° and the animal was trembling about the flanks, the eyes watered freely, at 10:35 the animal was excited and the temperature was $104\frac{1}{4}^{\circ}$, at 10:45 the animal fell over on its side in a spasm. The eyes were drawn and the muscles were rigid and contracted violently.

Before death an attempt was made to revive the animal by giving it spirits of turpentine. This caused the calf to recover its feet and walk about. A second

dose of turpentine and milk were given and the animal remained standing until 11.30, then it fell down as before. A full dose of aconite was given, but no results noticed and then a hypodermic injection of nitro-glycerin was given. Death occurred at 11.45 and spasms were almost continuous.

The most interesting discovery made by Prof. Hedrick was in regard to the apparent harmlessness of the roots while growing during the summer.

Prof. Hedrick says concerning the roots fed to an animal early in May.

The bulbs were of the same lot used in the first experiments, but had been growing for a month in a green house. It was expected that growth would remove some of the dangerous properties of the bulbs, but it was a surprise to find that an animal eating many times as much of them as had killed the cows in the previous experiment suffered no ill effects whatever. The conclusion is obvious; the bulbs are only dangerously poison when in the dormant state, or for a short time after growth begins in the spring. Cattle are likely, then, to be poisoned only from the first of January to the middle of May.

Cicuta Bolanderi Watson.

Leaves bipinnate, leaflets narrowly lanceolate, long acuminate, acutely serrate, lower leaflets petiolate and often deeply lobed; involucre of several linear leaflets, fruit 2 lines long, nearly orbicular, strikingly ribbed, and broad oil tubes.

Distribution. In salt marshes along the Pacific coast in California.

Poisonous properties. Like those of the preceding species.



Fig. 375. Water Hemlock (*Cicuta bulbifera*)
A powerfully poisonous plant. (Ada Hayden.)

Cicuta bulbifera L. Bulbous Water-Hemlock

A slender perennial from 1-3 feet high; leaves 2-3 times pinnate, leaflets linear, sparingly toothed, 2 inches long; upper axils bear clustered bulblets fruit small, ovate.

Distribution. In swamps Nova Scotia to Delaware, Indiana, Iowa, Nebraska and Manitoba.

Poisonous properties. Very poisonous, like the other species described above

3. *Sium* (Tourn.) L. Water Parsnip

Smooth perennial herbs with stem leaves pinnate, basal leaves with pinnatifid leaflets, flowers white in large umbels, calyx teeth minute, short styles, fruit ovate to oblong with prominent ribs, 1-3 oil tubes. A small genus of 8 species in temperate regions. The *S. latifolium* is poisonous especially to cattle. It produces stupor, excitement and gastro-enteritis

Sium cicutaefolium Schrank. Water Parsnip

An erect stout marsh herb from 2-6 feet high, lower leaves with long petioles, leaflets 3-8 pairs, segments linear or lanceolate, sharply serrate flowers in umbels, white, fruit ovate, prominently ribbed.

Distribution. From Nova Scotia across the continent to Florida and California.

Poisonous properties. The water parsnip is reported as poisonous from several different sources. Hyams reports it poisonous in North Carolina

4. *Aethusa* L. Fool's Parsley

Annual glabrous herbs. Leaves 2-3 ternately compound, dissected, umbels compound without involucre, involucre long and narrow, flowers white, calyx teeth obsolete, fruit globose, ovoid, glabrous, flattened dorsally. One species native to Europe and Asia

Aethusa Cynapium L. Fool's Parsley

An erect leafy branched annual. Leaves 2-3 times pinnate petiole dilated at the base, umbels long peduncled

Distribution. In cultivated grounds and waste places from Nova Scotia New England to Pennsylvania and New York

Poisonous properties. Contains the alkaloid *cynapin*, and a comin-like alkaloid. One physician in England regards the plant as non-poisonous, even recommending it as a pot herb or for salad uses. On the other hand numerous cases of poisoning have been recorded. The following may serve as an illustration which is recorded by Dr. Millspaugh from a statement made many years ago. He records the experiments made on animals. Seven ounces of the juicy leaves were given to a strong dog and the oesophagus tied. twenty minutes thereafter the dog became sick. He stretched out his limbs and lay on his stomach and it was impossible to arouse the animal. The pupils were scarcely dilated, the pulsations of the heart were slow and strong. The extremities were agitated by convulsive movements; the animal threw himself from one side to the other, and died an hour after taking the poison. The post-mortem examination showed that the heart was contracted and the stomach was found to be full of the poison. In domestic animals it causes stupor, paralysis and convulsions. The common name indicates that it is sometimes mistaken for parsley with injurious results.



Fig. 376. Fool's Parsley (*Aethusa Cynapium*). Flowering branch supposed to be poisonous to animals. (After Paguet.)

5. *Angelica* L. *Angelica*

Stout perennial herbs with ternately or pinnately compound leaves in large ample umbels with white or greenish flowers, involucre none or of few small bracts, calyx teeth wanting, fruit strongly flattened with prominent lateral wings or tubes solitary or several, the 30 or more species found chiefly in the northern hemisphere and New Zealand.

Angelica atropurpurea L. Great Angelica

A stout smooth perennial from 3-5 feet high, large ample, ternately divided

leaves, pinnate segments, ovate-lanceolate and broad dilated petioles, flowers in large umbels 9-25 rayed, fruit broadly oval, oil tubes 25-30.

Distribution. Common in swamps from Labrador, Wisconsin, Minnesota, Illinois to Delaware.

Poisonous principles. Supposed to be poisonous. Although no cases have been reported to the writer, it is known that the Indians of Canada used the fresh roots for suicidal purposes. On drying, however, the roots lose their poisonous properties. Dr. Millspaugh says that they are considered carminative, diuretic, emmenagogue and stimulant. The dried root was often used, especially in combination with other and better known diuretics, in anasarca and various diseases of the urinary organs, and alone in flatulent colic and suppressed menstruation. Dr. Schell claims that doses of 15 to 20 grains of the dried root will cause a disgust for all spirituous liquors. The stems were often made into a candied preserve in some sections of the country—a practice now nearly extinct.

6. *Oxypholis* Raf. Cowbane

Perennial glabrous marsh herbs, with clustered tuberous roots, leaves reduced, pinnate or ternate, flowers white, in compound umbels, calyx teeth acute, fruit ovate, dorsally flattened, compressed, dorsal ribs slender, the lateral broadly winged, oil tubes solitary in the intervals, 2-6 on the commissure. Species 4, native to North America. Poisonous.

Oxypholis rigidior (L.) Coult and Rose Cowbane

A slender marsh perennial from 2-5 feet high. Roots tuberous, clustered. Leaves simply pinnate, petioled, leaflets thick, ovate-lanceolate or oblong-entire or denticulate, involucre of 1-4 bracts or none. Flowers white, oil tubes small.

Distribution. In swamps from New York to Florida, to Missouri and Minnesota.

Poisonous property. The roots and leaves are known to be poisonous. Said to poison cattle.

7. *Pastinaca* L. Parsnip

Tall, branching biennial herbs, pinnate leaves, thick conical roots, compound umbels with yellow flowers, involucre and involucrels commonly absent, obsolete calyx teeth, fruit smooth, oval, flattened, the lateral ribs extending into broad wings, oil tubes solitary, 2-4 on the commissure. Six or seven species native to Asia and Europe.

Pastinaca sativa L. Wild Parsnip

Tall branched biennial or annual herb with thick conical roots, pinnate leaves smooth or somewhat pubescent, calyx teeth obsolete, petals yellow, fruit oval, glabrous, flattened dorsally, seeds flat.

Distribution. Common in the northern states and on the Pacific coast, where it has escaped from cultivation.

Poisonous properties. Frequently confused with cowbane and said to produce poisoning. The writer has not received a single specimen where Wild Parsnip was said to have produced the poisoning, that the plant did not prove to be cowbane. The wide spread belief of the poisonous nature of the cultivated parsnip running wild is entertained by a large number of people and also to some extent by the medical fraternity. A few years ago Professor Fred-

erick B. Power and one of his pupils (Mr J T Bennett) undertook some experiments to determine whether the cultivated parsnip running wild had any toxic properties. Mr Bennett failed to detect the presence of any poisonous principle in the root of the true wild parsnip (*Pastinaca sativa*) and when the boiled roots were fed in considerable amounts to a cat, no symptoms of poisoning were manifest. We may add as a further



Fig. 327. Wild Parsnip (*Pastinaca sativa*). A plant with yellow flowers and its feral form of ordinary parsnip. Often mistaken for Cowbane (Charlotte M. King.)

testimony that Professor Power reports that his associate, Dr Kreiners, insisted upon eating one-half of one of the raw roots obtained from Mr. Rynning of West Salem, Wisconsin, which were supposed to cause cases of the poisoning. Dr Kreiners reported no ill effect whatever. One of the largest of the fresh raw roots weighing three and one half ounces avoirdupois was chopped fine, mixed with some meat and fed to a small dog. The animal ate it greedily and without disturbance. There were no symptoms whatever of poisonous action. I have on different occasions eaten the wild parsnip (*Pastinaca sativa*), without any ill effect, so that the above results are corroborated. I will admit that I had some hesitancy at first and that Mr Sexton, the foreman of the horticultural department at the Iowa State College did not expect to see me alive by evening. I must confess also that the roots were somewhat woody and not very palatable.

Dr J J. Brown, Sheboygan, Wisconsin, as quoted by Dr Power, states that he had prepared and dug enough wild parsnips for a good dinner which he ate and can testify that he could discover but little difference cooked or raw, from cultivated parsnip and those which had run wild for about 50 years.

That the parsnip does at times produce dermatitis has been proven. The following very interesting letter from Professor F. C. Stewart, botanist of the New York Agricultural Experiment Station, is of interest in this connection:

I recollect that some fifteen years ago you were much interested in the reputed poisonous properties of the wild parsnip, and that you reached the conclusion that wild parsnip is not poisonous. I think you may be interested to know of an instance which has recently come to my attention. Henry Vandreser, a prominent lecturer on poultry in this state, last season had a very serious injury to his face and eyes. His face became badly swollen and his eyes were in a terrible condition. It was feared at the time that they would be ruined, but the sight was not lost although it was considerably impaired. The physician in charge diagnosed it as a case of poisoning, due probably to the flowers of wild parsnip. Very shortly before the trouble appeared Mr. Vandreser had been mowing a large patch of wild parsnip which was in bloom. It was a hot day, so that he perspired profusely. He gathered bunches of the wild parsnip plants in his arms and carried them. This brought the plants in contact with his face. Both Mr. Vandreser and the physician feel confident that the wild parsnip was the cause of the trouble. Another gentleman who heard of this case told me that some years ago he lost a little girl with poisoning of a somewhat similar character, and it was attributed to the parsnip blossoms, among which the little girl had been playing innocently before he attacked.

The writer has also known of a few cases of dermatitis produced by this plant. Poisoning similar to the above may be produced by other members of his family notably the cowbane. A young high school lad in Boone lost his life in a way similar to that from *Pastinaca sativa*.

B. *Heracleum* L. Cow Parsnip

Tall stout perennials with large ternately compound leaves. Flowers in large umbels, involucre deciduous and involucre of numerous linear bracts; calyx teeth obsolete or small, petals white, clawed, the outer flowers dilated, fruit broadly obovate, flattened, lateral ribs broadly winged, the intermediate narrow oil tubes extending to about the middle. About 60 species, of cooler regions.

Heracleum lanatum Michx. Cow Parsnip

A stout, hairy, pubescent perennial from 4-8 feet high, leaflets broad and large, irregularly cut-toothed, flowers white, in broad umbels.

Distribution. From the Atlantic coast, Newfoundland through the northern states and Allegheny Mountains to California. Common in the Rocky Mountains.

Poisonous properties. Said to be poisonous although the leaves of the fresh plant are eaten by the Indians. It contains the bitter principle *heracalin* $C_{25}H_{38}O_6$. The roots have a disagreeable flavor. Hyams states that it is poisonous and Hasted says that cow parsnip will produce blisters.

Daucus Tourn. Carrot

Bristly annual or biennial herbs with pinnately decompound leaves, umbels compound, of white or reddish flowers, involucre of numerous entire or toothed bracts, calyx teeth obsolete, petals obovate with apex turned in, the outer flowers often dilated, fruit oblong, flattened with 3 primary ribs, secondary ribs 4 winged, each bearing a row of prickles. About 25 species of wide distribution.

9. *Daucus Carota* L. Wild Carrot

Biennial or annual bristly herbs and conical tap root, lower leaves 2-3 times pinnate, segments dentate lobed or pinnatifid, bracts of the involucre of linear divisions, flowers white or pinkish.

Distribution. Native to Europe. Common in fields and waste places in the north to the Pacific coast. It is the original of the cultivated carrot.

Injurious properties. The fruit often clings to wool, making it of inferior quality. The leaves of the plant cause vesication. This is especially true when

the plant is wet with dew. In Australia, according to Maiden, the *Douglas brachiatum* is injurious to sheep, the hooks sometimes entering the skin.

CORNACEAE Link Dogwood Family

Shrubs or trees, with opposite or alternate leaves, calyx tube adnate to the ovary, lobes small, petals as many as the stamens, 4-5, inserted on the disk; stamens as many as the petals or more numerous, ovary inferior, 1-2 celled, style 1, ovules 1, in each cavity, fruit a 1-2 seeded drupe.

A small family of 16 genera and about 85 species, containing several plants cultivated for ornamental purposes, such as the red osier (*Cornus stolonifera*), native from Nova Scotia south to Virginia, Kentucky, northeastern Iowa, Nebraska, the Rocky Mountains, California and Alaska.

The *Aucuba japonica*, with handsome red berries, is cultivated in greenhouses. It contains the glucoside aucubin $C_{13}H_{18}O_8 + H_2O$.

Flowering dogwood (*Cornus florida*) is a small tree or large shrub, cultivated for its large corolla like bracts. It is native in dry woods from southern New England to Ontario, Illinois, Missouri, and Texas. The western dogwood (*Cornus Nuttallii*) of the Pacific, is a beautiful tree from 20-75 feet high and 12 feet in diameter. It produces a close grained wood that is capable of receiving a high polish and is used for furniture, cabinet work, and water wheels. Tupelos (*Nyssa aquatica*) and (*N. sylvatica*) produce wood of some importance.

The dogwoods, especially the flowering dogwood and the round-leaved dogwood (*Cornus circinata*) are used in medicine, the bark contains the bitter principle *cornin*. Some species, like panicled dogwood (*C. paniculata*) are occasionally troublesome in newly cultivated fields.

Poisonous properties. At various times, members of this family have been suspected of being poisonous. Two species, *Cornus Amomum* Mill. and *C. paniculata* L'Her (= *C. canadensis* Mill.) are thus listed in Part I of this Manual. Actual poisoning from species of *Cornus* has not, so far as is known to the present author, been observed. Greshoff, however, lists as poisonous *C. Amomum* Mill., *Alangium Lamarchii* Thwait., and *Garrya Fremontii* Torr.

METACILLAMYDEAE

Petals partly or wholly united, rarely separate or wanting, tubular or funnel-form. This includes the important plants classed as *Gamopetalae*.

ERICACEAE

Flowers complete, regular with lobed or distinct calyx; corolla cleft; stamens free from corolla, ovary compound. This order includes such families as *Perisactae* and *Ericales*.

ERICACEAE Heath Family

Shrubs, trees or occasionally herbs, with simple leaves, flowers regular, or nearly so, generally perfect, calyx usually free from the ovary, 4-5 lobed, generally persistent, corolla of 4-5 petals or 4-5 lobed, anthers 2-celled, commonly appendaged or opening by pores, ovary 3-10 celled, style 1, pollen consisting usually of 4 united grains, embryo small.

A large family of wide distribution. About 1400 species. Several species of the family are frequently cultivated for ornamental purposes—notably the rhododendron and the wild laurels (*Rhododendron catawbiense*, *R. maximum*) which are hardy out of doors in the south and east. The azalea (*Rhododendron indicum*) is frequently cultivated in greenhouses, flowering in late winter and early spring. The garden azalea (*Rhododendron sinense*) is cultivated in gardens.

The California mountain laurel (*R. californicum*) is a very attractive ornamental shrub or small tree and one of the most handsome of the genus in North America. Among the other cultivated members of this family, mention may be made of the mountain laurel (*Kalmia latifolia*). The South African species of the genus are frequently cultivated in conservatories. The Scotch heather (*Calluna vulgaris*) of Europe has been found on Nantucket Island. The May flower or trailing arbutus (*Epigaea repens*) is well known to people living in the Eastern and Northern States. The rich, spicy fragrance makes it one of the most attractive of our early flowering plants. Several species of the genus *Vaccinium* are cultivated for their fruits. Among them are cranberries, the larger berry being *V. macrocarpon*, which is the most common species in North America, and is cultivated extensively in Wisconsin, Michigan, New Jersey and Massachusetts. The smaller cranberry *V. Oxycoccos* is also sold under the name of Lunon berries and is common in Sweden and Norway and far northward in North America. The berries of this fruit are smaller and more acid than are those of *V. macrocarpon*.

The blue berries and huckleberries are obtained from several native species of the genus *Vaccinium* of the East. Dwarf blueberries of our northern dry woods are (*V. pennsylvanicum* and *V. canadense*). Another blueberry (*V. vacillans*) produces a sweet edible berry and occurs from Maine to Missouri. The mountain blueberry of the southern Alleghenias is *V. corymbosum* var. *pauciflorum*. The tall blueberry (*V. corymbosum*) has a pleasant acid flavor, and attains a height of 5-10 feet. In the Rocky Mountain, Lake Superior and White Mountain regions, there are several species of blueberries as *V. caespitosum* with blue sweet berries and *V. membranaceum*, which produces large black berries, and grows from 1-5 feet high.

The huckleberry (*V. stamineum*) is not edible nor is the fruit of the small tree known as farkle berry (*V. arboreum*). The huckleberry (*Gaylussacia bacata*) is edible. These come up spontaneously in burnt areas. The fruit is gathered from Maine to Minnesota. The shallon (*Gaultheria shallon*) of the Pacific coast, is a shrub or small tree with stout erect branches, and produces edible fruit with a spicy aromatic flavor. There are several species of arbutus, one of which Madrona (*Arbutus Menziesii*) forms a tree 1-6 feet in diameter and 20-100 feet high, the wood being used for cabinet work. The strawberry tree (*Arbutus* [*uva*]) is a native of Southern Europe to northern Africa. The fruit when ripe resembles the strawberry but is not edible. Several members of the order are used in medicine. One species only, however, is generally recognized. The Pipsissewa (*Chimaphila umbellata*) used for cutaneous eruptions, contains *chimaphilin* $C_{24}H_{32}O_8$. The bear-berry (*Arctostaphylos Uva-ursi*) is found throughout the mountain regions of North America—Pennsylvania, Illinois, Missouri to Nebraska, the Rocky Mountains and northward. It is an astringent tonic used in diseases of the liver, and contains *arbutin arbutin arsons* $C_{12}H_{16}O_6$. The wintergreen (*Gaultheria procumbens*) is a small creeping shrub with evergreen leaves and edible fruit, and is used for the production of the oil of wintergreen which serves as an aromatic stimulant and also in rheumatism. The fragrance is due to a volatile oil.

Drs. Price and L. Engle quite recently mention a case of poisoning from the oil of wintergreen. In this case a child swallowed the contents of a bottle containing oil of Gaultheria. The symptoms of poisoning were increased pulse, labored and irregular respiration, hearing impaired, some hallucinations of vision, twitching of heart and weak diarrhoea. The child died.

Many of the *Ericaceae* contain the glucoside *ericolin* $C_{24}H_{36}O_{11}$, this in turn contains *ericiol* $C_{10}H_{16}O$, a peculiarly fragrant oil, *ericolin* occurs in the leaves of *Ledum*, *Erica*, *Calluna*, *Rhododendron*, *Gaultheria* and *Epigaea*. *Rhododendron* $C_{16}H_{22}O_7$, is found in the leaves of *Rhododendron chrysanthum*, it is converted by hydrolysis into *rhododendrol*, $C_6H_{12}O_2$. *Oxytocin* occurs in *Vaccinium Oxycoecus*.

Andromedotoxin or *acetotoxin*, found in the leaves of *Andromeda Pokifolia* and *A. japonica* and also in many other *Ericaceae*, as *Anura Rhododendron japonicum*, *A. javanicum*, and *Kalmia* is one of the most toxic substances known.

Asebolin, $C_{24}H_{36}O_{11}$, derived from *andromedotoxin*, and *asebogensin* $C_{18}H_{24}O_7$, are found in *Andromeda Japonica*. *Arbutin* $C_{12}H_{18}O_7$, occurring in the leaves of many of the *Ericaceae* was first detected in *Arctostaphylos Uva-ursi*, *quercetin* is found in *Calluna*, as is also *calluna tannic acid*.

In Lehmann's work on the poisonous plants commonly found in Germany only one species of this family is listed as poisonous, namely the Labrador tea (*Ledum palustre*) of the Arctic region. This pretty shrub, growing in swamps, has alternate entire leaves with rusty wool underneath, revolute margins white flowers in umbel like clusters followed by a many seeded capsule. The *L. groenlandicum* related to it and also occurring in swamps, thickets and on mountain slopes but farther southward has wider leaves and is a larger plant.



Fig. 378. Great Laurel (*Rhododendron maritimum*) a, flowering branch; b, fruiting capsule—both one-third natural size. Very poisonous. (U. S. Dept. Agr.)



The *L. glandulosum*, similar to the above but a taller plant, occurs in bogs of the Rockies. All are more or less poisonous, one species is known to contain *andromedotoxin*. Several species are used as a substitute for tea. The Russians use the first named species. It is also said to be used to protect clothes from moths.

Genera of *Ericaceae*

Fruit a septicidal capsule, anthers unappendaged.

Corolla somewhat irregular

Corolla slightly 2-lipped, leaves deciduous (Rhododendron-Azalea)

Corolla campanulate, leaves evergreen 1 Rhododendron

Corolla regular

Stamens 10, capsule 5-celled 2 Kalnia

Fruit a loculicidal capsule, anthers often awned.

Corolla cylindrical, capsule depressed 3 Leucothoe

Corolla globose or urn-shaped, capsule globular

Anthers awned 4 Andromeda

Anthers awnless 5 Lyonia

Rhododendron L. Rose Bay

Shrubs or trees, with deciduous or persistent leaves, flowers in corymbose or umbellate clusters from scaly cone-like buds, calyx 5-lobed, or 5 parted, corolla bell-shaped, 5-lobed, nearly regular or irregular, stamens 5-10, spreading, declined and slightly or not exerted, anthers opening by terminal pores, style slender occasionally exerted and somewhat declined, fruit a woody capsule, seeds numerous. About 100 species, native of cooler regions. Several species cultivated for ornamental purposes. The genus includes Azalea.

Rhododendron californicum, Hook. California Rhododendron

A shrub or small tree from 3-25 feet high, with stout erect branches; leaves evergreen, broadly oblong, 3-6 inches long, flowers large, in corymbose clusters, calyx 5-lobed, corolla bell-shaped, rose color, capsule 1 1/4 inches long.

Distribution. On the Pacific slope from San Francisco north to British Columbia.

Rhododendron maximum, L. Rose Bay or Great Laurel

A tall evergreen shrub or small tree, leaves 6-10 inches long, thick and leathery, oblong-lanceolate, dark green on both sides, with revolute margins, flowers in corymbose clusters, pedicels viscid pubescent, corolla bell-shaped, 1 inch broad, pale rose color or nearly white or spotted with yellowish orange dots; fruit an oblong capsule.

Distribution. In damp woods from Maine to Ohio and the Alleghenies to Georgia.

Rhododendron catawbiense Michx. Mountain Rose Bay

Shrub from 3-6 feet high, or rarely small tree 20 feet high, leaves oval or oblong, mucronate, dark green above, pale or glaucous beneath; petioles stout, pedicels rather stout, corolla broadly bell-shaped, lilac-purple, petals rusty-brown.

Distribution. On mountain slopes of Virginia, West Virginia, and Georgia. *Poisonous properties.* According to Chesnut, *Rhododendron californicum*

is said, in Oregon, to be poisonous to sheep. The substance *andromedotoxin* $C_{21}H_{34}O_{10}$, a bitter principle has been found in *Rhododendron maximum*. It is extremely poisonous, even more so than aconitin, and more emetic than emetin. *Rhododendron ponticum* contains the glucoside *arbutin*.

Rhododendron occidentale A. Gray California Azalea

Shrub 2-6 feet high, shoots glabrous or minutely pubescent when young, not bristly, leaves obovate-oblong, sometimes approaching lanceolate, bright-green and shining above minutely pubescent, glabrate the margins minutely recurved ciliate, scars of the flower bud somewhat canescent. flowers appearing after the leaves sepals distinct oblong or oval conspicuous corolla minutely viscid pubescent outside white with the upper lobe yellow inside the narrow funnel-form tube equalling the deeply 5 cleft slightly irregular limb, the lobes ovate, stamens and style much exserted moderately curved capsule oblong



Fig. 379. California Azalea. *Rhododendron occidentale*. Flowering stem. Leaves are said to be poisonous to stock. (From American Agriculturist.)

Distribution. In wooded districts along streams in California extending to the mountains east of San Diego and common northward.

Poisonous properties. The leaves of the species have been analyzed chemically and have been found to contain a poisonous substance, said in California to be poisonous to stock. Certain species of *Azalea* contain the glucoside *ericoha* $C_{24}H_{34}O_{11}$.

Kalmia, L.

Erect shrubs with entire evergreen coriaceous leaves, naked buds, flowers in umbels or corymba, or solitary, or 2-3 together in the axils, calyx 5-parted, corolla somewhat wheel-shaped, 5-lobed, with 10 pouches in which 10 anthers are lodged, filaments long, stamens 10, shorter than the corolla, anther-sacs opened by large terminal pores; ovary 3-celled, ovules numerous, capsule obscurely 5-lobed, 5-celled, many-seeded, seeds small. Species 7, 6 native to North America 1 in Cuba.

Kalmia latifolia L. Calico-bush. Mountain-laurel

A shrub 4-8 feet high, forming dense thickets, becoming a tree 10-30 feet high in the mountains, branches stiff, leaves alternate, bright green on both sides ovate-lanceolate or oblong petioled, flowers profuse, large, and showy, lip rose-color to nearly white, in terminal corymba, capsule depressed, globose, 5-lobed, glandular.

Distribution. Rocky hills, damp soil. Canada to Maine and the Allegheny Mountains, through West Florida, Ohio, Kentucky, and Tennessee.

Poisonous properties. This plant has long been known as poisonous, and is very much dreaded in the Allegheny Mountains. The substance *andromeda-karin* $C_{21}H_{26}O_{10}$, referred to elsewhere, has been found in all parts of the plant. Prof. Chesnut, in reference to the plants says:

Scores of cattle and sheep are poisoned annually by eating the shrub. Access to it is generally obtained by breaking away from inclosures, or through neglect or accident when cattle or sheep are being driven past laurel thickets to upland pastures in early spring. Laurel leaves (commonly used for decorative purposes in winter), or the flowering branches, are often carelessly thrown into inclosures where animals are kept. The older cattle are not so frequently killed by it, but they are by no means immune. Horses and even goats have died from eating the leaves and in May, 1895, a monkey was killed at the National Zoological Park, at Washington, D. C., by eating a few flowers and leaves offered to it by a visitor. Deer and grouse are said to be immune, although it is claimed that their flesh, especially that of the ruffed grouse, is poisonous when they have fed upon it. It is stated that chickens have been poisoned by eating the vomited matter from poisoned animals. Experiments show, however, that they are able to withstand considerable quantities of the pure poison when it is fed to them. In these experiments the chickens were killed with chloroform after dosing for a few days. The entrails were then cast aside, and the well-boiled meat was fed to cats with nearly fatal results. The honey derived from the nectar of the flower appears to be poisonous under some conditions. Cases of human poisoning occur indirectly in the ways indicated above, directly by overdoses or improper use in domestic medicine, probably by the secret and criminal use of the leaves to increase the intoxicating effects of liquors, and, in children, by their eating the young shoots by mistake for the wintergreen (*Gaultheria procumbens*).

Dr. Johnson refers to the poisonous properties of the plant as follows:

"The leaves of the plant as also those of *Kalmia angustifolia* (Sheep-Laurel, Lamb-kill), are said to be poisonous to sheep and calves; and cases are reported of men being poisoned by eating the flesh of partridges which had fed upon the buds and berries. On the other hand Wilson, the ornithologist, ate freely of the flesh of such birds without any ill effect whatever, and deer feed upon the leaves in winter, not only without ill effect upon themselves, but also without rendering their flesh unwholesome to man. And yet the common opinion is that leaves are very poisonous. The writer himself was cautioned by an old Massachusetts farmer against



Fig. 380. Mountain laurel (*Kalmia latifolia*) a, flowering spray, one-third natural size, vertical section of flower showing peculiar attachment of stamens, natural size c, fruiting capsules, natural size. Very poisonous (Chenut, U. S. Dept. Agr.)



Fig. 381. Mountain Laurel (*Kalmia latifolia*) 1, section of flower bud showing anthers lodged in the pockets of the corolla 2, expanded flower showing anthers in pockets, 3, sectional view (Gray.)

meddling with *K. angustifolia* because of its poisonous nature, disregarding the advice he ate several leaves in the presence of the farmer, without, however, convincing the latter, notwithstanding no ill effects were experienced. That persons have been poisoned seriously by eating the flesh of partridges in winter cannot be denied, that these birds may have fed upon *kalmia* buds and berries is also probable, but that their flesh is thus rendered poisonous does not seem as yet fully established. Some experimenters have reported effects produced in their own persons by strong decoctions of the leaves, similar to those of partridge poisoning others quite as worthy of credence have failed to observe any sensible effect from them. Taking these contradictory statements in connection with the negative results of the chemical analyses thus far made, one may reasonably conclude either that the poisonous character of these plants has been greatly exaggerated, or that the energy of the poison, whatever it may be, is greatly influenced by the personal idiosyncrasy of the individual experimented with.

Dr. Mulspaugh, after digesting the various opinions that have been given regarding the plant, makes this statement:

From the experience of nearly all persons who have experimented upon themselves with a tincture or decoction of the leaves, it is obvious that the effects produced on cattle after grazing on the leaves, and on persons eating of "poisoned" partridges, are due to the plant itself, not to indigestion or putrefaction. Dr. Bigelow's later observations agree in toto with our previous ones. He gives the following as its action: "The flesh of the bird impairs the functions of the brain and acts directly as a sedative poison, secondarily affecting the digestive and circulatory organs." The symptoms arising in those proving the drug are: Vertigo and headache, almost complete loss of sight, pale, somewhat livid countenance, salivation and difficult deglutition, thirst, nausea and vomiting, with oppression and pressure in the region of the stomach, difficult respiration with great palpitation, and fluttering of the heart, followed by an irregular, feeble, and slow pulse, weakness, weariness and pains in the limbs, coldness of the surface and great prostration."

There would seem to be no question then that this plant is poisonous.

Kalmia angustifolia, L. Sheep-laurel. Lambkill

Shrub from 1-3 feet high. leaves usually opposite, or in 3's, pale or whitish beneath, light green above, acute or narrowly oblong, petioled, flowers in simple or compound corymbs, purple or crimson, pedicels filiform, recurved in fruit, sepals ovate, acute, capsule depressed, globose.

Distribution. In moist soil from Eastern Canada to Newfoundland, from Hudson Bay south to Georgia and Michigan.



Fig. 382 Sheep-laurel (*Kalmia angustifolia*), showing flowering branch, one-third natural size. (Chesnut, U. S. Dept. Agr.)

Kalmia polifolia, Wang. Swamp laurel

A low shrub with 2-edged branchlets, opposite nearly sessile leaves, oblong and white-glaucous beneath, margins revolute, flowers in terminal corymbs, few flowered, flowers $\frac{1}{2}$ inch broad, purple, capsule depressed, glabrous, smooth.

Distribution. In bogs, Newfoundland to Alaska, Connecticut to Pennsylvania Michigan, Rocky Mountains, and California.

Poisonous properties. Both of these species are regarded as poisonous, the *K. angustifolia* especially so. The symptoms of poisoning are very much the same as those produced by the preceding species.



Fig. 383 Branch Ivy (*Lencothoe Catesbaei*) a, flowering branch, b, fruiting capsule (Chesnut, U. S. Dept. Agr.)

Lencothoe, D. Don

Shrubs with alternate petioled leaves, and small, usually white, flowers in axillary or terminal spiked, racemes sepals 5, distinct, corolla cylindrical, 5-toothed, stamens 10, included; anthers naked, or the cells with 1 or 2 erect awns at the apex, opening by a pore, capsule depressed, more or less 5-lobed, 5-celled, 5-valved, seeds mostly pendulous, minute. About 35 species, natives of the Western Continent a few in Asia.

Leucothoe Catesbaei, (Walt.) Gray

Shrubs 2-4 feet high; leaves ovate-lanceolate, taper-pointed, serrulate, ever green, with spinulose teeth, racemes dense, many-flowered; bracts borne at the bases of the short petioles, corolla narrowly cylindric, anthers awned, capsules depressed.

Leucothoe racemosa, Gray

Shrubs 4-10 feet high, branches and racemes mostly erect, leaves oblong to ovate, generally acute at each end, sepals ovate-lanceolate, anther cells 2-awned, stigma capitate, capsule depressed-globose, seeds smooth and wingless.

Distribution. Moist thickets and swamps from Massachusetts, to Florida, and Louisiana.

Poisonous properties. Both species are known to be poisonous in the Alleghany Mountain region to all kinds of stock; probably contain the same principles found in other plants of this order.



Fig. 384. Swamp *Leucothoe* (*Leucothoe racemosa*). A well known poisonous plant of the Alleghany Mountain region and the Southern states. (Charlotte M. King.)

Andromeda, L.

Shrubs or small trees, leaves evergreen, short petioled, flowers in panicles, racemes or umbellate clusters, calyx persistent, without bractlets, corolla globose, urn-shaped, 5-parted, stamens 10, included, anthers fixed near the middle and opening by a pore, ovary 5-celled with columnar style, capsule globular and 5-celled, many seeded; seeds smooth. A small genus of 13 species found in Eastern Asia, the Himalayas, North America and Europe.

Andromeda Polifolia L. Wild Rosemary. Fetter-bush

A glabrous shrub 6-19 inches high, coriaceous, leaves with strongly revolute

margins, glaucous beneath, flowers in few-flowered drooping umbels, bracts persistent, calyx lobes triangular corolla urn-shaped, stamens 10, anther with a slender terminal awn.

Distribution. From Newfoundland to New Jersey, Michigan to British Columbia and Alaska. Europe and Asia.

Andromeda floribunda, Pursh. Mountain Fetter-bush

A leafy shrub from 2-6 feet high, branches nearly erect, bristly strigose pubescent, leaves oblong to ovate-lanceolate, ciliate and glandular dotted be-



Fig. 385. *Andromeda* (*Andromeda Petiolaris*)
Common in the North, especially in bogs and swamps
(Ada Hayden.)

neath. flowers white in densely flowered panicles, corolla 5-angled stamens 10, anthers without appendages, capsule somewhat globose shorter than the style.

Distribution. In the mountains of Virginia to Georgia.

Lyonia, Nutt

Shrubs. Leaves coriaceous and evergreen or thin and deciduous. flowers in racemes or panicles, white, calyx of 5 nearly distinct sepals, stamens 5 filaments hairy and often toothed or appendaged, anthers unappendaged capsule 5-angled dorsal suture with a thickened ridge, seeds with a thin testa.

Lyonia mariana, B. & H. Stagger-bush

A glabrous shrub with black dots, from 2-4 feet high, oblong or oval leaves, smooth above, slightly pubescent on the veins; flowers in nodding fascicles, racemose on nearly leafless branches segments of calyx acute, corolla ovoid, white or pinkish stamens 10, anther filaments 2-toothed near the apex capsule ovate, as large as the sepals.

Distribution Low grounds from Rhode Island to Florida, Tennessee and Arkansas.

Poisonous properties. All three species are poisonous. The leaves of the fetter-bush contain a narcotic poison, the *andromedotoxin* $C_{21}H_{31}O_{10}$ and have been known to kill sheep. Sheep have also been poisoned by the mountain fetter-bush. The stagger-bush received its name because of the intoxicating effect of its leaves on sheep and cattle.

PRIMULALES

Herbs or shrubs, corolla mostly present, gamopetalous stamens borne on the corolla as many as its lobes or twice as many or more. The family *Plumbaginaceae* contains *Statice*, growing mostly in saline soil along the coast. It is used as an astringent, particularly in diarrhoea. The baycurn (*Statice brussensis*) is one of the most powerful astringents, and is used locally as a gargle.



Fig. 384. Stagger-bush (*Lyonia maritima*) showing flowering branch, one third natural size. It contains a narcotic poison (Chesnut, U. S. Dept. Agr.)

PRIMULACEAE Primrose Family

Herbs with simple leaves and regular flowers; calyx 5-parted stamens as many as the lobes of the gamopetalous corolla and borne upon it, ovary 1-celled, bearing several or many seeds, calyx free from the ovary or partly adherent. A small order with about 28 genera, and 350 species, of wide distribution in the northern hemisphere, many of them arctic.

Few of the members of this order are weedy, or at least they are not troublesome. The honeywort (*Lysimachia Nummularia*) has escaped from gardens into damp ground and occasionally some species of the loosestrifes, are troublesome as weeds. Several members of this family are cultivated for ornamental purposes, among which are the cowslips or primroses and cyclamens. Native species of *Streptema*, like *S. lanceolatum*, and *Lysimachia*, are pretty ornamental plants. One of the prettiest of the native species in the northern




FIG. 387. Long and short-styled flowers of *Primula*; a and b styles; c pollen grains, and d petal of short-styled flower. Modified by C. M. King, after Seeburger.

states is the American cowslip (*Dodecatheon Meadia*), which is stemless and has a 6-parted reflexed corolla and exserted stamens. The true primroses (*Primula*) are frequently cultivated, the *P. sinensis* and *P. obconica* in greenhouses. The English cowslip, *P. vulgaris*, is an interesting European plant, and is also found in high altitudes in North America. The beautiful *P. Parryi*, of the Rocky Mountains, is abundant along brooks and in moist meadows at high altitudes, and is one of the prettiest of the Rocky Mountain plants. The root-stock has a strong odor of musk and is said to be somewhat poisonous. The leaves of *P. obconica* are poisonous to the touch, being similar in effects to the poison ivy, although not so strong. The European cyclamen is a commonly cultivated plant in greenhouses. The pimpernel or poor man's weather-glass (*Anagallis arvensis*) is common in sandy soil and occasionally somewhat weedy. It is native to Europe. In *Primula* and cyclamen is found cyclamin, $C_{21}H_{33}O_{12}$ with which *primulin* is supposed to be identified.

Anagallis, L.

Herbs with alternate sessile, or short petioled whorled leaves. flowers solitary, on axillary peduncles, calyx 5-parted, corolla wheel-shaped, deeply 5-parted, longer than the calyx, stamens 5, filaments bearded, capsule globose, circumscissile, many-seeded, seeds minute, flat on the back. About 15 species, mostly in the old world.

Anagallis arvensis, L.

Annual, leaves ovate, or oval, membranous, sessile, or somewhat clasping, calyx-lobe keeled, shorter than the obovate corolla-segments, fringed with minute stalked glands, flowers scarlet or white, usually with a darker center, capsule smooth.

Distribution. Newfoundland to Minnesota, Mexico, and on the Pacific Coast. Naturalized from Europe.



Fig. 141 Persian cyclamen (*Cyclamen*). Frequently cultivated in greenhouses. (From American Agriculturist)

Poisonous properties. This plant is especially common on the Pacific Coast, where it is well known as "Poison weed," and according to Prof. Cheanut, is suspected of having caused the death of a horse at Santa Ana. It contains the glucoside *cyclamin* $C_{25}H_{37}O_{13}$, and a saponin like substance, also a pepsin like ferment. According to Hyams, the fluid extract in 4 drachm doses is fatal to dogs.

Primula, L.

Perennial scapose herbs, producing a tuft of basal leaves, flowers dimorphous, umbellate, or in bracted racemose whorls; calyx tubular, angled, 5-cleft, persistent, corolla funneliform or salverform, tube longer than the calyx; stamens 5, included; capsule many seeded, splitting at the top into 5 or 10 valves. About 150 species, mostly in the northern hemisphere. This species is rather rare in eastern North America, but one species, (*P. Parryi*), is abundant in higher altitudes of the Rocky Mountains, and is suspected of being poisonous.

Primula obconica. Hance. Primrose

Leaves all radical and ovate-cordate, slender scapes 6-12 inches long, flowers bluish-lilac or purple, frequently drooping; calyx saucer-shaped and shallow, petals deeply notched, obconical.

Distribution. Native to China, but frequently cultivated in greenhouses.



Fig. 389. Poison weed (*Anagallis arvensis*) Flowering branch
 Flower c, corolla, s, sepals. Poisonous to horses and dogs. (From
 Vaseux's *Traité de Botanique*.)



Fig. 390. Parry's Primrose (*Primula Parryi*). This primrose is common in high altitudes. The roots have a more or less musky odor and it is regarded as poisonous. (Charlotte M. King.)

Poisonous properties. *Primula obconica* and *P. sinensis* both contain *oxalic acid* $C_2H_2O_2(OH)_2C_{11}H_{22}$. It has long been known that some people are sensitive to poisoning by coming in contact with the leaves of *P. obconica*. The hairs act as an irritant poison. Dr. White calls attention to the injury as follows:

The cutaneous disturbance therein described, dermatitis venenata, subsided in a short time, and the skin of the three affected persons has remained in a healthy condition until recently. Within the last two or three weeks, however, they have all manifested a recurrence of the same symptoms, an exzematous inflammation of the hands and face, and in about the same degree as last year. They now feel assured that the trouble is caused by *Primula obconica*, for the condition did not develop until a few days after this plant was first offered for sale in the shop, where it was freely handled by them. The proprietor informs me that his hands and face became affected immediately after making it up into dinner-table decorations.

Dr. Thurber refers to a case of poisoning quoted by Mr. Meehan:

A person, who after potting a lot of *Primula obconica*, had his face so swollen that he remained completely blind for a day. This was the worst case yet reported, but it should be stated that Mr. Cliffe was at that time suffering from poisoning by *Euphorbia pulcherrima*.

Mr. Thurber says:

Some have likened the effects produced by *Primula obconica* to those of *Rhus Toxicodendron*, the Poison Ivy, but it is a well established fact that many persons are poisoned by *Rhus* without coming in contact with the plant. Merely passing near it or being near a fire when it is being burned is sufficient to cause serious poisoning, indicating that the active principle is volatile.

EBENALES

Shrubs or trees, with alternate simple leaves, flowers generally regular, calyx more or less adnate to the ovary, corolla gamopetalous or sometimes polypetalous; stamens as many as the lobes, with corolla opposite them or more numerous.

FAMILIES OF EBENALES

Stamens as many as the corolla lobes	Sapotaceae
Stamens twice as many as the corolla lobes	
Flowers dioecious or polygamous	Ebenaceae
Flowers perfect	Stryaceae

SAPOTACEAE Sapodilla Family

Shrubs or trees with milky juice; simple and entire, alternate leaves, small and perfect, regular flowers, persistent sepals 4-7, corolla gamopetalous tube 4-7 lobed, frequently with as many or more lobe-like appendages, stamens as many as the proper lobes of the corolla, fruit a fleshy drupe, generally 1-celled and 1-seeded, seed large, coat hard, endosperm fleshy or none.

A small order of 35 genera and 400 species, of tropical regions. But few of these occur in the United States, the most common in the southern states being the Southern buckthorn (*Bumelia lycioides*) and *B. longinosa*, both being common in the woods from Illinois southward. The star apple (*Chrysophyllum Camuto*) with a fruit about the size of an apple, has a pleasant flavor but is said to be very constipating. The sapodilla plum (*Achras sapota*) also native to the West Indies, is widely cultivated in the tropics for its fruit, which is about the size of a pear and inferior only to the orange. The narmalade (*Lucuma mammosa*) of tropical America, from 3 to 5 inches long, is much esteemed for its pulpy fruit. The leaves, however, contain *prunic acid*, which is often used to adulterate cola nut. The fruits of the camno (*Lucuma obovata*) a native of Peru and Chili, are also eaten, as are the fruits of the black bully

tree (*Dipholis salicifolia*) of Jamaica. Seeds of the buttertree (*Beana butyracea*) furnish an oil which is an article of some commercial importance in India. The seeds of the buttertree (*Butyraspermum Parkii*) of South Africa, furnish an oil which is used as food by the natives. Gutta percha (*Dichopora gutta*), of Borneo, and the Malayan Peninsula is the source of some of the Gutta-percha of commerce which is used in surgical dressing the same sub-



Fig. 391. Gutta-percha Tree (*Dichopora Gutta*)
 obtained from the latex of this tree. (After Faguet.)

The gutta-percha of commerce is ob-

stance being also obtained from *Mimusops Kauh*, a large tree, native of British Guiana. The former contains *gutta* $C_{18}H_{16}N$, *fluorol* $C_{20}N_1O$, and *alban* $C_{40}H_{44}O_2$. *Sapotin* $C_{28}H_{32}O_{10}$ occurs in the bark of *Achras Sapota* and of *Symplocos racemosa*. In lotus bark occur three basic substances *loturin*, *col-loturin* and *loturidin*. In the seeds of *Ilpe Macclayana* a toxic glucoside, $C_{17}H_{33}O_{10}$, is found.

The *Sapotaceae* or Sapodilla family contains a number of plants of economic importance in the tropics, among these are *Mimusops Kauh*, *Dickopsia Gutta*, *Paquinum oblongifolium*, of the East Indies, and other species which furnish gutta percha. This is obtained from the latex of these plants and because of its plasticity, is particularly valuable in the manufacture of surgical instruments and for temporary filling of teeth. The same substance is produced by *Sideroxylon tomentosum* and *Paysona*. The Sapodilla tree (*Achras Sapota*) of Yucatan produces an aromatic gum resin which is largely used in the manufacture of chewing gum. Vegetable butter is obtained from the seeds of *Bassia butyracea*. The Sapodilla or Sapote (*A Sapota*) is commonly cultivated in the tropics, especially the West Indies and Central America, for its wild fruit. The star apple, *Chrysophyllum Cainito*, also produces a valuable fruit. Hard iron woods are produced by *Mimusops Kauh* of Australia and *Argania Sideroxylon* of Morocco. Saponin like substances are produced by several plants of the family such as *Omphalocarpum procerum*. The *Lucuma melleifolia* and *Paysona latifolia* contain hydrocyanic acid.

EBENACEAE. Ebony Family

Trees or shrubs with very hard wood, alternate entire leaves and polygamous regular flowers. Stamens from 2-4 times as many as the lobes of the corolla. ovary 3-12 celled. ovules 1-3 in each cavity. fruit a several-celled berry. seeds mostly single in each cell, large and flat, with a smooth coat.

About 6 genera and 275 species. The most widely distributed tree in the United States is the common persimmon (*Diospyros virginiana*) a tree from 20-70 feet high, found in woods and old fields from Rhode Island to New York and southern Iowa. The fruit is commonly eaten and the unripe fruit is used in medicine for internal hemorrhage. The Japanese persimmon (*D. Kaki*) with small flowers and a greenish-yellow fruit of variable size and shape, is cultivated in the Southern States and California for its fruit which is delicious. In Egypt it is cultivated under the name of Lotus tree.

The Indian persimmon (*D. Embryopteris*) is a middle sized evergreen tree native from India to Java, and is used in India in medicine, especially for chronic dysentery. Ebony wood comes from several species like the Ceylon ebony, Indian ebony (*D. Ebenum*) and (*D. melanoxylon*), and is highly prized for furniture since it is hard and black. The calamander wood (*D. quersia*) of Ceylon is a hard wood prized for making boxes. The mangosteen or mabola (*D. discolor*) a native of the Philippine Islands, is now commonly cultivated in the tropics for its fruit, which is about the size of a quince and has a pleasant flavor. Other hardwooded trees are *D. Cargulia* and *D. pentamera* natives of New South Wales.

Ebony wood is of various colors, the black from *D. ebenum*, red ebony from *D. rubra* native of Mauritius, white ebony from *D. chloroxylon*. Other species of *Diospyros* furnish a striped ebony, and others a green.

STYRACACEAE. *Styrax* Family.

Mostly tropical shrubs or trees with regular perfect, or polygamo-dioecious flowers, calyx free or adherent to the ovary corolla gamopetalous, or polypetalous, the lobes or petals 4-8, stamens twice as many as the lobes of the corolla, or more, inserted on its tube or base the filaments monadelphous, or in 4-5 sets, style 1, ovary 2-5-celled, fruit a berry or drupe.

About 75 species and 7 genera, few of them native to the United States. Several species are cultivated for ornamental purposes, especially the snow drop or silver bell tree (*Illicium carolinum*) which is a small tree native to the southern states. *Styrax Benzoin* is obtained from Sumatra and Java, benzoin being obtained by cutting incisions into the plant, from which a resin exudes and soon hardens by exposure to the air. *Benzoin* is used chiefly as incense in the service of the Greek Church.



Fig. 292. Black Ash (*Fraxinus nigra*). 1. Flowering branch of staminate tree. 2. Flowering branch of pistillate tree. 3. Pistillate flower showing rudimentary stamens, enlarged. 4. Longitudinal section of ovary, enlarged. 5. Fruiting branch. 6. Longitudinal section of fruit. 7. Embryo. 8. Winter branchlet. 9. 1/2, 5, 6, 8, one-half natural size. (M. M. Cheney in Green's Forestry of Minnesota.)

Styracin $C_{10}H_{10}O_7$, one of the ketones, occurs in *Styrax Benjoin* also resin, benzoic acid $C_7H_6O_2$, and cinnamic acid $C_9H_8O_2$. Siam benzoin contains vanillin.

CONTORTAE

Herbs, shrubs or trees, leaves opposite or rarely alternate, flowers regular, corolla gamopetalous, rarely polypetalous or wanting, stamens generally borne on lower part of corolla when present, as many as the lobes or fewer. Ovaries 2, distinct.

FAMILIES OF CONTORTAE

Corolla none. stamens usually 2	Oleaceae
Corolla present regular, stamens as many as its lobes	
Ovary 2, stigmas and sometimes styles united, fruit a follicle	
Stamens distinct	Apocynaceae
Stamens monadelphous attached to a stigmatic body	Asclepiadaceae
Ovary compound, 2-celled	Loganiaceae
Ovary 1-celled	Guttiferaceae

OLEACEAE. Olive Family

Trees, shrubs, or some nearly herbaceous, leaves opposite or rarely alternate simple or pinnate without stipules. flowers in panicles, cymes or fascicles. calyx small, inferior, sometimes wanting, stamens 2-4, filaments separate. ovary superior 2 celled, ovules a few in each cavity, style short or none. fruit a capsule, berry or drupe. About 500 species. Contains the lilac (*Syringa vulgaris* & *perica*), the jasmine (*Jasminum grandiflorum*) cultivated in France as the source of a valuable perfume, the fringe tree (*Chionanthus virginica*), well known in cultivation, and the ashes which make valuable timber trees like green ash (*Fraxinus pennsylvanica* var. *lanceolata*), white ash (*F. americana*), black ash (*F. nigra*) and manna of Europe, *F. Ornus*. The sweet exudate is called manna which contains mannitol and a glucoside resembling saccharin.

The olive (*Olea europaea*) of the Mediterranean region, is cultivated extensively in southern Spain, France, Italy to Arabia, California and other warm countries. The unripe fruit is used for table purposes by steeping in water containing lime and ash, and then pickling in a salt solution. The nearly ripe or ripe olives are used in a similar way. The seed as well as fruit contains an oil consisting of olein $C_{57}H_{104}(OC_{18}H_{33}O)_2$, the greenish color of which is due to the chlorophyll. It is nutritive and laxative. The hard wood takes a fine polish.

Edwards and Power have found an alkaloid in the bark of *Fraxinus americana*, one in the bark of *Olea glandulifera*, in the leaves and bark of *Ligustrum robustum* and in the leaves of *Jasminum*. The glucoside phillyrin, $C_{20}H_{32}O_9$, occurs in *Osmanthus fragrans* and *Forsythia suspensa*, and may be converted into phillygenin, $C_{20}H_{32}O_8$, which may again be readily converted into ugenol and vanillin. Ibo $C_{20}H_{32}O_{10}$, a saponin-like glucoside, occurs in *Ligustrum* and *Ibota chionanthi* is found in the bark of *Chionanthus virginica*, and is intensely bitter. It may also contain saponin. The plant is probably somewhat poisonous.

Ligustrum (Tourn.) L.

Shrubs or small trees with opposite entire leaves, flowers in a terminal thyrsus or panicle, corolla gamopetalous, funneliform, tube short, the limb 4-



Fig. 191. Large-flowered white jasmine (*Jasminum grandiflorum*). A native of Japan but cultivated in France as a source of perfume. (From Vauque's *Traité de Botanique*.)

lobed, stamens 2, filaments short, ovary 2-celled, with 2 ovules in each cavity, fruit a 1-3 seeded berry. About 35 species, natives of the Old World

Ligustrum vulgare, L. Privet

A shrub with long and slender branches, firm leaves, smooth, lanceolate or oblong, and short petioled, panicles dense, minutely pubescent, pedicels very short, flowers white

Distribution. Commonly cultivated for ornamental purposes, but escaped here and there from cultivation.

Poisonous properties. The leaves and fruit of the plant are said to be poisonous. Prof Chesnut states that accidents have been occasioned to children who have eaten the fruit or the leaves. The plant contains syringin and the bitter glucosidal principle *syringopictin* $C_{28}H_{34}O_{17}$. These also occur in the lilac. Edwards and Power have also found an alkaloid in the leaves and bark of an allied species, *Ligustrum robustum*, and it was discovered also to be in the seeds of *L. ibota*

LOGANIACEAE. Nux Vomica Family

Herbs, shrubs, vines, or in a few tropical genera, trees, simple, opposite leaves, stipules or small membranes, flowers perfect and regular in axillary or

cymose clusters, calyx of 4 to 5 parted sepals, corolla gamopetalous, 4 to 5 parted, stamens as many as the lobes of the corolla or alternate with them. Fruit follicles or drupes.

About 30 genera and 400 species, in warmer regions, several of them very poisonous. *Strychnos Nux-vomica*, native to India and the islands of southern Asia, is a small straggling tree, in its native country known as the Koocha tree. The fruit resembles an orange but with a hard rind and numerous fla-



Fig. 294. *Strychnos* (*Strychnos Crocousiana*). A curee plant of French Guiana. (After Faguet.)

seeds covered with silky hairs. The seeds contain two deadly alkaloids, *strychnin* $C_{21}H_{22}N_2O_7$ and *brucin* $C_{23}H_{26}N_2O_5 + 5H_2O$. Strychnin is bitter, used as a tonic and to stimulate the circulation. The bark and root are also bitter. The natives of India use it for snake bites and fevers. This alkaloid is also obtained from other plants of this order, being extracted by water acidulated with hydrochloric acid. It has an intensely bitter taste perceptible in very dilute solution (1 in 700,000).

Deafoy has shown that starving frogs are much more sensitive to strychnin than are normal frogs¹.




Fig. 395. Yellow Jessamine (*Gelsemium sempervirens*). Flowering branch, bud, dehiscent fruit, longitudinal sections of fruit and flowers. Root contains poisonous alkaloid. (After Faguet.)

¹ In a recent paper by Reid Hunt "The Effects of a Restricted Diet and of Various Diets Upon the Resistance of Animals to Certain Poisons," Bulletin Hygienic Laboratory, Public Health and Marine Hospital Service, of the U. S. Treasury Department he reports that diet has a marked effect upon resistance of animals to certain poisons, the resistance of some

It contains the glucoside *loganin* $C_{25}H_{41}O_{11}$ especially in the pulp. The *S. toxicaria* of the Orinoco, used as an arrow poison and called "curare," or curare, is obtained from the stripped bark, which is steeped in water. It contains the alkaloid *curarin* $C_{18}H_{23}NO$. Other alkaloids found in curare are *protocurarin* $C_{22}H_{33}NO$ which is poisonous, *lubocurarin* $C_{19}H_{27}NO_4$ and *curin* $C_{14}H_{17}NO_2$. The dried seeds of *S. potatorum* are used for clearing water. The *curatle* of Java is a strong poison causing tonic and clonic convulsions and there is no known antidote. It is used as an arrow poison. *S. Craccaiana* of French Guiana is used as a curare poison.

Dr. Winkler describes the toxicological effect of strychnin on a dog as follows:

One-twentieth of a grain of strychnin nitrate injected subcutaneously by the writer into a dog weighing 25 lbs., caused uneasiness and excitement, with protrusion of the eye-balls, and in the space of ten minutes, tetanic convulsions. The breathing was shallow and almost imperceptible, the pulse rapid and irregular, the lips were covered with foam, the tail was stiff and extended, the ears laid back and there was general muscular rigidity, the animal lying on his side in a state of quagitation. This condition lasted about three minutes and was followed by a period of relaxation. But the slightest noise or irritation of the skin brought on convulsions. The convulsions became less frequent and violent and ceased altogether within half an hour. The same animal was given a 1/60 of the alkaloid on the following day, but without producing any perceptible effect. One fortieth of a grain given on another day and in the same manner caused muscular uneasiness and restlessness, and within ten minutes induced a severe convulsion lasting a little over an hour, in which the animal was so rigid that he could be lifted but with great bending. The ears were soon laid back, the legs were extended and stiff, the tail was straight and rigid and the eye was protruding. The jaw muscles of the jaw and limbs. The corners of the mouth were drawn back and was anastomotic. The mouth was covered with foam and there was some tremor. The breathing was more or less increased owing to tetanic spasm of the respiratory muscles. Following this contraction the jaw dropped, he refused to swallow and another attack could not be produced by pressure on external carotid artery. Some twitching of the temporal muscles persisted. Later in the second stage of poisoning had ensued, and the motor nerves and cord of the inferior extremities had become paralyzed. Death occurred in general paralysis within half an hour, and without any recurrence of convulsions or tetanic rigidity. Death takes place more commonly in strychnin poisoning from asphyxia during a convulsion, and is caused by spasm of the respiratory muscles, or more rarely by spasm of the glottis. Sometimes a fatal result occurs from exhaustion, between the convulsions and occasionally death appears to follow the interference with the nervous system and depression of the heart.

The lethal dose for dogs has been stated as 1/50 gr. to 1/30 grain. A 1/20 is usually in large as estimated by the experiments mentioned above. The fatal amount varies greatly in accordance with the weight of the animal, probably less than 1/50 would be very serious, and cases are reported where they have been destroyed by 1/60 of the alkali. The therapeutic dose should, therefore, be proportioned, as advised, to the weight of the animal. Five to eight grains of nuxvomica will kill a dog.

animals must be increased forty fold by changes in diet. It was found that diet causes diet but not very marked differences in resistance to strychnin. The experiments further show that foods such as enter largely into the daily diet of man have most pronounced effect upon resistance of an animal to strychnin poisoning. They produce changes in metabolism but are not readily detectable by means ordinarily used in metabolism studies. Diet causes diet but not very marked differences in resistance to strychnin. It was found by Hunt that the effect of oatmeal, by increasing the resistance of certain animals to acetone is probably due in part to a specific effect of the diet upon the thyroid gland. The poisonous action of acetone is according to Hunt is largely due to the formation of hydrocyanic acid from it. This is the view expressed not only by Dr. Hunt, but also by Heymans and Mascoe.

In the paper by Dr. Hunt referred to above it is evident to learn from the work of Mansfield that starving rabbits are much more susceptible to strychnin, than well fed, and more than that well fed rabbits are much more resistant to strychnin. Mansfield and Rogers found the resistance of rabbits to strychnin to be directly proportional to the weight of the animal. In his connection it is of interest to state that Hunt found that strychnin has an important effect upon the resistance of animals to certain poisons. In some cases these effects seem to depend upon metabolic variations in the activity of the thyroid.

Power, who experimented with dogs and mice, found that when dogs were kept upon a high protein nutritious diet, they were much more resistant to strychnin. On a medium or low plane of protein nutrition, two survived this dose.

The minimum fatal dose of strychnin for man is one-half a gram. Usually four to seven grams constitute a lethal quantity, but recovery has ensued following the ingestion of 22 grams, after a full meal.

Horses—The toxic symptoms in horses resemble those already described in the dog. They do not appear for some time (20 minutes to 4 hours), depending on the rapidity of absorption when the drug is swallowed, and include excitement, muscular spasm and convulsions, increased frequency of the pulse, and difficult respiration. Death occurs in convulsions or in the interval between them. The minimum fatal dose of strychnine when given under the skin, is about $1\frac{1}{2}$ to 3 grams, and when swallowed, 3 to 5 grams of the alkaloid, or 1 to 2 ounces of max vomica.

Cattle are similarly affected with horses and dogs. There are exhibited muscular spasms, frequent pulse, difficult respiration, sensitiveness to light, sounds and external stimuli, protrusion of the eye-balls and convulsions. The fatal dose, by the mouth, varies greatly owing to difficulty of absorption in the complicated and capacious digestive apparatus of these ruminants. This is true of all medicines. When given under the skin the lethal dose is a little larger than that for horses. The fatal dose for same is said to be from gr. 1.6 gr. 36. Chickens are comparatively insusceptible, also guinea pigs and some monkeys.

Strychnin poisoning differs from tetanus in the fact that muscular rigidity is continuous in the latter but disappears to a considerable degree, if not completely, in the periods between the convulsions, in the case of strychnin poisoning. Moreover, in tetanus the body and limbs are bent, and the jaw more affected, while in strychnin poisoning the position is reversed.

Treatment—The treatment embraces the use of chemical antidotes, as ipecac or em. tals., or tannic acid, animal charcoal and emetics on the stomach tube before absorption has occurred. The best physiological antidote is chloral in large doses per rectum. Chloroform and amyl may also be given by inhalation, and quail and ice are indicated. Artificial respiration is of no service on account of the muscular spasm, unless air be forcibly driven into the trachea through a cannula. Calabar bean and gelsemium both depress the inferior cornua, but neither is of much value in strychnin poisoning.

A large number of the cases of poisoning are fatal according to Schauenstein 62 out of the 130 reported by him proved fatal. According to Falk, the minimum lethal dose is as follows: 6 mgm. *strychnin nitrate* for rabbit per kilogramm, rats, 75 mgm. frogs, 2 mgm. In one case $\frac{1}{2}$ grain *strychnin sulphate* produced death in 20 minutes, in another case 7-10 grain was required. The so-called vermin killers contain *strychnin*.

Strychnin is the best remedy to stimulate the action of the heart and to promote appetite and digestion. It is valuable in chronic complaints of cattle.

The St. Ignatius or False Angustura Bark (*S. Ignatii*) of the Philippine Islands, much used in medicine in the Philippines, contains the same substance as *S. Nux-vomica*. The *S. malaccensis* of Tonquin is used as a remedy for leprosy. Pink root (*Spigelia marilandica*) is a perennial herb, with opposite sessile leaves, tubular funnel form corolla, red outside yellow within 6 stamens slender style short 2-celled capsule found from Ohio to Florida and Texas. It contains a volatile alkaloid *spigelia* resembling *coniin* and a bitter acrid principle. It is a powerful anthelmintic. According to Hyams, when taken in overdoses it excites the circulation, causes dimness of vision, vertigo, dilated pupils, spasms of facial muscles and general convulsions, followed by death. It is especially fatal to children. Dr. True reported the plant to be poisonous. According to Dr. Stockberger this is frequently adulterated with *Ruellia ciliosa* (East Tennessee Pink-root). The prepared drug may contain, as impurities, roots of golden seal, serpentaria, soapwort, wild yam, and stone-wort.

Gelsemium, Juss. Yellow Jasmine

Smooth vines with opposite or whorled leaves, flowers in axillary cymes, regular, perfect, calyx deeply 5-parted, corolla funnel-form, 5-lobed, stamens 5, inserted on the tube of the corolla, pistil with a 2-celled ovary, style slender,

or filiform, 4-cleft, fruit a capsule, flattened, seeds winged. A small genus of 2 species, one in North America and the second in Asia.

Geisemum sempervirens (L.) Ait. Yellow Jessamine.

A well known poisonous woody vine climbing over shrubs and trees to the height of thirty feet or more with opposite or entire ovate or lanceolate leaves, shining and evergreen, flowers showy and fragrant in short axillary clusters, yellow.

Distribution. This plant grows in woods and low grounds from eastern Virginia to Florida, and as far west as Mexico to Guatemala.

Poisonous properties. It contains a poisonous alkaloid, *gelsemin* $C_{34}H_{54}N_2O_4$. The root is frequently used in medicine and poisoning has been caused by overdoses. It is used in febrile and inflammatory infections, and on the nervous system it is an antispasmodic sedative. It is, however, a powerful drug and should be used with great caution. In overdoses it produces nausea, pain in brows and eye balls, dilation of the pupils, paralysis and dimness of vision. When overdoses have been administered, stomach pump and coffee should be used at once in conjunction with whiskey. *Gelsemin* is a colorless, crystalline, bitter principle. This plant also contains the amorphous, bitter alkaloid *gelseminin* $C_{22}H_{37}(OH)N_2O_7$.

Dr Winslow gives the toxicological effects on animals as follows. Muscular weakness, especially in the fore legs, staggering gait and falling. These symptoms are followed by convulsive movements of the head, fore legs, and sometimes of the hind legs. The respiration is slow and feeble, temperature reduced, and there is sweating. Death occurs because of respiratory failure. Morphine subcutaneously has proved a good antidote.

Ryth records that 10 mgrs killed a frog in 4 hours, 8 mgrs. killed a cat in 15 minutes, 1.6 gram killed a woman in 7½ hours.

GENTIANACEAE. Gentian Family

Smooth, bitter herbs with opposite and sessile entire mostly simple leaves without stipules, flowers regular and perfect in clusters or solitary, calyx persistent 4-12 lobed, corolla gamopetalous, 4-12 lobed or parted, mostly persistent but withered, stamens as many as the lobes of the corolla, alternate, inserted in the throat of the corolla, pistil with a 1-celled or partly 2-celled ovary, ovules numerous, anatropous or amphitropous, fruit a capsule, mostly dehiscent by 2 valves, seeds small, with a minute embryo and fleshy endosperm.

About 600 species of wide distribution. Several plants of the family are used in medicine. The American centaury or rose pink (*Sabatia angularis*) furnishes a simple bitter tonic formerly much used in intermittent fevers. An allied species contains the glucoside *erythrocentaurin*, $C_{27}H_{24}O_8$. The Chirata (*Swertia Chirata*) a bitter tonic, contains the glucoside *chiratin* $C_{26}H_{40}O_{16}$, a very bitter substance, and *ophelic acid* $C_{22}H_{20}O_{10}$. The root of the American columbo (*Fraxea corymbosa*) is used as an emetic and cathartic or as a tonic. The *F. speciosa* is common in the Rocky Mountains. Several species of the gentian, as *Gentiana Andreuxii*, *G. puberula*, are used as bitter tonics.

The gentians are largely used in the preparation of the so-called stock foods, the chief constituents of the root being a bitter crystalline glucoside, *gentiopicrosin* $C_{20}H_{30}O_{12}$ and *gentiolic acid* $C_{14}H_{10}O_8$. The plants act as stomachics and bitter tonics.

Gentians are pretty flowers, but few of them are cultivated. The prettiest of these are the closed gentian (*G. Andromeda*), with large blue flowers, the white closed gentian (*G. flavida*), the fringed gentian (*G. crinita*), and the Texas lily (*Eustoma Russellianum*), native from Texas to Nebraska and introduced into Colorado, with handsome large blue and white flowers.



Fig. 396. Yellowish Gentian (*Gentiana flavida*). A plant with bitter properties frequently used in connection with stock foods. (Lous Pammel.)

Menyanthes (Tourn.) L.

Smooth marsh herb with creeping rootstocks, leaves 3-foliate, flowers conspicuous white or purplish, in racemes or panicle; calyx 5-parted, corolla short, funnel-shaped, 5-cleft, stamens 5, inserted on the tube of the corolla, ovary 1-celled, capsule oval, indehiscent.

Menyanthes trifoliata L. Buckbean. Bog-bean

Perennial from scaly rootstocks, leaves petioled, sheathing at the base, leaflets oblong or obovate, raceme 10-20 flowered, calyx shorter than the white or purplish corolla.

Distribution. Europe and Asia, and North America, especially northward from New England to Greenland and Alaska, Iowa and Minnesota.

Poisonous properties. Buckbean is bitter and suspected of being poisonous. It causes nausea, and contains the substance *menyanthin* $C_{33}H_{50}O_{14}$, in the green parts of the plant.



Fig. 397. Buckbean (*Menyanthes trifoliata*). This plant is said to be more or less poisonous. It is very common in low lands. (Charlotte M. King.)

APOCYNACEAE. Dogbane Family

Perennia, herbs, shrubs or vines with acrid or milky juice, entire, mainly opposite leaves, without stipules, and perfect, regular 5-parted flowers, the 5 lobes of the gamopetalous corolla convolute and twisted in the bud, stamens as many as the lobes of the corolla, alternate with them and inserted on the tube, or throat, pollen granular, ovary superior, with 2 distinct ovaries, fruit follicles or drupes, seeds with a large straight embryo, often bearing a tuft of hairs. About 25 genera and 1000 species. Very widely distributed, chiefly in the tropics.

The best known representative, perhaps, under cultivation is the oleander (*Nerium Oleander*) from the Levant, naturalized in Southern Europe, the Southwest and Mexico, and the *Nerium odorum*, of India. The periwinkles, *Vinca minor* and *Vinca major*, are common in cultivation, the most widely distributed being the common periwinkle (*Vinca minor*) erroneously called myrtle, found especially in country gardens and running wild in cemeteries and shady places. Some species of *Gynopogon* have a cumarin-like fragrance. Rubber is obtained from *Ochrosia*, *Landolphia*, *Kickxia*, *Hancornia*, *Urceola* and others. Of the medicinal species, the *Alstonia*, or Dita Bark, is a handsome forest tree from 50 to 90 feet high, common throughout the Indian Peninsula.

to Ceylon. The Dita Bark contains the alkaloid *ditamen* $C_{16}H_{11}NO_3$, recommended as a tonic and an antiperiodic, also the alkaloids *echitamine* $C_8H_{13}N_2O_4$ and *schitemin* $C_{10}H_{11}NO_4$. In the bark of *Alstonia spectabilis*, besides *ditamen* and *schitemin* there is found a third alkaloid *alstonamin*. An allied species native of eastern Australia, is used for ague. In North America the dogbanes are also used in medicine. The ordeal tree of Madagascar (*Cerbera manghin*) produces a very poisonous narcotic seed. It is used by the natives to poison persons, and formerly criminals were put to death by being pricked with a lance dipped in the juice of the kernel. A single seed is said to contain enough poison to kill 20 people.

Both species of *Apocynum* are used in medicine. The Indian hemp used as an emetic cathartic and diuretic contains the glucoside *apocynin* and a bitter resin like extractive, *apocynin*, both of which Schmedeberg isolated. The *Thevetia peruviana* of tropical America contains a powerful heart poison, *thevetin* $C_{34}H_{48}O_{12}$, and *thereverin*. *Strophanthus Kombe* of East Africa is a woody climber, and contains *strophanthin* $C_{16}H_{14}O_4$ which acts like *digitalin* being valuable in cardiac disease, it is also used as an arrow poison.* The *A. hispidus* contains the same glucoside and an alkaloid *masen*. The action of this drug is similar to *Digitalis* although not as efficient. The *Querbracho* (*Aspidosperma Querbracho blanco*) of Argentina furnishes a valuable tanning bark containing 3-4 per cent of tannin, it also contains alkaloids, six of which have been discovered among them being *aspidospermin* $C_{22}H_{26}N_2O_2$, *aspidospermatin* and *querbrachin* $C_{21}H_{30}N_2O_2$, *aspidosamen* and *querbrachamin*. The yellow flowered Nighshade (*Fraxinea*) of the West Indies is a cardiac poison not unlike *Digitalis*.

Apocynum, Dogbane

Perennial herbs with opposite entire leaves, small white or pink flowers, calyx 5-parted, with acute lobes and corolla bell-shaped, 5-cleft with 5 small triangular appendages below the throat opposite the lobes, stamens 5, inserted on the base of the corolla, with arrow-shaped anthers, style none, stigma large, ovoid, slightly 2-lobed. Fruit of 2 long slender follicles, seeds numerous, small, each with a long silky coma. A small genus of about 15 species, found chiefly in the north temperate zone.

Apocynum androsaemifolium L. Spreading dogbane

Rootstock horizontal, smooth, or rarely soft tomentose, branched above.

*There are also many snake poisons belonging to different families.

In the family *Acanthaceae* are *Andrographis paniculata*, *Rhinacanthus communis*, *Barleria cristata* and *B. lupulina*, the latter of which contains not only a large amount of potassium but some alkaloids also.

The common names of a great many plants indicate their supposed virtues as antidotes for snake bites. In the last edition of Gray's Manual a few of the common names of this class are among the *Compositae*, as Rattlesnake weed (*Hieracium venosum*), Rattlesnake root (*Prenanthes alba*, *P. serpentaria*), Rattlesnake master or Button snakeroot (*Frygiopsis yuccifolia*), Black snake-root (*Sanicula marylandica*); among the *Orchidaceae* Rattlesnake plantain (*Epipactis pubescens*), of the *Gramineae*, Rattlesnake grass (*Glyceria canadensis*), perhaps so named because of the rattling of the spikelets, Snakegrass (*Eragrostis megastachya*), so called because of the odor of the grass, of the *Ophioglossaceae*, Rattlesnake fern (*Botrychium Virginianum*).

spreading, leaves ovate, petioled; cymes loose, spreading, both terminal and axillary, calyx segments shorter than the tubes of the corolla, the latter pale rose color, open, bell-shaped.

Distribution. Common borders of thickets from Canada to British Columbia to Arizona and Georgia.

Apocynum cannabinum L. Indian Hemp

Glabrous or more or less soft-pubescent, 2-3 feet high, smooth, terminated by an erect, close, many-flowered cyme; corolla lobes nearly erect the tube not longer than the lanceolate segments of the calyx, greenish white appear in July and August.

Distribution. Common species eastward and troublesome as a weed in northern Mississippi Valley.

Poisonous properties Said to be poisonous to live stock



Fig. 398. Indian Hemp (*Apocynum cannabinum*). Flowering and fruiting branches. Bundle of fibers from stem. Section of flower. Furnishes a good bast fiber. (Dodge, U. S. Dept. Agr.)

Nerium. Oleander

Shrubs, leaves coriaceous, rigid, closely and transversely veiny; flowers showy, in terminal cymes; corolla salverform or tube narrow, funnellform; stamens attached to the middle of the tube; style 1; ovaries 2, forming pods; seeds tufted. Native to the Levant and India.

Nerium Oleander L.

Leaves lanceolate coriaceous, rigid, closely and transversely veiny flowers in terminal cymes, rose-color or white anthers scarcely protruding

Distribution Native to the Levant, naturalized in Southern Europe, and Southwest United States Frequently cultivated in greenhouses.

Poisonous properties John Smith, in his Domestic botany says, with reference to its poisonous properties

It grows abundantly in the valley of the Jordan, and when in flower is very beautiful. The whole of the plant is poisonous, and it is recorded that soldiers in Spain were poisoned through their meat being roasted on spits made of the peeled stem.

Prof. Chesnut states that stock are occasionally poisoned by eating the leaves, as the plant grows wild in northern Mexico and is abundant in the Southwest. The oleander is a heart stimulant acting like digitalis. Dr. S. Waterfall recently reported a case in which gastro-enteritis occurred, nausea, vomiting and irritation were prominent symptoms. He also reports the odor of the flower as poisonous.



Fig. 379 Oleander (*Nerium Oleander*). At the left, a stamen. *H* thread enlarged on cells of anther elongated end of connective. At the right flowering branch. Whole plant poisonous. (From Veague's *Traité de Botanique*.)

Prof. F. W. Wilson of the Arizona Station¹ has recently brought together the literature on the subject of oleander poisoning, also giving some of his own experiments.

Prof. Wilson finds that both the pink and white varieties are poisonous. He conducted some experiments with two horses, a cow, a mule, and three sheep

¹ *Bull. Ariz. Exp. Sta.* 59:38

The amount of oleander necessary to cause death in horses ranges from 15 to 20 gm. of green leaves, and from 15 to 30 gm. of dry leaves. This depends on the condition of the animal at the time the poison is obtained. A full stomach will necessitate more poison. In the case of cows it is safe to say that from 10 to 20 gm. of green leaves and 15 to 25 gm. of dry leaves are sufficient to cause death. For sheep the fatal dose of either green or dry leaves is from 1 to 5 gm. There is little danger in the bark, roots, or flowers since live stock would hardly obtain sufficient poison in that way. . . .

The general symptoms are increased temperature and pulse, coldness of the extremities, warm body temperature, dilation of the pupils of the eyes, and discoloration of the mouth and nostrils followed by vomitings. The body becomes wet with sweat due to the exertion caused by the powerful heart stimulation. The animal generally refuses to eat or drink during the 24 hours preceding death. This is usually due to soreness of the mouth and throat, making it painful to masticate and swallow food. The bowels act often and feces are usually greenish in color. The action of the kidneys is increased slightly and color of urine is normal. There is little doubt that numerous cases of oleander poisoning have never been brought to light because of death being attributed to other sources. It is safe to say, however, that many hundreds of animals have been lost in southern Arizona from this shrub.

ASCLEPIADACEAE. Milkweed Family

Perennial herbs, vines, or shrubs, with milky juice and opposite or whorled leaves, entire flowers in umbels, regular, calyx inferior, corolla bell or urn-shaped, rotate or funnel form, 5-lobed or 5-cleft the segments generally reflexed, a crown between the corolla and stamens, stamens 5, inserted on the corolla, generally monadelphous, anthers connivent around the stigma or more or less united with each other, commonly bearing an erect or inflexed member, pollen collected in masses generally 10, known as pollinia, ovary consisting of 2 carpels in fruit of 2 follicles, seeds flattened, usually appendaged by a long tuft of hairs called a coma.

About 200 genera, and 1800 species, of wide distribution, many members of the family being weedy, some are medicinal, but few are economic. Several plants of the family are cultivated. The *Periploca græca* of the Old World is an ornamental climber and produces granular pollen in place of pollinia. The *Stapeia* several species of which are cultivated, are natives of the Cape of Good Hope. They produce flowers of dull purple color with transverse stripes, exhaling a very disagreeable odor not unlike that of putrid meat.

The wax plant (*Elaeagnus*) is a well known house plant of India with rooting stems, thick, fleshy, cal leaves, and flesh colored flowers. The *Lonicera* is a European climber sometimes cultivated in the Eastern States. Several species native from South America are sometimes cultivated for ornamental purposes. Some species of the order are used in medicine. The Indian *Pariparika* (*Hemidesmus indicus*) growing in the Indian peninsula and Ceylon, has a medicinal root which is used as a tonic and diuretic. The root has the odor of the tonka bean or of sweet clover. The mulder (*Calotropis procera*) also a native of India from Ceylon and the Moluccas is common in waste grounds. The bark contains one bitter principle *madazin* used as a tonic and diaphoretic and in large doses as an emetic. It produces a strong fiber, the silk being exported as "kapok," or tree cotton. The Indian *specacanthia* (*Tylophora asthmatica*) is a climbing perennial of India and Mauritius, anciently much used by the Hindoos in dysentery. The pleurisy root or butterfly weed (*Asclepias tuberosa*) is used as a diuretic and an emetic. A *Curassavea* of the West Indies is a vermifuge. The flowers are said to produce excellent honey. Generally, however, honey bees are killed by becoming entangled in the pollen masses of some of the species of *Asclepias*. Many other plants of the order have an acrid poisonous juice. Two of our common species of milk-

weed (*A. tuberosa*) and (*A. incarnata*) and the (*Phaceloxylum officinale*) contain the bitter glucoside *asclepidin* which is an amorphous, bitter, yellow emetic substance. The root especially, which acts as an emetic, is recorded as poisonous in Europe and cattle and other domestic animals will not eat it. Friedberger and Fröhner state that it causes diabetes and general weakness. The caustic bush *Sarcostemma australe* of Australia is regarded in that country as poisonous. The Condurango (*Marsdenia Condurango*) a South American vine is used as an alterative. It is bitter and acid. The root of *Asclepias stellifera* of South Africa, according to J. Burtt Davy yields an excellent rubber.

Asclepias (Tourn.) L. Milkweed

Perennial herbs with milky juice, entire leaves, flowers in umbels, calyx 5-parted, persistent, the lobes spreading, corolla deeply 5-parted, reflexed during flowering deciduous, the crown consists of 5 hooded processes each containing an incurved horn, enclosing the stamen tube, stamens 5, inserted at the base of the corolla, anthers adherent to the stigma, each with 2 cells and containing a pair of pear shaped masses of pollen (pollinia), ovaries 2, follicles 2, the stigma 5-angled or 5-lobed, numerous seeds with a coma. About 85 species, mostly of the new world. Several members of the order are quite weedy, especially in grain fields and pastures.

Asclepias vestita Hook and Arn.

Densely floccose woolly, the white wool deciduous in age, leaves from ovate to oblong-lanceolate very acute or acuminate, often subcordate, short petioled or the upper sessile 4-6 inches long, umbels 1-4, the terminal usually peduncled the lateral all sessile, corolla greenish white or purplish, the lobes ovate, 3 inches long, column very short, hoods nearly erect verticillate, slightly surpassing the anthers, entirely at the back of the somewhat truncate summit, auriculate extended at the inner angle, the auricles or angles revolute; the great not horn-shaped attached up to the summit of the hood blunt not exserted, an interior crown of 10 tooth like processes in pairs between the hoods, ovaries glabrous, follicles at first canescent.

Distribution. California and adjacent regions.

Poisonous properties. Said to be poisonous. The juice of this species and of *A. eriocarpa* are irritating.

Asclepias mexicana Cav.

Stem 3-5 feet high, leaves in whorls of 3-6 or uppermost and lower opposite, sometimes also in axillary fascicles, linear or narrowly lanceolate (3-6 inches long, 2-6 lines broad), umbels corymbose, densely many flowered, on peduncles longer than the pedicels, flowers greenish white, sometimes tinged with purple, corolla lobes oblong, hoods broadly ovate, entire, shorter than the anthers, exceeded by the stout-umbulate incurved horn.

Distribution. In the southwest.

Poisonous properties. Said to be poisonous.

Asclepias speciosa Torr. Showy Milkweed

A perennial from 1-4 feet high, white tomentose or canescent, leaves thick, broadly ovate or oval, petioled, pedicel glabrate above, flowers greenish purple,



Common - known to *Scirpus setosus* - Dist. by Lelundary as monogonous. (Bull. Ia. Agr. Exp. Sta. 74)

borne in dense umbels or rarely solitary follicles, erect or spreading on the recurved pediceis.

Distribution. This weed is found in moist soil from Minnesota and northern Iowa to Kansas, in the Rocky Mountain region, common in Colorado, Wyoming and Utah. Troublesome not only in our meadows but occasionally also in our grain fields and gardens.




Fig. 400. Showy Milkweed (*Asclepias speciosa*). A well known troublesome weed with milky juice. This plant and other species are known to be poisonous. It is common from Western Missouri and Iowa westward (Ada Hayden.)

Asclepias syriaca L. Milkweed

A perennial herb with a stout stalk from 2-5 feet high, finely soft, pubescent leaves ova-oblong, or ovate, obtuse or roundish at the base the young leaf somewhat pubescent above, soon becoming glabrate, petioles stout, flowers borne in umbels, from a few to many, peduncles pubescent or tomentose, corolla dull purple or whitish in color, follicle borne on erect or recurved pediceis.

Distribution. This species is widely distributed in the north, occurring in waste places from New Brunswick to the Saskatchewan, along the Atlantic coast to North Carolina and south and west to Missouri and Kansas.

Poisonous properties. Said to be poisonous to live stock. It contains *asclepin*. Bees often become entangled in the pollen masses of this and preceding species and are unable to extricate themselves. Lehmann and other European writers list this species as poisonous.

TUBIFLORAE

Mostly herbs, rarely shrubs or trees, corolla generally gamopetalous, irregular or regular, stamens adnate to the corolla tube and generally alternate with them, ovary 1-celled, occasionally deeply 4-lobed.

Among the important families of this order are the *Polemoniaceae*, containing the ornamental *Phlox Drummondii*, the perennial *P. maculata*, *P. divaricata*, and *P. pilosa*, many cultivated varieties of *Phlox* and of *Gilia* of which *G. aggregata* and *Columba gracilis* are examples. According to Greshoff *Gilia aggregata* contains saponin. It is regarded as a poisonous plant. The order also includes the *Pedaliaceae*, including the sesame (*Sesamum indicum*) which furnishes the valuable sesame oil, the plant being indigenous to the East Indies, the *Orobanchaceae* which contains the troublesome brown rapeseed, (*O. ramosa* and *O. minor*), the *Gesneriaceae* containing the *Gloxinias*, native to Mexico but cultivated in this country as greenhouse plants, the *Lentibulariaceae* which includes the bladderwort (*Utricularia vulgaris*) and other insectivorous plants which are sometimes destructive to fish, and the butterwort (*Pinguicula vulgaris*) found in northern regions, the leaves of which are used by the Lapps to curdle reindeer's milk and also to thicken fresh warm milk so that it will neither curdle nor form cream afterward but makes a palatable tenacious mass, a small portion of which will act similarly upon another quantity of fresh milk, and the *Acanthaceae* containing several plants cultivated as ornamentals like the *Thunbergia alata* and the *Ruellias*. The *Ruellia ciliosa* is used as a substitute for *Spigelia* and is common in the central states to Iowa and Southward. The *Harleria Priankis* of Siam is used for snake bites. The *Strobilanthes callosus*, a shrub found in India, is an irritant poison, according to Major Kirtkar, these irritant properties being due to hairs on the leaves.

FAMILIES OF TUBIFLORES

Corolla generally regular

Ovary not 4 lobed, ovules 2 or more.

Style 1, ovary 2-celled, fruit a berry or capsule

S. 2011-1154

Style 1, entire 2-cleft or 2-parted, frequently twining plant.

Caryophyllaceae

Style 1, 2-lobed or 2 parted herbs not twining

Hydrophyllaceae

Ovary generally 4-lobed

Вот так nasce

Corolla generally irregular

Ovary generally 4 lobed

Labalgae

Ovary not 4-lobed!

Placentae axillary

Scrophulariaceae

Placentae parvulae

Figoniaceae

CONVOLVULACEAE

Chiefly twining or trailing herbs, shrubs or trees, some with milky juice; alternate leaves without stipules; flowers regular and perfect; calyx inferior, 5-parted or 5-divided; a 5-lobed or plaited corolla, convolute or twisted in the bud; stamens 5, inserted on the tube of the corolla and alternate with its lobes; a 2-celled, rarely 3-celled ovary with a pair of erect ovules in each cell. About 900 species of wide distribution, but chiefly in warm regions. Contains a number of important economic plants, among them the sweet potato (*Ipomoea*

Batatas), which has been widely cultivated in all tropical and sub-tropical countries and is a well known and cultivated plant in the United States. There are many varieties. In the South the large varieties are called yams, but these should not be confused with the Chinese yam (*Dioscorea*), which forms an important article of food in tropical countries, in the islands of the Pacific and in New Zealand. The man of the earth (*Ipomoea fastigiata*) was used by the Indians as food. It has properties similar to Jalap, for which it is sometimes substituted. The *Ipomoea Jalapa*, native to Mexico, produces a large root which is also a purgative and contains a glucoside *convolvulin* $C_{41}H_{56}O_{16}$. Several plants of the order, like the moon-flower (*Ipomoea Bona-nox*), the morning-glory (*Ipomoea purpurea*) and cypress vine (*Ipomoea Quamoclit*) are cultivated for ornamental purposes. Several, like Jalap (*Ipomoea Purga*) and (*I. congesta*) are used in medicine and are strong purgatives. The scammony (*Convolvulus Scammonia*) of Western Asia is also used as a purgative. the milky juice being collected when hard, it contains *scammonin* $C_{16}H_{28}O_{10}$. *Convolvulus scoparius* and *C. floridus*, furnishing the oil of rhodum, are small shrubby species of the Canary Islands. The wood is strongly scented and the oil is used to adulterate attar of roses.



Fig. 401. Sweet Potato (*Ipomoea Batatas*). A well known cultivated plant in tropical and sub-tropical countries. (From Vauquelin's *Traité de Botanique*.)




Fig. 402. Jalap (*Ipomoea Purga*). Plant and tuber. Purifies a strong purgative (From Vosque's Traité de Botanique).



Fig. 403. Man-of-the-Earth (*Ipomoea fastigiata*). Used by the Indians as food. (M. Spangh & Co. by)




Fig. 404. Dodder. To the left—Field dodder (*Cuscuta arvensis*). a, flower, b, flower spread apart, c, capsule with stamens and styles, d, seed. To the right—A. falla dodder (*C. spithameum*). a, flower, b, flower spread apart to show stamens and corolla, c, capsule showing styles, d, seed. (Dewey, I. S. Dept. Agr.).

The dodders (*Cuscuta*) belong to the family *Cuscutaceae* and are of interest in this connection mainly because they are parasitic on clover, alfalfa and other plants. Those growing on the above named plants are *Cuscuta arvensis* and *C. Epithymum*, the flax dodder (*C. Epilinum*) occurs on flax. The plants contain *cuscutin*. From two independent sources in this country there have come reports that clover dodder is injurious. Dr J. L. Taylor of Lisbon, Ohio, wrote Prof. A. D. Selby that when dodder was fed to horses it caused bowel trouble. Dr E. H. Jenkins of Connecticut states that cattle were seriously troubled with scours when clover hay in which this plant occurred was a part of the ration. He could not determine whether it was the dodder or the mould which had developed because the dodder had so matted with the clover the hay was not cured well. When this hay was cut out from the ration the trouble ceased.

Ipomoea L. Morning Glory

Twining or erect herbs. flowers large, showy, axillary, calyx 5 parted, corolla broadly bell-shaped, 5-cleft, stamens included, pistil with 2-4 celled ovary 4-6 ovules style undivided, stigmas capitate, 1, 2 or 3, fruit a globose capsule, 4-6 seeded. About 350 species, tropical, or of warm regions.

The *Ipomoea leptophylla* of the plains of Nebraska to Kansas and the Rocky Mountains produces a large root weighing from 10-100 pounds. It has an erect non-twining stem from 2-4 feet high, narrow and long leaves. large pink purple flowers.

Ipomoea fastigiata Sweet. Man of the Earth. Wild Potato

A deep rooted, smooth, trailing or twining perennial with a stout large root occasionally weighing 30 pounds, leaves heart shaped, acuminate or occasionally fiddle-shaped, peduncles 1 5-flowered, sepals smooth, ovate, oblong, very obtuse; corolla funnel-shaped, 3 inches long, tube purplish, pistil with a 2-celled ovary, stigma 2-lobed, each cell 2-seeded.

Distribution. Common in fields, dry or alluvial grounds from New England to Florida to Ohio, Michigan, Illinois, to Texas.

Poisonous properties. The large root of this plant is said to be poisonous. It contains the glucoside *ipomoein* and is purgative. The same is true of *I. leptophylla*. The *I. Purga* is regarded as somewhat poisonous.

Convolvulus (Tournef.) L. Morning Glory or Bindweed

Herbs or somewhat shrubby plants, twining, erect or prostrate, leaves generally cordate or sagittate and petioled, flowers large and axillary, calyx bractless or with a pair of bracts, sepals nearly equal, corolla funnel-form or campanulate, stamens included, inserted on the tube, style undivided or 2-cleft at the apex, stigmas 2, filiform, oblong or ovoid, ovary 1 or 2 celled with 4 ovules, fruit a capsule, 1-4 celled, 2-4 valved. About 175 species of wide distribution.

Convolvulus sepium L. Hedge Bindweed

Smooth, occasionally pubescent, twining around supports or trailing, leaves triangular, halberd or arrow-shaped, the tip acute or pointed, the basal lobes obliquely truncate or sinuate lobed, the flowering peduncles 4-angled, with 2 leaf-like bracts which are commonly acute, corolla is white or tinged with rose purple.

Distribution. The weed is common in the Mississippi Valley in the moist alluvial bottoms along streams and in fields. It is also found extensively creeping over weeds and other herbaceous vegetation and roadsides throughout the state. The species is native to North America and is common from Nova Scotia to Maine, south to North Carolina and Texas to Kansas, Utah, Montana, Minnesota and eastward. It also occurs in Europe and Asia.




Fig. 405 Bindweed (*Convolvulus sepium*) Supposed to be poisonous to swine. (After Vasey.)

Poisonous properties. The plant has a somewhat disagreeable odor. Dr. Schaffner states that it is supposedly poisonous to swine. It is more than likely that some of the same substances are found in hedge bindweed that occur in Jalap. Jalap contains the glucosides *convolvulin* $C_{31}H_{50}O_{10}$, *jalapin* $C_{34}H_{56}O_{10}$, *surpethin* $C_{31}H_{50}O_{10}$, *tampicin* $C_{24}H_{44}O_{11}$. The rootstock is rich in starch.

Convolvulus arvensis L. European Bindweed

A deep-rooting perennial, procumbent stem, twining or creeping propagates freely by underground rootstocks, the leaves from 1 to 2 inches long ovate-oblong, arrow-shaped, the lobes at the base running to a point, the flowers are borne in 1-flowered peduncles with very small leaf like bracts some distance from the flowers. flowers an inch or less long, short broadly funnel-shaped white, or commonly of a rose tinge.

Distribution. This weed has been known for a considerable length of time in eastern North America, where it has been sparingly naturalized for some time. Its distribution may be given as Nova Scotia to Ontario, New Jersey, Nebraska and Kansas.

Poisonous properties. Probably the same as the preceding. European authorities list this and the common morning glory, especially the latter as somewhat poisonous because of their purgative properties.

HYDROPHYLLACEAE Water-leaf Family

Herbs, generally hairy, with alternate leaves; perfect, regular 5-parted flowers, calyx inferior, deeply cleft or divided, corolla gamopetalous, stamens 5, inserted on the corolla, ovary superior, 2-celled, with 2 parietal placentae, styles 2-cleft or partially united, fruit a capsule, seeds generally reticulated or pitted.

A small family of 17 genera and about 160 species, chiefly in western North America. Very few plants of the family are ornamental. Some of the western species are occasionally cultivated. Of these the *Phacelia* is the most important. A few of the plants are weedy. The Yerba Santa (*Eriodictyon crassifolium*) is an evergreen shrub with funnel-shaped, white or purple flowers in cymose clusters, it contains a yellow acrid resin, a crystal in principle, *eriodictyonic acid* and *eriodictin*.

Phacelia Juss.

Mostly hirsute, hispid, or scabrous herbs, leaves alternate or the lower opposite, flowers blue or purple, violet, or white, inflorescence hispid, calyx naked at the cymes, deeply 5-parted, stamens attached near the base of the corolla, ovary 1-celled, capsule 1-celled or falsely 2-celled, seeds reticulated. About 80 species, natives of the New World.



FIG. 406. Hairy *Phacelia* (*Phacelia sericea*). The *Phacelia* is common in the Rocky Mountains. The bristly hairs produce mechanical injuries. (Charlotte M. King.)

Phacelia sericea A Gray

Herbs 6 inches to a foot high from a branching caudex, silky pubescent or canescent, or the simple virgate stems and inflorescence vilous-hirsute rather easy to the top, leaves pinnately parted into linear or narrow-oblong numerous and often again few-cleft or pinnatifid divisions, silky canescent or sometimes greenish, the lower petioled, the uppermost simpler and nearly sessile, short spikes crowded in a naked spike-like thyrsus, corolla violet-blue or whitish, stamens long exserted, capsule a little longer than the calyx.

Distribution. Common in the mountains from Colorado and northward in rather dry soil.

Phacelia Menziesii Torr

Herbs 6 inches to a foot high, at length paniculate-branched, hispid or roughish-hirsute, leaves mostly sessile, linear or lanceolate and entire, or some of them deeply cleft, the lobes few or single, linear or lanceolate, entire, spikes or spike-like racemes thyrsoid-paniculate, at length elongated and erect, corolla bright violet or sometimes white, stamens about the length of the corolla, capsule shorter than the calyx.

Distribution. Common in the Rocky Mountains from Montana to Utah and westward.

Poisonous properties. The stiff bristles upon these plants certainly produce mechanical injuries. A form of dermatitis venenata occurs after handling the plants. The writer has had abundant experience in contact with these plants in the Rocky Mountains.

BORAGINACEAE. Borage Family

Herbs, shrubs or trees, with alternate entire, rough or frequently scabrous or setose leaves; flowers perfect, usually regular, generally blue, borne in one-sided spikes; racemes, cymes or scattered, calyx inferior, mostly 5-lobed or 5-cleft; corolla short, bell or wheel-shaped 5-cleft or 5-parted, stamens as many as the lobes of the corolla and alternate with them, ovary superior, deeply 2-lobed, style entire; capsule globular, fruit forming 4 seed-like, 1-seeded nutlets, or into two 2-seeded or four 1-seeded nutlets.

A large family of wide distribution, consisting of about 80 genera and 1500 species. Some of the members are ornamental and are frequently cultivated, like the common heliotrope (*Heliotropium peruvianum*) native to Peru used for bedding and in greenhouses. Borage or horage (*Borago officinalis*), used in old gardens for ornamental purposes is an excellent honey plant. Lungwort (*Mertensia virginica*), an early spring blooming plant of the North, with handsome blue flowers, is occasionally cultivated. The Rocky Mountain *M. sibirica* is an equally handsome species. The forget-me-not (*Myosotis scorpioides*) with small but pretty blue flowers, native of Europe, is occasionally cultivated. Alkanet (*Achæna tinctoria*—*Anchusa tinctoria*) a native of southern Europe, yields a red dye used for coloring oils and wax. Other plants of this order yield a similar product and one species is known to color the wool of sheep. The roots of the common puccoon (*Lithospermum*), also yield a dye. The comfrey (*Symphytum officinale*) is used as a forage plant in Europe, but in the United States is seldom cultivated. The wood of *Cordia allba* is used by military authorities in San Juan in the manufacture of gun carriages, carpenters' benches, vases, etc.

Genera of *Boraginaceae*

Ovary not lobed	<i>Heliotropium</i>
Ovary deeply 4-lobed.	
Flowers regular	
Nutlets horizontally radiate, covered with prickles	<i>Cynoglossum</i>
Nutlets erect, armed with prickles.....	<i>Lappula</i>
Corolla irregular	
Stamens exserted .	<i>Echium</i>

Heliotropium (Tourn.) L. Heliotrope

Herbs or shrubs with entire alternate leaves, the small blue or white flowers, scattered or borne in scorpioid spikes, corolla salver or funnelliform without appendages, more or less plaited in the bud, stamens not exserted, with nearly sessile anthers, stigmas conical or capitate. Fruit 2-4 lobed, separating into 4, 1-seeded nutlets or into 2, 2-seeded carpels.

About 115 species, mostly tropical. The best known representative in the North is the cultivated *H. peruvianum*. The seaside heliotrope (*H. curassavicum*) occurs in saline soil along the Atlantic coast, to Maine, and from Mexico to Illinois and westward.

Heliotropium indicum L. Indian Heliotrope

An erect, hairy annual with hirsute or hispid stem, leaves petioled, ovate or oval and somewhat heart-shaped, flowers blue, borne in spikes, fruit deeply 2-lobed, smooth, 2 seeds in each cell.

Distribution. Common in waste places from North Carolina to Southern Indiana and Missouri. Naturalized from India.

Heliotropium europaeum L. European Heliotrope

An erect annual branched, roughish pubescent herbs, leaves oval, long petioled, narrow at the base, flowers borne in 1-sided bractless spikes, the terminal spikes in pairs, calyx spreading, the segments shorter than the corolla tube; anthers distinct, obtuse.

Distribution. Common in waste places southward. New York to Pennsylvania and Florida.

Poisonous properties. The European heliotrope contains *heliotropin*, a bitter, volatile, alkaline, poisonous alkaloid.

Cynoglossum (Tourn.) L. Hound's Tongue

Hirsute or hispid, rarely smooth herbs, alternate or entire leaves, purple, blue or white flowers in paniced racemes, calyx 5-cleft or 5-parted, corolla funnelliform or salverform, the throat clothed with 5 obtuse scales; ovary deeply 4-lobed fixed near the apex to the base of the style, roughened with short barbed or hooked prickles. About 75 species, of wide distribution.

Cynoglossum officinale L. Hound's Tongue

A coarse biennial herb, clothed with short, soft hairs, lower leaves oblong or oblong lanceolate, the upper closely sessile with a slightly heart-shaped base; racemes nearly bractless, elongated in fruit, divisions of the calyx ovate, lanceolate, acute, corolla reddish-purple, rarely white; nutlets flat on the broad upper face, splitting away at maturity.

Distribution In fields and waste places, especially eastward from New England to Quebec, Ontario, Minnesota, Manitoba, and Kansas.

Cynoglossum virginicum L. Wild Comfrey

Perennial hirsute herb with simple stem, 2-3 feet high leafless above, stem leaves lanceolate oblong, clasping by a heart-shaped base, flowers long peduncled pale blue, small, fruit broad, nutlets not margined, convex on the upper surface.

Distribution. Common in woods of the central Mississippi Valley states from New Brunswick to Ontario, Florida, Louisiana to Texas.

Poisonous properties. The common hound's tongue is suspected of being poisonous. The European species, *C. officinale* contains a powerful alkaloid, *cynoglossin* which resembles curare in its action. It also has the principle *consolidin*.



Fig. 407 Hound's-tongue (*Cynoglossum officinale*) Suspected of being poisonous. (From Da Lingdon's Weeds and Useful Plants.)

Lappula (Rivinus) Moench. Stickseed

Roughish pubescent or hairy herbs with alternate narrow or entire leaves, small blue or white flowers in racemes or spikes, calyx deeply 5-cleft or 5-parted with narrow segments, corolla salverform or funnelform.

About 40 species in north temperate regions, several of which are weedy in North America.

Lappula virginiana (L.) Greene Beggar's Lice

A coarse pubescent biennial from 2-4 feet high, lower leaves ovate, orbicular cordate, long petioled, stem leaves ovate-oblong or oval, flowers nearly white globose nutlets, flattened and barbed.

Distribution. Common especially in woods northward from New Brunswick to Wisconsin, Minnesota, Kansas to Louisiana.

Lappula echinata Gilbert. Stickseed

An erect annual from 1-2 feet high, pale, leafy, hispid with erect branches, leaves linear or linear-oblong racemes 1-sided, bracteolate, calyx segments lanceolate, corolla blue, nutlets rough-granulate or tuberculate on the back, the margins with a double row of slender prickles.

Distribution. Abundant in waste places along roadsides from eastern Canada and New England to Minnesota, Kansas and British Columbia. Weedy also in Europe, where it is native.

Lappula floribunda (Lehm.) Greene Large-flowered Stickweed

An erect perennial or biennial, rough-pubescent 2 feet or more high leaves oblong to linear-lanceolate, the lower tapering on margined petioles racemes erect or nearly so, pedicels deflexed in fruit, flowers pretty, blue, or occasionally white, nutlets scabrous on the margin with a row of flat prickles.

Distribution. Common in Saskatchewan and Minnesota and abundant in the Rocky Mountains.

Poisonous properties. perhaps the fruits of all the species are somewhat injurious to animals, causing inflammation. They are frequently found in wool.

Echium L. Blueweed

Bristly annual, biennial, or perennial, erect stems, corolla with a funnel form tube and a 5-lobed spreading border stamens many, exserted. The 30 species are natives of the old world.

Echium vulgare (Tourn.) L. Blueweed

An erect, rough, bristly biennial, 18 inches to 2 feet high, with an erect, mostly simple stem leaves of the stem linear-lanceolate sessile, flowers in cymose clusters, corolla reddish purple, changing to blue, tube funnelform, border unequal, spreading, 5-lobed, stamens 5, inserted on the tube; style threadlike nutlets roughened or wrinkled.

Distribution. A common weed along roadsides, fields, and meadows from New England to Indiana.

Poisonous properties. Probably poisonous. According to Friedberger and Frohner it causes slavering.

VERBENACEAE. Verbena Family

Herbs, shrubs or trees, with opposite or alternate leaves; perfect, regular, or somewhat irregular flowers borne in spikes, racemes, cymes or panicles, calyx generally persistent, inferior 4-5-lobed or cleft, corolla 4-5-cleft, tube cylindrical; stamens 4, didynamous or 2, ovary superior 2-4-celled or more, fruit dry or drupaceous, splitting into 1-4 nutlets.



Fig. 408. Blueweed (*Echium vulgare*). Probably poisonous. (Vasey-Schuy)

A large family which is found chiefly in tropical regions, there being about 1200 species. The wood of some of the tropical members of the family is valuable. Among these is the Indian teak tree, (*Tectona grandis*), native to India. The wood is hard, heavy and very durable and is used in ship-building and for railway cars. The *Uiter litoralis*, a large tree, native of New Zealand, produces a hard and heavy wood. The fragrant leaves of *V. Negundo* are used to stuff pillows.

This family includes many ornamental plants, the most commonly cultivated in green houses being the *Clerodendron Thompsonae*, which has a showy crimson corolla and white calyx. The lemon verbena, (*Lippia citriodora*), a stiff, branching shrub from Chili, is commonly cultivated in the south on account of its very fragrant leaves, an oil, from it, contains *citral*. The fog-fruit, (*L. nodiflora*), is valued in Egypt as a lawn plant, proving a success where all grasses or other plants tried for that purpose have failed, the lawns lasting five or six years without renewal. The *Lippia mexicana*, used in medicine, contains *lippiol*. The extensively cultivated verbena of our flower gardens is the *V. Aubletia*, which is produced in many colors, the species being native from southern Indiana to Texas. Lantanas are also cultivated. The French mulberry, (*Callicarpa americana*), is said to be poisonous. The vervains (*Verbena stricta* and *V. hastata*) with blue flowers, are common pasture weeds. The white vervain (*V. urticifolia*) contains a bitter glucoside.

LAMIATAE. Mint Family

Chiefly aromatic herbs, some shrubs and trees, with square stems; opposite leaves without stipules. Flowers with cymose inflorescence, perfect, irregular, more or less 2-lipped, calyx 5-toothed or 5-lobed, corolla 4-5-lobed, commonly 2-lipped, upper 2-lobed or entire, the lower usually 3-lobed, stamens borne on the corolla tube, ovary superior, deeply 4-lobed or 4-parted, in fruit forming 4 small seed-like nutlets or achenes surrounded by the persistent calyx, usually exalbuminous or with some albumin. A large family of about 3000 species

of very wide distribution. Many of these plants are used in medicine. Lavender (*Lavandula vera*) of the mountain regions of northern and eastern Europe, is cultivated for the oil of lavender, much used in perfume and contains *linalool* and *geraniol*. Spearmint (*Mentha spicata*) a fragrant perennial plant of Europe is used by confectioners and in the manufacture of perfumed soap. The volatile oil contains *carvol* $C_{10}H_{14}O$. Peppermint (*Mentha piperita*) a native to Europe and naturalized in North America, is cultivated in New York and Michigan for the manufacture of peppermint oil. It contains a volatile oil and *menthol* $C_{10}H_{20}O$, and is used for flavoring mutton and sweetmeats and as a cordial. It is a stimulant. The Japanese peppermint is obtained from *M. arvensis* var. *piperascens*. Pennyroyal (*Mentha Pulegium*), native to Europe, is used for the same purposes for which peppermint is used. Garden thyme (*Thymus vulgaris*) is a small erect woody shrub of southern Europe, it is fragrant, has a pungent taste and contains *thymol* $C_{10}H_8ON$, which is used as an antiseptic. It also contains *cymene*, *borneol* and *linalool*. Rosemary (*Rosmarinus officinalis*), an evergreen shrub of Europe, is chiefly used as a perfume. This and *Lavandula Spica* both contain *borneol* $C_{10}H_{18}O$, one of the pinene group of terpenes and *camphene*. Germander (*Taenidium canadense*) is a stimulant and has aromatic properties. Horse mint (*Monarda fistulosa*) is a stimulant and is used to remove colic pain. Oswego tea (*Monarda didyma*) is used as a substitute for tea, and catnip is used for the same purpose. *M. punctata* contains *thymol*, *carvacrol*, etc. Horehound (*Marrubium vulgare*), a perennial weed native to Europe, is a stimulant and tonic and is also slightly laxative. It contains the bitter principle *marrubidin*. Sweet basil (*Ocimum Basilicum*) of India is a strong aromatic herb used for culinary purposes and in the manufacture of Chartreuse liquors while the mucilaginous



Fig. 409. Horse Mint (*Monarda fistulosa*). A common roadside weed with pungent properties. (Charlotte M. King).

'seeds" are used for genito-urinary troubles. Savory (*Satureia hortensis*) an annual herb of Europe, sparingly run wild in western United States, is used as a pot herb. The patchouli (*Pogostemon Heyneanus*) is used by the Arabs to scent mattresses and shawls. Sage (*Salvia officinalis*) is cultivated as a pot herb and contains *pinene*, *cineol*, *thujone* and *borneol*. Marjoram (*Origanum majorana*), a perennial plant native to South of Europe, and cultivated as an aromatic herb, is also an excellent honey plant containing *citral*. Many species of mints, like the brilliantly colored South American sage (*Salvia coccinea*, *S. splendens* etc.) and others native to the southern states and the west, are cultivated for ornamental purposes. The lance leaved sage (*S. lanceolata*) is a troublesome weed in the West. *S. officinalis* of Europe, a stimulant and tonic contains *cineol* and *safrol* $C_{10}H_{12}O$, and is commonly used with meat and sausage in German communities. The Japanese potato (*Stachys Sieboldii*), is used as the Jerusalem artichoke is. Sweet balm (*Melissa officinalis*) a pot herb, contains a bitter principle.

Cymol, $C_{10}H_{14}O$, one of the *benzol* derivatives, occurs in many of the Labiales, especially in *Thymus officinalis*, *Monarda punctata*, and *Micromeria punctata*. *Thymol*, a *benzol* derivative, is also found in *Origanum floribundum* and *Monarda citriodora*.

Genera of Labiales

Calyx rigid, spiny, pointed	3 Leonurus
Calyx not spiny toothed.	
Anthers approximate.	
Fertile stamens 2	.. 2 Hedysarum
Fertile stamens 4	
Calyx tubular curved	1 Nepeta
Calyx bell-shaped.	
Aromatic herbs, stamens exserted	5 Mentha
Not aromatic herbs ascending under the upper lip	4 Lamium

1. *Nepeta* L. Catnip

Herbs with dentate or incised leaves, flowers white or blue, in clusters, calyx tubular corolla 2 lipped, lower lip spreading, 3-lobed. About 150 species native to Europe and Asia.

Nepeta Cataria L. Catnip

A perennial erect herb, 1-3 feet high leaves ovate, cordate, coarsely serrate, petiole, whitish, downy underneath, flowers in cymose clusters corolla whitish, dotted with purple.

Distribution Native to Europe, widely naturalized in northern states.

Nepeta hederacea (L.) Trevisan. Ground Ivy

A creeping, trailing perennial, with leaves all alike, petioled, round, kidney-shaped, crenate smooth green on both sides, flowers light blue in axillary whorls of about 6, appearing in early spring and summer.

Distribution Native to Europe, widely naturalized in the northern states, especially in shady places.

Poisonous properties. According to Dr. Schaffner, Ground Ivy is said to be poisonous to horses. Contains a volatile oil and a bitter principle. The common catnip also contains a volatile oil and bitter principle.



Fig. 410. Ground Ivy (*Veronica hederacea*). A somewhat weedy plant naturalized from Europe. Said to be poisonous to horses. (From Johnson & Med. Bot. of N. A.)

2. *Hedeoma* Pers.

Aromatic pungent herbs, leaves small, flowers in axillary clusters, crowded into terminal spikes or racemes. calyx ovoid or tubular, bearded in the throat, 2-lipped, the upper lip toothed. corolla 2-lipped, the upper 2-lobed, the lower spreading, 3-cleft. fertile stamens, 2, the upper pair reduced; sterile filaments or wanting, nectaries ovoid, smooth.

Hedeoma pulegioides Pers. American Pennyroyal

An erect, branching, hairy annual, leaves ovate to ovate-oblong, petioled, sparingly serrate. whorls few flowered. upper calyx teeth triangular, gibbous, corolla bluish, rudimentary stamens, evident but not usually anther-bearing.

Poisonous properties. It has been regarded with suspicion. It has the odor and taste of true Pennyroyal. *Hedeoma pulegioides* contains a volatile oil, hedeomol $C_{10}H_{16}O$.

Leonurus L. Motherwort

Tall herbs with palmately cleft or dentate leaves, flowers small, white or blue, in axillary clusters, calyx tubular, 5-nerved and 5 rigid teeth. corolla 2-lipped. About 10 species in the old world.

Leonurus Carduica L. Motherwort

Tall perennial herb with erect stem, 2-6 feet high, leaves long petioled, the

lower round and palmately lobed, the upper crenate at the base, 3-cleft flowers pale purple in close axillary whorls, corolla bearded.

Distribution. Native to Europe, widely naturalized in the northern states.

Injurious properties. The stiff bristles of the calyx are often injurious, producing mechanical injuries. It contains a volatile oil with an unpleasant odor, a bitter principle, etc.

Lamium L. Dead Nettle

Decumbent herbs, leaves usually cordate, doubly toothed flowers small axillary and terminal clusters, calyx tubular, 5-toothed, the teeth nearly equal, the upper ones larger, corolla dilated at the throat, upper lip ovate or oblong arched, narrowed at the base, the middle lower lip spreading, the lobe notched at the apex stamens 4, small, the anthers nearly in pairs, nutlets truncate. About 40 species in the old world some troublesome weeds like *L. amplexicaule*.

Lamium amplexicaule L. Dead Nettle

An annual or biennial herb with rounded, deeply toothed crenate leaves, the upper leaves small, clasping, calyx tubular, 5-toothed, flowers small, corolla purple, upper lip bearded, the lower spotted. The *L. album* is perennial, has larger flowers, and has slender calyx teeth.

Distribution. Common in the eastern states to Missouri. The *L. album* escaped and not infrequent westward.

Injurious properties. The dead nettle is regarded as injurious.



Fig. 411. Dead Nettle (*Lamium, album*), frequently escaped from cultivation. (Ada Hayden).



Fig. 412. Dead Nettle (*Lamium amplexicaule*). Regarded as poisonous. (Selby).

Mentha (Toora.) L. Mint

Herbs with the odor of mint. Leaves usually with punctate spots, flowers small in whorled clusters, pink or white, calyx bell-shaped or tubular, 5-toothed, corolla tube shorter than the calyx, limb 4-cleft, stamens equal, erect, included or exserted, filaments smooth, anthers ovoid, smooth. About 30 species of temperate regions. Our native species (*M. arvensis*, var. *canadensis*) is common in low marshy ground.

Mentha crispata contains *hnacal*, $C_{10}H_{16}O$, one of the terpenes. It may be mentioned in passing that this same substance recurs in *Orimum*, *Thymus* and *Darwinia*. A ketone carbon, $C_{10}H_{14}O$, is found in several species of the genus.

Mentha piperita L. Peppermint

Smooth, erect perennial herb with creeping rootstocks from 1-2 feet high, leaves petioled, ovate, oblong to oblong lanceolate, acute and sharply serrate, flowers whorled in interrupted loose, leafless spikes, purplish or whitish.

Distribution. Commonly escaped from cultivation and troublesome in the East.



Fig. 413. Peppermint (*Mentha piperita*). One of the sources of the peppermint of commerce. (From Vesque's *Traité de Botanique*.)



Fig. 414. Tomato (*Lycopersicon esculentum*). An important food plant (W. S. Dudgeon.)

SOLANACEAE. Nightshade Family

Herbs rarely shrubs, vines, a few of the tropical species, trees with alternate leaves without stipules, flowers regular or nearly so, borne in cymes, calyx inferior, 5-lobed, corolla gamopetalous, generally 5-lobed, stamens as many as the lobes of the corolla and alternate with them, inserted on the tube, generally equal, style and stigma 1, ovules numerous, fruit a berry or capsule. A large family chiefly tropical, consisting of 70 genera and 1600 species. Several of these are important medicinal plants and several important food plants. Many plants of the order have poisonous properties.

The red or Cayenne pepper (*Capicum annuum*), native to North America, is much used in tropical countries to stimulate the appetite. It seems to run to many forms, but has been shown by Prof. Irish that many of the so-called species belong to *C. annuum*. A shrubby species, *C. frutescens* is native to western Texas and Mexico. The tobacco (*Nicotiana Tabacum*), was undoubtedly native to America. It was used by the Indians in North America at the time of the discovery by Columbus, and was introduced into England in 1585 by Lane who was a deputy of Sir Walter Raleigh. Tobacco is now cultivated in many civilized countries, as Cuba, Philippine Islands, Sumatra, also in Florida, the Carolinas, Connecticut, Kentucky, and Wisconsin in the United States. It is used for making cigars, snuff and for chewing. It is an important article in commerce. The potato (*Solanum tuberosum*) is indigenous to Peru and Chili but was introduced into Spain about the beginning of the 16th century, and into England from Virginia in 1586 by Sir Walter Raleigh. The greatest yields occur in irrigated districts. It is one of the most important food




Fig. 414a. Tobacco Plant (*Nicotiana glauca*). a. Flower, b. corolla, cut open; c. ovary; d, e young fruit. a, b, c natural size; d, e, x 2. (After Strasburger, Nail, Schenck, and Schimper).

plants. Other species of tuberous *Solanum* occur, as the *S. Jansoni*, in southwestern United States, and several others in Mexico and South America. The egg-plant (*Solanum Melongena*) is native to India, but is now widely cultivated in tropical countries and temperate regions, the fruit being used for culinary purposes. The tomato (*Lycopersicon esculentum*), a native of tropical America, of which there are many varieties, is now widely distributed in tropical and temperate regions. The fruit is eaten fresh or canned, or made into

various products. The currant tomato (*L. pimpinellifolium*), a fruit about the size of a currant, is cultivated as a curiosity. Ground cherries are much cultivated. The strawberry tomato (*Physalis Alkekengi*), a perennial native to southern Europe, has a large fruiting calyx which turns red. The fruit is eaten, but it is not so palatable as the fruit of the Cape gooseberry (*P. peruviana*) which is native to Peru. The fruit of a native species (*P. pubescens*) is also eaten, but this also is inferior to the Cape gooseberry.

Beladonna is derived from *Atropa Belladonna*, a tall glabrous or slightly downy herb, with a perennial rootstock, native to southern Europe and eastward to Asia Minor. This furnishes the *atropin* of commerce used for dilating the pupil of the eye. The earliest investigations of the alkaloids of belladonna were made by Hauberlein in 1809, who first determined their presence. Kise was the first to find *atropamin* $C_{11}H_{17}NO_3$ in the roots of the plant, although Schmidt denies the presence of *atropin*, affirming the presence of *hyoscyamin* only. *Belladonnin*, an isomer of the above, is probably also present. The root of *Atropa Belladonna* sometimes contains from 4/10-1% of the alkaloid and the leaves about half as much. The greatest amount of the alkaloid occurs during the flowering period. *Pseudo-hyoscyamin* occurs in the roots of *Mandragora*. The alkaloids *mandragorin*, $C_{11}H_{15}NO_2$, and *monacin* $C_{22}H_{33}N_2O_{10}$, occur in *Brunfelsia Hopeana*. Several other undetermined alkaloids, such as *purpurebin* have also been found in the family.



Fig. 415 Ground Cherry (*Physalis peruviana*) Cultivated for its fruit. (W. S. Hodgeon)

The mandrake (*Mandragora officinarum*) of the ancients was, at one time, supposed to have medicinal virtues. The flowering tobacco (*Nicotiana glauca*) is a well known cultivated plant of Brazil. The thorn apple (*Datura Metel*) is much cultivated, as is the *D. meteloides*, which has large, sweet-scented flowers. The bittersweet (*Solanum Dulcamara*), the hardy annual (*Schizanthus pinnatus*) the *Brunfelsia latifolia*, which has fragrant flowers, and the *Cestrum*



Fig. 416. Sleeping or Deadly Nightshade (*Atropa belladonna*). Top of flowering and fruiting branch; entire fruit; cross section of fruit; corolla cut open and spread out. Source of the belladonna of commerce. (From Vauque's *Traité de Botanique*.)

regions are frequently cultivated. The odor from the flowers of the latter is very overpowering. The berries of *C. pallidum* are said to be poisonous but birds have scattered the plant very widely in the tropics. *Petunia violacea*, *Lycium halimifolium* and *L. chinense* are cultivated. The *Duboisia myoporoides* of Australia is a tall shrub, its leaves having narcotic properties and containing the substance *duboisin*, a mixture of *hyoscyamine* and *atropine* producing an action like that of *hyoscyamine* but more hypnotic. According to Maiden this plant is poisonous to stock. Other species like *D. Leichardtii* contain the same substance. The pituri (*D. Hopwoodii*) contains a liquid volatile alkaloid *piturin* C_8H_9N resembling *nicotine*. The natives mix the pituri leaves with the ashes of some other plant and chew them like tobacco. In its action it resembles *nicotine*.

The *scopola* (*Scopola carnioleica*) of Austria and Hungary, is a perennial herb used like *Belladonna* in medicine. The leaves and rhizomes of this species and *S. japonica* are poisonous. The *S. carnioleica* plant contains *atropine* $C_{17}H_{21}NO_3$, *hyoscyamine* and *scopolamine*. The latter substance is broken up into *scopoline* $C_{14}H_{17}NO_2$ and *tropic acid* $C_8H_9O_3$. The *hyoscin* $C_{17}H_{23}NO_2$ is impure *scopolamine*. *Scopoline* causes dilation of the pupils, the heart action is at first diminished, then increased, due to the stimulation of the inhibitory nervous apparatus. The pichu used in kidney troubles is the dried leafy twigs of the Chuban shrub (*Fabiana imbricata*). The tree tomato (*Cyphomandra betacea*) produces a fruit similar, in taste, to that of the common tomato, if eaten when raw, but after it is stewed, provided the skin and seeds have first been removed, an apricot-like flavor is produced. It is much used in carts and pastry in the mountainous districts of the tropics.



Fig. 4-7 Cuckoo tree (*Dubautia myosotoides*). A shrub, bearing leaves that possess narcotic qualities. (After Fournet)

Genera of Solanaceae

Fruit a berry

Corolla wheel shaped.

 Anthers opening by uplifted valves

 Anthers opening longitudinally, widely spreading

Corolla not wheel shaped

Corolla funnel form

Fruit a capsule.

Calyx urn shaped somewhat irregular

Calyx prismatic, corolla funnel form

Calyx tubular.....

..1 Solanum

2 Capsicum

5 Nicotiana

7 Lycium

4 Hyoscyamus

6 Datura

5 Nicotiana

1 *Solanum* (Tourn.) L. Nightshade

Herbs or shrubs, often with stellate pubescence, flowers in cymose, umbellate, or racemose clusters; calyx bell shaped or rotate generally 5-parted or 5-cleft, corolla rotate 5-lobed or cleft, plaited in the bud, stamens exserted, filaments short inserted on the throat, anthers converging around the style opening at the top ovary usually 2-celled, stigma small, berry with persistent calyx at the base or enclosing it, seeds numerous. About 900 species of wide distribution. Several are troublesome weeds as horse nettle (*Solanum carolinense*) and buffalo bur (*S. rostratum*). The potato (*S. tuberosum*) and egg plant (*S. Melongena*), are cultivated.



Fig. 412. Common potato (*Solanum tuberosum*). The potato under some conditions is very poisonous, especially when the tops are green. (Linn Fammel)

Solanum Dulcamara L. Bittersweet

A more or less pubescent perennial, stem climbing or straggling, somewhat woody below. leaves petioled ovate or hastate, the upper usually halberd shaped. flowers purple or blue in cymes, berry globose, red.

Distribution. Naturalized from Europe, New Brunswick to New Jersey, Pennsylvania, Kansas to Minnesota.

Poisonous properties. The plant contains the alkaloid *solanin* $C_{28}H_{47}NO_8 + H_2O$, with a hot, bitter taste, *dulcamarin* a bitter principle $C_{28}H_{47}O_{10} + 20H_2O$ and the alkaloids *solanidin* $C_{40}H_{61}NO$ and *solanine* $C_{52}H_{83}NO_{11}$.

Chestnut says with reference to this plant

Resides *solanin* (0.3 percent), this plant contains another less poisonous compound *dulcamarin*, which gives it its peculiar bitter-sweet taste. Neither of the compounds is abundant. The berry, though its taste is not remarkably disagreeable, is somewhat poisonous, and it has been shown that an extract of the leaves is moderately so. The plant has nevertheless caused some effect. The treatment is the same used in case of the above species.

According to Schimpfky the berries of this plant have been used to poison dogs and the juice of the fruit acts as a poison to rabbits. Fluckiger and Hanbury in their Pharmacographia, make this statement with reference to *Dulcamara*.

Dulcamara is occasionally given in the form of decoction, in rheumatic or cutaneous affections, but its real action, according to Garrod, is unknown. This physician remarks that it does not dilate the pupil or produce dryness of the throat like belladonna, henbane or stramonium. He has given to a patient 3 pints of the decoction per diem without any marked action, and has also administered as much as half a pound of the fresh berries with no effect.

Johnson in his Medical Botany of North America refers to the use of the plant as follows

Bittersweet in full doses, produces a certain amount of cerebral disturbance of a narcotic character, together with dryness of the throat, and sometimes an erythematous eruption of the skin, with a tendency to diaphoresis. It has been employed with benefit in a variety of cutaneous eruptions in muscular rheumatism, and in chronic bronchial and pulmonary affections.



1) Bittersweet (*Solanum Dulcamara*). a, Flowering spray, b, fruit—both one-third natural size. Berries somewhat poisonous. (U. S. Dept. Agr.)

Lehmann states that it is a narcotic poison when given in large doses even causing death in rabbits.

It will be seen from the above quotations that the plant is not a violent poison and yet ill effects are probably produced by it under some conditions.

Solanum nigrum L. Common Nightshade or Stubbleberry

Annual, low branched and often spreading, stem glabrous or hairy, hairs simple roughened on the angles, leaves ovate, petioled, flowers white in small, umbel-like drooping lateral clusters calyx spreading, the lobes obtuse, much shorter than the white corolla, berries glabrous, globose, black, occasionally large.

Distribution. Found in northern United States. Also occurs in Europe. Shady grounds and fields. A cosmopolitan weed.

Poisonous properties. Stubbleberries are occasionally cultivated for their fruit. They are sometimes sold as huckleberries and used for pies and pre-



Fig. 430. Black Nightshade (*Solanum nigrum*) Unripe fruit thought to be poisonous. (Charlotte M. King)

serves. The fruit should, however, be used with caution, especially before it is ripe. A decoction of the ripe stubbleberry when fed to cats does not appear to be poisonous according to Dr. Buchanan, who fed considerable quantities to cats in the writer's laboratory without injurious effects. Mr. Chesnut says

The amount of poison present in any part of this plant varies with the conditions of growth. The more musky-odored plants are the most poisonous. In some, the amount of alkaloid in the ripe fruit and leaves is so small that these parts may be, and are, consumed in considerable quantity without any ill consequences. Poisoning does sometimes follow, but it is not clear whether this is due to improper preparation or to careless selection of the parts used. The use of black nightshade for food is certainly not to be recommended. Cases of poisoning are recorded for calves, sheep, goats and swine.

The characteristic symptoms are about the same in man and animals. They are stupefaction, staggering, loss of speech, feeling and consciousness, cramps, and sometimes convulsions. The pupil of the eye is generally dilated. Death is directly due to a paralysis of the lungs, but fortunately few cases are fatal.

In June, Dr. Fackinger reported from Greenfield, Iowa, several cases of sheep poisoning probably due to eating plants of *Solanum nigrum*. A subscriber of the Iowa Homestead also says that for a number of years poisoning has affected his

sheep in a pasture where this weed has been abundant. Dr Koto, Iowa State Veterinarian, also reports cases due to this plant.

This Black Nightshade contains the alkaloid *solanin* $C_{25}H_{47}NO_{11}$ which is probably present in larger quantities in the fruit before it is entirely ripe, also *solanidin* $C_{25}H_{47}NO_{11}$ with stronger basic properties. From all the evidence I can get, I may say that the fruit should be eaten with caution. In Europe it has been looked upon with suspicion for a long time. The ancients held it in suspicion and many superstitious beliefs were connected with it. Schimpfky in discussing the poisonous and non-poisonous action of the plant states that the amount of poison produced depends upon climatic conditions and the character of the soil. In some places it may be entirely harmless and in other places poisonous. In Europe the plant is sometimes used as a salad plant, but the author above referred to, remarks that when the odor is unpleasant and disagreeable it should not be eaten.

According to Lehmann, Schreber and Haller the berries are poisonous to ducks and chickens. Cases of poisoning from the berries of this plant have been recorded in Europe by Hirtz,¹ Manners,² etc.



Fig. 421. Spreading nightshade (*Solanum triflorum*) one-third natural size. Suspected of being poisonous. (U. S. Dept. Agr.).

¹ Gas. Med. d. Strasbourg. 1842.

² Edin. Med. Jour. 1847.

For these and other references see Blyth "Poisons" 4th ed. 598. Poisoning by bitter sweet berries is recorded in Lancet, 1856. Berries of *S. tuberosum*, Brit. Med. Jour. 1895.

Solanum triflorum Nutt. Three-flowered Nightshade

Annual, low spreading, slightly hairy or nearly glabrous, leaves acute, pinnatifid, 7-9 lobed, peduncles 1-3 flowered, corolla white berries greenish or inclined to blackish, about the size of a small cherry pedicels reflexed in fruit.

Distribution. On the plains, and waste places from Nebraska and Kansas to Northwest Territory and Arizona, introduced eastward.

Poisonous properties. Prof. Chesnut says experiments on guinea pigs show that the berries are poisonous. No cases of human poisoning have been reported. The berry is not attractive to the eye, but has an agreeable odor and taste. It is therefore to be suspected in cases of poisoning which occur in localities where the weed is abundant. The writer has also received complaints of the poisoning by this plant from Nebraska and other western states. The active constituent is, no doubt, *solanum*.

Solanum rostratum Dunal. Buffalo Bur Sand Bur

Herbaceous, woody when old, somewhat hoary or yellowish 8 inches to 2 feet high covered with copious stellate pubescence, the branches and stems covered with sharp yellow prickles, leaves somewhat melon like 1-3 times



FIG. 422 Buffalo Bur (*Solanum rostratum*). a, branches of the plant with burs; b, yellow flowers c and d, seeds, c, very much enlarged; d, natural size. The prickles cause mechanical injuries to stock.

pinnatifid, lobes roundish or obtuse and repand, covered with soft pubescence. hairs stellate. flowers yellow, corolla gamopetalous, 1 inch in diameter, nearly regular the sharp lobes of the corolla broadly ovate, stamens, 5, declined. anthers tapering upward, linear lanceolate, dissimilar, the lowest much larger and longer with incurved beak, hence the technical name *rostratum*. style much declined. fruit a berry but enclosed by the close fitting and prickly calyx, which has suggested the common name buffalo bur or sand bur, pedicels 1. fruit erect. seeds thick, irregular, round or somewhat longer than broad, wrinkled, showing numerous small pits, seeds surrounded by a gelatinous substance. The related species *S. citrullifolium* of the southwest is glandular pubescent with slender yellow subulate prickles, lowest anther violet.

Distribution. The buffalo bur was undoubtedly a native to the region of the




FIG. 4. a. Horse nettle (*Solanum rostratum* L.) a plant with flowers and fruit, b. flower, c. seed enlarged. Considered poisonous by Bessey and others. (U. S. Dept. Agr.)

plains occurring in the bare places where grass is scant and in former times was most abundant around the "buffalo wallows." Its range is from New Mexico to Wyoming and across the plains. The general traffic from the west to the east has caused the weed to be distributed in various eastern and middle states, Iowa, Minnesota, Wisconsin, Missouri, Illinois, Indiana, Ohio, New York, Massachusetts and Tennessee.

Injurious properties. This plant is carefully avoided by stock, but so far as known is not poisonous although the prickles on the plant produce mechanical injuries. When the prickles enter the tissues of animals inflammation occurs and pus is formed.

Solanum carolinense L. Horse Nettle

Horse nettle is a deep rooting perennial, propagating freely by its underground roots these running roots are often 3 feet long, stem from 1-2 feet high somewhat straggling, half shrubby at the base, stems hairy or merely roughish with minute hairs which are usually numerous, leaves oblong or sometimes ovate, obtusely sinuate, toothed or lobed or deeply cut, 2-4 inches long flowers borne in racemes which later become 1 sided the outer part of the flower the calyx consists of slender lobes, the corolla is light blue or white an inch or less in diameter and resembles that of the common potato. the flowers are followed by yellow globose berries, 1-2-3-4 inch in diameter, the small seeds are yellowish, a little less than 1/12 of an inch long, minutely roughened.

Distribution. Its distribution in North America is from Connecticut through New York, Pennsylvania and New Jersey, West Virginia along the Atlantic seacoast to Florida, west along the Gulf Coast to Texas, through Kansas, Nebraska, Iowa, Illinois and Michigan.

Poisonous properties. The root as well as the fruit of the plant has a very disagreeable narcotic odor according to several authors, the plant is poisonous. Dr. Bessey reports it as possibly poisonous. It contains *solanin*, according to Kraemer, 0.8 per cent in the berries.

Solanum elaeagnifolium Cav. Horse-weed. Bull Nettle

A deep-rooted spreading perennial from 1-3 feet high stem silvery canescent, finely pubescent leaves lanceolate, oblong or linear, petioled entire or repand-dentate flowers in cymose clusters, peduncle stout and short corolla gamopetalous, blue, calyx lobes lanceolate, berry yellow, smooth globose.

Distribution. Common on the prairies of Kansas to Texas and New Mexico.

Poisonous properties. The berries of this fruit are used to curdle milk in northern Mexico and southern Texas. They are crushed into a powder, put into a muslin bag, suspended in the milk until coagulation occurs. It is also used as a medicine by the Mexicans.

Solanum tuberosum L. Potato

An erect herb, cultivated as an annual for the esculent tubers leaves pinnate of several ovate leaflets and smaller ones between, flowers blue or white, berries round, green.

Distribution. Native to Chile north to Mexico and Arizona. Introduced into Europe between 1580 and 1585.

Poisonous properties. The wilted green stem and leaves are poisonous, containing the alkaloid *solanin*. The water from boiled potatoes contains a poisonous substance. Some persons cannot eat potatoes because poisonous to them. According to Kassner, healthy potatoes do not contain *solanin* but diseased potatoes contain this substance, from 150 gms. he separated 30-50 mgrs. of *solanin*. It is probable that this substance occurs in other species of *Solanum*. Thos. Maiden states that the *S. eremophilum* poisons sheep and cattle when they eat the tops. Friedberger and Fröhner state that potato tops are injurious, that the diseased animals show symptoms resembling foot and mouth disease.

Dr Doerr has recently reported a case of poisoning in a cow that had been fed exclusively upon refuse from the Club-house kitchens near the Iowa State College Campus. These contained potato parings among other things and to these the poisoning was charged. The trouble was diagnosed as gastro-enteritis. The post-mortem revealed diffuse intestinal hemorrhages with enlarged liver and spleen.

2. *Capsicum*. Pepper

Herbs or shrubs with sharp taste, leaves fleshy, flowers white, corolla wheel shaped, 5-lobed, tube short, stamens separate with filaments longer than the heart shaped anthers which open longitudinally, fruit a berry. The Guinea pepper and the Indian goat pepper (*C. frutescens*) are much more powerful stimulants than cayenne and often produce violent pain and purging. This shrub is native to the Southwest. The genus *Capsicum* has two species. A monograph by Prof Irish published in the Missouri Botanical Garden Reports describes many of the varieties.

Capsicum annuum L. Cayenne Pepper

Annual. Leaves ovate entire; flowers with truncate calyx and white corolla; fruit a berry oblong or globular, red or green.

Distribution. Widely cultivated. Native to the Southwest



FIG. 424. Red Pepper (*Capsicum annuum*). The fruit of this as well as the leaves are powerfully pungent. (W. S. Dudgeon).

Poisonous properties. The peppers are often used in domestic practice in making a stimulating plaster; if its action is continued long enough, however, a vesicular formation makes its appearance. In domestic animals it causes

gastro-enteritis. Death has even occurred where too much of the Cayenne pepper has been used. Thresh isolated a principle to which he gave the name of *capsaicin* $C_8H_{14}NO_2$, however, the bitter pungent substance has been called by Morbitz, *capsacutin*, a crystalline nitrogenous compound containing the substance $C_{23}H_{31}N_4O_4$. This is so powerful that 1 part in 11,000,000 will impart the pungent taste. A volatile alkaloid resembling *cocaine* has been found in small quantities.

3. *Nicotiana glauca* Adans. Apple of Peru

Tall smooth annual with alternate leaves, calyx 5-parted, angled; corolla wheel shaped or somewhat funnel shaped with widely spreading border, tube short anthers 5 connivent, fruit a somewhat 3-5 celled berry

Nicotiana Physalodes (L.) Pers. Apple of Peru

A tall smooth annual, 2-5 feet high, leaves ovate angled or sinuate toothed; flowers solitary corolla pale blue rather large, fruit a globular dry berry calyx, 5-parted, 5-angled, enlarged and bladder like in fruit

Distribution. Native to Peru but sparingly naturalized in the United States.

Poisonous properties. Said to be poisonous used as a fly poison in parts of the United States.

4. *Hyoscyamus* (Tourn.) L. Henbane

Clammy-pubescent, fetid, narcotic herbs leaves alternate, mostly lobed or pinnatifid, flowers large, calyx bell-shaped or urn-shaped, 5 lobed; corolla funnelform, oblique 5-lobed, the lobes unequal, capsule enclosed in the persistent calyx, 2-celled

About 15 species native to the Mediterranean region Medicinal and poisonous plants.

Hyoscyamus niger L. Black Henbane

Biennial or annual. Stem 1-3 feet high leaves ovate, sinuate toothed and angled the upper clasping flowers short pedicelled in one sided leafy spikes, corolla dull yellowish, reticulated, with purple veins capsule globose oblong.

Distribution. Common only eastward in waste places from Nova Scotia to Michigan, also in Montana, Utah, Idaho and the Pacific Coast

Poisonous properties. A well known medicinal plant from which *hyoscyamine* is obtained. *Hyoscyamine* is an anodyne and hypnotic and is poisonous. Dr. Cheesnut says

One or two cases are recorded in foreign literature in which stock have been poisoned by eating the plant of their own accord, but there is very little danger from it, on account of its ill odor and harsh texture.

It contains *hyoscyamine*, $C_{17}H_{23}NO_3$ causing a dilation of the pupils and having a sharp and disagreeable taste. It also contains *pseudo hyoscyamine* $C_{17}H_{23}NO_3$ another alkaloid, and *hyoscin* $C_{17}H_{21}NO_3$ the latter of which also dilates the pupils. The *hyoscyamine* resembles *atropin* in its composition and action and is obtained from the *Hyoscyamus* seed. When damp the alkaloid has a tobacco like odor and a bitter taste. According to Dr. Winslow, the *hyoscyamine* is practically *atropin* except that its mydriatic action is shorter. *Hyoscin* is a powerful depressant to the cerebral respiratory center spinal reflex centers, and motor tract. It is a cerebral sedative. According to Winslow

The tetanic stage succeeding spinal paralysis observed in *atropin* $C_{17}H_{23}NO_3$ poisoning, does not ensue with *hyoscin*. The latter alkaloid slightly depresses and slows the

heart, and does not paralyze the vagus terminations, nor depress the motor and sensory nerves or muscles. The circulation is but slightly influenced, and vasomotor depression only occurs in the latter stage of lethal poisoning. Death occurs from paralysis of the respiratory centers. Poisoning in animals is exhibited by loss of muscular power, slowing and failure of respiration, dryness of the mouth, stupor and asphyxia. The pulse may be infrequent the pupils are dilated and the skin is moist rather than dry. Delirium and convulsions sometimes occur in man. The effect of the combined action of *Hyoscyamin* and *Hyoscin* in *Hyoscyamus* is shown when we compare the drug with belladonna. *Hyoscyamus* is more of a cerebral sedative and hypnotic and less of a heart and respiratory stimulant. It is said to possess more power in overcoming spasm, and giving of cathartics, and in aiding intestinal movement. *Hyoscyamus* is also thought to exert a more pronounced antispasmodic action than *belladonna* upon the smooth muscles of the bladder and urethra.

Blyth gives the action of *hyoscyamin* as follows:

Thirty-two mgrs. ($\frac{3}{4}$ gr.) begins to act within a quarter of an hour, the face flushes, the pupils dilate, there is no excitement, all muscular motion is enfeebled and the patient remains in this state for many hours. 64.8 mgrs. would possibly be a fatal dose. The root is




Fig. 425. Black Henbane (*Hyoscyamus niger*). At the left, open corolla, and flowering branch. At the right, longitudinal section of flower. A well-known medicinal plant. (From Vesque's *Traité de Botanique*.)

more poisonous than the leaves, and the seeds of *Datura* contain a considerable quantity of *hyoscyamin*, they are often mistaken for other seeds such as poppy.

Many cases of children being poisoned by this seed are recorded. One instance is given by Schumsky where of two children who had eaten the seeds of the plant, one died before purgative action could be produced. The second child slowly recovered but growth was checked.

5. *Nicotiana* (Tourn.) L. Tobacco

Rank, viscid-pubescent narcotic herbs or shrubs; leaves alternate, entire, flowers borne in paniced racemes; calyx tubular, bell-shaped. 5-cleft, corolla funnel-form or salver form, the limb with five separating lobes, stamens 5, inserted on the tube of the corolla, dehiscence of the anthers longitudinal; ovary 2-celled, stigma capitate, capsule 2-valved, seeds numerous, small. About 50 species nearly all native of North America.

Nicotiana Tabacum L. Common Tobacco

A coarse annual from 4-6 feet high, leaves lanceolate, ovate, decurrent,

1-2 feet long, flowers paniced, rose-purple, corolla funnel-form, 2 inches long, lobes short, somewhat inflated

Distribution. Native to South America but widely cultivated introduced into Europe by the Spaniards shortly after the discovery of America. The best types of tobacco are cultivated in Cuba and Porto Rico, and this forms an extensive industry in North America, especially in Connecticut, Wisconsin, Virginia, etc.

Poisonous properties. Various opinions are expressed in regard to its poisonous properties. It is known however, that an alkaloid occurs in *N. tabacum*, *N. macrophylla*, *N. rustica* and *N. glutinosus* which apparently does not occur in any other plant.

The active principle of the tobacco leaf is the alkaloid *nicotin* $C_{10}H_{14}N_2$ which is easily extracted from tobacco by means of alcohol or water. It occurs to the extent of 6 per cent in the dry leaves. It has a sharp burning taste, is very poisonous and is said to have sixteen times the toxic power of *conium*. On application of heat *nicotin* is changed into *pyridin*, C_5H_5N , and other similar alkaloids like *picotin*, $C_{11}H_{11}N$. *Pyridin* depresses the spinal motor tract and causes paralysis of respiration. Moderate doses cause contraction of the pupil. *Nicotin* $C_{10}H_{14}N_2$ was found by Pieter and Rotshy in leaves of tobacco, also *nicotemin* $C_{10}H_{14}N_2$ and *nicotidin* $C_{10}H_{14}N_2$. According to more recent investigations the seeds of *Nicotiana* are free of *nicotin*. The following statement is made with reference to the toxicology of *nicotin* by Dr Winslow.

Nicotin is one of the most powerful and rapidly acting poisons. When swallowed, it causes, in animals, local irritation and pain in the throat and stomach, muscular tremors and weakness, on account of which the animal falls. These symptoms are followed, first by severe tonic and clonic convulsions, and then by abolition of voluntary motion and quiescence. The pupils are contracted, and there is vomiting (in the case of some animals), purging and micturition. The respiration is at first shallow and rapid, but becomes weaker and slower, and death occurs from respiratory failure and general collapse. The pulse is primarily slow and intermittent, but later becomes rapid. The treatment of poisoning consists in evacuation of the stomach, the use of tannic acid, respiratory and heart stimulants, as *strychnin*, *atropin*, and *alcohol*, together with external heat and artificial respiration. The maximum lethal dose is about one drachm of tobacco, or one minim of *nicotine* for small dogs. For horses, five to ten drops of *nicotin* or one half pound of tobacco.

Friedberger and Fechner state that animals have convulsive spasms, great muscular weakness, with acute paralysis.

The chronic effects of the use of tobacco, according to Millsbaugh, are as follows:

The effects produced upon smokers are almost nil, in the study of the drug itself, and it is only in that class of chewers who swallow the juice, that positive data could be pooled for, still here as well, we are at a loss to determine facts, for in manufacturing the narcotic, processes are used which alter the product greatly. nevertheless some few symptoms seem to be more or less common to all who have been, for protracted periods, subjected to the drug. Mental anxiety and irritability with at times confusion of ideas, distention of the pupils, ringing in the ears, increased secretion of saliva, uncertainty of speech, dryness of the throat, at times weakness of the stomach and bowels, increased secretion of urine, dry cough especially at night, precordial oppression with palpitation of the heart and at times an irregular pulse, trembling of the extremities when held long in one position, general anemic condition of the blood, spasmodic contraction or relaxation of single muscles, sensations of exhaustion and especially lassitude, sleepiness, profuse perspiration and sensitiveness to cold.

A writer in the London *Lancet* quoting from the *Therapeutic Gazette** states that the injury from tobacco smoke comes largely from the inhalation of carbon monoxide. Cigarette smoking is more harmful than smoking a pipe because more of the gas is inhaled.

* Vol. 33 790.

According to Zalazkas, *eserin* and *strychnin* are not antagonistic to *nicotin* but the juice of *Nasturtium officinale* counteracts it.

Nicotiana rustica L. Wild Tobacco

Annual, with obovate, petioled leaves, flowers greenish yellow paniced, longer than the calyx capsule globose

Distribution In fields and waste places from Canada to Florida, and Minnesota. Cultivated by the Indians.

Poisonous properties Probably the same as those of the preceding species.

Nicotiana glauca Link & Otto

An annual, pubescent plant from 3-4 feet high leaves lanceolate, flowers large tubular, tube 5-6 inches long the limb deeply 5-cleft

Distribution Native to Brazil but a frequent escape in gardens in the east

Nicotiana quadrivalvis Pursh Wild Tobacco

An annual 1 or more feet high leaves oblong or the upper lanceolate and the lower obovate lanceolate, acute at both ends flowers few corolla white tubular funnel-form; tube 1 inch long.

Distribution Oregon to the plains. Often cultivated by the Indians for tobacco. The allied species, *N. attenuata*, Torr., is found from Colorado to Nevada and California.

Poisonous properties The poisonous properties are probably the same as those of common tobacco. According to Maiden the *N. glauca* is poisonous to stock in New South Wales.

6. *Datura* L. Thorn Apple Jimson Weed

Rank-scented, tall, narcotic herbs, or a few tropical shrubs or trees with alternate petioled leaves large flowers calyx 5-cleft, corolla funnel-form, 5-lobed the limb plaited, stamens generally included, inserted at or below the middle of the corolla tube, ovary 2-celled, forming a capsule which is globular and prickly

A small genus of about 12 species of wide distribution, 2 being cosmopolitan weeds. Several of the species are used for ornamental purposes. Among these are the common white-flowered thorn apple (*Datura Metel*) which is native to tropical America, also the *Datura meteloides*, native to New Mexico and cultivated for its large sweet-scented flowers. Several tropical American tree-like shrubs like *Datura suaveolens* are often cultivated in conservatories. The seeds of *D. fastuosa* are used in India as a poison according to Gimlette and are commonly used on the Malay Peninsula. The *D. alba* is common in India, about Madras, and *D. atrox* occurs on the coast of Malabar. The *Daturas* are all important in India from the point of view of poisoning. The seeds of *D. alba* are often mistaken and eaten for the seeds of *Capsicum*.

Datura Tatula L. Purple Thorn-apple or Purple Stramonium

A glabrous annual from a few inches to 5 feet high, stem purplish, leaves thin, ovate, acute or acuminate, flowers consisting of a 5-toothed calyx and a 5-lobed funnel-form corolla, with stamens included, filiform filaments inserted below the middle of the corolla tube, capsule globular, prickly, 4-valved and 2-celled

Distribution. Abundant in fields and waste places from New England to

Ontario to North Dakota, Nebraska, Texas, and Florida, naturalized from tropical America.

Poisonous properties The poisonous alkaloids found in this plant are *atropin* $C_{17}H_{23}NO_3$, *hyoscyamin*, and *hyoscin*. The *daturin* is a mixture of *hyoscyamin* and *atropin*.

Professor Cheanut, in his work on the Poisonous Plants of the United States, referring to the jimson weed, says

The poisonous alkaloids, *atropin* and *hyoscyamin*, the active constituents of belladonna are found also in both of the jimson weeds. *Hyoscyamin* is the poison of the henbane and as it is identical in its physiological action with *atropin*, the above-named plants present the same symptoms of poisoning, which must be met in the same manner. The alkaloids exist in all parts of the two *daturas*. The seeds are especially poisonous.



Fig. 426. Jimson Weed (*Datura stramonium*). a, leaf and flowers, b, fruiting capsule.

Cases of poisoning arise in adults from excessive use of a stimulant or a medicine. Children are sometimes tempted to eat the seeds, if they are permitted to play where the weed is to be found. Several cases of this kind were reported to the Department during the fall of 1897. At Alpena, Michigan, five children were badly poisoned in August by eating the seeds of the purple-flowered species, which was cultivated in a garden as a curiosity under the fanciful trade name of "Night-blooming Cactus." In Sept. a boy was killed in New York by eating the seeds of a jimson weed, which was permitted to grow in a vacant lot, his brother poisoned at the same time was saved only with difficulty. In October two other cases occurred in New York. Four children were playing in one of the public parks of the city where jimson weeds were growing luxuriantly. The boys imagined themselves Indians and roamed about and ate parts of various plants. Three of them ate the seeds of the jimson weed. One died in a state of wild delirium, another was saved after heroic treatment with chloral hydrate and morphine;

the third, who ate but few of the seeds, was but little affected. Children are also poisoned by sucking the flower, or playing with it in the mouth. The fresh green leaves and also the root have occasionally been cooked by mistake for other wild edible plants. One or two instances are recorded in which cattle have been poisoned by eating the leaves of young plants which were present in grass hay, but these animals generally either avoid the plants or are very resistant to its poison.

The symptoms of the poisoning are about the same in all cases, those characteristic of large doses being headache, vertigo, nausea, extreme thirst, dry burning skin and general nervous confusion, with dilated pupils, loss of sight and of voluntary motion, and sometimes mania, convulsions, and death. In smaller amounts the effects are like those of the ordinary anæsthesics. As vomiting is not a common symptom, the contents of the stomach must be quickly removed by the use of the stomach tube or emetics. It is well then to wash out that organ thoroughly with strong tea, tannic acid, or an infusion of oak bark, and to administer stimulants, such as brandy and hot, strong coffee. *Pilocarpus* is recommended by physicians to counteract the drying effect upon the secretions (sweat is very useful) and prolonged artificial respiration must often be resorted to to maintain the aëration of the blood.

As nothing has been said in regard to the *atropin* which is found in the Jimson weed, it might be said that the commercial *atropin* is derived from the root of belladonna and when used externally it is a local anodyne. Dilute solutions of *atropin* paralyze and stop the corpuscular movement in the blood and large doses give rise to slowing of the pulse. In poisoning it causes a paralysis of the vascular motor centers and stimulates the brain, large doses produce restlessness and excitement and delirium in man and organs only delirium in lower animals. With reference to the spinal cord, large doses cause complete loss of motion. Its action upon the nerves is very important and on this depends much of the value of the drug. Dr Winslow says:

The peripheral motor nerve terminations, and to a less extent, their trunks, are depressed and paralyzed. This is never so complete, however, but that there is some voluntary power left in an animal fatally poisoned.

Dryness of the mouth is one of the first symptoms following the use of belladonna because of the paralysis of the peripheral terminations of the secretory nerve. The muscular nerves are not affected by moderate doses of belladonna. The motor nerves ending in the voluntary muscles are paralyzed by poisonous doses of belladonna. Small doses do not affect the respiration, large doses make it quicker and deeper. Fatal doses cause asphyxia. Moderate doses cause a rise of temperature, but fatal doses lessen the body heat. Dr Winslow gives the following summary of the action of the drug:

It will be observed that belladonna, generally speaking, first stimulates and then depresses the nerve centres, while it chiefly paralyzes the motor nerve terminations, including the inhibitory (vagus and splanchnic), the secretory (salivary, vaginal, etc.) and, to a less extent, the sensory nerves. Secondary depression of the cerebrum is not so profound as that of the great medullary centres, especially the respiratory centre and there is sometimes a slight and brief stimulation of the motor nerves of the smooth muscles, viz. vagus, splanchnic, and possibly vaso-motor nerves.

Full medicinal doses depress the peripheral filaments of the inhibitory and secretory nerves and those of the striated muscles, lessen the functional activity of the voluntary motor system, and, to a less degree, that of the afferent nerves. The pulse becomes quickened because of paralysis of the peripheral vagus endings and stimulation of the heart or its ganglia, the blood tension is augmented because of the increased cardiac action and stimulation of the vasomotor centers, and the respiration is accelerated because of excitation of the respiratory centers. The temperature is elevated owing to the circulatory excitation and stimulation of the heat-producing centers. Slight delirium may be present from the exciting action of the drug upon the cerebral motor centers.

The spinal cord is unaffected by therapeutic doses. Locally applied belladonna is a direct paralyzant to nerves, muscles, vessels and cells.

Toxic doses of belladonna cause in animals dryness of the mouth, increased frequency of the pulse and respiration, elevation of temperature, dilation of the pupil and partial blindness, restlessness, nervousness, delirium, twitching of the muscles (occasionally erythema) and

frequent micturition. These symptoms are succeeded, in fatal poisoning, by fall of temperature, retention of urine, muscular weakness, staggering gait, partial anesthesia, convulsions and paralysis (one preponderating over the other), weak, slow, irregular respiration feeble, rapid pulse, paralysis of the sphincters, stupor and death. Death occurs mainly from asphyxia, but is due in part to cardiac failure. The physiological test consists in placing a drop of urine (secreted by the poisoned animal) into the eye of a healthy animal when mydriasis should follow if the case be one of belladonna poisoning. Three-quarters of a grain of atropin under the skin has proved fatal to dogs. Two grains of atropin produce mild toxic symptoms in the horse. Small dogs are slightly poisoned by gr 1.80 of atropin medium sized dogs by gr 1.60, given hypodermatically. Cattle are as susceptible as horses, although herbivora are not so easily influenced as carnivora. The pulse in dogs is greatly accelerated, sometimes as high as 400, while the pulse rate of the horse is not generally more than doubled. Rodents, as guinea pigs and rabbits, and pigeons, are particularly susceptible to belladonna, in regard to its effect upon the pupil, circulation, etc.

The treatment for poisoning includes the use of the stomach pump, emetics, cardiac stimulants, and *pilocarpin* under the skin. Also external heat, general friction and artificial respiration.

Datura Stramonium L. Jamestown or Jimson Weed

Much like the last, an annual glabrous or the young stem somewhat pubescent, stout and green, branches and leaves sparingly pubescent, leaves thin ovate, sinuate toothed or angled calyx less than $\frac{1}{2}$ the length of the corolla, corolla white, 3 inches long, the border 5-toothed capsule ovoid, prickly the lower prickles mostly shorter.

Distribution. Nova Scotia, New England to Minnesota Iowa Nebraska, Texas and Florida. Naturalized, native to tropical regions of the Old World, probably Asia. De Candolle says that it is probably native to the borders of the Caspian Sea.

Poisonous properties. The *Datura Stramonium* has been used in medicine since the close of the sixteenth century. Earlier than this it was used by the people of western Asia and eastern Europe. Miss Henkel describes the method of collecting as follows:

The leaves are collected at the time of flowering, the entire plant being cut or pulled up and the leaves stripped and dried in the shade. The unpleasant narcotic odor diminishes upon drying. The leaves are poisonous, causing dilation of the pupil of the eye, and are used principally in asthma.

All three species of *Datura* are poisonous, the seeds being especially poisonous. Dr. Hasted records a case of poisoning of a boy five years old in New York, New Jersey who ate freely of a half grown capsule of this species and died the next morning. The seeds of *D. Stramonium* are known to have poisoned a child in eastern Iowa. They contain *hyoscyamine atropin* and *scopolamine*. In some analyses, as much as 0.33 per cent of the alkaloid *atropin* has been found in the seeds, and about 0.2 per cent in the leaves.

Datura Metel L. Thorn Apple

A clammy pubescent annual 3-4 feet high leaves ovate, entire or obscurely angular toothed, rounded at the base flowers large, white, calyx about $\frac{1}{2}$ as long as the corolla capsule globose prickly.

Distribution. Native to tropical America. Naturalized from New England to Florida and westward.

Datura meteloides DC. Wright's Datura

A spiny pubescent annual, pale in color, leaves obovate entire flowers large showy white or pale violet, sweet scented corolla with a 5-toothed border capsule nodding spiny.

Poisonous properties. The writer a few years ago saw a notice in a local

paper of a child being poisoned by sucking the nectar of a flower of Wright's Datura. Professor Chesnut, in speaking of the poisonous properties of the same species, says

Datura meteloides is a very large-flowered species, which is native from southern California to Texas, and in some localities is common in cultivation. No cases of poisoning have yet been recorded against it, but it is largely used as an intoxicant by Indians, and is used in general for the same purposes as jimson weed. It undoubtedly contains the same poisons.

7. *Lycium* L. Matrimony Vine

Shrubs or woody vines, often spiny, leaves small, entire, alternate, with smaller ones between, calyx 3-5 toothed or cleft, persistent, corolla funnel-form or salver-shaped, usually 5-lobed, the lobes imbricated, stamens 5 rarely 4, anthers opening lengthwise, style slender, stigma capitate, ovary 2-celled, berry globose, ovoid or oblong.

About 75 species of wide distribution.

Lycium baumifolium Mill. Common Matrimony Vine

A glabrous spiny or unarmed shrub, slender climbing or trailing stem, leaves lanceolate, oblong or spatulate, petioles short, peduncles filiform, calyx lobes ovate, corolla short, funnel-form, greenish-purple, stamens slightly exserted, berry oval, orange red.

Distribution. In thickets and waste places, escaped from gardens from Canada to Texas. Introduced from Europe.

Poisonous properties. Supposed to be poisonous.

SCROPHULARIACEAE, Figwort Family

Mostly herbs, shrubs or rarely trees, leaves without stipules, flowers perfect, regular or irregular, calyx 4-5 toothed, cleft or divided, corolla irregular 2-lipped or nearly regular, stamens 2-5, didynamous or nearly equal, inserted on the corolla, pistil 1, 2-celled, many ovuled, fruit a capsule, seeds numerous, with a small embryo in copious albumen.

About 2500 species of wide distribution. Few plants of the family are of economic importance. Several species are medicinal. The most important are foxglove (*Digitalis purpurea*), mullein (*Verbascum Thapsus*) used as a stimulant because of its mucilaginous properties, and speedwell or Culver's root or Culver's physic (*Veronica virginica*), used for digestive disorders, when fresh being a violent emetic-cathartic and containing a glucoside *leptandrin*. Several species of the order are cultivated for ornamental purposes. The *Paulownia tomentosa*, native to Japan, is hardy in the south. The foxglove is also much cultivated. Several species of monkey flowers (*Mimulus luteus*) and musk flower (*M. moschatius*) are cultivated. The best known of all is the snap dragon (*Antirrhinum majus*), native to Europe. The genus *Casteolaria*, of which there are numerous species, is native to Chili and other parts of South America, and is cultivated. The *C. crenatiflora* is a showy herbaceous plant cultivated for its pretty slipper-shaped, sac-like flowers. The *Maurandias* are Mexican climbers with heart shaped or halberd-shaped leaves and open-mouthed, somewhat bell-shaped, purple, rose colored or violet corollas. The *Torenia asiatica*, native to Asia, is cultivated for its handsome pale violet or purple flowers. The turtle head (*Chelone glabra*) is occasionally cultivated and has large, white or rose-tinted corollas that are very pretty. The beard-tongue (*Penstemon*) contains many species, found mostly in western America and Mexico, the most beautiful of our western species being *P. grandiflorus*, with

large, lilac-purple flowers collected in ample racemes. Many of the Rocky Mountain forms are handsome perennials. The Mexican *Russelia juncea* is a showy bedding and greenhouse plant with carmine flowers and leaves reduced to scales. The painted cup (*Castilleja coccinea*) is a pretty species, native of the northern states. The roots of most species are parasitic. The common lousewort (*Pedicularis canadensis*) is an early spring blooming plant of northern prairies. *P. groenlandica* of Europe, is found in the colder regions of North America also and has handsome purple flowers borne in spikes. The flowers of *Lythris atropurpurea* or Cape Saffron resemble true saffron very greatly in odor, taste and drying qualities.

Vanquelin isolated the glucoside *gratiolin* $C_{16}H_{18}O_{11}$ from *Gratiola of ficoides*. This species is poisonous to stock, strong medicinal doses are poisonous to man as well. Some species, according to Maiden, are often poisonous to stock in Australia. The row wheat of Europe (*Melampyrum arvense*) causes colic and sleepiness.

Genera of Scrophulariaceae

Flowers regular or nearly so.

Flowers racemose, stamens 5.

1. *Verbascum*

Flowers axillary or racemose, stamens 2

2. *Veronica*

Flowers irregular

Stamens 4 not in pairs.

Corolla spurred

2. *Lamaria*

Corolla not spurred.

Tubular

3. *Digitalis*

Bell-shaped

4. *Gerardia*

Stamens in pairs.....5. *Pedicularis*

1. *Verbascum* (Tourn.) L. Mullen

Biennial or perennial, generally tall herbs with alternate leaves. flowers in spikes, racemes or panicles, calyx 5-parted, corolla flat with 5 broad rounded or slightly unequal divisions, stamens 5, inserted on the base of the corolla, unequal filaments of all of the stamens woolly or only the 3 upper style flattened at the apex, fruit a capsule 2-valved, seeds rough.

About 125 Old World species. Several naturalized in North America.

Verbascum Thapsus L. Common Mullen

A tall, densely woolly annual from 2-6 feet high. leaves oblong, thick, covered with branched hairs, the basal leaves margined petioled. flowers in long dense spikes. corolla rotate, yellow or rarely white, stamens unequal, the 3 upper shorter woolly with short anthers the 2 lower smooth with large anthers.

Distribution. From Nova Scotia north across the continent; south to Missouri and Kansas and west to Utah.

Verbascum Blattaria L. Moth Mullen

Stem round, sparingly branched, biennial with smooth leaves the lower petioled, oblong ovate, lanceolate, lacinate, serrate, upper clasping, flowers in loose racemes, yellow or white with a tinge of purple, all the stamens bearded with violet hairs, capsule nearly globose, numerous seeds.

Distribution. Common eastward, rarely in the Mississippi Valley. Abundant in the west in Salt Lake basin.

Poisonous properties. The common mullein produces an irritation but is probably not very poisonous to stock.

2. *Linaria* (Tourn.) Hill. Toad Flax

Herbs or shrubs, with alternate leaves, or those of the sterile shoots opposite or whorled, flowers in racemes or spikes, calyx 5-parted, corolla per-




Fig. 437. *Verbascum Thapsus*. (From Johnston's Medical Botany of N. A.)



Fig. 438. Toad Flax (*Linaria vulgaris* n. seed. Regarded with suspicion, (Selby

sonate and with a spur at the base; the upper lip erect, 2-lobed, the lower 3-lobed; stamens 4, didynamous, not exerted, fruit a capsule, opening by 1 or more holes in the top, seeds small, numerous.

About 150 species of wide distribution. One native species in the northern states.

Linaria vulgaris Hill. Ramsted. Butter and Eggs.

A pale green perennial with erect, leafy globose or sparingly pubescent stem; leaves sessile, entire, upper, at least, alternate, flowers in dense racemes; calyx segment oblong, spur subulate, corolla orange color, nearly erect, 1 inch long, spur subulate, nearly as long as the body of the corolla.

Distribution. Native to Europe. In fields and waste places from Nova Scotia to Kansas, North to Manitoba.

Poisonous properties. It is regarded with suspicion. The plant has a very disagreeable odor. A glucoside *linaria* $C_{14}H_{18}O_{10}$ has been isolated.

Veronica peregrina L. Purslane Speedwell, Neckweed

Glabrous, glandular, or nearly smooth, branching annual, 4-9 inches high.

leaves petioled, upper oblong, linear and entire, floral leaves like those of the stem but reduced, flowers axillary and solitary, white, capsule orbicular.

Distribution. A common weed in fields in eastern North America from Nova Scotia to Florida, Missouri, Kansas, Texas, British Columbia and California. Also found in Mexico, South America and in Europe, almost cosmopolitan.

Veronica virginica L. Culver's Root. Culver's Physic

A tall, smooth, or occasionally somewhat hairy perennial simple stem, leaves lanceolate pointed in whorls finely serrate, flowers in panicked spikes, small, nearly white, salverform, tube longer than the calyx, stamens 2, exerted, pistil 1, style 1, fruit a capsule, oblong-ovate.

Distribution. From western New England to Minnesota, Manitoba to Nebraska and Kansas.

Poisonous properties. The *V. peregrina* has been reported as poisonous. The root of *V. virginica* contains *leptandrin*. It is a violent emetic, cathartic and, according to Johnson, cannot be used with safety in medicine.

3. *Digitalis* L. Foxglove

Tall herb, leaves alternate, large purple yellowish or white flowers borne on 1-sided racemes, calyx 5-parted, corolla irregular, tube contracted, upper lip 2-cleft, lower lip 3-lobed, middle largest, stamens 4 didynamous, style slender; fruit a capsule, seeds numerous, roughened. About 20 species in Asia and Europe.

Digitalis purpurea L. Purple Foxglove

Biennial or annual pubescent herb with stout stem, lower leaves ovate or ovate-lanceolate, slender petioled, upper leaves smaller sessile, flowers borne in long drooping racemes, corolla spotted.

Distribution. Native to Europe but widely naturalized in the Pacific Northwest.

Poisonous properties. The plant has long been used in medicine. For this purpose the leaves of the second year's growth are collected. The active constituents are such glucosides as *digitoxin*, $C_{41}H_{64}O_{16}$. The most poisonous are active, *digitalin*, $C_{28}H_{44}O_{14}$, *digitalen*, an amorphous bitter substance soluble in water, *digitonin*, $C_{27}H_{44}O_{14} + H_2O$ and *digitophyllin* $C_{32}H_{52}O_{16}$. The leaves also contain *luteolin* which occurs in mignonette.

In large doses digitalis is a gastro intestinal irritant and in poisoning causes nausea. It causes the pulse to become slower, fuller and stronger and more regular. It causes stimulation of the heart muscles. In poisonous doses it is rapid, weak and irregular. The respiratory centers are unaffected except by toxic doses. The temperature is reduced by toxic doses. Dr Winslow in speaking of the cumulative action of digitalis and the toxic action says:

Digitalis and strechnon are said to be cumulative in their action. Evidence is stronger in the case of the former drug than in that of the latter. Its cumulative action is marked sudden transition from a therapeutic to a toxic effect. This may be due to three causes: 1. Tardy absorption. 2. Increasing susceptibility. 3. Delayed elimination and accumulation of the drug in the system. The cumulative action of digitalis is chiefly due to the latter cause. It should never be administered in full medicinal doses uninterruptedly for any considerable length of time.

Toxicology.—Poisoning may occur from large single doses within 3 to 6 hours of their ingestion, and last for 16 or more hours with a fatal result, or may appear suddenly after the administration for several days of large medicinal doses (therapeutic action). A minimum fatal dose for the horse is about 2 vi. of digitalis, or gr. i. as. of Hanolle's digitalin. For dogs,

3 i. of digitalis, or gr $\frac{1}{4}$ of digitalin. The symptoms exhibited are chiefly concerned with the digestion and circulation. They consist in dulness, lassitude, loss of appetite, nausea, flatulences, diarrhoea, infrequent, full pulse (reduced 6-10 beats in the horse), and contracted pupils. There is vomiting in dogs. In fatal cases these symptoms are followed by severe colic and tympanites, rapid, feeble, diabolic, irregular or intermittent pulse (120-140 in horses), while the heart may be heard and felt beating wildly and strongly, and a systolic blowing murmur can frequently be detected. This is due to mitral or tricuspid regurgitation caused by irregular contraction of the columnæ carneæ. The pulse is imperceptible because of the failure of the heart to fill the vessels. The extremities are cold, the eye is protruding, and salivation occurs. Bloody diarrhoea is very often present and the urine may be suppressed. The breathing finally becomes difficult and death ensues within a few hours, or as late as several days.

Treatment.—Evacuation of the stomach and bowels. Tannic acid, as a chemical antidote, alcohol, opium, and aconite, which is the physiological antagonist in depressing the action of the heart and lowering blood tension. In addition, external heat should be applied and complete quiet and rest secured.



Fig 429 *Digitalis purpurea*. Flowering branch—diminished. Flower natural size. A well known medicinal and poisonous plant. (From Vesque's *Traité de Botanique*).

4. *Gerardia* (Plumier) L. *Gerardia*

Erect herbs or a few shrubs, leaves generally opposite and sessile or the upper alternate, flowers showy purple or yellow, calyx bell-shaped, 4 toothed or 5-lobed, corolla somewhat irregular, bell-shaped, 5-lobed, slightly 2-lipped, stamens 4, somewhat unequal, filaments pubescent, capsule many seeded. About 40 species native to America.

Gerardia tenuifolia Vahl. Slender Gerardia

A glabrous annual, leaves narrowly linear acute, calyx teeth very short acute corolla light purple spotted $\frac{1}{2}$ inch long

Distribution. In low grounds Quebec to Minnesota, Iowa, Kansas and Louisiana.

Gerardia grandiflora Benth

A common pubescent annual or biennial 2-4 feet high, leaves ovate lanceolate coarsely toothed or cut, the lower pinnatifid pedicels shorter than the calyx, calyx-lobes oblong or ovate, corolla longer than calyx-lobes.

Distribution. In dry woods Wisconsin, Minnesota to Texas and Tennessee

Poisonous properties. The first species said to be poisonous to sheep and calves. Other species probably poisonous.

5. *Pedicularis* (Tourn.) L. Lousewort

Perennial herbs with pinnately lobed or cleft or pinnatifid leaves calyx tubular corolla 2 lipped, the upper lip arched, frequently beaked at the apex, lower lip erect, stamens 4, under the upper lip, anthers transverse, capsule generally oblique several-seeded. About 125 species mostly in the Northern Hemisphere.

Pedicularis canadensis L. Lousewort

Herb simple stemmed plant from 6 inches-1 foot high, leaves scattered, the lowest pinnately parted flowers in short spikes, calyx split in front, oblique, corolla greenish yellow and purple, upper lip of the corolla hooded, 2-toothed under the apex.

Distribution. In woods and prairies from eastern Canada to Florida, to Missouri and New Mexico, to Manitoba.

Pedicularis lanceolata Michx. Swamp Lousewort

An upright glabrous perennial from 1-3 feet high, leaves opposite and alternate, lanceolate or linear-lanceolate, pinnately lobed, the lower petioled, flowers in spikes, calyx 2-lobed, leafy-crested, corolla pale yellow, bearing a short truncate beak, the lower lip nearly erect capsule ovate, about as long as the calyx.

Distribution. In swamps from Ontario to Connecticut Virginia, Iowa, Nebraska, Minnesota to Manitoba.

Poisonous properties. *P. canadensis* is said to be poisonous to sheep and *P. lanceolata* is also suspected. Rocky Mountain species such as *P. groenlandica*, *P. racemosa* and *P. bracteosa*, are frequently eaten by sheep without any ill effects. Lehmann lists three European species as poisonous, the *P. palustris*, *P. sylvatica* and *P. rudetica*. A decoction made from these European plants is used to destroy animal parasites. In cattle these plants cause anemia.

Dr. Landley, in speaking of the European species, says, "they are acrid but are eaten by goats." The European *P. palustris* was formerly official and is much used in Europe as a domestic remedy. The glucoside, *rhamanthin*, is found in the different species of the genus. It is identical with the material found in the common Butter and Eggs (*Linaria vulgaris*) $C_{24}H_{34}O_{10}$

BIGNONIACEÆ. Bignonias Family

Woody plants trees, shrubs or woody climbers or some exotic herbs, leaves opposite or rarely alternate, flowers mostly large and showy calyx 2-

lipped, 5-cleft or entire, corolla tubular bell-shaped, 5-lobed, somewhat irregular, stamens inserted on the corolla, some of the stamens sterile or rudimentary, inserted on the tube of the corolla, anther bearing 2 or 4 ovary usually 2-celled, fruit a 2-valved capsule, seeds flat, winged, cotyledons broad and flat

About 500 species mostly tropical. The trumpet creeper (*Tecoma radicans*), native from Pennsylvania to Minnesota and southward, produces large scarlet or orange flowers, and is much cultivated as an ornamental plant. It contains narcotic principles. *T. jasminoides*, much cultivated in greenhouses for its pretty white, pinkish or purple flowers, is a native of Brazil. *Bignonia capreolata*, from Virginia to southern Illinois and southward, produces pretty orange red flowers and is cultivated southward. *B. venusta* is a greenhouse plant native to Brazil. The leaves of Caroba (*Jacaranda procera*) furnish a valuable astringent. The *Newbonidia laevis* is used in dysentery.

Catalpa Scop. Catalpa

Trees or shrubs, leaves opposite or verticillate, simple, petioled, flower large in terminal panicles, calyx deeply 2-lipped, corolla bell-shaped with spreading margin, some stamens with fertile anthers 2, the others sterile or rudimentary, capsule large and slender, 2-celled, seeds numerous, winged. Seven species in Asia, North America and the West Indies.

Catalpa speciosa. Warder. Common Catalpa

A large tree with thick bark, leaves large, heart shaped, long acuminate corolla 2 inches long white and mottled, capsule thick with numerous seeds.

Distribution. The common hardy catalpa (*C. speciosa*) is native to the United States from Illinois to Arkansas, is a tall tree largely planted for its wood which is used for posts, and railroad ties. It is hardy as far north as northern Iowa.

Catalpa bignonioides Walt

Tree with thin bark, spreading branches, leaves strongly scented and broadly ovate entire or 3-lobed acute or acuminate, deeply pubescent beneath, flowers in panicles, white mottled with yellow and purple corolla tube bell-shaped, the lower lobe entire, capsule rather thin walled, drooping.

Distribution. Commonly cultivated but less hardy than the preceding, native to the Gulf States.

Poisonous properties. The odor coming from the fragrant flowers is poisonous and Dr. White in his *Dermatitis Venenata* states that the flowers are irritating to many persons. Dr. Mollspaugh, also, says that it is considered to be dangerous to inhale the odor of the flowers for a long time, which, however, is probably not generally the case. The allied caroba (*Jacaranda procera*) contains the bitter principle *carobin*. The glucoside, *catalpin* comes from the bark and pod of *C. bignonioides*. The *Oroxylum indicum* contains *oroxylum* $C_{18}H_{16}O_8(OH)_2$.

PLANTAGINALES

This order contains but one family the *Plantaginaceae*.

PLANTAGINACEAE Lindl. Plantain Family

Mostly stemless herbs, leaves, in species with stems, opposite or alternate flowers small, perfect, polygamous or monoecious, calyx 4-parted persistent,

corolla 4-lobed, hypogynous, stamens 4 or 2 or only 1, inserted on the throat or tube of the corolla, ovary 1-2 celled, or falsely 3-4-celled, sessile, ovule 1, several ovules in each cavity from a gynae, circumscissile, at or below the middle of a whorl.

3 genera, 2 native to North America, and more than 200 species, of wide distribution. None of the plants are of any economic importance. The seeds of several species (*Plantago major* and *Plantago Rugelii*) are used to feed birds. The *P. ovata* is used in France as a salad. The seeds of *P. arvensis* of Europe and *P. indica* are used for sizing in the manufacture of muslin. All of the seeds of the genus have a mucilaginous testa.




FIG. 16. *Plantago major* (Plantain). 1. Pair of stamens. 2. Single pistil. 3. Single flower. 4. Seed. 5. Longitudinal section of seed. (After M. M. Cheney in Curtis, *Flora of North America*.)

RUBIALES

Leaves opposite or whorled. Flowers with gamopetalous corolla, separate anthers, stamens as many as the lobes of the corolla and alternate with them.

or occasionally fewer or twice as many, ovary compound, adnate to the calyx tube, ovules one or two in each cavity of the ovary. The important families are *Rubiaceae*, containing cinchona, coffee, asperula, and galium, *Adoxaceae* containing a single genus *Adoxa* the Musk-root, *A. Moschatellina*, *Valerianaceae* containing the corn salad (*Valerianella olitoria*, and *Valeriana officinalis*), native to Europe and North America, the roots of which are used in medicine, *Dipsacaceae*, containing Fuller's teasel (*Dipsacus fullonum*), whose rigid chaff hooked at the end is used for carding woollen cloth, and scabious (*Scabiosa maritima*) frequently cultivated for ornamental purposes, and *Caprifoliaceae*

Families of Rubiales

Herbs or shrubs; flowers regular; leaves with stipules.	<i>Rubiaceae</i>
Herbs or shrubs, flowers regular or irregular, leaves without stipules.	<i>Caprifoliaceae</i>



Fig. 431. Teasel (*Dipsacus sylvestris*). A common American weed at bed to Fuller's Teasel. (Millsap: Seely).

RUBIACEAE B. Juss. Madder Family

Herbs, shrubs or trees, with simple opposite leaves connected by stipules, or the leaves sometimes in whorls without stipules, calyx tube adnate to the ovary, flowers regular and perfect, often dimorphic, corolla funnel-shaped club-shaped, bell-shaped or rotate, 4-5 lobed, stamens as many as the lobes of the corolla and alternate with them, pistil with a simple or lobed style, ovary 1-10 celled, ovules one to many in a large cell, fruit various, capsule, berry or drupe, seeds small or large, the coat thin or hard, endosperm fleshy or horny.

A large order, chiefly tropical, consisting of about 350 genera and 5000 species. Only a few of them found in northern United States. Some species are abundant in southern United States, a few being weedy. Cinchona or Peruvian Bark, from which quinine is derived, is found in several species of the genus *Cinchona*, a tree with evergreen leaves. Quinine is derived chiefly from *Cinchona officinalis*, which is a native of South America, *C. lanceifolia* is native to Peru. The Cinchona is now, however, extensively cultivated in India. Its use in fevers has been recognized since its earliest introduction from the wild plants gathered in the Andes Mountains by the Indians. The Red cinchona is obtained from *C. succirubra* and Calisaya bark from *C. Ledgeriana*.

Cinchona contains a large number of alkaloids of which the following are more important. *Cinchonin*, $C_{19}H_{27}N_2O$, *quinamin*, $C_{19}H_{25}N_2O_2$, *quinin*, $C_{20}H_{25}N_2O_2$, *hydroquinin*, $C_{20}H_{25}N_2O_2$, *arcin*, $C_{22}H_{29}N_2O_4$. Another alkaloid belonging to this group is *disinchonin*, $C_{37}H_{44}N_4O_2$. *Jaronin*, $C_{28}H_{36}N_2O_4$, occurs in Calisaya bark. The Cuprea bark (*Remya pedunculata*) from the U.S. of Colombia is also used in the manufacture of quinine and contains *cinchonanin*, $C_{19}H_{25}N_2O$. The partridge berry (*Mitchella repens*) is a tonic. Gambier (*Laccaria*) of the East Indies is used for tanning. The root of ipecac (*Psychotria Ipecacuanha*) of Brazil is a systemic emetic used as a remedy in dysentery and contains *emetin*, $C_{15}H_{21}N_3O_5$ and *cephallin*, $C_{14}H_{20}NO_2$. Madder (*Rubia tinctorum*) of the Levant and Southern Europe is used for dyeing and contains a red coloring matter, *alizarin*. The *Morinda* contains a yellow coloring principle *morindin*. The cape jasmine (*Gardenia jasminoides*) also contains a yellow coloring resembling crocin.

Coffee obtained from the *Coffea arabica* and other species, contains the chemical principle *caffein*, $C_8H_{10}N_4O_2$. This is the same as *thein*. This substance occurs in a large number of plants including cocoa (*Theobroma Cacao*), cola (*Cola acuminata*), yopon (*Slex Cassiae*), maté (*I. paraguayensis*), *Sterculia peltatifolia*, *Paulinia Cupano*.

Green seeds of *Coffea arabica* contain 1.22 per cent of *caffein* or *thein*, $C_8H_{10}N_4O_2$, the young leaves of Chinese tea 2.12 per cent. *Caffeidin*, $C_8H_8N_2O$ is obtained from *caffein*. *Caffein* causes the heart to beat more




Fig. 432. *Cinchona* (*Chinchona lancifolia*). A native of Peru and one of the species furnishing the Peruvian bark of commerce. (From Strasburger, Naegeli, Schenck and Schimper.)

forcibly. It is a cerebral stimulant, producing wakefulness and restlessness. In lower animals it produces excitement and mania. From a toxicological point it is a spinal and muscle poison to the frog. In dogs and mammals it causes restlessness, and in dogs it produces vomiting. The minimum fatal dose according to Winslow is 1 gr. to 1 lb. of live weight.



Fig. 4.3 Coffee-plant (*Coffea arabica*). Fruiting branch. Furnishing the soft berry (country) (After Engel).




Fig. 4.4 1. Flower or branch. 2. Berry. 3. Transverse section of berry showing seeds. 4. Seeds.

The coffee plant is a small evergreen tree native to the tropical mountain districts of Africa but now cultivated in all warm countries. It was introduced into Arabia early in the 15th century or perhaps earlier. Brazil supplies a large amount of coffee. Large amounts of coffee also come from Ceylon, Java and the Celebes. It is also grown in Puerto Rico and Cuba. Its first introduction is said to have occurred in the middle of the 16th century. The Mocha coffee comes from southwestern Arabia.

The sweet-scented bedstraw (*Galium triflorum*) is used in making an aromatic drink, especially in German communities in this country. It contains coumarin. In Europe the sweet woodruff (*Asperula odorata*) is used like *G. triflorum* and when added to wine, the drink is known as "Mai-trunk."

Cephalanthus L. Button Bush

Shrubs or small trees, leaves opposite or verticillate, flowers in spherica, penicillate heads, white or yellow, calyx tube obpyramidal with 4 obtuse lobes corolla tubular 4-toothed stamens 4, inserted on the throat of the corolla ovary 2-celled ovules solitary in each cavity, style thread like stigma capitate fruit dry obpyramidal 1-2 seeded.

About 7 species native to America and Asia.

Cephalanthus occidentalis L. Button Bush

A shrub or small tree, leaves petioled ovate or lanceolate-oblong pointed opposite or whorled with small petioles, flowers borne in globular head sessile white, style longer than the corolla.

Distribution. In swamps and low grounds from Canada to Mississippi, Texas and Florida.

Poisonous properties. The leaves contain a poisonous, bitter glucoside *cephalanthin* $C_{12}H_{24}O_6$. It has been used in medicine on account of its bitter properties.

CAPRIFOLIACEAE Vent. Honeysuckle Family

Shrubs, trees or vines, or rarely herbs, with opposite leaves, stipules absent or present, flowers perfect, mostly cymose, calyx adnate to the ovary, 3-5 toothed or 3-5 lobed, the gamopetalous corolla with a 5-lobed limb or 2-lipped, stamens 4-5, inserted on the tube of the corolla and alternate with its lobes ovary 2-5 celled style slender, stigma capitate fruit a berry drupe or pod, seeds with a membranous or hard coat.

About 275 species. Generally found in the northern hemisphere. The plants of this order are of small economic importance. Several are used in medicine as the feverwort (*Triosteum perfoliatum*).

Many members of the order are used for ornamental purposes. The most important are members of the genus *Lonicera*. Of the native species, the trumpet honeysuckle (*Lonicera sempervirens*) is widely cultivated also Sullivan's honeysuckle (*L. Sullivanii*), Fraser's honeysuckle, (*L. flava*) and the western honeysuckle (*L. involucrata*). Some of the *Loniceras* are possibly poisonous. Of the European and Asiatic species, the *L. tatarica*, *L. japonica*, *L. fragrantissima* and *L. Periclymenum* are cultivated. The elders (*Sambucus canadensis* and *S. racemosa*) are likewise cultivated in the North. The former is often weedy. Several species of the snowberries, like the wolfberry (*Symphoricarpos occidentalis*) and the snowberry proper *S. racemosus* are native to the northern states. The Indian currant (*S. orbiculatus*) is sometimes weedy in Iowa and Missouri. Suckers used in Arkansas for making

baskets. The *Linnæa borealis* or twin-flower is native in the cool damp woods of the North.

Several of the viburnums are cultivated. The hobblebush (*V. alnifolium*), a native from Ontario and southward, is frequently cultivated in the East. The cranberry-tree or guelder rose (*V. Opulus*) found along streams from New Brunswick to northeastern Iowa, is cultivated both in its native and cultivated forms. The well-known snowball is a cultivated form of the cranberry tree. The fruit of this plant is used in the North. The root of horse gentian (*Triosteum perfoliatum*) has a bitter taste and is used as a cathartic. The flowers of the elder berry (*Sambucus canadensis*) are sudorific. The bark of *Viburnum prunifolium*, the black haw, is official, an antispasmodic, nervine and an astringent. It is also used as an uterine sedative, contains viburnin, oxalic acid, tannic acid, etc. The bark of *V. Opulus* is said to be antispasmodic. *Xylosteum* occurs in *Lonicera Xylosteum*. According to Greshoff the leaf of *Viburnum macrophyllum* and of *Symphoricarpos* are common saponary.

Genera of Caprifoliaceae

Flowers in compound cymes, corolla rotate	<i>Sambucus</i>
Flowers not in cymes, tubular to campanulate	
Erect perennial herbs	<i>Triosteum</i>
Shrubs	<i>Symphoricarpos</i>

Triosteum L. Horse Gentian. Feverwort

Coarse hairy perennial herbs with simple stems; leaves connate perfoliate or sessile, flowers axillary perfect sessile, calyx tube ovoid with a 5-lobed limb, persistent, corolla tubular, gibbous at the base, 5-lobed, stamens 5, anthers linear, ovary 3-5 celled, ovule 1 in each cavity, style bifid, fruit a dry drupe orange or red, enclosing 2-3 or rarely more 1-seeded nutlets, embryo minute.

Triosteum perfoliatum L. Feverwort Wild Coffee

A soft hairy perennial 2-4 feet high, leaves oval, abruptly narrowed below, downy beneath, flowers brownish purple, clustered, corolla purplish, fruit orange in color.

Distribution. In rich woods New York to Minnesota, Kansas and Alabama.

Poisonous properties. Some species of the genus were used by the Indians as a cure of fevers and early practitioners in this country used the root as an emetic. In early days, the berries of this plant were used as a substitute for coffee. The physiological action of the plant is to produce vomiting. It has a bitter nauseous taste.

Sambucus (Tourn.) L. Elder

Shrubs, trees or occasionally herbs, leaves opposite pinnate, flowers small in compound cymes, calyx-lobes minute or obsolete, corolla rotate or somewhat campanulate, regular with a 5-cleft wing, stamens 5, inserted at the base of the corolla, stigmas 3, ovary 3-5-celled, ovules 1 in each cavity, fruit a berry like drupe with 3-5 1-seeded nutlets, endosperm fleshy.

About 20 species of wide distribution.

Sambucus canadensis L. Common Elder

A shrub from 5-10 feet high, wood with large lenticels and large pith, leaflets 5-11 ovate or oval acuminate or acute, short stalked smooth above, sharply serrate, flowers white.

Distribution. From Canada to Manitoba, Kansas, Texas to Florida.

Poisonous properties. In regard to the poisonous properties Dr. Rusby says:

The common black elder or *Sambucus canadensis* L., a plant very common throughout the entire eastern and central United States, and represented by other species, apparently with similar properties, upon the Pacific Coast and in the old world, has dangerous properties which have remained unrecognized, or, in any the least, very obscure, to the present time.

Of the last mentioned, Dr. Robert Christian reports in the *Edinburgh Medical and Surgical Journal*, 1830, page 73, as follows:

Two boys in the vicinity of Edinburgh encountered a clump of the *S. Ebuus*, and one of them ate freely of the flowers, the other of the leaves. The boy who had eaten the leaves was attacked with enteritis, the abdomen at length becoming so sore that it could scarcely be touched. There was enormous vomiting, the matter containing blood. (Hemorrhagic colic) existed throughout. The boy was saved by vigorous treatment. The one who had eaten the flowers suffered considerably, and for a considerable time from vertigo with some headache, but the symptoms were not very serious.

Dr. Christian observed that both the berries and the flowers were known to kill fowls which fed upon them and that when berries were freely eaten they often caused giddiness. He also quotes a report of a case of a woman who dressed the shoots with vinegar and ate them as a salad, and who was promptly seized with violent purging, forty times in two days, coma resulting on the third day. Of our own species, *S. canadensis*, Dr. Johnson states that the bark and the root are actively cathartic and hydragogue when freely used. There is little doubt that he refers in this instance to the bark and the root in the green condition, since it is well known that the properties become much less active upon drying and keeping.

Our most direct evidence bearing upon the poisonous character of the elder-berry comes rests upon a case which occurred in the spring of 1894, at the Institution of Mercy, a Roman Catholic institution for children at Tarrytown, on the Hudson, and which attracted a great deal of attention at the time in the public press. The grounds of this institution were comparatively new and ditching and fencing were still in progress at the time stated. A workman in digging a drain, discovered a large number of roots to which the children took a fancy and which they began eating. Within a very few minutes, and while still engaged in eating, a large number of the boys were seized with convulsions and several of them died. One of them had the remainder of the root, the marks of his teeth upon it still clutched in his hand after death. The symptoms correspond in most features to those of the acute poisoning above described and in that regard the accident was described in the public press. Several months later I visited the institution in company with Mr. Frederick A. Coker, the botanist of the United States Department of Agriculture and Prof. Edward L. Coker, Professor of Botany to the Catholic University at Washington. At this time and subsequently through correspondence a pretty thorough investigation of the case was made. We found that it was not a locality where *Cicuta* would be apt to grow and no evidence existed that any had grown there. Three poisonous plants grew upon the spot, viz. the least pokeweed and elder. The workman who had dug the drain, the surviving boys and the Sisters in attendance were positive that it was the elder root which had occasioned the poisoning. They did not know the name of the plant, but had accepted the statements of the parents that it was elder, but they positively identified it by its appearance and by the young purple shoots and coarsened leaves which they had observed carefully while still attached to the pieces of root which had been taken from the hands of the boys poisoned. Their story was so clear, connected and positive that it was difficult to doubt that the elder root was the poisoning agent. Furthermore the least would not have produced the symptoms that were observed and the bark should have at once been distinguished by even a casual observer. Nevertheless, since the root was described as "like a carrot or turnip," and since the symptoms in some respects resembled those of poke-root poisoning the question cannot be regarded as settled beyond a doubt. In the case of so large a number of victims it is even possible that both of the roots were concerned. The attending physician, Dr. Luke Fleming, does not believe that the poisoning was caused by Elder. The active constituent of the elder is not known farther than that a report has been received from the twigs and leaves of the related European species, *S. nigra*. This would of course explain the very similar symptoms to those of *Cicuta* poisoning. The chemistry of the plant is now receiving thorough investigation in the division of pharmacology in the United States Department of Agriculture.

In Part I attention was called to Treub's theory in regard to the role of hydrocyanic acid in plants. Prof. Treub in a recent paper¹ reiterates his former conclusions that it performs some part in the products of assimilation. It was found that the amount of hydrocyanic acid in plants of *Sorghum* increases during the day because of its relation to the products of assimilation of carbon. It had previously been shown in the case of *Pongium edule* and *Phaseolus lunatus* that light plays no part in the formation of this substance, except as it favors photosynthesis. The same results have now been obtained with *Prunus japonica*, *Pastiflora foetida*, and some other plants. The results of the investigation with these plants show a direct proportion between the formulation of hydrocyanic acid and the function of the chlorophyll. The amount of acid is usually greatest in the young leaves and gradually diminishes as the leaves grow older. Leaves about to fall contain very little hydrocyanic acid. *Sambucus nigra*, according to Guignard is one of the exceptions to the rule, and Treub has also found this to be true for *Indigofera galeoides*. Hydrocyanic acid is probably the first recognizable product of the assimilation of nitrogen "and perhaps the first organic nitrogen compound formed." The amounts of the acid in the plants could be increased and decreased in proportion to the amount of nitrate used. Ravenna and Peli think that the nitrates are necessary for the formation of the acid. Treub agrees with this and adds that dextrose is especially essential. The acid probably occurs in the form of a glucoside and is liberated by an enzyme or by boiling water.

The investigations of Em. Bourquelot and Em. Danyon² with the glucosides found in various plants show that they are not identical. In *Sambucus nigra* they find *sambunigrin*. They also studied the character of glucosides from which hydrocyanic acid is derived in *S. racemosa* and *S. libanica*.

The flowers have long been used in domestic practice. The physiological action recorded for the drug by Dr. Millspaugh are as follows:

Dr. Uebelacker's experiments with from 20 to 30 drops of the tincture gave the following symptoms of physical disturbance: Drawing in the head, with anxious dread, flushed and blotched face, dryness and sensation of swelling of the mucous membranes of the mouth, pharynx, and trachea, frequent and profuse flow of clear urine, heaviness and constriction of the chest, palpitation of the heart, pulse rose to 100, and remained until perspiration ensued, sharp, darting rheumatic pains in the hands and feet, exhaustion and profuse perspiration, which relieved all the symptoms.

Prof. Hyman states that the young buds of the American elder are especially poisonous.

The European *Sambucus nigra* contains the alkaloid *sambucin*, according to Sanctis the same plant contains *convall.* *Sambunigrin* has also been isolated and hydrocyanic acid is known to be present.

Symphoricarpos (Dill.) Ludwig. Snowberry

Low, branching shrubs, leaves oval, short petioled entire or wavy-toothed, downy underneath, flowers white tinged with purple, calyx tube nearly globular, teeth short, corolla bell-shaped 4-5 lobed, stamens 4-5 inserted in corolla; ovary 4-celled, 2 with a fertile ovule; fruit a berry 4-celled and 2-seeded, embryo minute.

About 10 species of North America. One species frequently cultivated for ornamental purposes. The wolf-berry (*S. occidentalis*) northward is com-

¹ Nouvelles recherches. Ann. Jard. Bot. Botanique, II. 2:84-118. 1910. See R. Catin Rose, Bot. Gazette 30:156.

² Soc. Biol. Paris. 7 July, 1905. 1. c. Oct. 9, 1905. Jour. de Pharm. et de Chimie, Aug. 16, and Sept. 1, 1905. Compt. rend. Acad. Sci. Paris, July 8, 1905.

mon in the Rocky Mountains and frequently cultivated. The snowberry (*S. racemosus*) is common in rocky woods, and abundant along river course from Minnesota to Arkansas and westward.

Symphoricarpos orbiculatus Moench. Indian Currant Coralberry

A shrub 2-4 feet high, purplish, usually pubescent, branches, leaves oval or ovate entire or undulate, nearly glabrous above, pubescent underneath; flowers in short axillary clusters, corolla bell-shaped sparingly bearded, pinkish, stamens included; fruit a purplish berry.

Distribution. Rocky woods and along streams; from New Jersey, Illinois, Southern Iowa, South Dakota, Nebraska, Texas to Georgia.

Poisonous properties. It is suspected of being poisonous, but there is no direct evidence to support this view.

CAMPANULATAE

Herbs or rarely shrubs, corolla gamopetalous, petals occasionally separate; stamens as many as the lobes of the corolla, anthers united, ovary inferior. It contains the families *Cucurbitaceae*, *Candolleaceae* (mostly Australian), and *Compositae*, this last order including *Cichoriaceae*, *Compositae* and *Ambrosiaceae* of some authors.

Families of Campanulatae

Herbs or rarely shrubs; flowers in an involucrate head	Compositae.
Flowers not in involucrate heads.	
Flowers monoecious or dioecious, generally vines	Cucurbitaceae.
Flowers perfect	Lobeliaceae.



Fig. 435. Water melon (*Citrullus vulgaris*). (W. S. Dudgeon).

CUCURBITACEAE B. Juss. Gourd Family

Herbaceous vines, usually with tendrils, leaves alternate, petioled, palmately lobed or dissected, flowers dioecious, monoecious or rarely perfect, calyx tube adnate to the ovary, 5-lobed, petals usually 5, inserted on the limb of the calyx, stamens 13, 2 of them with 2-celled anthers, the other with a 1-celled anther, filaments short, frequently monadelphous, ovary 13-celled, stigmas 2 or 3, fruit indehiscent or rarely dehiscent, seeds flat in the large embryo, exalbuminous.

About 650 species, mainly in tropical regions. A few of the species are medicinal. The squeezing cucumber (*Leballium Elaterium*) is a fleshy decumbent herb used for making elaterium, a powerful hydragogue cathartic and an irritant poison. Its poisonous nature was known to Pliny. It contains *elaterin*, $C_{20}H_{28}O_8$. *E. officinale* contains *prophetin*, another glucoside. The colocynth (*Citrullus Colocynthis*) a slender scabrous plant with perennial roots, is native to the dry regions of the Old World, Palestine and North Africa. Its gourd is about the size of an orange and used as a purgative, while the seeds are roasted and boiled and used as food by some of the tribes of the Sahara. Tea made from this fruit is used by the people of the Nile to smear their water bags to prevent camels from cutting them. It contains *colocynthin*




Fig. 436. Bryony (*Bryonia dioica*). A, a branch with flowers, B, female flowers, C, male flowers, D, stamens, E, fruits, F, section of fruit. (After Strasburger, Noll, Schenck and Schimper).

$C_{40}H_{40}O_{10}$ The plant is intensely bitter. The towel or sponge gourd (*Luffa aegyptiaca*), a native of Egypt, produces a fruit one foot or more long filled with a spongy fiber, which when the outer part is removed is used to rub the skin, and for many other domestic purposes. The fruits of many plants of the family are economic. The nara (*Acanthosicyos horrida*) in Southern Angola is used as food and has medicinal virtues. The chayote (*Serchium edule*) is cultivated in the West Indies for its fruit. The green and ripe fruit



Fig. 437. *Colocynthis* (*Chenille colocynthis*). An intensely bitter plant of economic importance. (After Faguet).

of the cucumber (*Cucumis sativus*) native to India, has long been used for food, especially for pickles, and the West Indian gherkin (*C. Anguria*) is also cultivated for the same purpose. According to Greshoff the foliage of *Cucumis metuliferus* contains saponin. He also states that he found saponin in the seeds of *Lagenaria vulgaris*, and *Cucurbita maxima*. The *C. myriocarpus* contains the toxic alkaloid myriocarpin. The musk melon (*Cucumis Melo*), a native to British India, is now widely cultivated. Sugar and nutmeg melons are well known everywhere in North America. The water melon (*Citrullus vulgaris*), native to tropical Africa, where large areas of wild plants occur, has long been cultivated in Mediterranean countries, and is well known everywhere in North and South America. The citron is a form of the water melon. The common pumpkin (*Cucurbita Pepo*) is native to southwestern North America, Arizona and Mexico. It is used for stock food and for culinary purposes. The nest-egg gourd, bush scalloped squash, crookneck squash and common pumpkin are all from the same species. The seeds of pumpkin and squash are used in North Africa and Egypt much as peanuts are in this country and have taenifuge properties. The winter squash (*C. maxima*) is probably also of




Fig. 438. Muskmelon (*Cucumis Melo*)
A part of a plant of the common melon.
(W. S. Dudgeon)




Fig. 439. Pistillate flower of cucumber (*Cucumis*, a, sepal, b, petal, c, pistil, d, stigma, e, style, f, ovary. (Pieters, U. S. Dept. Agr.)

American origin. The Hubbard squash is a well known representative of the species and is used in various food preparations. Other species like cushaw (*C. moschata*) are also cultivated. The Missouri gourd (*C. foetidissima*), with a large root sometimes a foot in diameter, occurs from Nebraska to Texas. The wild cucumber or wild balsam apple (*Echinocystis lobata*) is frequently used as a cucumber and is sometimes weedy. The star cucumber (*Sicyos angulatus*) is used in a similar way. The cocoon antidote, (*Pe villen cordifolia*), a native of Jamaica, has seeds which are used as a cure for snake bites and as an antidote against *Entada scandens*. They also contain an oil. The seeds of *Tefaria pedata*, a native of the East coast of Africa and Zanzibar, contain an excellent oil. The bryony of Europe (*Bryonia dioica*) is a climbing plant producing a pretty colored fruit. It is a drastic purgative and is poisonous. It contains the alkaloid *bryonin* $C_{10}H_{11}NO_7$, and the root of *Bryonia alba* contains the glucoside *bryonin* $C_{42}H_{82}O_{11}$. Dr. Halsted states that a friend of his has been repeatedly poisoned by handling the star cucumber (*Sicyos angulata*). Friedberger and Frohner state that poisoning has occurred from *Cucurbita Pepo*, causing symptoms of dullness and in some cases the animals showed excitement. Major Kirtikar says that the pulp of a cucurbitaceous plant of India *Trichosanthes palmata*, a perennial herb, is used in India to poison cattle, and that the *T. cucumerina* also of that country is an emetic and a drastic purgative. *Echinocystis macrocarpa*, according to Trimble and Sayre contains a glucoside. The marimbo or dipper gourd (*Lagenaria vulgaris*) yields gourds which are edible when small.

LOBELIACEAE. Lobelia Family

Herbs with acid milky juice, leaves alternate, flowers perfect, irregular, 5-lobed, gamopetalous corolla, stamens 5, free from the corolla, united into a tube, stamens monadelphous and syngenesious, flowers proterandrous, the stigma of the single style often fringed with hairs, fruit a capsule with numerous small seeds, embryo minute and straight.

About 600 species of wide distribution, comparatively few in northern United States. Some of the tropical species shrubby. Several species are medicinal, among them Indian tobacco (*Lobelia inflata*). The *Lobelia erinus* of the Cape of Good Hope is frequently cultivated in the conservatories and in gardens. It has small azure blue flowers. It has escaped on the Pacific Coast.

Lobelia (Plummer) L. *Lobelia*

Herb or occasionally shrub with alternate or radical leaves, flowers racemose or spicate, calyx 5-cleft with a short tube, corolla irregular, with a straight tube split down on one side, the upper lip of 2 erect lobes, the lower lip spreading and 3 cleft; stamens 5, free from the corolla tube, monadelphous, two of the anthers or all of them bearded at the top, ovary 2-celled, fruit a 2-celled pod, many-seeded.

About 200 species, of wide distribution. Some 25 species native to the United States.

Lobelia inflata L. Indian tobacco

A pubescent or hirsute, much branched annual from 1-2 feet high, leaves dentate or denticulate, the lower larger the upper small, bract like, but longer than the pedicels of the flower, flowers pale blue; calyx tube ovoid, capsule ovoid, inflated.

Distribution. In fields, especially clay soils, from Labrador to Georgia, Tennessee, Arkansas, Kansas, Iowa to Northwest Territory.

Poisonous properties. It is used medicinally for laryngitis and spasmodic asthma. In full doses it produces nausea, vomiting and great prostration, in overdoses it produces prostration, stupor, coma, convulsions and death.

We quote from Dr. Millipugh in regard to poisoning.

Thanks to much reckless prescribing by many so-called eclectic physicians, and to careless intent, as well as to experimentation and careful provings, the action of this drug is pretty thoroughly known. *Lobelia* in large doses is a decided narcotic poison, producing effects on animals generally, bearing great similitude to somewhat smaller doses of tobacco, and *lobeline* in like manner to nicotine. Its principal sphere of action seems to be upon the pneumogastric nerve, and it is to the organs supplied by this nerve that its toxic symptoms are mainly due, and its "physiological" cures of pertussis, asthmatic asthma, croup and gastralgia gained. Its second action in importance is that of causing general muscular relaxation, and under this it records its cures of strangled hernia (by enema), tetanic spasms, convulsions, hysteria, and mayhap, hydrophobia. Its third action is upon mucous surfaces and secretory glands, increasing their secretions.

The prominent symptoms of its action are: great depression, exhaustion, and mental depression, even to insensibility and loss of consciousness; nausea and verigo; contraction of the pupil; profuse clammy salivation; dryness and prickling in the throat; pressure in the esophagus with a sensation of vermiform motion, most strong, however, in the larynx and epigastrium, sensation as of a lump in the throat; increase and violent nausea, with pain, heat and oppression of the respiratory tract; vomiting, followed by great prostration; violent and painful cardiac constriction; griping and drawing abdominal distress; and urine, easily decomposing and depositing much uric acid; violent racking cough, dry and cough withropy expectoration; small irregular slow pulse; general weakness and oppression, more marked in the thorax; violent spasmodic pains, with paralytic feeling especially in the left arm; weariness of the limbs, with cramps in the gastrocnemii; and sensation of chill and fever. Death is usually preceded by insensibility and convulsions.

It contains *lobelic acid*, *lobelacrin inflata* and the alkaloid *lobelin* $C_{12}H_{21}NO_2$ has been isolated which according to Lloyd is a powerful emetic. *Lobelia nicotianifolia* of India and *L. purpurascens* contain the same alkaloids. According to the late Baron Ferdinand von Muller the *L. Brevia* of Australia and other species are poisonous.

Lobelia siphilitica L. Great Lobelia

A somewhat hairy stout, perennial herb from 1-3 feet high, leaves thin, acute or acuminate at the apex, dentate or crenate-dentate, sessile or the lower petioled, flowers large, spicate, racemose, leafy bracts, calyx hirsute, corolla bright blue or occasionally white.

Distribution. In moist soil near springs and in marshes from New England to South Dakota, Kansas, Louisiana and Georgia.

Poisonous properties. It is suspected of being poisonous. Johnson in his manual says of the action of the species of *Lobelia*

In full doses lobelia produces severe nausea, obstinate vomiting, and great prostration. In overdoses the prostration becomes extreme, there is failure of voluntary motion, followed by stupor, coma, and not infrequently convulsions and death. Though formerly much used for emetic effect by empirics, dangerous effects were so often produced that it is now seldom employed in this manner. It is chiefly employed in spasmodic affections of the air-passages, as spasmodic laryngitis and spasmodic asthma. In the latter disease it often produces the expectorant effects.

The great lobelia is probably not as poisonous as *L. inflata* to which the above remarks chiefly apply.

Lobelia cardinalis L. Cardinal-flower

A tall smooth or slightly pubescent perennial 2-4 feet high, leaves thin, oblong, lanceolate, smooth or slightly pubescent, crenulate, flowers racemose, bright scarlet or red.

Distribution. In moist soil, usually alluvial bottoms, from New Brunswick to Manitoba, Kansas, Texas, to Florida.

Poisonous properties. Reported as poisonous.

Lobelia spicata Lam. Spiked Lobelia

A perennial or biennial, smooth or pubescent herb; leaves smooth or minutely pubescent, leaves thickish, the lower obovate or spatulate, the upper linear or club-shaped bracts, entire or dentate or crenulate, flowers in a racemose spike, pale blue. calyx tube short, obconical or nearly hemispherical.

Distribution. Prairies or dry sandy soil. From Nova Scotia to Manitoba, Louisiana to North Carolina.

Poisonous properties. Reported as poisonous.

COMPOSITAE Adans. Thistle Family

Herbs or rarely shrubs, flowers borne in a close head on the receptacle, surrounded by an involucre of a few or many bracts, anthers usually united into a tube. syngenesious, sometimes caudate, calyx adnate to the ovary. Limb crowning the summit in the form of capillary or plumose bristles or chaff called the pappus. corolla tubular or strap-shaped, when tubular, usually 5-lobed, ligulate or biciliate in one small division of the family, the flowers of a head may be all alike when they are called homogamous, or of two kinds, heterogamous, bracts or scales on the receptacle are often present, flowers inside of the rays are disk flowers and a flower without rays is said to be discoid, stamens 5 or rarely 4, style 2-cleft at the apex or in sterile flowers usually entire, fruit a dry, indehiscent acheneum containing a single seed without endosperm.

A large family consisting of 840 genera and 13,000 species, found in all parts of the world. This is the largest order of flowering plants. Sometimes it is divided into the families *Cichoriaceae*, *Ambrosiaceae* and *Compositae*.

A few of the plants of the family are medicinal. Inulin is obtained from




Fig. 38a. Illustrating structure of Compositae (Thistle) (*Cirsium altissimum*). 1, head; 2, leaf; 3, outer bracts of head; 4, inner bracts; 5, single flower; a, achene; b, pappus; c, tubular corolla with 5 lobes; d, anthers; e, style; 6, style enlarged with two stigmas and pollen grains on style; 7, single pollen grain cut lengthwise to show pollen grains and tailed appendages; 8, single pollen grain. (Charlotte M. King)

the *Inula Helenum*, native to Europe and occasionally naturalized in the northern states. It is a mild tonic and contains *inulin* $C_6H_{10}O_5$, *helennin* and a volatile oil. The pebbly root (*Chrysanthemum Pyrethrum*) contains *pyrethrin* with a pungent taste, which, according to Dunstan, is apparently identical with *piperoytin* $C_{15}H_{21}NO_2$, used for toothache, the flowers of Roman chamomile (*Anthemis nobilis*) are used as a tonic and stomachic. However, German chamomile (*Matricaria chamomilla*) is sometimes substituted for the preceding and contains *anethidin* and a deep-blue volatile oil. Santonica, a species of wormwood (*Artemisia Cina*), contains *santonin* $C_{15}H_{21}O_3$, and *cinerol* $C_{15}H_{21}O_3$, is

found in the volatile oil, an anthelmintic. *Artemisia maritima*, *A. pontica*, *A. Absinthium*, *A. biennis* and *A. Abrotanum* are also used for the dislodgement of worms. The latter contains the alkaloid *abrotanin* $C_{21}H_{22}H_2O$. Sage brush (*Artemisia tridentata*) and other species produce sneezing.

Wormwood (*Artemisia Absinthium*) is a stimulant and tonic, the volatile oil produces cerebral disturbances and enters into the familiar composition of absinthe, made by the French, and contains *absinthin* $C_{18}H_{26}O_4$. Arnica root (*Arnica montana*) native to arctic Asia and America, is used as a popular remedy for chilblains and bruises and contains the bitter principle *arnicin* $C_{30}H_{50}O_8$. The tincture is liable to produce a form of dermatitis. Dandelion root (*Taraxacum officinale*) is used as a mild laxative and tonic and contains a bitter principle *taraxacin*, and *taraxacerin* $C_9H_{16}O$. *Lactucarium*, the milky juice from several species of the genus *Lactuca* occurs in lettuce, in which is also found *lactucoparin*, a bitter acid substance and *lactucol* $C_7H_{14}O$. Colts foot (*Tussilago farfara*), a bitter astringent containing much mucilage, is used for asthma. The cistus (*Santarea Lappa*) produces flowers with thistle-like heads and large roots, the latter of which are used as a perfume and an incense, according to Kraemer, it contains a ketone. The musk tree (*Olea argophylla*) of Tasmania, whose leaves emit a musk-like odor, grows to a height of 20 feet and is often 1 foot in diameter. The wood takes a nice polish. Other trees of the order are found in the genus *Senecio* (*Senecio Forsteri*). The genus *Baccharis*, found along the sea coast, is shrubby. The *B. cordifolia* contains *baccharin*, an alkaloid poisonous to sheep. It is the Mio Mio of South America.

The button snake root (*Liatris spicata*) has been used as a remedy for snake bites, but it probably contains no antiodotal properties whatever. Boneset (*Eupatorium perfoliatum*) and other species much used in domestic medicines for colds and in large doses are emetic. They are tonics, emetics, cathartics, and diaphoretics containing the bitter glucoside *eupatorium*. Horserweed (*Erigeron canadensis*) and other species are used as tonics and astringents.

Golden-rod (*Solidago odora*) is used to relieve colic, and guaiweed (*Grindelia squarrosa*) is beneficial in catarrhal affections. It is said to contain an alkaloid known as *grindelin*. Mada oil is obtained from tarweed (*Medicago sativa*). The tiger seed, the fruit of *Limnium abyssinica* is an important source of oil in Abyssinia and India.

The root of the burdock (*Arctium Lappa*) used by the laity as a remedy in skin diseases, contains a bitter glucoside, *lappin*. Chicory (*Cichorium Intybus*) is used to increase the appetite and to aid digestion. Rattlesnake weed (*Hieracium venosum*) is a popular antidote to the bites of poisonous snakes. Rattlesnake root (*Frenanthes alba*) is used as a remedy for toothache. The ragweeds (*Ambrosia artemisiifolia* and *A. triphylla*) are stimulants and astringents, the larger weed being also supposed to cause hay fever.

Durbar has demonstrated that the producing cause of hay fever may be pollen, and that pollen of all grasses, lilies of the valley, asters, and certain other plants may produce an irritation similar to that accompanying hay fever. He isolated an active principle which is believed to be a tox-albumin. A very interesting account of his experiments is given by Rochussen in the twenty-sixth volume of the Pharmaceutical Review.

According to Maxden, the *Helichrysum apiculatum* of Australia causes death from irritation and from the formation of hair balls. The African mari-

gold is poisonous to stock and the may weed (*Anthemis Cotula*), when applied to the surface, causes vesication. The European prickly lettuce (*Lactuca virosa*) contains a bitter principle, *hyoscyamina*, the prepared milky juice is called *Lactucarium*. It is listed as poisonous by Lehmann, a small amount of *hyoscyomin* also occurs in the cultivated lettuce.

The seeds of the sunflower (*Helianthus annuus*) are said to be diuretic. The oil cake from these seeds is used as stock food. Sneezeweed (*Helenium autumnale*) is used by the Indians to produce sneezing, a decoction made from it is used as a tonic. The mayweed (*Anthemis Cotula*) acts like chamomile, and is used as a tonic and stimulant in colic, when applied to the skin, it causes vesication. Yarrow (*Achillea Millefolium*), a stimulant and tonic, contains *achillein*, $C_{24}H_{34}N_2O_{12}$. The oil of tansy (*Tanacetum vulgare*) is used as an abortifacient, in many cases with fatal results. It contains the substance *thujone*, found in *Thuja occidentalis*. This has commonly been called *tanacetin* $C_{11}H_{14}O_4$, and is identical with *absinthol* obtained from *absinthum*, and with *salvol* from *salvia*. The *Cnicus benedictus* contains *cnicin* $C_{25}H_{37}O_{10}$.

Of the many cultivated plants of this order used for ornamental purposes, the best known in the northern states are probably the bachelor's button (*Centaurea Cyanus*), a native of Europe, ageratum (*Ageratum conyzoides*), Chinese aster (*Callistephus hortensis*), garden daisy (*Bellis perennis*), and dahlia (*Dahlia variabilis*), native to Mexico, running into many varieties. The last named produces thickened roots, that contain a great deal of *inulin*, $C_6H_{10}O_5$. Among the species of the order which are common in the gardens are the zinnia (*Zinnia elegans*) golden glow (*Rudbeckia laciniata*) sunflower (*Helianthus annuus*), coreopsis (*Coreopsis tinctoria*), gaillardia (*Gaillardia puchello*), dusty miller or cineraria (*Senecio Cineraria*), common cineraria (*S. cruentus*) from the Tenerife, and the purple ragwort (*S. elegans*) from the Cape of Good Hope, yarrow (*Achillea Ptarmica*), whiteweed (*Chrysanthemum Parthenium*), several species of the genus *chrysanthemum* from Japan (*C. sinense* and *C. indicum*), Marguerite (*C. frutescens*) and summer chrysanthemum (*C. coronarium*), marigold (*Tagetes erecta*), the pot marigold (*Calendula officinalis*), cosmos (*Cosmos bipinnatus*) of Mexico. The marigold (*Calendula officinalis*) cultivated in country gardens contains *calendulin* $C_4H_{12}O_5$.

Of the economic plants, the following are the more important: the cultivated lettuce (*Lactuca sativa*), probably native to Asia, slightly narcotic, comprising many varieties; the safflower or saffron (*Carthamus tinctorius*), native to Egypt, used for dyeing; marigold flowers (*Calendula officinalis*) sometimes used as an adulterant for saffron; wormwood (*Artemisia Absinthium*), used for making absinthe; tarragon (*Artemisia Dracunculus*) used as a pot herb and in making vinegar, a native of Asia, but cultivated now in Holland and England; the Jerusalem artichoke (*Helianthus tuberosus*), with thickened roots, cultivated as food for hogs and stock, native to North America, the great sunflower (*Helianthus annuus*) the seeds of which are eaten in Russia, and from which an oil is obtained; the chicory (*Cichorium Intybus*), the burdock (*Arctium minus*), cultivated for its thick root which is sometimes dried and mixed with coffee and also used as a forage plant, the endive (*Cichorium endivia*) cultivated as a winter salad plant; the cardoon (*Cynara Cardunculus*), also known as the European artichoke, used as a vegetable; the thickened scales being used as food, salsify (*Tragopogon porrifolius*) a vegetable, and the dandelion (*Taraxacum officinale*) used like lettuce.

The Compositae are divided into two sub-families the *Tubuliflorae* with corolla tubular in all the perfect flowers, 5-lobed, rarely 3-4-lobed, ligulate only in the marginal flowers, called the ray flowers, which are absent in some species the *Liguliflorae* with corolla ligulate in all the flowers of the head and all the flowers perfect, herbs with milky juices and alternate leaves. The former contains the tribes *Vernoniaeae*, with the large genus *Vernonia* of 450 species mostly of the tropics, the *Eupatorieae*, containing *Eupatorium*, *Asteraceae*, containing *Solidago*, *Aster*, *Frigeron*, *Grindelia*, *Bigelovia*, *Inuleae*, containing *Inula*, *Antennaria*, *Helianthus*, containing *Siphium*, *Parthenium*, *Helianthus*, *Bidens*, *Coreopsis*, the *Heleniaceae* with *Helenium*, *Actinella*, the *Anthemideae* with *Anthemis*, *Achillea*, *Tonacetum*, the *Senecioneae* with *Senecio*, *Tussilago* the *Cynaraeae* with *Archium* *Cnicus*, *Cirsium* and *Centaurea*. The *Liguliflorae* contain but one tribe, the *Cichorieae*, which contains the genera *Taraxacum*, *Cichorium*, *Lactuca* *Hieracium*, *Sonchus* and *Scorzonera*.

Genera of Compositae

Corolla ligulate in all of the flowers of the head, flowers perfect

LIGULIFLORAE CICHORIEAE

- | | | |
|--|----|-------------------|
| Pappus none | 1 | <i>Cichorium</i> |
| Pappus composed of capillary bristles. | | |
| Flowers yellow achenes not beaked | 2 | <i>Sonchus</i> |
| Flowers yellow, purplish or cream colored. | | |
| Achenes beaked | 3. | <i>Lactuca</i> |
| Achenes not beaked | 4. | <i>Lygodesmia</i> |

Corolla tubular in perfect flowers, 5 or rarely 3 or 4-lobed, ligulate only in the marginal flowers.

TUBULIFLORAE

Stamens distinct or nearly so.

AMBROSIEAE

Staminate and pistillate flowers in the same head

5. *Iva*

Staminate and pistillate flowers in a separate head.

Involute of pistillate heads with several tubercles

6. *Ambronia*

Involute of pistillate flowers forming a bur

7. *Xanthium*

Stamens generally united by their anthers into a tube around the style.

Anthers not tailed.

Style branches thickened upward, papillose

EUPATORIEAE

Achenes 3-5 angled, flowers discoid

8. *Eupatorium*

Achenes 8-10, ribbed or striate.

Bracts of involucre in several series.

9. *Lactis*

Bracts of involucre in 2 or 3 series

10. *Trips*

Style branches of perfect flowers flattened with terminal appendages.

ASTEREAE

Ray flowers yellow

Pappus of scales or awns

11. *Grindelia*

Pappus of numerous capillary bristles

12. *Solidago*

Ray flowers not yellow, bracts nearly equal

14. *Erigeron*

Bracts in several unequal series.

13. *Aster*

Style branches truncate or hairy appendaged.

HELIANTHEAE

Pappus wanting or minute crown

15. *Rudbeckia*

Pappus of 2 persistent downwardly barbed awns or tubes

16. *Bidens*

- Style branches truncate or with hairy tips. HELENÆÆ
 Bracts of the involucre spreading.
 Receptacle naked; bracts of involucre spreading or reflexed at maturity. 17 *Helenium*
 Bracts of the involucre united. 18. *Dysodia*
 Style branches mostly truncate with brush hairs on the tip ANTHEMIDÆÆ
 Pappus of short scales or a crown.
 Scales of involucre scarious and imbricated
 Receptacle chaffy.
 Heads small, involucre obovoid or campanulate, achenes flattened. 19. *Achillea*
 Heads large achenes tetecle 20. *Anthemis*
 Heads solitary or corymbose
 Receptacles not chaffy
 Ray flowers usually present, conspicuous 21 *Chrysanthemum*
 Ray flowers inconspicuous 22. *Tanacetum*
 Heads small spikeate or racemose paniculate 23 *Artemisia*
 Style branches truncate or triangular with brush hairs.
 Heads radiate or discoid.
 Pappus of capillary bristles. SENECIONIDÆÆ
 Heads showy, leaves opposite 24. *Arnica*
 Heads usually showy, leaves alternate 25. *Senecio*
 Style branches short or united, anthers caudate CYNARÆÆ
 Achenes base fixed 26. *Arctium*
 Involucral bracts pointed or prickly
 Receptacle densely bristly 27 *Cirsium*
 Receptacle honey-combed 29 *Silybum*
 Achenes attached laterally.
 Pappus of short scales 28. *Centaurea*

LIGULIFLORÆ

Herbs with milky juice. Corolla ligulate in all of the flowers of the head, and all of the flowers perfect.

1 *Cichorium* (Tourn.) L. Chicory

Erect branching perennial or biennial herbs with alternate leaves, involucre of two series of herbaceous bracts, the inner of 9-10 scales, the outer of 5 short spreading scales, receptacle flat, naked or slightly fimbriate. Flowers bright blue purple or pink, rays 5 toothed. Achenes striate pappus of numerous small chaffy scales forming a crown.

About 8 species of the old world

Cichorium Intybus L. Chicory or Succory

A branching perennial with deep roots and alternate leaves basal leaves spreading on the ground stem leaves oblong or lanceolate, partly clasping.

Distribution. Common along roadsides and in fields and waste places from New England to Canada and Nebraska, especially where chicory has been cultivated. It has become a troublesome weed in Wisconsin and Minnesota. It is allied to endive (*Cichorium Endivia*), cultivated as a salad plant.

Poisonous properties. When fed in large quantities it imparts a bitter flavor to milk and butter. It contains the bitter glucoside *chicarin* $C_{32}H_{44}O_{10}$. Chicory root is used as an adulterant of coffee.



Fig. 439a. Dandelion (*Taraxacum officinale*). 1, Single head during flowering; single head after flowering. 2, Single flower with corolla, stamens and style. 3, Achensum. 4, Receptacle and single achene. (After Strasburger, Noll, Schenk and Schimper.)

2. *Sonchus* (Tourn.) L. Sow Thistle

Annual or perennial herbs with alternate, mostly amplexicaule-clasping, entire dentate, lobed or pinnatifid leaves with soft prickly margins; flower heads in corymbose or paniculate clusters, involucre bell-shaped, scales imbricated in several rows; receptacle flat and naked; achenes oblong, more or less flattened, 10-20-ribbed; pappus of soft white capillary bristles.

About 45 species of the old world.

Sonchus oleraceus L. Annual Sow-thistle

Annual or perennial succulent herbs with leafy stems, smooth and glaucous with corymbose or umbellate heads of yellow flowers. Stem leaves dentate, runcinate-pinnatifid, terminal segments large and triangular, heads numerous, flowers pale yellow, occurring in summer and fall.

Distribution. Common in fields and waste places throughout North America, except far northward. Also from Mexico to South America.

Sonchus arvensis L. Field Sow-thistle

A glabrous perennial, producing deep creeping root stock; stem leafy, branched, basal leaves runcinate-pinnatifid, spiny toothed, clasping by a heart-shaped base; flowers yellow; achenes transversely wrinkled.




Fig. 439c. Chicory (*Chicorium Intybus*) a, part of plant with several heads, b, single head side view c, single flower with strap shaped corolla, d, achene with small chaffy scales. (U. S. Dept. Agr.)

Distribution. Common in eastern states, Canada (Manitoba) and occasionally in Wisconsin, Illinois, Iowa and Utah.

Poisonous properties. The plants are more or less bitter and not liked by cattle. Their milky juice probably contains some active principle.

3. *Lactuca* (Tourn.) L. Lettuce

Tall, leafy-stemmed herbs with milky juice and alternate leaves, flowers white, yellow or blue, in panicle heads; involucre cylindrical, bracts imbricated in two or more series, receptacle flat naked; anthers sagittate at the base; achenes oval, oblong or linear, abruptly contracted into a beak, dilated at the apex, bearing a soft white capillary or brown pappus.

About 90 species natives of the northern hemisphere. Garden lettuce (*L. sativa*), native to Europe, is cultivated.

Lactuca Scariola L. Prickly Lettuce

Tall, erect herbs, annual or winter annual, 2-6 feet high, simple or branched except the lower part of the stem which has stiff bristles, leaves glaucous, green, smooth except the midrib which is beset with weak prickles; lanceolate

to oblong in outline with spinulose, denticular margins occasionally sinuate toothed, sometimes pinnatifid, base sagittate clasping, leaves becoming vertical by a twist, the leaves are not twisted in shady situations, flowers in small open panicle heads, each head has from 4-18 yellow flowers, achenes flat, striate nerved, obovate, oblong, produced in long filiform beak which is paler in color than the achene, pappus consists of delicate white bristles arising at the end of the beak.

Distribution. Prickly lettuce is a native of temperate and southern Europe, Canary Islands, Maderia, Algeria, Abyssinia, and the temperate regions of eastern Asia. It was introduced into North America about 1863.



Fig. 440. Prickly Lettuce (*Lactuca Scariola*). After Fitch.

Lactuca canadensis L. Wild Lettuce

A tall, leafy smooth or occasionally somewhat hairy biennial 4-9 feet high, leaves 6-12 inches long, stem leaves sessile or auriculate clasping, the upper leaves smaller, lanceolate acuminate and entire, heads with about 20 flowers in spreading panicles involucre cylindrical, rays yellow, achene somewhat longer than the beak.

Distribution. In moist places, borders of thickets and in fields from Nova Scotia to Manitoba south to Arkansas, Louisiana and Georgia.

Lactuca fulchella (Pursh.) D. C. Blue Lettuce

A glabrous perennial, with milky juice, simple stem from 1-2 feet high, sessile, oblong or linear lanceolate entire leaves the lower runcinate-pinnatifid heads corymbose paniculate, peduncles with scaly bracts, scales of the involucre imbricated in 3 or 4 ranks, flowers blue, achenes oblong lanceolate, somewhat flattened.

Distribution. Native to the plains from western Iowa north to Manitoba and west to the Great Basin and California, and rare as far east as Michigan. A troublesome weed in grain fields of Montana and Utah.

Poisonous properties. The wild blue lettuce is common in the west but is not liked by cattle although sometimes eaten by sheep. It has been regarded

with suspicion. The cultivated lettuce contains several active principles as follows: *Lactucorol* $C_{16}H_{30}O$, *lactucol* $C_{21}H_{42}O$, a small amount of *hyoscyamin*. The bitter taste of *L. canadensis* is due to *lactucrim* $C_{22}H_{44}O_8$, and *lactucopicroin* $C_{44}H_{88}O_2$. I quote Dr. Millspaugh in regard to the physiological action of *L. canadensis*: "Lactucarium, in large doses, causes delirium, confusion of the brain, vertigo and headache, dimness of vision, salivation, difficult deglutition, nausea and vomiting, and retraction of the epigastric region, with a sensation of tightness, distension of the abdomen, with flatulence, urging to stool followed by diarrhoea, increased secretion of urine, spasmodic cough, oppressed respiration, and tightness of the chest, reduction of the pulse ten to twelve or more beats, unsteady gait, great sleepiness, and chills and heat, followed by profuse perspiration." The *L. virora* a wild lettuce of Europe, and occasionally in the Mississippi Valley but never abundant westward, is regarded in Europe as poisonous.

4. *Lygodesmia* D. Don.

Low smooth perennial herbs with linear leaves or the lower somewhat pinnatifid, the upper of scales, heads 3-12 flowered, a single one terminating the branch, flowers purple or pink, achenes smooth or striate, pappus of copious, somewhat unequal simple bristles.

About 6 species of western North America.

Lygodesmia juncea (Pursh.) D. Don

A tufted smooth, frequently glaucous perennial, a foot or more high, coming from a thick woody root, copious milky juice and stems lower leaves rigid, linear lanceolate, entire, the upper scale-like; heads erect with purple flowers; achenes narrow-ribbed, pappus light brown.

Distribution. Common on the plains from the Missouri river to western Montana, Northwest Territory and east to the St. Croix river in Wisconsin. This has been reported as a troublesome weed in corn fields in northwest Iowa, troublesome also in Colorado and Montana.

Poisonous properties. The plant is bitter like many others belonging to sub-family *Cichoriaceae*. The milky juice no doubt contains some deleterious properties. The plant is not liked by stock.

Professors Chesnut and Wilcox say with reference to the species in Colorado

"This species, sometimes known as prairie pink, grows abundantly on dry prairies and plains in Park, Sweet Grass, Gallatin, Meagher, Lewis and Clarke, Choteau, and Teton counties. The general distribution of the plant is from Minnesota to New Mexico and Nevada. It has been suspected by stockmen both in Montana and Utah of being poisonous to stock. The plant was not investigated, but it was ascertained that the milky orange-colored juice of nonflowering plants gathered at Toston was extremely bitter and disagreeable to the taste."

TUBULIFLORÆ

Corolla tubular in all the perfect flowers, 5-lobed, rarely 3-4-lobed, or ligulate only in marginal flowers, called the ray flowers which are absent in some species.



Fig. 440a. Boneset (*Eupatorium altissimum* L.) Common in the Mississippi valley. At the right: Mountain Sunflower (*Helianthus Magnusii* Connor) in Mississippi valley. (Charlotte M. King).

Iva L. Marsh Elder

Herbaceous or some shrubby plants, pistillate and staminate flowers in the same head. The lower leaves opposite, the upper alternate, flowers greenish, rays absent, subtended by an involucre of hemispherical or cup-shaped bracts, achenes obovoid or lenticular, without pappus.

About 12 species of western and southern America.

Iva axillaris Pursh. Small flowered Marsh-elder

A smooth or sparingly pubescent perennial with herbaceous stems, from 1-2 feet high, with woody roots. Leaves sessile, entire or nearly so, obovate, oblong or linear-oblong the lower opposite, the upper smaller and alternate, heads generally solitary in the axils of the leaves, short petioled, involucre hemispherical, pistillate, flowers with tubular corolla.

Distribution. Common especially in the saline soils from Nebraska to the Dakotas. British Columbia, California and New Mexico.

Iva xanthifolia Nutt. Marsh Elder

An annual, from 1-8 feet high, stem frequently pubescent when young, all the leaves opposite, rhombic, ovate or the lowest heart-shaped, doubly serrate or cut-toothed, obscurely lobed, the upper surfaces minutely scabrous, canescent beneath, especially when young. Petiole frequently ciliate at its upper end, flowers born in spike-like clusters forming a compound panicle, heads small, crowded. Outer bracts of the involucre broadly ovate, greenish, inner membranaceous, achenes glabrate.

Distribution. In alluvial ground or along streams, Saskatchewan and Nebraska to New Mexico, Utah and Idaho. In the western part of the state of Iowa this weed is extremely common as in Woodbury, Harrison, Monona and Fremont counties. It occupies not only the vacant lots but is found in the streets and cornfields. To the west in Nebraska it becomes increasingly abundant and in the irrigated fields of portions of Colorado it is frequently 8 feet high. It is common in the Red River Valley of the North and other parts of Minnesota and Dakota, and Manitoba. It is a most aggressive weed.

Poisonous properties. This plant and the preceding produce an unusually large amount of pollen and have been looked upon as in part responsible for hay fever, the pollen being simply an irritant of the nasal mucous membrane.



Fig. 441. False ragweed (*Iva axanthifolia*)
The pollen is possibly a cause of hay fever (Dewey, U. S. Dept. Agr.)

6. *Ambrosia* (Tourn.) L. Ragweed

Herbs, leaves alternate or opposite, lobed or dissected, flowers in heads, fertile, 1-3 together, sessile in the axils of leaves or bracts, involucre of the pistillate flowers top-shaped, ovoid or globose, closed, 1-flowered, armed with 4-8 tubercles or spines, corolla none, pappus none, involucre of the staminate flowers saucer-shaped, 5-12 lobed, many flowered, receptacle nearly flat or with

filiform chaff, corolla funnelliform, 5-toothed, anthers but slightly united, achenes ovoid.

The twelve species are mostly native of North America.

Ambrosia trifida L. Great Ragweed, Ironweed

A stout scabrous, hispid or nearly glabrous annual, 3-12 feet high, leaves opposite and petioled, 3-nerved, deeply 3-5 lobed, the lobes ovate, lanceolate and serrate, the upper leaf sometimes ovate and unlobed, flowers monoecious,



Fig 442 *Ambrosia* showing flowering stalk, pistils, stamens, and longitudinal sections of stamens in perianth and ovary. (After Engelm.)

staminate borne in spikes surrounded by the larger bract-like leaves, the involucre is turbinate to ovoid, 5-7 ribbed, beaked, each rib bearing a tubercle near the summit, the involucre enclosing a single achene.

Distribution The greater ragweed is distributed from Quebec to Florida throughout the Atlantic region to Texas, common throughout the Mississippi Valley west to Colorado and Northwest Territory, in Manitoba and Saskatchewan.




Fig. 443. Tall ragweed (*Ambrosia trifida*).
(U. S. Dept. Agr.)



Fig. 444. Small Ragweed (*Ambrosia artemisiifolia*). In the lower left hand corner is a bur that contains the seed, upper right hand corner a flower. In middle, branch and leaf. Probably poisonous. (Holm-Britton.)

Ambrosia artemisiifolia L. Hogweed. Bitterweed

A puberulent or hirsute branched annual 1-3 feet high, leaves thin, once to twice pinnatifid, the upper alternate, the lower usually opposite, pale or canescent beneath. flowers monoecious, the staminate above and the pistillate in the lower axils of the leaves, the fertile heads are obovoid or globose, short beaked and 4-6 spined.

Distribution. A troublesome weed in northern states. Its distribution in North America is from Nova Scotia to Florida throughout the Atlantic states and Mississippi Valley, in the Rocky Mountain region and west to British Columbia, also in Mexico and the West Indies and South America.

Poisonous properties. The greater ragweed is regarded as especially

troublesome as an exciting cause with reference to hay fever. The following from Dr. Johnson on this point is of interest:

These plants are said to be stimulant, tonic, and astringent. A decoction has been used, chiefly in domestic practice, as a topical astringent in chronic catarrhal affections.

Of late years *A. arvensifolius* has attracted considerable attention on account of its real or assumed agency in the production of hay-fever. The plant produces pollen in great abundance, which is extremely irritating to the air-passages of many people, and is capable of exciting asthmatic attacks in susceptible persons. Now as the weed is so very abundant, and its time of flowering coincident with the greatest development of hay-fever, the relation of cause and effect has been asserted by many writers. That it may be so in a certain proportion of cases is quite probable, but that its influence in this direction has been overrated is still more probable. The pollen of all plants is irritating to the air-passages of sensitive people, but probably little more so than any other dust of an organic character, and the proportion which ragweed pollen in the air of any specified locality bears to that of all other plants combined must be very small indeed. Much less still must its proportion be to other pollen and organic dust in the air of cities, where this affection has become endemic—and fashionable.

The ragweeds, marsh elder, goldenrod, and chrysanthemum, as well as the pollen of some grasses, are said to produce hay fever. In recent years a toxic substance has been isolated which belongs to a class of poisonous substances known as toxalbumin.

The fact that this troublesome disease is caused by a poisonous toxin has led to a study of serum treatment by Dunbar* who has produced an antitoxin which he calls "pulantin," and Weichardt** another called "grammol." Pulantin is obtained from the blood serum of horses which have been immunized with the pollen toxin. Dunbar's hay fever serum is sold both as a powder and as a fluid. There are those, however, who believe that hay fever is not due to poison by pollen toxin so that this treatment can be of no use. A Wolf-esser*** attributes the action of the serum not to antitoxins but to colloidal substances. Weichardt prepared the serum from the blood of herbivorous animals. Statistics of The Hay Fever Union of Germany for 1906 indicate that the results with the treatment of "grammol" were favorable.

A correspondent from Nebraska sent to the writer a specimen of the annual ragweed stating that it was abundant in his pasture and that where cattle used it as forage, the mouths of these animals became very sore. The plant is bitter and possibly may be irritating at times. There were no parasitic fungi on the specimens sent us.

7 *Xanthium* (Tourn.) L. Cocklebur Clothur

Coarse low branching annual herbs with alternate toothed or lobed petioled leaves, monoecious flowers. staminate flowers with a short involucre of several distinct bracts, receptacle cylindrical, pistillate flowers with a closed involucre covered with hooked prickles, 2-celled, 2-flowered, in fruit forming a bur, achenes oblong flat, without pappus.

The 12 species are widely distributed

Xanthium spinosum L. Clothur

A pubescent branched annual herb with slender yellow 3-parted spines in the axils, leaves lanceolate or ovate lanceolate white downy underneath, bur oblong cylindrical, armed with single short beak and numerous glaucous prickles.

* Deutsch. Med. Woch. 1903 340.

Berlin Klin. Woch. 1903 4 25, 26, 28.

** Klinisch-ther. Woch. 1903 1452.

*** Das heufieber, sein weesen und seine Behandlung. München 1906. In this connection the very excellent review on Serum Therapy in E. Merck's Annual Report, 1902, Vol. 23 should be consulted.

Distribution. Common in waste grounds from Ontario to Missouri and Texas to Florida. Native to the old world.



Fig. 445. Thorny Cuckoo-bur (*Xanthium spinosum*). After Hochstein.

Xanthium Strumarium L.

A low, rough, branching annual from 1-2 feet high, leaves slender, petioled, broadly ovate or orbicular, 3-5-lobed, both surfaces rough, but oblong, smooth or nearly so, with two straight or nearly straight beaks.

Distribution. In waste places along the Atlantic seacoast. Native to Europe.

Xanthium canadense Mill. Cockle Bur

A coarse rough annual from 1-3 feet high, stem marked with brown punctate spots, leaves alternate, cordate or ovate, 3-nerved, long petioled, flowers monoecious, staminate and pistillate flowers in different heads, the pistillate flowers clustered below, the involucre of the staminate flowers somewhat flattened of separate scales, receptacles cylindrical, scales of the fertile involucre closed in fruit, 2-beaked, containing 2 achenes the bur is densely prickly and hispid, achenes oblong without pappus.

Distribution. In Iowa this species is very common along the sandy bottoms of our streams and river courses. It is less troublesome in uplands but here and there it does occasion some trouble even in the central and southern part of the state. However, in southern and southeastern Iowa the weed is often quite troublesome in cornfields, coming up in enormous quantities. In Texas, too, it appears in the very richest soil. Its distribution in North America is

from Nova Scotia to North Carolina, southwest to Texas and west through Colorado, Utah and Nevada, and north to Saskatchewan and Manitoba. Most abundantly developed in the Mississippi Valley region from Minnesota to Texas.

Poisonous properties. The injury from this plant probably comes largely from its mechanical action. As the involucre is indigestible, its barbs sometimes injure an animal feeding upon it. Stock will probably not eat very much of it, but on account of the hooked awns of the involucre the animal may have considerable difficulty in removing them. The hairs of the plant cause itching. Several cases of poisoning of hogs, probably due to this plant, have been reported to me. The plant contains the poisonous glucoside *xanthostrumarin* which resembles *datiscin*. On heating the odor of succinic acid is given off. According to Chesnut, the young seedlings of three species of cockle bur, among them our Canadian cockle bur (*Xanthium canadense*) are poisonous to hogs. Dr. Bitting was unable to find a poisonous substance in the growing plants. He thinks the injurious properties are largely mechanical.



Fig. 446. Cockle-bur (*Xanthium canadense*)
Cause of mechanical injuries to animals.
(Dewey, U. S. Dept. Agr.)

A writer in the *American Agriculturist* says:

When the seed is ripe the bristles are very hard and sharp, and in the stomach and intestines of the animal, mat or felt together by the aid of their barbs, forming large balls, which

obstruct the intestines and prove fatal. In these cases, death is evidently not from any poisonous quality of the hay, but is due to mechanical causes. While Cocklebur may not be poisonous, it may in a similar manner kill swine by mechanical obstructions. When the burs are ripe, they readily leave the plant, and attach themselves to the coat of any passing animal. The plant is especially annoying to sheep owners, as these burs become entangled in the wool. Those who have traveled in Texas and Mexico, soon make acquaintance with the cockle bur. Horses and mules, while grazing for a single night, will have their tails clotted with these burs, and converted into a useless club. It can be understood, that if swine eat the burs in considerable quantity, the pieces of them can, by means of the prickles, form masses which may prove fatal. Both on account of its probable danger to swine and its injury to sheep, there should be an united effort to destroy the plant. Being an annual, its extermination would not be difficult. If the plants are cut down before the seed is ripe, new ones can not appear unless the ground is re-seeded.



Fig. 447. Boneset (*Eupatorium alticarpum*). A common plant in woods. It is supposed by some to cause milk fever or trembles. Branches with numerous small heads. (Low Pam. mel.)

B. *Eupatorium* (Tourn.) L. Boneset

Erect perennial herbs with opposite, whorled or alternate leaves, often sprinkled with resinous dots, flowers in corymbose heads, white, bluish, or purple bracts of the involucre in two series, receptacle naked, corolla regular tube short 5-lobed, branches of the style slender, thickened upward or club-shaped, very minutely and uniformly pubescent, with stigmatic lines indistinct; achenes 5-angled, truncate, pappus of numerous fine capillary bristles, arranged in one row.

About 450 species in warm temperate and tropical regions. An oil is made from the southern *E. capillifolium*, which has an aromatic, pepper-like odor.

The boneset (*E. perfoliatum*), is commonly used as a domestic remedy, employed in catarrhal affections and fevers. The Joe-Pye weed (*E. purpureum*) is used in urinary affections.

Eupatorium urticifolium Reichenb. White Snake-root

A smooth or nearly smooth branching herb from 2-3 feet high leaves opposite, thin rounded, cordate at the base or abruptly narrowed into a slender petiole, coarsely and sharply toothed, heads in ample and loose corymbose clusters flowers white, involucre narrowly campanulate bracts linear, acute or acuminate.

Distribution. Common in rich woods from New Brunswick to South Dakota and Nebraska to Indian Territory and Louisiana.

Poisonous properties. This plant is said to produce the disease known as trembles in cattle, horses and sheep, and milk sickness in people. Mr E. L. Moseley states that:

Milk-sickness is known to be due to the use of milk butter cheese or meat of animals afflicted with the trembles, but what causes the trembles has not been well understood. It has long been known that only the animals allowed to run in the woods were affected, and experiments showed that certain woods were very dangerous, while others were safe. People who came from Pennsylvania with a view to settling here returned to their own State on learning of the peril of pasturing animals in Ohio. To this day many woods in this district are not pastured, because animals would soon die if turned into them.

The Eupatoriums are not palatable. Anyone who has tasted boneset will admit that this is true of *Eupatorium perfoliatum*. In the South I have observed that animals leave *Eupatorium spirale* untouched even where they have been compelled to go to eat almost every other green thing in reach. In northern Ohio I have found *Eupatorium urticifolium* the white snake-root, growing abundantly in a number of woods where animals were pastured but no sign of their having eaten it. But if the pasture becomes more tame some are likely to eat it.

On the 8th of last October I visited a piece of woods in Sandusky county where there was nothing fit for an animal to eat the principal herbs being cattle white snake-root poke and black nightshade, with some rich weed, lamb and hellebore. Every plant of snake-root had been nipped off so that I did not see one more than about half the normal height. This had probably been done by cattle from the adjoining pasture who were denied the accustomed spending a portion of his sunny days in the shade of the woods. A few weeks before my visit a man and his wife who had been using butter made from the milk of cows in this pasture had milk-sickness and the wife died.

Joshua Huff Townsend township, Sandusky county, did not think trembles were due to any weed, until he found that western sheep which he raised got sick and die the white snake-root and died of trembles. Sheep whose ancestors had long been in the region did not eat it, and did not have trembles. Since that he has been destroying the weed.

Mr William Hamelitt of Bloomingville informs me that about 1840 when there was so much discussion of the subject the boys of the neighborhood used to assemble evenings at the old lime-kiln southeast of Castalia and experiment on dogs. They would boil or steep the white snake-root and putting the extract in milk give it to the dogs, in which it would induce the trembles a large number were killed in this way. Some one experimented on sheep with the same result. He informs me also that a Mr. Richmond (who did not live) saw that the weed was the cause of trembles) chewed some of the weed and died after suffering for about four weeks.

On November 26th my pupil Oscar Kuback, using snake-root I had recently gathered, broke up the stems and leaves of two plants and soaked them over night in about a pint of milk, of which he gave about a gill at about 9 A. M. to his tom-cat. The cat took about one half of .

About 9-10 it seemed to take effect and he tried very hard to cough but could not. He took long, deep breaths. He was quiet and wanted to sleep very hard. All of a sudden he would tremble very hard, then again very little. A watery fluid passed from his eyes and mouth. He chose a spot in the sun and when driven away walked back in a staggering manner. He had no appetite. His senses seemed to be duller as he did not care for anything. He went to sleep about 10-30 but did not sleep sound. He seemed to be in an unconscious state for the rest of the day. The next morning about 10-30 he walked about three rods and there died about noon.

Mr Mosely estimates that 5000 animals have succumbed from the disease in a small area in northern Ohio. There are many cases commonly in this same region from milk sickness.

Dr Albert C. Crawford of the Bureau of Plant Industry, investigated the so-called milk sickness occurring in Alton, Illinois, which resulted in the death of about 50 head of cattle, and gives the following conclusions:

To sum up, it certainly can not be said that it has been proved that milk sickness is due to any constituent of *E. verticillatum*. The transmission of the disease by eating some quantity of meat or milk of animals sick with the "trembles" and the fact that cooked meat or boiled milk does not produce this disorder point primarily rather to a parasitic origin while the fact that *Eupatorium verticillatum* is abundant in areas where the disease is not known and absent in some milk-rich districts also indicates that the plant has no relation to the disease. If it does, it would be only an accidental carrier of some pathogenic organism. According to reports the same flora may be in areas in which "trembles" occur as in those free from it and milk sickness is also said to occur where no vegetation grows (improved pastures). The disease also has disappeared from an area after simply clearing the woodland where it occurred and turning it into pasture. Again, severe epidemics have occurred in winter when the foliage has disappeared, which would tend to exclude the higher non-herb green plants as the cause of this disorder. In fact, all the evidence in hand is against the causation of this disease by such plants, and certain analogies with cases of botulism suggest a somewhat similar cause. If there is any truth in the statement that cattle caused in pasture to night air exposure contract the disease this fact might suggest the more or less direct connection of some night-gathering as a carrier of the bacillus, and certain parasites are supposed to be associated with certain locusts. Very little is known chemically of *Eupatorium verticillatum*.

And it is seen to confirm the investigation by Dr. Bitting of the Indiana Agricultural Experiment Station. It seems very doubtful that this plant causes milk sickness, since it is very common in many pastures in the west where trembles does not occur.

In regard to boneset (*E. purpureum* and *E. perfoliatum*) Dr. Johnson states as follows:

Of domestic remedies few are better known or more largely used than boneset. It is tonic, diaphoretic, emetic and cathartic, the different effects depending largely upon the size of the dose and mode of administration. The infusion, taken cold in moderate doses, is tonic and is employed in debility of the digestive organs and in convalescence. Taken warm in large doses, the infusion or decoction produces copious diaphoresis, and is employed in the acute stages of catarrhal affections and in fevers, especially those of an intermittent or remittent type. In still larger doses the warm infusion or decoction produces emesis or catharsis. These effects are however seldom sought.

E. purpureum or gravel root is said to be diuretic and to have been employed in urinary affections, but it has not attained an established reputation and is seldom used.

Boneset (*E. perfoliatum*) contains the glucoside *eupatorium*, the *E. purpureum* contains the glucoside *euparin* $C_{12}H_{21}O_5$.

Dr. T. H. M. gives an extended account of the medicinal qualities accompanied by the anatomical structure of this plant. *Eupatorium perfoliatum* according to Dr. H. M. is prescribed as a tonic and in large doses is an emetic.

9. *Trilisa* Cass. *Trilisa*

Erect perennial herbs, fibrous roots, leaves alternate and simple, heads in terminal corymbose panicles, discoid, 5-10-flowered, flowers white, receptacle flat, corolla regular 5-lobed, achenes nearly terete, 10-ribbed. A small genus closely allied to *Liatris*. Two species native to eastern North America.

Trilisa odoratissima Cass. Vanilla Plant

A rather stout, somewhat glabrous perennial, leaves pale obovate-spatulate,

* Merck's Rep. XVII:326-328. L. 1-11.

or oval, thickish and clasping head in corymbose clusters, achenes glandular pubescent.

Distribution. From Virginia to Louisiana.

Trilisa paniculata Cass. Hairy Trilisa

Viscid-hairy perennial, leaves entire, base lanceolate, narrowly oblong, acute, or obtusish, those of the stem small, heads paniculate achenes finely pubescent.

Distribution. From Virginia to Georgia and Florida.

Poisonous properties. The former plant has the odor of vanilla and contains the substance *cumarin* $\text{C}_9\text{H}_7\text{O}$. Dr. Johnson says

Odonatum deserves much more attention from the fact that it is largely used as an adulterant of smoking tobacco, than from any demonstrated medicinal virtues. There is abundant evidence to show that the leaves of this plant enter largely into the manufacture of many grades of smoking tobacco, especially those employed in our domestic cigarettes. And the author is convinced, from personal experience and observation, that the deleterious effects produced by smoking tobacco thus adulterated are much greater than those produced by the consumption of pure tobacco in even great excess. The inhalation of a few whiffs of the smoke from a cigarette made from this adulterated material produced the sensations are made in quick succession produces a train of cerebral sensations of an intoxicating character as much different from any effect of tobacco alone as could be imagined and prolonged use of such cigarettes inevitably produces great derangement of the digestive organs, very little resembling the dyspepsia induced by excessive use of tobacco, together with cardiac symptoms often of a distressing character. And again, the habit of smoking *cumarin* in this form appears to become more inveterate, more exacting, than that of the use of tobacco alone, so that the unhappy victim—for such he should be called—is never comfortable except when indulging. Hence it happens that cigarette-smoking in this country in its effects upon adolescents especially, is assuming the proportions of a great national evil, and is producing far more deleterious effects than in other countries where it is practiced to a greater extent but with different material.

10. *Liatris* Schreb. Button Snakeroot. Blazing Star

Perennial herb, usually from a corm-like tuber, leaves alternate entire, narrow, flowers spicate or racemose, discoid, scales of the involucre few or many imbricated in several series, the outer shorter, corolla regular, 5-lobed or 5-cleft, branches of the style exerted, achenes 10-nubbed, slender tapering to the base, flowers rose-purple, rarely paler in color.

Distribution. About 20 species in North America.

Liatris spicata (L.) Willd. Snakeroot

Smooth or somewhat hairy perennial, leafy stem, leaves near the lower 3-5 nerved, heads crowded in a long spike, 8-12 flowered, involucre cylindrical, bell-shaped, flowers purple, pappus not very plumose.

Distribution. In moist soil from New England to Florida, Kentucky, Arkansas, and South Dakota.

Liatris pycnostachya Michx. Snakeroot

Hairy or smooth perennial with a stout stem 3-5 feet high, leaves linear-lanceolate, the upper very narrowly linear, spikes 6-20 inches long, densely flowered, flowers purple, pappus not very plumose.

Distribution. Prairies, Indiana to Minnesota and southward.

Liatris punctata Hook. Western Snakeroot

A glabrous or sparingly pubescent perennial, stout rootstock, leaves rigid punctate, spike many flowered, heads 3-6 flowered, purple, bracts of involucre oblong, often ciliate on the margins, pappus very plumose.

Distribution. From Minnesota, Western Iowa, to Montana, Texas, New Mexico and Sonora.

Poisonous properties. Several species of *Liatris* are powerful diaphoretics, formerly these species were used as antidotes against snake bite. Dr. Johnson says that they probably possess no antidotal properties whatever and beneficial effects attributed to them are doubtless due to the diaphoresis induced by the administration of large quantities of the hot decoction.

11. *Grindelia* Willd. Gum Weed

Coarse perennial or biennial herbs, occasionally woody at the base, leaves alternate, sessile or clasping, spinulose serrate, involucre hemispherical; scales imbricated in several series, heads large, terminating leafy branches, radiate or rayless ray flowers yellow, pistillate, disk flowers perfect or staminate, achenes short, thick, compressed or turgid pappus of 8 awns, soon falling.

About 25 species, from western Minnesota and Iowa westward and southward to Peru and Chile.

Grindelia squarrosa (Pursh.) Gum Weed

A resinous, viscid, glabrous perennial from 1-3 feet high, leaves alternate, spatulate to linear oblong, sessile or clasping, spinulose serrate, heads many flowered, ray flowers yellow, pistillate, scales of the involucre hemispherical, imbricated in several rows with green tips, achenes short and thick, pappus consisting of 2 or 3 awns.

Distribution. Common west of the Missouri river from Mexico, Nevada and Texas to British America and east to Minnesota, Illinois and Missouri, occasionally naturalized eastward.

Poisonous properties. Very abundant in the west, is not liked by stock. The *G. robusta* and *G. squarrosa* are used in medicine in moderate doses to stimulate the mucous membrane and are beneficial in catarrhal affections. They are also antispasmodics. An alkaloid has been isolated from *G. robusta*. It contains *grindelin*, a bitter alkaloid, and two glucosides which resemble the saponins of *Polygala*.

12. *Solidago* L. Golden-rod

Perennial erect herbs, simple or branched, leaves alternate, simple toothed or entire, heads small, in terminal or axillary panicles. cymose corymbose ray flowers yellow, rays few or many pistillate, disk flowers yellow perfect involucre hemispherical or bell shaped, bracts appressed, destitute of green tips, achenes many-ribbed, terete or nearly so, pappus of simple capillary bristles.

A genus of about 100 species mostly in North America. About one-half of the number found in northern states east of the Missouri river. A few of the species are handsome and ornamental, like *Solidago speciosa*, *S. Drummondii*, *S. Missouriensis* and *S. odora*. The latter species is used as a stimulant and carminative. From this species there is derived an aromatic oil. An oil is also derived from *Solidago canadensis*, which strongly resembles the oil obtained from pine needles. The leaves of the fragrant golden-rod (*S. odora*) are often used as a substitute for tea. It contains an aromatic volatile oil.

Solidago canadensis L. Golden-rod

A perennial, with rough stem, from 3-6 feet high, leaves hairy beneath rough above, lanceolate and pointed, sharply serrate heads small, few flowered,

rays yellow, short pistillate, scales of the involucre appressed, not herbaceous, receptacle small, not chaffy, achenes ribbed; pappus simple, of capillary bristles.

Distribution. Widely distributed from New Brunswick to Florida. Common in the Mississippi Valley, especially along fences and in pastures and borders of fields in the Rocky Mountains to the Northwest Territory, British Columbia to Arizona.

Solidago rigida L. Rigid Golden-rod

An erect perennial from 2.5 feet high, rough and somewhat hairy, with minute pubescence, leaves oval or oblong, thick and rigid, not 3-nerved, the upper sessile, slightly serrate, heads in a compound corymb, large, 3 or more flowered, rays large, 7 to 10, yellow.

Distribution. Common in the upper Mississippi Valley, especially on the prairies, occurs east to New England.

Poisonous properties. It is thought by some that hay fever is caused by this and other species. A disease of horses in Wisconsin a few years ago was attributed to the eating of golden-rod, but Chesnut thought this might be caused by a rust fungus, *Uromyces solidaginis*.

Mr. J. L. Scott, who made an investigation of this disease, reported in Garden and Forest as follows:

During the past four years a large number of horses have died in the northern part of this state from the ravages of a disease which has baffled the skill of veterinarians, and I have been called upon to make investigations as to the cause and nature of the malady. At first it was thought to be anthrax and samples of the blood and sections from the spleen and other internal organs were sent to the Bureau of Animal Industry and to the Faculty of the State University, for bacteriological examination. Numerous bacteria were found, but the bacillus anthracis was not present.

The horses affected were in the majority of cases heavy draught horses from the lumber camps. These animals were brought from the woods in the winter, usually in good condition, and turned out in pasture. Most of them were fed grain while in pasture. On the farm of Mr. C. P. Reynolds, Hayward, Wisconsin, over seventy horses died during the past four years from this peculiar malady. The pasture contained about four hundred acres, three hundred acres of which had been broken and seeded to timothy. Adjoining this was one hundred acres of "slash oak" or land from which the timber had been cut, but which had never been broken. This was thickly covered with golden-rod. On one side of the farm is a lake with a clean gravel bottom and shore. The lake is fed by springs. There is no marsh or low land on the farm. Upon investigation I became convinced that the cause of the trouble was to be found either in the food or water and watched the horses closely for several days, and saw them eating the golden-rod greedily. Some of them, especially those affected, seemed to prefer the plant to anything else.

I also visited the farm of Peter Traux, near Eau Claire. There is no golden-rod to be found on this farm and the disease has not made its appearance. During the past summer, Mr. Traux placed ten horses in pasture near by, where the plant was plentiful and eight of them died during the summer and the remaining two were affected. When the healthy horses are taken from the pasture in the fall the disease disappears. None of the animals attacked by the malady have recovered, and medicinal treatment does not seem to produce any beneficial effect.

Symptoms. The animal appears dull, ears drooped, temperature elevated, ranging from 103° to 107° Fahrenheit, during the entire course of the disease. The visible mucous membranes are pallid. On the mucous membranes of the vulva small petechial spots are seen. Occasionally the legs swell and oedematous enlargements appear under the abdomen. The appetite remains fairly good during the entire course of the disease. Emaciation takes place rapidly as the disease advances. Loss of coordination with staggering gait. Death takes place in from two weeks to two months from the onset.

Post mortem. On cutting open the body the blood appears to be completely disintegrated, resembling ordinary blood serum. Intestines bloodless, with numerous petechial spots on the mucous membrane. Spleen enlarged, weighing from six to ten pounds. No structural changes

apparent to the naked eye. The lungs and kidneys apparently normal. The brain and spinal cord were not examined.

I am fully convinced that this disease is due either to some poisonous principle in the plant or some parasitic fungus upon the surface of the same. It is now too late in the season for any investigation to be carried on in this direction this year, but I intend to have the matter thoroughly investigated next summer.

13. *Aster* (Tourn.) L. Aster

Herbs generally perennial with corymbose panicled or racemose heads, heads many flowered, radiate, the ray flowers in a single series, fertile. Bracts of the involucre more or less imbricated, generally with herbaceous tips, receptacles flat, achenes somewhat flattened, pappus simple, consisting of capillary bristles. A large genus of 275 species, chiefly in eastern North America. Flowering in the autumn. A few of the species are cultivated for ornamental purposes.

The New England aster (*A. novae-angliae*) occurs in moist ground. The *A. laevis* with somewhat clasping leaves is common in dry soil or prairie regions of the West. The small white aster (*A. multiflorus*) with small leaves is common along roadsides in dry soil. Very few of the species have deleterious properties.

Aster Parryi Gray. Parry's Aster or Woody Aster

A somewhat hoary perennial with deep woody roots and a short more or less branched stem, hoary leaves, spatulate linear, cuspidate, heads solitary, bracts of the involucre oblong lanceolate long acuminate pubescent, rays white over half inch long, achenes white villous. This is the *Xylorrhiza Parryi* Gray.

Distribution. Common in saline soils. In Colorado, West Wyoming and Utah.

Poisonous properties. This plant has recently come into prominence in western Wyoming where the disease "grub in the head" has been attributed to the plant but Dr. Aven Nelson¹ has attributed this disease to a fungus *Puccinia xylorrhizae* which according to Dr. Nelson abundantly occurs on the plant. He says

If it should prove to be true that the malady is due to the eating of the aster, then it may be the aster itself that is the source of the trouble, but the chances are rather better that the specific poisonous qualities are due to the fungus. Some other parasitic fungi have been proved poisonous and we may well, in the absence of evidence to the contrary, also suspect this one.

Dr. Nelson states that the trouble is not a new one and that similar outbreaks have occurred in previous years, the disease recurring in the same localities. The disease was prevalent near Medicine Bow and northward to the Shirley Basin as well as Natrona county. The farmers in the vicinity where this disease occurs speak of the area in which the plant occurs as the "poison patch." Dr. O. L. Prien, Wyoming Agricultural Station, and Dr. Frederick of the Utah Agricultural Station, are making a careful study of this disease. To prevent the trouble, sheep should be kept, so far as possible, away from areas in which this plant occurs.

14. *Erigeron* L. Fleabane, Daisy

Branching or scapose herbs with entire or toothed leaves, heads in corymbose, paniculate or solitary peduncled heads, scales of the involucre narrow nearly equal, not foliaceous or green-tipped, flowers radiate, white violet or purple, numerous, fertile, disk flowers yellow tubular and perfect, branches of

¹ Press Bull. Wyoming Agr. Exp. Sta. No. 10.

the style more or less flattened with short appendages; achenes pubescent and 2-nerved, pappus a single row of fine capillary bristles.

About 120 species of wide distribution, most numerous in North America. Several species of fleabane (*E. philadelphicus*, *E. annuus* and *E. canadensis*) are used in medicine. They are reputed diuretics, tonics and astringents.



Fig. A. Small Horseweed (*Erigeron discoloratus*). From Indiana to Minnesota to Nebraska and southward. Common in sterile grounds. (Charlotte M. King).



Fig. B. Horseweed, Mares' Tail (*Erigeron canadensis*). A common weed throughout the United States, especially northward, also in Europe. Acid and to be irritating. (Charlotte M. King).

Erigeron canadensis L. Horseweed

Bristly herb, stem hairy or somewhat glabrate, 1-6 feet high, simple or panicately branched, leaves usually pubescent or ciliate, the lower spatulate, incised or entire obtuse or acutish, the upper generally linear and entire. Heads numerous, with inconspicuous white ray flowers shorter than the pappus, achenes small, flattened, pappus of numerous small fragile bristles.

Distribution. Common throughout the eastern part of North America except far northward. Common also in the Rocky Mountain region, and in

waste and cultivated grounds along the Pacific coast appearing as an introduced weed. Also an introduced weed in Europe and South America.

Erigeron romosus (Walt.) BSP. White Weed. Fleabane

Stem and leaves somewhat hirsute and hairy, roughish, leaves entire or nearly so the upper lanceolate, the lower oblong or spatulate, heads borne in corymbose panicles, ray flowers white and twice as long as the scales of the involucre, achenes small, pappus double, the inner of fragile bristles.

Distribution. From Nova Scotia to Florida, west to Louisiana and Texas to Northwest Territory.

Erigeron annuus (L.) Pers. Fleabane. White Weed.

A sparingly pubescent annual from 3-5 feet high, leaves thin, coarsely and sharply toothed, the lower one ovate, or ovate-lanceolate, acute and entire on both ends; heads corymbous and rays white, tinged with purple.

Distribution. A common weed in meadows, pastures and woods in northern United States.



Fig. C. White weed (*Erigeron annuus*). Common in clover and timothy meadows. A troublesome weed. Leaves and stem with a somewhat bitter flavor (Charlotte M. King).

Poisonous properties. Common horseweed (*E. canadensis*) abundant in many places has been looked upon as being suspicious. It contains an oil, mainly a terpene $C_{10}H_{18}$. It has an acrid taste and is neutral in its reaction. It causes smarting of the eyes, soreness of the throat, aching of extremities, and colic. It causes irritation to people handling the plant. When in hay it

may be injurious because the leaves are very bitter and cause much dust. The other species are very abundant in meadows in central Mississippi Valley states, and are regarded with much disfavor.

15. *Rudbeckia* L. Cone-flowers

Annual, biennial or perennial herbs, usually rough or rarely smooth, with alternate leaves, flowers in terminal clusters, involucre hemispherical, bracts imbricated in 2-4 series, scales leaf-like, in 2 rows, spreading, receptacle conical or columnar with chaffy scales, ray flowers neutral, disk flowers perfect, fertile, achenes 4-angled, obtuse or truncate at the apex, pappus wanting or a minute crown.

A small genus of some 30 species in North America and Mexico. Several species are common in cultivation.

Rudbeckia hirta L. Cone-flower Nigger-head

A rough, bristly hairy biennial from 1-2½ feet high with stems simple or branched, near the base bearing a long pedunculate head, leaves nearly entire, the upper sessile, oblong or lanceolate, the lower petioled and spatulate heads many-flowered, radiate, the rays about 14, neutral, receptacle columnar or conical chaff hairy at tip, acutish ray flowers yellow, disk dull brown achenes four-angled; pappus none.

Distribution. Common in dry soils and on the prairies and meadows in the northern Mississippi Valley, also a common weed in meadows in the eastern states. Introduced largely with clover seed.

Rudbeckia laciniata L. Golden Glow or Tall Cone-flower

A tall branching smooth perennial leaves thin, minutely pubescent, the upper leaves irregular ray flowers yellow disk flowers dull greenish, bracts of the involucre unequal pappus a short crown. Improved forms of this species are cultivated under the name of golden glow.

Distribution. From Canada to Florida, Kentucky, Kansas and New Mexico to Manitoba and Montana.

Rudbeckia occidentalis Nutt. Western Cone-flower

Nearly glabrous and smooth, or somewhat scabrous-puberulent, leaves undivided, ovate or ovate-lanceolate, acuminate, entire or irregularly and sparingly dentate, 4-8 inches long upper sessile by a rounded or subcordate base; lower abruptly contracted into a short winged petiole, rarely a pair of obscure lateral lobes, disk in age becoming 1½ inches long, and achenes 2 lines long.

Distribution. Common in the Rocky Mountains from Utah, Wyoming and Idaho.

Poisonous properties. Dr. Schaffner states that the *R. laciniata* is poisonous to sheep. It is eaten by cattle. The *R. occidentalis* is eaten by sheep but not relished. May be suspicious. A correspondent in Kansas and one in Iowa state that the plant is poisonous to cattle.

16. *Bidens* L. Bur Marigold or Beggar-ticks

Annual or perennial herbs with opposite leaves or the upper alternate, involucre double, campanulate, the outer scales commonly foliaceous, heads many-flowered, receptacle chaffy, rays present or absent when present neutral, in our species yellow disk flowers perfect, fertile, achenes flattened or 4-sided; pappus of 2-6 teeth, upwardly or downwardly barbed.

About 70 species of wide distribution.

Bidens discolor (T. & G.) Britton Small Tickseed

A diffusely branched annual from 1-2 feet high; leaves petioled, ternately divided, leaflets ovate-lanceolate, pointed and serrate, rays wanting, disk flowers yellow, achenes narrow wedge-shaped, bearing a pair of short and stout upwardly or downwardly barbed awns.

Distribution. Common on wet banks throughout the northern Mississippi Valley

Bidens frondosa L. Beggar-ticks Stick-tights

A branching hairy or smooth annual from 2-6 feet high, leaves petioled, 3-5 divided, the stalked leaflets lanceolate, pointed, coarsely toothed rays small and yellow, involucre double, the outer foliaceous, longer than the head, receptacle flattish with deciduous chaff, achenes wedge-obovate, 2-awned, the awns downwardly barbed.



Fig. 447a. Spanish Needle. Foot Jacks (*Bidens frondosa*). The flattened achenes with downwardly barbed awns more or less irritating. (Charlotte M. King)

Distribution. Widely distributed in moist places throughout the northern states, often a very troublesome weed in gardens. Widely scattered because of the "seeds" adhering to clothing, pelts of animals, etc.

Bidens bipinnata L. Spanish Needle

A smooth branched annual from 1-5 feet high leaves petioled and 1-3

pinnately dissected, heads many flowered, small, long peduncled involucre double, outer of linear scales, ray flowers yellow or none, achenes linear 4-angled, narrowed upward into a beak, nearly smooth, the four downwardly barbed awns usually spreading, shorter than the achenes.

Distribution. Native to Mexico and tropical America and the southern states now occurring as far north as Rhode Island to Ohio, southern Iowa and Nebraska. A weed also in southern Europe and Asia.

Poisonous properties. All of the species are local irritants but especially *B. frondosa*.

17. *Helianthus* L. Sneezeweed.

Erect branching annual or perennial herbs, plants often sprinkled with resinous matter, leaves alternate decurrent on the stem branches terminated by a single or corymbose heads, yellow or purple in color, involucre short and broad but small, bracts in 1 or 2 series, linear or subulate reflexed or spreading, ray flowers pistillate and fertile or neutral, rays 3-5 cleft, disk flowers perfect, fertile, corolla 4-5 toothed, style branches dilated and truncate at the apex, achenes top-shaped and ribbed pappus of 5-8 thin curved chaffy scales extended into a point.

About 25 species in North America. All of the species of the genus more or less poisonous. Among these are the *H. Bigelovii* of Oregon, California and *H. nudiflorus* from Texas to Kansas, Illinois and Florida.

Helianthus Hoopesii Gray

A stout perennial from 1-2 feet high, tomentose or pubescent when young becoming smooth with age leaves rather thick, entire oblong lanceolate the lower spatulate and tapering base, somewhat nerved heads large single or several, ray flowers fertile yellow, an inch long, soon reflexed bracts of the involucre in 2 series.

Distribution. Common in the Rocky Mountains to eastern Oregon to California. It covers large stretches between 6,000 and 8,000 feet in the Uintah mountains.

Poisonous nature. It is said to be poisonous like other species of the genus. Sheep carefully avoid it, feeding on the grass and other herbaceous plants, leaving the plant standing.

Helianthus autumnale L. Sneezeweed

A smooth or pubescent perennial from 1-6 feet high, leaves toothed, lanceolate to obovate, oblong decurrent on the stem, acuminate at the apex, heads numerous, many flowered, radiate rays yellow, wedge shaped, 3-5 cleft, fertile disk flowers yellow, involucre small, reflexed scales linear or awl-shaped receptacle glabrous or oblong achenes pubescent on the angles scales of the pappus ovate or acuminate, the variety *grandiflorus* with larger heads scales of the pappus narrowed and longer awned.

Distribution. Common in low meadows and alluvial banks from eastern Canada to South Dakota, Florida to Alabama, and especially common in the northern Mississippi Valley to Kansas. Occurs also in the Rocky Mountains, Wyoming to Montana. The variety in Washington to California.

Poisonous properties. This plant is considered poisonous to cattle and sheep. Numerous complaints have been made in various parts of the state of Iowa in regard to the poisonous nature of this weed. This plant is more or less bitter acrid and pungent, especially the flowers. Parts of the plant

rubbed were used by the Winnebago Indians for colds and to produce sneezing. Dr. Chestnut says:



but at 3 p. m. the calf was resting easy. Next day he seemed to be nearly as well as usual, and took his feed.

Experiment 2.—October 13th, same calf. A hot infusion of six ounces of the drug was prepared in one quart of water and given between 8 a. m. and noon. Soon after the first dose some weeping began but no other marked effect was observed. After second dose the calf laid down. Third dose, he continues down, but somewhat restless and sick. Fourth dose, becomes very restless, bowels loose, staggering, and very sick, holding head down, spitting wind, left flank tympanic but not very tense, right flank contracted, trying to put head between forelegs, and very sick greater part of the day. movements weak and uncertain, purposeless, hyperæthemia very marked, especially along the back, shivering from least touch but by persistent handling submits to touch, respiration and pulse quickened, appearance of slight spasms along the back neither eats nor drinks of food and water and milk in his stall, a little straw from floor in his mouth, and although the jaw is in constant motion the straw is undisturbed and he seems unconscious of its presence. Next day apparently well.

Experiment 3. October 30th another calf about four months old. One pound of the dry plant was taken up in hot water poured on it and boiled down to a quart & given by 10 35 a. m. After third dose the left flank tympanic, feces and urine discharged, slight jerking as though choked, tremor of muscles of hindquarters, lying on right side over an hour breathes with some noise and distress, very sensitive from n. h. to sternum, at 11 o'clock fullness of left flank subsided, limbs showed weakness and later stiffness, very sick foaming at mouth and flow from nose. Next day the calf was well.

The large quantity of the drug given this calf producing little more effect than the smaller dose on the former one leads me to suppose that the prolonged boiling may have dissipated a portion of the poisonous principle. Knowing that none of the symptoms observed would lead on to death no antidote was administered in any of the experiments. The weeping and movement of the jaw were probably due to the pungent effect of the drug on the throat and stomach, and I incline to the opinion that the nervous symptoms were not caused by direct action of the drug on nerve centers, but were reflex from the irritated throat and stomach.

In the sheep the spasms in severe cases are violent form and a sheep may have many such convulsions and yet recover without treatment and after many hours rise up and walk.

The horse and mare succumbed to the injurious effects of the poison quicker and more completely than other animals. The effect is manifested very soon after ingestion and with great violence. The animal cannot control his motions, plunges about blindly, falls dead or perhaps breaks his neck in falling forward with the head under the body.

An antidote which has proved effective is a pint or two of melted lard poured down the animal's throat. The relief is very prompt. But it must be given before the horse loses control of his limbs, or I cannot be assured of a cure. The relief is so prompt and complete that it is difficult to be sure that it is the direct effect of the lard on the nerve centers, but as a local emollient applied directly to the burning throat and stomach. The local trouble being quickly relieved, the violent reflex effect on the nervous system suddenly ceases.

When we consider the wide distribution of this plant and that in some places it is so abundant as to hide the earth from view over whole acres it is remarkable that so few poisonings occur. But animals have no special fondness for it. When they have been long on the road and deprived of green food, stopping where this weed is found and other green plants scarce they are supposed to eat a little of it. It is under such conditions that the plant is most frequently eaten and the poisoning occurs.

Animals, notably sheep once intoxicated by the plant and recovering, seem to acquire a mania for it and when finding themselves in localities where it grows hurry to and devour it.

Within 200 miles of the Gulf of Mexico the *H. quadridentatum* takes the place of the preceding, is very abundant, harmless to animals early and dies by August.

Helenium tenuifolium Nutt. Narrow leafed Sneezeweed.

A perennial, erect branching herb from 1-2 feet high, leaves alternate decurrent on the stem, linear oblanceolate. flowers borne in corymbed clusters, heads many flowered, ray flowers yellow, fertile disk-shaped, ribbed, pappus of 1-nerved chaffy scales, the nerve extended into a point.

Distribution. Common from Texas to Missouri, Florida and Virginia, especially in pastures and meadows and along roadsides. It is said to have been widely scattered in the south after the war of the rebellion.

Poisonous properties. It is often the cause of bitter milk in the south. Dr. Chesnut says

The fine-leaved Sneesweed has been reported from several of the Gulf States, where it is a troublesome weed, fatal to horses and mules. It is not known to what extent cattle may feed on the plant with impunity, but the bitter principle in milk and meat sometimes met with in the Southern States is quite generally supposed to be due to these plants.

It contains a narcotic poison.



Fig. 449. Fine-leaved sneesweed (*Helenium tenuifolium*). Often the cause of "bitter-milk" (Dewey, U. S. Dept. Agr.)

18. *Dioscorea* Lag. Fetid Mangold

Mostly annual or biennial herbs with strong scent, dotted with large pellucid glands, leaves mostly finely dissected, heads many-flowered, small, of both tubular and ray flowers, involucre cylindrical or hemispherical, bracts in one series united into a cup, receptacle flat, not chaffy but with short bristles, ray flowers pistillate, short, disk flowers perfect, achenes slender, 4 angled, pappus a row of chaffy scales dissected into rough bristles. A single species

Dyssodia papposa (Vent.) Hitchc. Fetid Marigold.

A nearly smooth branched annual from 6 inches-2 feet high, leaves opposite, sessile, pinnately parted, bristly-toothed with large pellucid glands, which give to the plant a strong odor. Heads many-flowered, rays small, yellow pistillate, disk yellow, involucre with a few scales at the base, one row of scales united to form a cup, achenes slender, 4-angled, pappus a row of chaffy scales finely divided into numerous rough bristles.

Distribution. Common from western Iowa and Minnesota to Illinois and the southwest. In this section of the country it occupies the same ground occupied by the Mayweed.

Poisonous properties. It is a very disagreeable smelling herb. Not relished by stock. It is probably injurious.



Fig. 450. Fetid Marigold (*Dyssodia papposa*). Pungent odor said to be injurious. (Charlotte M. King.)



Fig. 451. Dog Fennel (*Anthemis cotula*). Contains a pungent principle.

19. *Achillea* (Vaill.) L. Yarrow

Perennial herbs with erect leafy stems and finely dissected alternate leaves; heads many flowered, radiate, in corymbose clusters, ray flowers few white

or pink, disk flowers perfect, fertile; corollas yellow and 5-lobed, receptacle nearly flat; achenes oblong or obovate, slightly compressed, pappus wanting.

About 75 species mostly of the Old World. The yarrow (*Achillea Ptarmica*) occasionally cultivated.




Fig. 451 Yarrow (*Achillea Millefolium*). Structure of flower: a, stigma; b, papilion; c, style pollen; d, single flower; e, f, g and h, flowers in different stages. (Müller).

Achillea Millefolium L. Yarrow

Perennial herb from 1-2 feet high, simple stems, leaves twice pinnately parted, consisting of crowded linear divisions. flowers in compound flat-topped corymbs, heads many-flowered, radiate rays 4-5, white, scales of the involucre with scarious margins, receptacles chaffy, disk flowers small, tubular, achenes oblong, flattened, pappus none.

Distribution. A common weed from the Atlantic to the Pacific. In the East it is found in old fields and dry hills and is abundant in open parks and rocky dry soils in the Rocky Mountains up to the timber line. It is native also to Europe, where it is used as an ornamental plant. Sheep will eat the weed. The form with rose-colored ray flowers is frequently cultivated.

Poisonous properties. The alkaloid *achillein* $C_{20}H_{29}N_2O_5$ has been isolated from the plant. Millsbaugh describes the action of the drug as follows: "Yarrow seems to have a decided action upon the blood vessels, especially in the pelvis. It has been proven to be of great utility in controlling hemorrhages, especially of the pelvis, viscera, where hemorrhage is caused by it. Its common



Fig. 451a. Yarrow (*Achillea Millefolium*). 1, plant with leaf and flowers. 2, A single head. a, ray flowers. b, a single flower. (Charlotte M. King)

European name *Nosebleed*, was given from the fact that the early writers claimed hemorrhage of the nose followed placing its leaves in the nostrils; this may have been either due to its direct irritation, or the use of *Achillea Ptarmica*, its leaves being very sharply serrate and appressed-toothed. *Millefolium* causes burning and raw sensations of the membranes with which it comes in contact, considerable pain in the gastric and abdominal regions with diarrhoea and enuresis. An alkaloid having the same formula as *achucerin* has been isolated in *A. moschata*, a second alkaloid, *moschatin*, $C_{21}H_{27}NO$, is said to occur in the same plant. In Europe sometimes regarded as a forage plant.

20. *Anthemis* (Mich.) L. Mayweed

Annual or perennial herbs with finely dissected leaves and a strong scent; heads peduncled, involucre hemispherical, bracts imbricated in several series; ray flowers white or yellow, 2-3 toothed; pistillate and fertile, style branches of the disk flowers truncate, achenes oblong angled, ribbed; pappus none or short crown. There are about 60 species in the Old World. They are strong scented or aromatic herbs.

Anthemis Cotula L. Mayweed Dog Fennel

An acriid branching scented annual from 1-2 feet high, leaves thrice pin-

nately dissected, heads solitary, many flowered, ray flowers white, pistillate, fertile or neutral, involucre of numerous small imbricated dry and scarious scales, disk flowers yellow, small, tubular, achenes terete or ribbed, smooth, truncate, pappus none or a minute crown.

Distribution. A common weed in Europe where it is a native. Widely scattered in the Northern States from the Atlantic to the Dakotas and Nebraska. Occurs also in Washington and Oregon. In the north Mayweed is common in yards and along roadsides.

Poisonous properties. The strong odor and acrid taste makes it extremely undesirable for stock. The Mayweed blisters the skin.

Antennaria arvensis L. Corn Chamomile

A much branched annual or biennial pubescent herb, not all ascending, leaves sessile, finely 1-2 pinnately parted, less divided than the preceding; heads numerous, bracts of the involucre oblong, obtuse, usually somewhat pubescent with



Fig. 452. *Antennaria nobilis*. Cultivated as an ornamental plant. Faguet.

broad scarious margins ray flowers white, pistillate, mostly 2-toothed, chaff of the receptacle obtuse achenes oblong, obtusely 4-angled pappus a small border

Distribution. In fields and waste places in the Eastern States. Nova Scotia to Virginia, Missouri, and Wisconsin to the Pacific coast.

Poisonous properties. The oil of *Anthemis nobilis* contains the terpene anthemene $C_{11}H_{16}$, angelic acid $C_5H_8O_2$, tiglic acid $C_5H_8O_2$. The oil produces abdominal pain, increase of heart action and freeness of the bowels. It is not known that our species is poisonous but stock carefully avoid the plant.

21 *Chrysanthemum* (Tourn.) L. Ox-eye Daisy

Perennial or annual herbs, leaves alternate, dentate, pinnatifid or dissected, heads single or corymbose, or both ray and tubular flowers rarely wanting, involucre with broad imbricated scales, receptacle flat or convex, ray flowers pistillate and fertile, the rays white, style, branches of the pistillate flowers, truncate and pectinate achenes angled or terete, striate with pappus.

About 100 species of wide distribution. The common feverfew (*Chrysanthemum Parthenium*) is frequently cultivated in Germany and is rarely an escape in eastern North America. It contains the oil of feverfew.

Insect powders are derived from several species of *Chrysanthemum*, the Dalmatian from *C. cinerarifolium* cultivated in Europe, North Africa, New York and California, the *C. coccineum* Willd and *C. Marshalis*, found in the region of the Caucasus, furnish the Persian insect powders.

Chrysanthemum Leucanthemum L. Ox-eye Daisy

A perennial herb with erect stem and spatulate, petioled root leaves, those of the stem partly clasping all the leaves cut or pinnatifid toothed, the nearly simple stem bears a large many flowered head with numerous white rays, scales of the involucre with scarious margins disk and ray flowers pistillate and fertile, achenes marked with longitudinal lines, pappus alveol.

Distribution. Common in fields and meadows of the eastern states and eastern Canada, less common in the middle and western states.

Poisonous properties. This abundant weed probably contains an active principle. The *C. coccineum* contains the alkaloid chrysanthemin $C_{14}H_{21}N_3O_6$. The flowers are more or less irritating.

22 *Tanacetum* L. Tansy

Bitter acid strong scented aromatic annual or perennial herbs, leaves alternate, 1-3 pinnately dissected and corymbed heads involucre hemispherical, depressed, scales imbricated into several series, rays wanting or present, yellow marginal flowers pistillate and fertile corollas 5-toothed, style branches truncate and pectinate achenes angled or ribbed, pappus a short crown. A small genus of 35 species, natives of the northern hemisphere. The *Tanacetum balsamita* contains a balsamic oil.

Tanacetum vulgare L. Tansy

A bitter acid strong scented simple or branched perennial herb from 2-4 feet high leaves pinnately divided into linear pinnatifid divisions, lobes serrate, heads corymbose, many flowered ray flowers few disk flowers yellow marginal flowers fertile, scales of the involucre imbricated in several series, receptacle flat or convex naked, branches of the style brush like at the summit, achenes 5-angled or 5-ribbed, truncate or obtuse, pappus none or a short crown.

Distribution. Native to Europe, frequently escaped from gardens to roadsides along the Atlantic Coast as far west as Iowa and Kansas.

Poisonous properties. The oil of tansy is obtained from this plant. The

bulk of the oil consists of *anaceton* or *thujon* $C_{10}H_{16}(OH)$. This oil has been used as an anesthetic since the middle ages. Tansy is a very poisonous, producing a condition similar to rabies. From this plant many serious cases of poisoning in the human race occur, but poisoning of animals is very infrequent. It is an irritant narcotic poison. Dr. Millsbaugh cites a few cases as follows:

A young woman had been in the habit of using tansy tea, made from the herb, at nearly every menstrual period, for difficult menstruation. On this occasion about two and a half drachms of the oil was poured into half an ordinary tin cupful of water (this, with the exception of a small portion of the water containing about one-half drachm of the oil, was taken at one dose). Convulsions were almost at once produced, and when Dr. Bailey was sent for the patient was unconscious, foaming at the mouth and in violent tonic spasms with dilated pupils, frequent and lock-jaw pains. Constant knocking on the stomach had produced partial emesis, and then spiced mustard, and a large draught of hot water emptied the stomach. Two drachms of magnesium were then given, and a full dose of acetate of morphine consciousness returned, no untoward symptoms followed, and, after thirty-six hours, without additional medication she was entirely restored.

A woman took half an ounce of the oil: the most violent rigid kind of tonic spasms occurred once in about twelve minutes, coming on generally and instantly and continuing about one minute. They were attended with sight, if any means of the arms it might be called a trembling. The arms were peculiarly affected, and even when in the same way they were thrown out forward of and at right angles with the body. The hands at the wrists bent at right angles with the forearm supinated, the points of the fingers nearly in contact, the fingers straight and slightly bent at the metacarpophalangeal joints. The movements of respiration were strikingly affected during each paroxysm: the air was forced from the chest slowly but steadily and made a slight hissing noise as it escaped from between the patient's lips. During the intermission of spasms, the muscles were perfectly flexible, and the transition seemed very sudden. The jaws were the only exception to this rule: they were for the first hour and a quarter rigidly closed and were with difficulty opened, but after that they were subjected to the same action as the rest of the body—when the spasms were on they were rigid, when off they were relaxed. After the patient grew weaker the spasms were more frequent but had about the same severity and length. Death ensued in two hours.

On animals the symptoms are as follows according to a statement condensed by Dr. Millsbaugh:

Dr. Hy Van DeWarker records cases of the action of the oil upon dogs. In one case two drachms were given, causing salivation, vomiting, dilation of the pupils, muscular twitchings, followed by clonic spasms, and a cataplectic condition from which the animal recovered. Recovery was inaugurated a half ounce after the same cause of symptoms, but however on repeating the dose the already poisoned animal was plunged into a long and fatal convulsion. Postmortem examination disclosed the cerebral veins and spinal cord itself highly congested, a serious effusion had taken place in the meninges. The lungs were found to be engorged, the left heart empty and the right distended with dark liquid blood. Congestion of the kidneys had also taken place, and the bladder was found contracted.

The safe maximum dose of the oil is indeterminable: a few drops only sometimes proving fatal.

The symptoms occurring in a number of cases of poisoning and experiments were substantially as follows: Mental confusion, loss of consciousness, vertigo, with cerebralgia at first contraction, then wide dilation, of the pupils, staring, prominent eye balls, ringing in the ears, face congested, roughness of the mouth and throat, difficult deglutition, eructations, nausea, fre vomiting and burning of the stomach, sharp cutic pains in the abdomen, diarrhoea, constant desire to urinate, urine at first suppressed, then profuse, respiration hurried and laboured, pulse at first high, then very low and irregular, numbness of the extremities, tonic and clonic spasms and nervous tremblings, drunkenness and cold sweat. Death appears to ensue from a paralysis of the heart and lungs.

23. *Artemisia* L. Wormwood, Sage Brush

Bitter and aromatic herbs or shrubs with alternate leaves, heads discoid, few or many flowered, collected in panicles, racemes or spikes, with greenish or yellow flowers, involucre ovate or oblong, scales imbricated, the outer shorter, receptacle small, naked, marginal flowers patulate, disk perfect, or



Fig. 453. *Artemisia*. Sage Brush. Absinthe and Wormwood belong to this genus. (After Engelm.)

perfect and fertile with branches of the style truncate or sometimes sterile with ovaries abortive, styles undivided, marginal flowers usually pistillate and fertile, or flowers all perfect and fertile in some species, anthers often tipped with subulate appendages, achenes obovoid and no pappus.

About 200 species, mostly native of the northern hemisphere. A few in southern South America. The oil of Levant wormseed (*A. maritima* var. *Stechmanniana*) contains the substance *cineol*. Santonin $C_{15}H_{16}O_4$ is the active principle found in the flowers of *A. cuneata*. The *A. maritima* contains *artemisin* $C_{15}H_{16}O_4$. Absinthe is furnished by *Artemisia absinthium* of Europe. The *A. barbatula* furnishes the Algerian absinthe. Several species of sagebrush (*Artemisia tridentata* and *A. cana*) are common in the west and much used as forage for sheep. The alkaloid *abrotanin* $C_{21}H_{23}N_2O$ is obtained from *A. Abrotanum*.

Artemisia biennis Willd. Wormwood

An aromatic, somewhat bitter smooth annual or biennial, from 1-3 feet high, with leafy stems and erect branches, lower leaves twice pinnately parted, the upper pinnatifid, the lobes linear or linear-oblong, serrate or cut-toothed; ray flowers absent, heads numerous in short axillary spikes, the bracts of the involucre green, scarious, margined.

Distribution. Common in the northern Mississippi Valley, now widely scattered east to Nova Scotia and south to Kentucky.

Artemisia tridentata Nutt.

From 2-12 feet high, much branched, silvery canescent, leaves cuneate, obtusely 3-toothed or 3-lobed, or 4-7 toothed at the summit, upper leaves cuneate-linear, heads homogamous, flowers all perfect and fertile, heads densely paniculate, 5-8 flowers.

Distribution. From Montana and Colorado westward covering immense areas. The characteristic sage brush, similar species are the *A. trífida* Nutt. a smaller plant common westward, and the *A. arbuscula*, Nutt., also a dwarf plant found in the high mountains and elevated plains, the *A. cana*, Pursh from 1-2 feet high silvery canescent leaves, found from Saskatchewan to Dakota, Montana and Colorado.

Poisonous properties. The sage brush is consumed by sheep and cattle,



Fig. 454. Wormwood.
(*Artemisia biennis*). A com-
mon weed in the Mississippi
Valley. Ada Hayden.

although the latter do not relish the plant. The inhalation of the dried powder of the plant causes violent sneezing. Chesnut and Wilcox, in speaking of the sage brush of Montana, say that some species are recognized by stockmen as valuable forage plants for the late fall and winter grazing. Various species are suspected in Montana of being poisonous to stock, but no specific cases have been reported to the department. At Toston in May 1900, it was noticed that the terminal branchlets of *A. tridentata* had all been eaten off from a considerable number of plants.

A. Absinthium L. Common Wormwood Absinthe

A shrubby, silky, hairy plant from 2-3 feet high. Leaves 2-3 pinnately parted, the lobes lanceolate, obtuse, heads in racemose panicles, nodding yellow, marginal flowers pistillate, fertile or sterile, central flowers fertile, involucre hemispherical outer bracts linear.

Distribution. Native to the old world but escaped from gardens and found along roadsides. From Newfoundland to Massachusetts and North Carolina, occasionally westward to Wisconsin and Montana.

Poisonous properties. Wormwood is a stimulant and tonic and has been employed chiefly for digestive disorders, although seldom used medicinally. The volatile oil of wormwood in large doses produces cerebral disorders, convulsions and even death. The well-known absinthe is made from this plant and used with alcoholic drinks. For this purpose the plant is extensively cultivated in Europe and to some extent, in Nebraska, Michigan and Wisconsin. Absinthe contains the principle substance $C_{15}H_{24}O_4$. Dr. Rusby, in speaking of absinthe says that absinthium is in a general way similar in composition and properties as a poison to *Tanacetum* though the symptoms are more largely cerebral. Neither is it liable to be taken except for medicinal purposes or as an addition to spirits. As a result of the latter, poisoning is usually chronic and extremely difficult to cure. Dr. Millspaugh reports the experience of a clerk who took about a half ounce of the oil. He was found on the floor perfectly insensible convulsed, and foaming at the mouth. Shortly afterward the convulsions ceased, the patient remained insensible with the jaws locked, pupils dilated, pulse weak, and stomach retching. After causing free emesis and applying stimulants, the man recovered, but could not remember how or when he had taken the drug. According to Dr. Legrand, the effects prominent in absinthe drinkers are: Derangement of the digestive organs, intense thirst, restlessness, vertigo, tingling in the ears, and illusions of sight and hearing. These are followed by tremblings in the arms, hands, and legs, numbness of the extremities, loss of muscular power, delirium, loss of intellect, general paralysis, and death. Dr. Magnan, who had a great number of absinthe drinkers under his care, and who performed many experiments with the liquor upon animals, states that peculiar epileptic attacks result, which he has called "absinthe epilepsy."

Post-Mortem. Great congestion of the cerebro spinal vessels, of the meninges of the brain, extreme hyperaemia of the medulla oblongata, injection of the vessels of the cord, with suffusion of the cord itself. The stomach, endocardium, and pericardium show small ecchymoses.

Absinthe is sometimes added to hops to make beer more exhilarating.

A. vulgaris L. Common Mugwort

Tall, branching perennial with fine and closely appressed hairs. Leaves mostly

smooth and green above, but quite woolly underneath; parastich, the divisions cut-toothed, linear-lanceolate; heads small in open leafy panicles.

Distribution. In waste places from Canada to North Carolina to Pennsylvania and occasionally westward.

Poisonous properties. The common mugwort was used medicinally by the physicians of the 16-18 century, but in modern times the plant is not used to any extent. Mugwort causes epileptic spasms, profuse sweat with the odor of garlic, violent contractions of the uterus, labor-like pains, prolapsus and rupture of the uterus, miscarriage, metrorrhagia and increase of lochial discharges.

24. *Arnica* L.

Perennial herbs, erect, simple or branching, leaves opposite, or rarely with the upper alternate, heads many-flowered, radiate on long peduncled heads. Scales of the involucre bell-shaped, bracts in 1-2 series, the ray flowers pistillate and fertile, disk flowers perfect and fertile achenes linear, 5-10 ribbed, pappus of roughened-denticulate bristles.

About 30 species native to the Northern Hemisphere. The tincture of *arnica* is a well-known home remedy used for bruises and chailbains. *A. montana* contains a bitter principle *arnica* $C_{21}H_{33}O_5$. The value of *arnica* as a germicide has been largely overestimated. The germicidal action is probably due to the presence of alcohol. Dr. White in his *Dermatitis Venenata*, records several instances of injurious effects from the use of *arnica*. In one case a gentleman descending the stairs to mount his horse for a ride, slipped and scraped the lower part of his back. A handkerchief dipped in the tincture was applied to the bruised skin and worn in contact with the part during the ride. During the ride he felt considerable itching and upon his return home found that the skin was greatly congested. On the next day I was called to see him. The skin of the back, nearly to the shoulders, was in a state of active hyperaemia, and already covered with innumerable papules. The inflammatory process extended rapidly downwards nearly to the knees, and forward upon the abdomen and genitals. In a few days these parts presented all the characteristic appearances of acute eczema in its various stages of progression—general hyperaemia, papules, vesicles, excoriated and exuding surfaces, and crusts. The subjective symptoms were intense itching, stinging and burning in the parts. Scarcely any clothing could be borne in contact with the skin by day, and sleep was for a few nights almost impossible, but the system generally was only slightly disturbed. Dr. White records two other cases. All three seem to have been old gentlemen and he states: "In all of them we have an acute inflammatory process, confined to the upper dermal layers, and manifesting itself, according to the stage reached, by the following appearances—hyperaemia, papules, vesicles, excoriations, crusts, and scales, in regular sequence. The local sensations were intense itching and some degree of burning in the parts affected."

Arnica cordifolia Hook. *Arnica*

A low perennial, 1-2 ft. high or less, pubescent, or the stem hirsute and peduncles villous, the root and leaves deeply cordate or ovate at the base, on slender petioles, stem leaves opposite, in 1-3 pairs, dentate or denticulate, upper small, sessile, heads few, solitary, involucre pubescent or villous, rays an inch long; achenes more or less hirsute.

Distribution. Higher altitudes of the Rocky Mountains, from Colorado to California and British Columbia.

Arnica mollis Less. Hook.

A leafy branching perennial, tomentose or villous-pubescent or nearly smooth, leaves denticulate or dentate, acute or obtuse, lower leaves tapering into a margined petiole, upper broader at the base and clasping, flowers much smaller than the preceding, ray flowers yellow, achenes hirsute-pubescent.

Arnica monocephala Olin.

An erect simple, branching perennial, from 1-2 feet high, leaves opposite, long peduncled heads, leaves narrow, oblong and lanceolate, the upper linear, ray flowers, yellow.

Distribution. Grows in woods and moist localities. Montana to New Mexico and westward.

Poisonous properties. This well known arnica has, for some time, been suspected of being poisonous. When bruised all the plants exhale the odor of arnica. Chesnut and Wilcox state

It has an odor and taste much like that of the official drug, and, in fact, an extract made from the heads is considerably used locally as a liniment. No specific cases of poisoning caused by the plant have been reported from Montana, and we are not aware that it is eaten but it has, nevertheless, been suspected of being poisonous to stock.

Dr. S. B. Nelson states that a sheep ate 2 pounds (of the leaves?) of *A. alpina* within a day without experiencing any bad effect. The plants had been gathered eighteen hours.

25. *Senecio* (Tourn.) L. Groundsel

Perennial or annual herbs, shrubs or trees. Leaves alternate, heads many flowered, solitary, corymbose or paniculate, involucre cylindrical or bell shaped, principal bracts in one series, distinct or united at the base, usually with some shorter rays, outer ones pinnate or none. Disk flowers perfect, corolla tubular; achenes terete or those of the marginal flowers compressed, 5-10 ribbed, pappus of numerous capillary bristles. A large genus of 1000 species of wide distribution. Some ornamental. Some species have been looked upon as poisonous.

Dr. Day in the Twenty-third Annual Report of the Bureau of Animal Industry, 1906, gives the following in regard to a disease supposed to be associated with a species of this genus.

At the present stage of the work there appears reason for associating this malady with an affection of cattle known in Nova Scotia as Pictou disease, with Winton disease of horses of New Zealand, and with Molteno cattle disease of South Africa. The lesions examined present practically the same lesions as those described for the above-mentioned affections—namely, uterubular connective tissue hyperplasia, fatty infiltration of the liver cells, and hemorrhages into the spleen.

Pictou disease, Winton disease, and Molteno cattle disease have been investigated time and again in an endeavor to find a causative agent. All inoculation experiments have failed, and it has only been found of late that feeding experiments with certain weeds have given seemingly reliable results.

Pictou disease, occurring in Pictou and Antigonish counties, Nova Scotia, is now ascribed to the ingestion of a weed called ragwort. Winton disease is thought to be caused by a member of the ragwort family, *Senecio Jacobaea*. The Molteno disease of South Africa is probably induced, according to recent investigations, by still another ragwort, *Senecio Burckellii*. At least these weeds when fed to animals in their respective localities have produced diseases followed by a train of symptoms corresponding to those seen in the naturally induced disease. The post mortem findings were also quite similar. It was also proved that these plants were most dangerous in the young state. Yet another variety known as *Senecio latifolius* was found to be deadly to stock.

In some parts of the states in the west *Senecios* are common but they are not

commonly found when this forage is cut for hay. It is extremely doubtful whether the *Senecio* occurring in the western part of Iowa is responsible for this disease. The groundsel or mug-wort (*Senecio Balsamifera*) blooms in May



Fig. 454a. Burdock (*Arctium Lappa*). 1, general view of plant, 2, single flower, 3, scales of involucre hooked. These heads annoying to stock. (L. S. Dent Agr.)

and after blooming disappears except a few of the root leaves. The stinking Willie (*S. Jacobaea*) is naturalized in the east and of course may be responsible for the Picton disease.

26. *Arctium* L. Burdock

Large coarse biennial, rough or pubescent herbs, leaves broad, alternate and petioled, heads large, racemose, corymbose or paniculate, flowers purple or white, all tubular perfect and similar, involucre globose, the scales imbricated, lanceolate, tipped with hooked points, receptacle flat, bristly, achenes oblong, somewhat compressed and 3-angled and wrinkled, pappus short, of numerous short bristles.

A small genus of 5 or 6 species in the old world. The root of the common burdock is used in medicine, while in Japan it is also used for food.

Arctium Lappa L. Burdock

A coarsely branched biennial from 1-3 feet high, hairy, leaves large, roundish or heart-shaped, thin, obtuse, entire or dentate, floccose tomentose beneath, petioles deeply furrowed, heads of purplish or whitish flowers, clustered or somewhat corymbose, the involucre lengthened into hooked tips, glabrous or slightly cottony.

Distribution. Common from New Brunswick westward in the Canadian provinces in Canada, southward to Alabama, general in the Mississippi Valley, Rocky Mountain region, and Utah.

Poisonous and injurious properties. Where the burdocks are allowed to grow freely the burrs cling to the fleece of animals, often forming large balls which may be found on the tails of cattle and horses. When they are common where sheep pasture, the burdocks work into the wool which soon becomes of little value. Formerly burdock was used medicinally more than it is at the present time. The root has diuretic properties. Dr. Johnson gives the chief medical properties as follows:

Burdock, though chiefly used by the laity, is highly esteemed by some of the profession as a diuretic and alterative. It is, perhaps, more frequently used in rheumatism than in any other disease, but is also recommended in chronic cutaneous diseases, catarrhs, affections, syphilis, and scrofula. In the form of an ointment or liniment it is used as an application to burns, ulcers, etc.

Some forms of the burdock are used for food, the young stems when stripped of their rind may be eaten raw or boiled, or in a pottage with vinegar.

27. *Cirsium* (Tough.) Hill. Thistle

Perennial, annual or biennial, erect, branched or simple, herbs, leaves sessile, alternate, often pinnatifid, prickly, heads usually large, many flowered, solitary or clustered, heads discoid, flowers purple, yellow or white, tubular perfect and fertile or dioecious, involucre ovoid or globose, scales imbricated in many rows, often glutinous, tipped with a prickle or unarmed, receptacle flat or convex, bristly, corolla tube slender, the limb deeply cleft, anthers sagittate at the base, style elongated, branches short or obtuse, unappendaged, often with a pubescent ring below, achenes obovate or oblong compressed or obtusely four-angled, glabrous, often ribbed; pappus plumose or minutely serrulate.




Fig. 455. Woolly Thistle (*Cirsium canescens*). 1, Head, 2, Flower and pappus, 3, Achene, 4, Anthers and style cut open, 5, Pollen grain, 6, end of style. (Charlotte M. King)

About 200 species of the north temperate regions. Some of them are cultivated for ornamental purposes, many of the species are pleasantly scented; many of them are troublesome weeds. *Cnicus* $C_{20}H_{32}O_{16}$ has been found in a related genus (*Cnicus benedictus*). Horses are fond of the heads of many species like *C. eriocephalum*, *C. Drummondii*, *C. undulatum* and others that occur on the Rocky Mountains. None of the species are probably poisonous but many of them have irritating properties. The Canada thistle is used in domestic practice.

Cirsium arvense (L.) Hoffm. Scop. Canada Thistle

Smooth perennial herbs, spreading by creeping rootstocks, 1-3 feet high, corymbose-ly branched at the top, stem smooth, leaves lanceolate, sessile and deeply pinnatifid, lobes and margins of leaf with spiny teeth, heads small, $\frac{1}{4}$ -1 in. high; bracts appressed, the outer with a broad base inner narrow, all with an acute tip, never spiny. flowers purple, dioecious. in staminate plant flowers exerted with abortive pistil in pistillate less so, scarcely exceeding the bracts. long stamens with abortive anthers, tube of the corolla 6 lines long, anther tips acute, filaments minutely pubescent, all of the bristles of the pappus plumose.

Distribution. The Canada thistle is found in waste places from Newfoundland, Nova Scotia, various provinces of Ontario, to New York, Virginia, southwest to Missouri and Kansas, Colorado to Idaho, Montana and Oregon.

Cirsium lanceolatum (L.) Hill. Bull Thistle

Branching biennial, 3-4 feet high, tomentose, becoming dark green and villous or hirsute with age, branchlets bearing large heads, leaves lanceolate,



Fig. 456 Canada Thistle (*Cirsium arvense*). Common in the North, causes inflammation. (Charlotte M. King).

decurrent on the stem with prickly wings, deeply pinnatifid, the lobes with rigid prickly points, upper face roughened with short hairs, lower face with cottony tomentum, heads $1\frac{1}{4}$ to 2 inches high, bracts of the involucre lanceolate rigid when young, more flexible with age, long attenuated prickly pointed spreading tips, arachnoid woolly flowers perfect anther tips acute, filaments pubescent, achenes smooth, pappus of numerous plumose bristles.

Distribution. Distribution in North America in fields and waste places from Newfoundland to Georgia, Missouri, Kansas to Nebraska, the Dakotas, Montana, Washington, Oregon and Idaho. Naturalized from Europe, native also to Asia.



Fig. 457 Bull thistle (*Cirsium lanceolatum*). A common pasture weed. Causes mechanical injuries. (U. S. Dept. Agr.)

Cirsium canescens Nutt. Woolly Thistle

Branching perennial 2-4 feet high, woolly throughout bearing single medium sized heads, stem angled, white woolly, leaves radical 8 inches to 1 foot long, the divisions usually 2-lobed, prominently ribbed, ending in stout spines, stem leaves except the lower, 1-4 inches long, pinnatifid, the upper sessile, slightly roughened, with a slight cottony down, the lower white, woolly, heads $1\frac{1}{2}$ to 2 inches high, bracts of the involucre somewhat arachnoid, lower scales with a broad base, glutinous ridge, and ending in minutely serrated spine, inner scales long, attenuated, tips straw colored; flowers purple. This is *Carduus Flodmannii* Rydb.

Distribution. This species is distributed from Mason City, Iowa, to southwestern Minnesota, west to the Rocky Mountains. Collected by Charles A. Geyer in 1839, and described by Nuttall. The writer has seen it very abundant in Wyoming, Montana and Colorado.

Cirsium discolor (Muhl.) Spreng. Prairie Thistle

Tall, branching, leafy biennial, 5-7 feet high, with heads larger than the Canada thistle, stem striate, slightly hirsute, leaves radical 12-14 inches long, deeply pinnatifid, the divisions frequently divided, prickly toothed, the upper surface smoothish, and the lower white, woolly single heads terminating the branches, with purple flowers, heads $1\frac{1}{2}$ inches high, bracts of the globose involucre slightly arachnoid, lower bracts ovate, with a broad base and a weak prickly recurved bristle, slight dorsal gland, inner linear lanceolate with a nearly colorless entire appendage, flower purple, lobes of the corolla terminating in clavate tips, anther tips acute, filaments pubescent, bristle of pappus plumose; achenium smooth, upper part yellow.

Distribution. In fields and along roadsides from Quebec, Ontario, south through New England, New York and Georgia, west to Missouri, Nebraska and South Dakota.

Cirsium ochrocentrum Gray

Biennial, 2-8 feet high, white tomentum, leaves commonly deeply pinnatifid and armed with long yellowish prickles heads 1-2 inches high involucre scales with a viscid line on the back, with a prominent spreading yellowish prickle; corolla purple, rarely white.

Distribution. From Arizona to Colorado and Utah.

Cirsium undulatum (Nutt.) Spreng.

Biennial 1-2 feet high, white tomentose leaves pinnately parted, somewhat prickly, heads about 2 inches high, the outer bracts thickened by glandular viscid ridge, corolla rose-color, purple or pale purple, the variety *megacephaleum* with larger heads.

Distribution. West of the Missouri River to Oregon and New Mexico.

Injurious properties. None of the species is poisonous so far as the writer knows. The spiny involucre and spiny leaves inflict inflammation and cause the formation of pus.

28. *Centaurea* L.

Perennial or annual herbs, with leaves alternate, entire, dentate, incised or pinnatifid; involucre ovoid or globose, bracts in many series, marginal flowers usually neutral and larger than the central ones in some species, all the flowers perfect and fertile, heads middle-sized, tubular, purple-violet, or white or rarely

yellow; receptacle flat and bristly, corolla-tube slender and 5-cleft or 5-lobed; anthers sagittate at the base, achenes oblong or obovoid, pappus of several series of bristles and scales.

About 350 species mostly of the old world where they are troublesome weeds. The common bluebottle or corn flower is frequently cultivated in gardens and is a frequent escape from cultivation.




Fig. 458. Flowers of *centaurea*. 1. Flower 2. anther 3. style At the left. Achene and Pappus.

Centaurea solstitialis Lam. Knapweed

Erect, branched cottony, stems winged, lower leaves lyrate, upper linear entire, decurrent, spines of upper bracts long spreading with a few smaller ones at the base, pappus soft.

Distribution. Fields in California, also in Europe and England, rare, introduced with alfalfa. Introduced into Iowa with alfalfa seed.

Injurious properties. Because of the spiny character of the plant it is often troublesome to man and animals.

29 *Silybum* (Vahl.) Adans. Milk Thistle

Annual or biennial, much branched herbs leaves large, alternate, clasping, white-bleached heads discoid, solitary at the end of the branches, involucre large; bracts rigid, some armed with large spreading or recurved spines; flowers all tubular, corolla-tube slender top 5-cleft anthers sagittate achenes obovate-oblong, pappus-bristles in several rows.

A single species, native to the Mediterranean region.

Silybum Marianum (L.) Gaertn. Lady's Thistle

A tall branched glabrous annual with striate stem, leaves oblong lanceolate, prickly clasping.

Distribution. Occasionally found eastward especially in ballast and common on the Pacific coast, from British Columbia to Southern California.

Injurious properties. The spiny leaves and involucre have been troublesome, produces mechanical injuries.

A CATALOGUE OF THE POISONOUS PLANTS OF THE WORLD

In the following catalogue an attempt has been made to record all of the species that have been in some way or another regarded as poisonous or injurious to man. This catalogue, however, is based mainly on the plants listed by Greshoff, Cornevin, Chesnut, Ernst, Lyons, Maiden, Miquel, Radkofer, Rusby, Smith, White in "*Dermatitis venenata*," but includes also a few from other scattered sources. The poisonous fungi were listed from Atkinson, Hard, Clements, and some from Fries, Peck, Farlow, and Bulliard. It was not thought best to list the fungi from the works of Ratzburg and Phoebeus entirely because of the uncertainty as to the poisonous qualities. It should be stated that many of the plants in this list are not poisonous nor are they always given as poisonous by the authors credited to them after the species, in fact, many of them on the whole must be regarded as plants simply having medicinal virtues. It was thought best, however, to list them, to permit future investigators to take up the problem of the chemistry and poisonous qualities of these plants from an experimental standpoint. It is simply an indication of the lines along which work should be done. We have also indicated the action of the plant or in some few cases the important substance found in the plant. The alphabetical arrangement has been adopted except as to the lower forms. In the last column the distribution is indicated usually only where the plant is indigenous, but in some cases North America is added where the plant has been introduced.

I have been greatly aided in this work by the excellent treatise of A. B. Lyons, "Plant Names, Scientific and Popular," published by Nelson Baker & Co., and also by the excellent work of Sayre, "Organic Materia Medica and Pharmacognosy."

In the preparation of this catalogue I have received very substantial help from Miss Harnette S. Kellogg, who has looked after the details in cataloguing the species and the synonyms. In some cases the synonyms could not be found in the Kew Index, and these have been allowed to stand as they were given by the authors. Some duplication of names may occur and possibly through an oversight some have been placed in the wrong orders, but I ask the reader's indulgence in errors of this kind.

Finally, I am indebted to Miss Bertha D. Herr for the laborious task of getting the copy ready for the printer.

L. H. FAMEL

The following abbreviations refer to the properties mentioned in the third column of the catalogue:

Abort.—abortifacient

Acr.—acrid poison

Alk.—alkaloidal poison

Amvg.—contains amygdalin

Androm.—contains anarommedotoxin

Anes.—anesthetic

Ant.—anthelmintic

Ant-sc.—antiscorbutic

Antisp.—antispasmodic

Astr.—astrigent

Berb.—contains berberin

Card.—cardiac poison

Cath.—cathartic

Con.—convulsive

Con.—contains coumarin

Cur.—curare poison

Cyt.—contains cytsin

Del.—deleriant

A. P.—arrow poison	Diaph.—diaphoretic
Ast.—asthenic	Diur.—diuretic
Dr. Em.—drastic emetic	Naus.—nauseant
Em.—emetic	Ord.—ordeal poison
Ent.—causes enteritis	Par.—paralyzing
Eser.—contains eserin	Pur.—purgative
Exp.—expectorant	Sap.—contains saponin
F. P.—fish poison	Sed.—sedative
Gastr.—causes gastro-enteritis	Sial.—salagogue
Hyd.—contains HCN	Sol.—contains solanin
Hyp.—hypnotic	Sop.—soporific
Ineb.—inebriant	Stim.—stimulant
Ins.—insecticide	Taen.—taenifuge
Intox.—intoxicant	Toxal.—Contains a toxalbumin.
Ir.—irritant	Tymp.—causes tympanites
Lax.—laxative	Ur.—causes uræmic poisoning
Loco.—causes loco poisoning	Urt.—urticarial
Mech.—causes mechanical injuries	Ver.—vermifuge
Nar.—narcotic	Ves.—vesicant

The abbreviations for the literature referred to are as follows

C.—Cornevin's Les plantes vénéreuses et ses empoisonnements qui elles déterminent.

Ches.—Chesnut's Preliminary Catalogue of Plants Poisonous to Stock

E.—Ernst's Ueber nachvergiftenden Pflanzen.

G.—Greshoff's The Distribution of Prussic Acid in the Vegetable Kingdom.

Phytochemical Investigations at Kew

K.—Kirtikar's The Poisonous Plants of Bombay

L.—Lyon's Plant Names, Scientific and Popular

M.—Maiden's Plants reputed to be Poisonous to Stock in Australia

Mi.—Miquel's Poisonous Plants of North Netherlands

R.—Radtkofer's Plants said to be Poisonous to Fish.

Rusby.—Rusby's The Poisonous Plants of the Vicinity of New York City.

Sm.—Smith's Poisonous Plants of all Countries.

Wh.—White's Dermatitis venenata.

ACANTHACEÆ

NAME OF PLANT	Authority	Properties	Locality
<i>Hygrophila hispida</i> Nees	G.	Em.	Cuba
<i>Justicia Gendarussa</i> Burm.	G.		Tr. As.
<i>Lepidagathis Wrightiana</i> Benth.	G.	Abort.	
<i>Paulownia polisperma</i> Benth.	G.	F. P.	Tr. Afr.
<i>Paulownia speciosa</i> Horbat.	G.	Em.	Abyss.
<i>Rhinacanthus communis</i> Nees.	G.		Ind. Malay
			Burma. Abyss.
<i>Ruellia patula</i> Nees.	G.	Em.	Arab. Ind.
<i>Ruellia strepens</i> L.	G.	Em.	N. Am. W. Ind.
<i>Ruellia tuberosa</i> L.	G. L.	Em.	N. Am. Tr. Am.

ADOXACEÆ

ALISMACEÆ

<i>Alisma Plantago</i> L.	G. M. L.	Ir.	Tem. N. Am. Austr.
---------------------------	----------	-----	-----------------------

AMARANTHACEAE

NAME OF PLANT	Authority	Properties	Locality
<i>Achyranthes bidentata</i> Bl.	G.	Sap.	Tr As
<i>Amaranthus gangeticus</i> L.	G.	Sap.	Trop. Reg
<i>Amaranthus hybridus</i> L.	L.	Astr	Eur N A
<i>Amaranthus hypochondriacus</i> L.	G.	Sap	N Am. Eur
<i>Amaranthus viridis</i> L.	G.	Sap. Ant	Tr Reg
<i>Celosia anthemina</i> Aschers.	G.	Ant.	Tr Afr
<i>Celosia trigyna</i> L.	G.	Tacn.	Tr Afr
<i>Deeringia celsioides</i> R Br	G.		As Tr Austr

AMARYLLIDACEAE

<i>Amaryllis Belladonna</i> L.	G.		S. Afr
<i>Buphane disticha</i> Herb.	G.C Sm.L.	A P	S. Afr
<i>Crinum asiaticum</i> L.	G.		Tr As.
<i>Crinum zeylanicum</i> L.	G.		Ind
<i>Curculigo scorzoneraefolia</i> Baker	G.		Tr Am.
<i>Furcraea gigantea</i> Vent.	G.	Sap	Tr Am.
<i>Galanthus nivalis</i> L.	G.		Eur
<i>Haemanthus</i>	G.		Afr
<i>Haemanthus coccineus</i> L.	L.	Diur	S. Afr
<i>Hippeastrum equestre</i> Herb.	G.		Mex
<i>Hippeastrum rutillan</i> Herb.	G.		Braz. I
<i>Hymenocallis</i> sp.	G.		Tr Am.
<i>Leucojum aestivum</i> L.	C.		Ort
<i>Leucojum vernum</i> L.	G.		Eur
<i>Lycoris radiata</i> Herb.	G.	Fan.	China
<i>Narcissus poeticus</i> L.	M.	Ir	Eur
<i>Narcissus Pseudo-Narcissus</i> L.	M G.C.L.	Ir	Eur
<i>Narcissus Taxetta</i> L.	M.		Med. Reg
<i>Pancratium Illyricum</i> L.	G.		Eur Austr
<i>Pancratium maritimum</i> L.	G.C.L.	Diur	Med. Reg N
<i>Pancratium zeylanicum</i> L.	G.		Am.
<i>Sprekelia formosissima</i> Herb.	G.		Ind
<i>Zephyranthes Atanasco</i> Herb.	Ches.		Mex
			N A

ANACARDIACEAE

<i>Buchanania</i> sp.	G.		Tr As Pac. Is
<i>Comocladia glabra</i> Spreng.	G.		Austr
<i>Corynocarpus laevigata</i> Forst.	G.	Con.	Cuba
<i>Gluta Benghas</i> L.	G.		N Zeal
<i>Holigarna caustica</i> Roxb.	G.		Java.
<i>Holigarna longifolia</i> Roxb.	G.		Ind
<i>Lithraea venenosa</i> Miers.	G.		Ind.
<i>Mangifera indica</i> L.	G.		S Am
<i>Oncocarpus vitiensis</i> A.	G.		Malay, Ind.
<i>Pistacia Khanjuk</i> Stocks	G.	Ir	Fiji Is.
<i>Pseudosmodium perniciosum</i> Engl	G.	Ir	Egyp. Pers.
<i>Rhus Cotinus</i> L.	C.		Him
			Mex
			Eur China
			Med. Reg.
* <i>Rhus glabra</i> L.	L.	Astr	N A.

* Just as this part of the work is going to press I have received two important papers on *Rhus* by L. E. Warren in which the statement is made that the poisonous substance of *Rhus* is a powerful escharotic, one milligram producing very severe blistering when

NAME OF PLANT	Authority	Properties	Locality
<i>Rhus Metopium</i> L.	L.	Emeto-Cath.	N. A. W. I.
<i>Rhus Toxicodendron</i> L.	G. L.	Ir.	N. Am.
<i>Rhus venenata</i> DC.	G.	Ir.	N. Am.
<i>Rhus vermicifera</i> DC.	G. L.	Ir.	Th. b. Jap.
<i>Scierocarya Caffra</i> Sond.	G.	Ir.	S. Afr. Austr.
<i>Semecarpus Anacardium</i> L.	G. L.	Ir.	As. Tr. Austr.
<i>Semecarpus heterophylla</i> Bl.	G.	Ir.	Java
<i>Spondias mangifera</i> Willd.	G.	Ir.	Trop. As.

ERUCACEAE

<i>Anona amara</i> Raeusch.	G.		Guiana
<i>Anona Cherimolia</i> Mill.	G.	F. P.	Tr. Am.
<i>Anona muricata</i> L.	G. L.	Astr.	Tr. Am.
<i>Anona palustris</i> L.	G. L.	F. P.	Tr. Am. & Afr.
<i>Anona reticulata</i> L.	G. L.	Acr.	Tr. Am.
<i>Anona spinescens</i> Mart.	G.		Brazil
<i>Anona squamosa</i> L.	G. L.	F. P.	Ind.
<i>Asimina triloba</i> Dun.	G. L.	Em.	N. Am.
<i>Gomothalamus macrophyllus</i> Hook.	G.		Malacca.
<i>Guatteria veneficiorum</i> Mart.	G.		Brazil
<i>Popowia pisocarpa</i> Endl.	G.		Java
<i>Xylopi odoratissima</i> Welw.	G.		Tr. Afr.
<i>Xylopi polycarpa</i> Oliv.	L.	Berb.	Tr. Afr.
<i>Xylopi saicifolia</i> HBK.			Tr. S. Am.

APOCYNACEAE

<i>Acokanthera Lamarkii</i> G. Don.	G.		S. Afr.
<i>Acokanthera venenata</i> G. Don.	G.		S. Afr.
<i>Adenium Boehmanum</i> Schinz.	G.		Afr.
<i>Adenium obesum</i> Roem. & Schult.	G.		Afr. Arab.
<i>Adenium somalense</i> Oliv.	G.	A. P.	Afr.
<i>Agaveasma carycina</i> A. DC.	G.	P. P.	Ind.
<i>Alamanda cathartica</i> L.	G.	Em.	Tr. Am.
<i>Astonia constricta</i> F. Muell.	L.	Bitter	Trop. Old World.
<i>Alstonia Scholaris</i> R. Br.	G. L.		Ind.
<i>Alstonia venetata</i> R. Br.	G.		Austr.
<i>Alyxia buxifolia</i> R. Br.	G.		

placed on the arm for fifteen minutes. When in contact with the air it produces the characteristic non-toxic varnish. The resinous products from the latex of the *Rhus* give black compounds with the alkali hydroxides. The irritating properties he thinks will be found to be connected with the presence or the relations of these hydroxyl groups.

This writer found that the milk sap of *Rhus vernix* is analogous in almost every particular to the Japanese lac. Warren agrees with nearly everyone else that the bacteria infection theory has very little to support it. Nor is there much to support the recently elaborated theory of immunity. He shows how a popular impression in regard to supposed immunity has become widespread even among scientists. It has not however been demonstrated. In the same way, contrary to the early published records, *Rhus Michauxii* has been shown by Warren and Trelease not to be poisonous. As a summary, Warren, after an extensive investigation of the subject, states that the great majority are empirical. "There is no specific for this troublesome complaint. Remedies which have alleviated the symptoms in one case have proved utterly useless or worse in others. The following symptoms are listed by Warren as poisonous:

Rhus venenata D. C. (*R. vernix* L.), *Rhus radicans* L., *Rhus Toxicodendron* L., *Rhus diversiloba* Torr. & Gray, *Rhus Rydbergii* Small, *Rhus Metopium* L., *Rhus Florida* Mearns, *Rhus littoralis* Mearns, *Rhus vermicifera* DC., *Rhus succedanea* L., *Rhus sylvestris* Sieb. & Zucc., *Rhus Wallichii* Hook. fil., *Rhus Griffithii* Hook. fil., *Rhus striata* R. & Pav., *Rhus peruviana* H. B. & K., *Rhus chinensis* Mill., *Rhus jamaica* L., *Rhus copallina* Hook. & Arn., *Rhus lucida* L.

NAME OF PLANT	Authority	Properties	Locality
<i>Axylia daphnoides</i> A. Cunn.	G.		Norfolk Is
<i>Amsonia Tabernaemontana</i> Walt.			N. Am.
<i>Apocynum androsaemifolium</i> L.	Ches. G.C. Rusby L.	Card.	N. Am.
<i>Apocynum cannabinum</i> L.	Ches. G.C. Rusby L.	F P Card.	N. Am.
<i>Apocynum venetum</i> L.	G.		Med. Reg. Ori-
<i>Aspidosperma quebracho-blanco</i> Schlecht.	G. L.		ent N. As. Ind
<i>Aspidosperma sessilifolium</i> Fries.	G. R.	P	S. Am.
<i>Cameraria latifolia</i> L.	G.		S. Am.
<i>Carissa ovata</i> R. Br.	G.		Ind.
<i>Carissa xyloperon</i> Thon.	G.		Austr.
<i>Cerbera Odollam</i> Gaertn.	G. C. R.	P	Mascar Is
<i>Cerbera tanghin</i> Hook.	C. L.		Ind. Malay
<i>Echites alexicaca</i> Mart. & Stadelm.	L.	Cath.	Braz.
<i>Echites maculata</i> A. DC.	G. C.		St. Doming. Is.
<i>Echites venenosa</i> Marr.	G. L.	A. P.	Brazil.
<i>Ellertonia Rheedii</i> Wight.	G.		Ind.
<i>Gelasospermum velosum</i> Allen.	G. L.	Bitter	Braz.
<i>Haplophyton curcuidum</i> A. DC.	G.	Jus.	Mex.
<i>Malouetia nitida</i> Spruce.	G. L.	A. P.	Braz.
<i>Melotinus monogynus</i> Roxb.	G. R.	F. P.	Him. Malaya
<i>Nerium odoratum</i> Soland.	G.	Car.	China
<i>Nerium Oleander</i> L.	G. C. Mi. Ches. L.	Car.	Per. Jap. Ind.
<i>Ochrosia Moorei</i> F. von Muell.	G.		Med. Reg. Ori
<i>Plumera rubra</i> L.	C. L.		Austr.
<i>Pottsia cantoniensis</i> Hook. & Arn.	G.		Trop. Am.
<i>Prestonia toxifera</i>	G.		Ind. Java China
<i>Rauwolfia serpentina</i> Benth.	G.	P	Ind. Java.
<i>Rauwolfia tripartita</i> Gaertn.	G.		Java.
<i>Rauwolfia verticillata</i> Baill.	G.	F. P.	
<i>Strobilanthes calons</i> Nees.	K.		Ind.
<i>Strophanthus Emmeri</i> Asch.	L.	A. P.	Cent. Afr.
<i>Strophanthus hispidus</i> DC.	G. L.	F. P. A. P.	S. Afr.
<i>Strophanthus Kombe</i> Oliver.	L.	F. P.	Trop. Afr.
<i>Strophanthus Pierrei</i> Herm.	G.		W. Afr.
<i>Tabernaemontana Borbonica</i> Lam.	G.		
<i>Tabernaemontana citrifolia</i> L.	G.		Ind. Mex.
<i>Tabernaemontana cotinifolia</i> Willd.	G.		Ind.
<i>Tabernaemontana malaccensis</i> Hook.	G.		Malacca
<i>Tabernaemontana Mauritanica</i> Poir.	G.	F. P.	Mascar. Is.
<i>Tabernanthe iboga</i> Baill.	G.		Trop. Afr.
<i>Thevetia Ahoua</i> A. DC.	G. R.	F. P.	Brazil.
<i>Thevetia nereifolia</i> Juss.	G. C. R. L.	F. P.	Tr. Am.
<i>Urechites suberecta</i> Muell.	G. L.	Sed.	St. Doming. Is.
<i>Vincetoxicum major</i> L.	G. L.	Abort.	Med. Reg.
<i>Vincetoxicum minor</i> L.	L.	Astr.	Eur.
<i>Vincetoxicum pusilla</i> Murr.	G.		Ind.

AQUIFOLIACEAE

<i>Ilex Cassine</i> Walt.	Sm. L.	Em. Cath.	N. Am.
<i>Ilex glabra</i> A. Gray.	L.	Astr.	N. A.
<i>Ilex verticillata</i> A. Gray.	L.	Astr.	N. A.

NAME OF PLANT	Authority	Properties	Locality
<i>Alocasia indica</i> Schott.	G.		Ind. Malay
<i>Alocasia montana</i> Schott.	G.		Ind. Malay
<i>Amorphophallus viridis</i> Bl.	G.		
<i>Anthurium Fernerense</i>		Ir	
<i>Arisaema curvatum</i> Kunth.	G.	Ir	H. m. Reg.
<i>Arisaema tortuosum</i> Schott.	G.	Ir	H. m. Reg.
<i>Arisaema triphyllum</i> Schott.	Rusby	Em	N. A.
<i>Arisaema vulgare</i> Targ.	G.	Ir	Med. Reg.
<i>Arum Dioscoridis</i> Sibth.			Greece As. M. n.
<i>Arum Dracontium</i> Schott.	L.	Acr	N. A.
<i>Arum italicum</i> Mill.	G. C. M.	Sap.	Eur.
<i>Arum maculatum</i> L.	G. M.	Ir	Eur.
<i>Arum triphyllum</i> Schott.	L.	Acr	N. A.
<i>Caadium bicolor</i> Vent.	G.	Ins	Am. Austr.
<i>Calla palustris</i> L.	G. C. L.	Acr	Eur. As. N. Am.
<i>Colocasia antiquorum</i> Schott.	G. L.	Acr	Tr. As.
<i>Colocasia gigantea</i> Hook. f.	G.	Acr	
<i>Colocasia vitosa</i> Kunth.	G.	Acr	Ind.
<i>Cryptocoryne spiralis</i> Fisch.	G.	Ins	Ind.
<i>Cyrtosperma Merkusii</i> Schott.	G.		Sumatra
<i>Dracontium asperum</i> C. Koch.	G.		Brazil
<i>Dracontium vulgare</i> Schott.	C.		S. Eur.
<i>Epipremnum mariae</i> Schott.	G. L.	Ant	Java
<i>Homalomena amabilis</i> Schott.	G.		Malay Ind.
<i>Lasia aculeata</i> sp. Lour.	G.	Hyd	Tr. As.
<i>Monstera pertusa</i> Schott.	G.	Ir	Tr. Am.
<i>Philodendron bipinnatifidum</i> Schott.	G.		Brazil
<i>Philodendron beneraceum</i> Schott.	G.	P.	S. Am.
<i>Philodendron imbe</i> Schott.	G.	P.	Argentina
<i>Philodendron Simsii</i> Sweet.	Sm.	Ir	Guyana
<i>Pnelia tuberosa</i> Ten.	G. L.	Ir	Japan
<i>Semdayas officinalis</i> Schott.	G.	Ant	Burma.
<i>Spathiphyllum candicans</i> Poepp.	G.		Trop. Am.
<i>Synplocarpus foetidus</i> Nutt.	Sm.	Ir	E. N. Am.

ARALIACEAE

<i>Aralia</i> sp.	G.	Sap.	N. Am. Asia
<i>Aralia hispida</i> Vent.	L.	Dur.	N. E. U. S.
<i>Aralia nudicaulis</i> L.	L.	Dur.	N. E. U. S.
<i>Aralia racemosa</i> L.	G. L.	Pan.	N. A.
<i>Aralia spinosa</i> L.	L.	Dur.	N. Am.
<i>Hedera Helix</i> L.	G. C. L.	Lin. Nar.	Eur. Afr. As.
<i>Heptapleurum emarginatum</i> Seem.	G.		Zeylan
<i>Heptapleurum scandens</i> Seem.	G.		Malay
<i>Heptapleurum venulosum</i> Seem.	G.		Austr.
<i>Panax</i> sp.	G.	Sap.	N. Am. As.
<i>Polyscias</i> sp.	G.	Sap.	Austr. N. Zeal.
			Afr. Madagas.
			Malay. Pac. Is.
<i>Polyscias nodosa</i> Seem.	G.	S. P.	Malay
<i>Trevesia</i> sp.	G.	Sap.	Java Sumatra.

ARISTOLOCHIACEAE

<i>Aristolochia angustida</i> Jacq.	G.	Diaph.	S. Am.
<i>Aristolochia antihysterica</i> Mart.	G.	Em.	Brazil.
<i>Aristolochia argentina</i> Griseb.	G.	Em.	Argent. Reg.

NAME OF PLANT	Authority	Properties	Locality
<i>Aristolochia brasiliensis</i> Mast. & Zucc.	G.	Em.	Brazil.
<i>Aristolochia Clematitis</i> L.	G. C. M.	Em.	Eur Caucas Reg As. Minor
<i>Aristolochia elegans</i> Mast.	G.	Em.	Brazil
<i>Aristolochia grandiflora</i> Arruda	G. C.		Pernamb
<i>Aristolochia indica</i> L.	G.		Ind.
<i>Aristolochia Kaempferi</i> Willd.	G.		Japan
<i>Aristolochia longa</i> L.	C. L.		Med Reg.
<i>Aristolochia pallida</i> Willd.	G.	F P Nar	Eur As. Minor
<i>Aristolochia Pistolochia</i> L.	C. L.		Med Reg.
<i>Aristolochia rotunda</i> L.	G. L.	Nar	Med Reg.
<i>Aristolochia Serpentaria</i> L.	L.	Diaph	E. U. S.
<i>Aristolochia</i> sp.	E.	F P	Italy
<i>Asarum albiventum</i> Regel.	G.	Em.	Japan
<i>Asarum arifolium</i> Mx.	L.	Ir	N A
<i>Asarum canadense</i> L.	G. L.	Ir	N Am
<i>Asarum caudatum</i> L.	L.	Ir	Calif
<i>Asarum europaeum</i> L.	M. Sm. L.	Ir Em-Cat	Eur
<i>Asarum virginicum</i> L.	L.	Ir	N A
<i>Bragantia tomentosa</i> Bl.	G.	F P	Ind. Malay
<i>Thottea dependens</i> Klotzsch.	G.	Ir	Malay

ASCLEPIADACEAE

<i>Asclepias sericifera</i> Brot.	G.	Em.	Peru
<i>Asclepias curassavica</i> L.	G. L.	Em.	S Am
<i>Asclepias etiocarpa</i> Benth.	L.	Em.	W N Am
<i>Asclepias incarnata</i> L.	Ches. C. L.	Ant	N Am.
<i>Asclepias mexicana</i> Cav.	Ches.	Em.	W N Am
<i>Asclepias syriaca</i> L.	Ches. C. L.	Em.	N Am
<i>Asclepias tuberosa</i> L.	C. Ches.	Em.	N Am
<i>Calotropis gigantea</i> R. Br.	G. L.	Em.	Ind.
<i>Calotropis procera</i> R. Br.	G.	Inch	Peru N Afr Ind.
<i>Ceropegia bulbosa</i> Roxb.	G.		Ind.
<i>Chlorocodon Whitei</i> Hook.	G.	Cou	S. Afr
<i>Choristigma Streckertianum</i>	G.		Arg.
<i>Cosmosigma racemosa</i> W. ght.	G.		Ind.
<i>Cryptostegia grandiflora</i> R. Br.	G.		Tr Afr Ind (Cult.)
<i>Cynanchum acutum</i> L.	G. C. L.	Cath.	Eur Ori. As
<i>Cynanchum sarcostemmonifolius</i> K. Schum.	G.		
<i>Gonolobus larvis</i> Mx.	G.	A P	N Am.
<i>Gonolobus obliquus</i> R. Br.	G.		N Am.
<i>Gymnema latifolium</i> Wall.	G.		Ind Burma
<i>Gymnema sylvestre</i> R. Br.	G. L.	Destroys sense of taste	Ind Tr Afr Austr
<i>Hemidesmus indicus</i> R. Br.	G. L.	Cou	Zeylan.
<i>Marsdenia Cundurango</i> Nich.	L.	Dog Poison	S. A
<i>Marsdenia erecta</i> R. Br.	G. L.		Syr
<i>Metabea venenata</i> Baill.	G. L.	Ord.	Madagas
<i>Metaplexis Stuntoni</i> Schult.	G.		China
<i>Morrenia brachystephana</i> Griseb.	G.		Argentina
<i>Periploca graeca</i> L.	G. L.	Em. Wolf- Poi	S. Eur
<i>Periploca vomitoria</i> Leschen.	G.	Em.	Java
<i>Sarcolobus carinatus</i> Wall.	G.		Ind. Burma
<i>Sarcolobus narcoticus</i> Miq.	G.	Nar	Java

NAME OF PLANT	Authority	Properties	Locality
<i>Sarcostemma australe</i> R Br	G. M.		Aust.
<i>Sarcostemma glaucum</i> HBK.	G.	Em.	S. Am. Venez.
<i>Solenostemma Argel</i> Del.	L.		Nabia.
<i>Stape</i> a sp.	C.		S. Afr.
<i>Tylophora asthmatica</i> W & A.	G. L.	Em.	Ind. Malay.
<i>Tylophora fasciculata</i> Ham.	G.	Em.	Ind.
<i>Tylophora laevigata</i> Decne.	G.	Em.	Maurit. Is.
<i>Vincetoxicum nigrum</i> Moench.	C.	Dr Em.	Eur. As. M. nor.
<i>Vincetoxicum officinale</i> Moench.	C. M. L.	Dr. Eo.	Eur. Can. C.

BALSAMINACEAE

<i>Impatiens amphorata</i> Edgew.	G.		Hon.
<i>Impatiens aurea</i> Muhl.	L.	Diur.	N. A.
<i>Impatiens biflora</i> Walt.	L.	Diur.	N. A.
<i>Impatiens Noli-tangere</i> L.	G. L.	1/2 Diur.	Eur. Siberia.

BEGONIACEAE

<i>Begonia gracilis</i> HBK.	G.	Em.	Mex.
------------------------------	----	-----	------

BERBERIDACEAE

<i>Berberis Aquifolium</i> Pursh.	G. C. L.		N. Am.
<i>Berberis aristata</i> DC.	G. L.	F. P.	Ind.
<i>Berberis vulgaris</i> L.	G. L.	F. P. fr.	Eur. Tem. As.
<i>Caulophyllum thactioides</i> Michx.	G. L.	Sap.	N. Am.
<i>Leontice Leontopetalum</i> L.	G. L.	Sap. for.	Italy Orient.
<i>Narcissa vomica</i> Thunb.	G.	Opium.	Japan China.
<i>Podophyllum Eriod. Walt.</i>	L.	Cath.	Hon. Reg.
<i>Podophyllum peltatum</i> L.	Sm. Rusby.	Pur.	N. Am.
	C. Chex.		

BETULACEAE

<i>Betula lenta</i> L.	Witthaus.	Vetr.	E. N. Am.
<i>Betula lutea</i> Michx.	Witthaus.	Aptr.	E. N. Am.

BIGNONIACEAE

<i>Bignonia capreolata</i> L.	G. R. L.	F. P.	N. Am.
<i>Crescentia cucurbitaria</i> L.	G.		Tr. An.
<i>Crescentia Cujele</i> L.	G.		Trop. An.
<i>Dolichanassa falcata</i> Seem.	G.	F. P. Abort.	Ind.
<i>Jacaranda procera</i> Spr.	R.	F. P.	Guana.
<i>Osmohydrophora nocturna</i>	G.	Hyl.	Brax.
<i>Tanaecium crucegerum</i> Seem.	G.		W. Ind.
<i>Tecoma ceramensis</i> Teijsm. & Binn.	G.		Ceram. Is.
<i>Tecoma leucoxylen</i> Mart.	G. R.	F. P.	Guayaq. W. Ind.
<i>Tecoma obtusata</i> DC.	G.		Brazil.
<i>Tecoma radicans</i> Juss.	G. R.	F. P.	N. Am.
<i>Tecoma speciosa</i> DC.	G.		Brazil.
<i>Tecoma stans</i> Juss.	G.		N. Am. S. Am.
<i>Tecoma toxophora</i> Mart.	G.		Brazil.

BIXACEAE

<i>Gynocardia odorata</i> R Br.	G. L.	F. P.	Ind.
<i>Hydnocarpus anthelminticum</i> Pierce.	L.	Ant.	
<i>Hydnocarpus castanea</i> H. F. & Thoms.	G.	F. P.	Andoman Is. Malacca.

NAME OF PLANT	Authority	Properties	Locality
<i>Hydnocarpus heterophylla</i> Bl.	R	F. P	Java.
<i>Hydnocarpus Kurzii</i>	L.	Ant.	Trop. As.
<i>Hydnocarpus pascidia</i>	Sm.		
<i>Hydnocarpus venenata</i> Gaertn.	E. G. L. R.	F. P	Zeylan.
<i>Hydnocarpus Wrightiana</i> Bl.	L. R.	Ant.	Ind.
<i>Pangium edule</i> Reinw.	G. R. E.	Hyd. F. P.	Java.
<i>Taraktogenos Blumei</i> Hassk.	G.	F. P.	Sumatra.
<i>Trichadenia zeylanica</i> Thw.	G.		Zeylan.

BORAGINACEAE

<i>Asperugo procumbens</i> L.	Sm.		N. As. Eur.
<i>Borago officinalis</i> L.	Wh. C.	Tr.	Eur. Afr. As.
<i>Pourreria Havanensis</i> Miers.	G.		Cuba.
<i>Cynoglossum officinale</i> L.	G. L.	Tr.	Eur. Orient.
<i>Echium vulgare</i> L.	G.	Tr.	Eur. N. Am.
<i>Hebentropium europaeum</i> L.	G.		Eur. Afr. Ori.
			Austr. N. Am.
			Ind.

BROMELIACEAE

<i>Ananas sativus</i> Schult.	G. L.	Ant.	Tr. Am.
<i>Bromelia Pinguin</i> L.	G. L.	Sap.	Tr. Am.
<i>Karatas Plumieri</i> Emorr.	G.	Ant.	Parana.

BOMBACACEAE

<i>Liquidambar punctatam</i> R. Br.	G. M.		Amer.
-------------------------------------	-------	--	-------

BURSERACEAE

<i>Canarium</i> sp.	G.	F. P.	Aust. Pac. Is.
			Philipp.
<i>Canarium commune</i> Lam.	L.	Cath.	Molucca Is.

BUXACEAE

<i>Buxus balearica</i> Lam.	G.	Cath.	Balearic Is.
<i>Buxus sempervirens</i> L.	Ches. G. C.	Tr.	Eur. Ori. Tem.
	L.		As.

CACTACEAE

<i>Anhalonium Lewinii</i> Henn.	G. L.	Card. I. vel. Nar.	Mex.
<i>Cereus Hoplandii</i> Parm.	L.	Card. Sed.	Brazil.
<i>Cereus flagelliformis</i> Mill.	G.	Ant.	S. Am.
<i>Cereus grandiflorus</i> Mill.	G. L.	Ir. Card. Sed.	W. Ind.
<i>Cereus McDonaldii</i> Hook.	L.	Card.	Honduras.
<i>Echinocactus</i> sp.	G.	Tr.	N. Am.
<i>Echinocactus Williamsii</i> Lem.	G. L.	Nar.	Mex. Tex.
<i>Mammillaria</i> sp.	G.		W. N. Am.
<i>Rhipsalis</i> sp.	G.		S. Am. Trop.
			Am.

CALYCANTHACEAE

<i>Calycanthus glaucus</i> Willd.	G. L. Ches.	Sheep Poi.	N. Am.
-----------------------------------	-------------	------------	--------

CAMPANULACEAE (Including LOBELIACEAE)

NAME OF PLANT	Authority	Properties	Locality
<i>Isotoma axillaris</i> Lindl.	G. M.		Austr.
<i>Isotoma Brownii</i> G. Don.	M.		Austr.
<i>Isotoma longiflora</i> Presl.	G. M.		W. Ind.
<i>Lobelia cardinalis</i> L.	C. Mi. L.	Ant.	E. N. Am.
	Rusby		
<i>Lobelia fulgens</i> Willd.	C. Mi.	Ac.	Eur.
<i>Lobelia inflata</i> L.	Sm. Ches.	Dep. Em.	N. Am.
	L. Rusby		
<i>Lobelia Kalma</i> L.	Ches.		N. A.
<i>Lobelia nicotianaeifolia</i> Heyne	G.	Ac.	Ind.
<i>Lobelia pratioides</i> Benth.	G. M.	Ac.	Austr.
<i>Lobelia thynchopetalum</i> Hemsl.	G.		(Tr. Afr.)
<i>Lobelia siphilitica</i> L.	Rusby L.	Ac.	E. N. A.
	Ches.		
<i>Lobelia spicata</i> Ruiz. & Pav.	Ches.		Peru.
<i>Lobelia Tupa</i> L.	G. R.	F. P.	Chil. Peru.
<i>Lobelia urens</i> Willd.	G. C.	Ac.	Eur.
<i>Pratia erecta</i> Gaud.	G. M.		Austr.
<i>Siphocampylus giganteus</i> Don.	G.		Equad.

CANELLACEAE

<i>Canela alba</i> Marr.	G. L.	Sap. F. P.	W. Ind.
--------------------------	-------	------------	---------

CAPPARIDACEAE

<i>Boscia urens</i> Welw.	G.		Tr. Afr.
<i>Cadaba ineica</i> Lam.	G.		Ind.
<i>Capparis cynophallophora</i> L.	G. Sm. L.	Ant. Diar.	S. Am.
<i>Capparis ferruginea</i> L.	L.	Ant. Diar.	W. Ind.
<i>Capparis globifera</i> Del.	G.		Tr. Afr.
<i>Capparis odoratissima</i> Jacq.	G.		S. Am. Venez.
<i>Cleome Chelidoni</i> Lf.	G.		Ind. Java
<i>Cleome frutescens</i> Aubl.	G.		Gaiana
<i>Cleome gigantea</i> L.	G.		Braz. S. Am.
<i>Cleome graveolens</i> Raf.	G.	Tr.	N. Am.
<i>Cleome psoralenifolia</i> DC.	G.		Brazil
<i>Cleome rosea</i>	G.	F. P.	Texas
<i>Cleome spinosa</i> Jacq.	G.		S. Am.
<i>Maerua angolensis</i> DC.	G.		Tr. Afr.
<i>Morisonia americana</i> L.	G.		S. Am.

CAPRIFOLIACEAE

<i>Abelia uniflora</i> R. Br.	G.	Hyd.	China.
<i>Diervilla japonica</i> DC.	G.	Sap.	Japan
<i>Diervilla trifida</i> Moench.	L.	Diar.	N. Am.
<i>Lonicera chrysantha</i> Turcz.	G.	Sap.	Siberia
<i>Lonicera involucrata</i> Banks.	G.	Sap.	N. Am. Ry. Mts.
<i>Lonicera japonica</i> Thunb.	G.	Sap.	Him. Reg.
<i>Lonicera Standishii</i> Hook.	G.		China
<i>Lonicera tatarica</i> L.	G.	Sap.	Siberia
<i>Lonicera tomentella</i> Hook. f. & Thoma.	G.	Sap.	Him. Reg.
<i>Lonicera Xylosium</i> L.	G.	Sap.	Eur. N. As.
<i>Sambucus canadensis</i> L.	C. Rusby L.	Em.	E. N. Am.
<i>Sambucus Ebulus</i> L.	G. C.	Sap.	E. Eur. Him. Reg.
<i>Sambucus mexicana</i> Presl.	L.	Em.	W. N. A.

NAME OF PLANT	Authority	Properties	Locality
<i>Sambucus nigra</i> L.	G.C. Rusby	Pur	Eur N As
<i>Sambucus racemosa</i> L.	G. C.		N Tem Reg.
<i>Symphoricarpos molle</i> Nutt.	G.	Sap.	W N Am
<i>Symphoricarpos racemosus</i> Michx.	G.	Ir	N Am
<i>Triosteum majus</i> Michx.			E N Am
<i>Triosteum perfoliatum</i> L.	G. Rusby L.	Calh	N Am
<i>Viburnum Lantana</i> L.	G. L.	Astr	Eur Cauc Reg
<i>Viburnum macrophyllum</i> Thunb.	G.	Sap	Japan
<i>Viburnum Opulus</i> L.	G. L.	Sed	Eur As. N Am

CARYOPHYLLACEAE

<i>Acaulophyllum C. A. Mey. sp.</i>	G.		
<i>Agrostemma Githago</i> L.	G. C. Sm.	Sap Ir	Eur As Int. N
	Ches.		& S. Am Austr
<i>Arenaria serpyllifolia</i> L.		Short	Eur
<i>Dianthus chinensis</i> L.	G.	Sap.	Eur Tem. As
<i>Gypsophila Struthium</i> L.	G. L.	Sap.	Spain
<i>Herniaria glabra</i> L.	G.	Sap.	Eur N As.
<i>Lychris chalcidonica</i> L.	G.	Sap.	Japan
<i>Lychris dioica</i> L.	G.	Sap.	Eur N As
<i>Lychris Flos-cuculi</i> L.	G.	Sap.	Eur N As.
<i>Lychris indica</i> Benth.	G.	Sap.	Him Reg
<i>Polycarpha sp.</i>	G.	Sap.	Austr Pac. Is.
			Trop.
<i>Saponaria officinalis</i> L.	G. C. L.	Ir Sap.	Eur, W As.
<i>Saponaria Vaccaria</i> L.	G.	Sap.	Eur As Min
			Siber
<i>Silene antirrhina</i> L.	Ches.		E. N Am
<i>Silene Cucubalus</i> Wibel	G.		Eur N Afr
			Him. Reg
<i>Silene Griffithi</i> Boiss.	G.		Tem. Reg.
<i>Silene macrodon</i> Steud.	G.		Abyss
<i>Silene virginica</i> L.	G.	Ant	N Am
<i>Silene viscosa</i> Pers.	G.		Eur N Am
<i>Stelaria crassifolia</i> Ehrh.	G.		N & Arct Reg

CELASTRACEAE

<i>Catha edulis</i> Forsk.	G.	Sum	Trop Afr
<i>Celastrus scandens</i> L.	Ches. L.	an	E N Am.
<i>Elaeodendron orientale</i> Jacq.	G.		Madag Is.
			Maurit
<i>Euonymus atropurpureus</i> Jacq.	G. L.	Em	N Am.
<i>Euonymus Europaeus</i> L.	G. C.	lar Pur	Eur As Min
<i>Euonymus latifolius</i> Mill.		Em	Eur As Minor
<i>Karrimia seylanica</i> Arn.	G.	Hyd	Zeyl.
<i>Lophopetalum pallidum</i> Laws	G.		Malacca.

CHENOPODIACEAE

<i>Atriplex Halimifolia</i> L.	G.	Sap.	N & S Afr
<i>Atriplex hortensis</i> L.	G. L.	Sap. Em.	Cosmop. Cult.
<i>Atriplex laciniata</i> L.	G.	Sap.	Eur N Am
<i>Atriplex Nuttallii</i> S. Wat.	G.	Sap.	W N Am
<i>Atriplex rosea</i> L.	G.	Sap.	Eur N Am N
			Afr As Minor
<i>Atriplex tatarica</i> L.	G.	Sap.	Cert As.
<i>Atriplex vesicaria</i> Hew.	G.	Sap.	Austr
<i>Chenopodium ambrosioides</i> L.	G. Rusby L.	Ant	Temp. & Trop

NAME OF PLANT	Authority	Properties	Locality
<i>Chenopodium anthelminticum</i> L.	Sm. L.	Ant	Reg. N. Am. Temp. & Trop.
<i>Chenopodium Bonus-Henricus</i> L.	L.	Ant	Reg. N. Am.
<i>Chenopodium Botrys</i> L.	G. L.	Ant.	Eur. N. Am. N. Reg. N. Am.
<i>Chenopodium californicum</i> S. Wat.	G.	Sap.	Eur.
<i>Chenopodium hybridum</i> L.	G.		Cal.
<i>Chenopodium mexicanum</i> Moq.	G.	Sap.	N. Temp. Reg.
<i>Chenopodium polyspermum</i> L.	R.	F. P.	N. Am. Eur.
<i>Chenopodium rubrum</i> L.	G.		Mex.
<i>Chenopodium Vavaria</i> L.	Sm. L.	Ant.	Eur. N. As.
<i>Eurotia ceratoides</i> C. A. Mey.	G.	Sap.	Eur. N. As.
<i>Halenia frutescens</i> D. Dietr.	G.	Ant.	Cal. Eur. N. As.
<i>Kochia arenaria</i> Roth.	G.	Sap.	W. N. Am.
<i>Kochia scoparia</i> Schrad.	G.	Sap.	S. Eur. N. Afr.
<i>Kochia trichophylla</i> Hort.	G.	Sap.	S. Eur. W. As.
<i>Salsola Kali</i> L.	G.	Ant.	Eur. W. As.
<i>Salsola Kali</i> var. <i>tenuifolia</i> G. F. W. Mey.		Tr.	N. Am.
<i>Salicornia tamariscifolia</i> Cav.	G.	Ant.	Am.
<i>Sarcobatus Maximilianii</i> Nees.	Ches.	Typ.	Eur. As. N. Am.

CISTACEAE

COMBRETACEAE

<i>Combretum bracteosum</i> , Brandis	G.	Sap.	
<i>Combretum erythrophyllum</i> Sond.	G.		S. Afr.
<i>Combretum phaneropetalum</i> Bak.	G.		S. Afr.
<i>Combretum racemosum</i> Beauv.	G.		Trop. Afr.
<i>Combretum trifoliatum</i> Vent.	G.	Ant.	Hab.
<i>Gustavia augusta</i> L.	G. R.	F. P.	N. Grana.
<i>Gustavia brasiliensis</i> DC.	G. R.	F. P.	Braz.
<i>Quisqualis indica</i> L.	G.	Ver.	Tr. As.
<i>Terminalia Belerica</i> Roxb.	G. L.	F. P.	Ind. Malay.
<i>Terminalia Chebula</i> Retz.	L.	Ant.	Tr. As.
<i>Terminalia tomentosa</i> Wight.	G.		Ind.

COMMELINACEAE

<i>Athyrocarpus persicariifolium</i> Hemsl.	G.		Panama
<i>Commelina deficiens</i> Herb.	G.	Sap.	Braz.
<i>Commelina nudiflora</i> L.	G.		Tr. Reg.
<i>Commelina tuberosa</i> L.	G. L.		Mex.
<i>Tradescantia crassifolia</i> Cav.	P.		S. Am.
<i>Tradescantia elongata</i> G. F. W. Mey.	G.		S. Am.

COMPOSITAE

<i>Achillea Ptarmica</i> L.	C. L.	Acc.	N. Temp. Reg.
<i>Anacyclus officinarum</i> Hayne.	G. L.	Hyd.	S. Eur.
<i>Anacyclus pedunculatus</i> Pers.	G.	Tr.	Barbar.
<i>Anacyclus Pyrethrum</i> DC.	Sm. L.	Tr.	N. Afr. Orient.
<i>Anthemis aetnensis</i> Schouw.	G.	Hyd.	Mt. Aetna.
<i>Anthemis altissima</i> L.	G.	Hyd.	Eur. Orient.

NAME OF PLANT	Authority	Properties	Locality
<i>Anthemis arvensis</i> L.	G.	Hyd.	Eur Orient
<i>Anthemis austriaca</i> Jacq.	G.	Hyd.	Eur Orient
<i>Anthemis Blancheana</i> Bossa.	G.	Hyd.	Syria
<i>Anthemis chia</i> L.	G.	Hyd.	Greece, As. Min.
<i>Anthemis Cota</i> L.	G.	Hyd.	S. Eur
<i>Anthemis Cotula</i> L.	G. L.	Ir	Eur N Afr Orient
<i>Anthemis elbuenensis</i>	G.	Hyd.	
<i>Anthemis montana</i> L.	G.	Hyd.	Eur Orient
<i>Anthemis nobilis</i> L.	L.	Naus Stim	Eu As
<i>Arnica alpina</i> Olin and Ladau.	L.	Nar	Eur N A
<i>Arnica Chamissonis</i> Less.	L.	Nar	Unalaschk
<i>Arnica montana</i> L.	G. Sm. L.	Pur Ir	Eur N A
<i>Arnica nudicaulis</i> Ed.	L.	Nar	N A
<i>Artemisia Abrotanum</i> L.	G. L.	Inch	Eur Temp. A
<i>Artemisia Absinthium</i> L.	G. C. L.	Inch. Nar	
<i>Artemisia arenaria</i> DC	Rushy	Ant	Eur.
<i>Artemisia maritima</i> L.	Sm.	Inch	Cau. Reg.
<i>Artemisia mexicana</i> Willd.	L. G.	Inch. Ant	Eur. Cauc. Sib
<i>Artemisia pontica</i> L.	G.	Nat	Mex
<i>Artemisia triffida</i> Nutt.	L.	Ant	Eur Cau Reg
<i>Artemisia trifida</i> Nutt.	L.	Diaph	W N A
<i>Artemisia vulgaris</i>	L.	Antisp	Tem. Reg
<i>Athanasia amara</i> L.	G.	Ant	
<i>Atractylis gummifera</i> L.	G. C. L.		Med. Reg.
<i>Baccharis coraifolia</i> DC	G. L.	Poa Alk	Brazil
<i>Cardopatum corymbosum</i> Pers	G.		As. Min. Med Reg.
<i>Carlinia aculeis</i> L.	G. L.	Diur	Eur
<i>Centaurea Calceitrapa</i> L.	L.	Ir	N Afr Eur Temp. As
<i>Centipeda orbicularis</i> Lour	G.	Ant	As. Austr Pac. Is
<i>Chrysanthemum Balsamita</i> L.	G. L.	Ant	W As.
<i>Chrysanthemum cinerariacifolium</i> Vis.	G. L.	Ins	Dalmatia
<i>Chrysanthemum Parthenium</i> Bernh.	G. C. L.	Ins. Ant	Eur
<i>Chrysanthemum pinnatifidum</i> L.	G.	Cuc	Madera Is.
<i>Chrysocoma Coma-aurea</i> L.	G.		S. Afr
<i>Cibadium Barbases</i> DC	R. F.	F P	
<i>Cibadium surinamense</i> L.	G. R. F.	F P	S. Am.
<i>Cosmos sulphureus</i> Cav	G.		Mex
<i>Crepis lacera</i> Ten.	G.		Italy
<i>Cynara Cardunculus</i> L.	G.	Diur	Med. Reg
<i>Dimorphotheca Ecklonia</i> DC.	G.	Hyd. Sap	S. Afr
<i>Dimorphotheca paviata</i> Moench	G.		S. Afr
<i>Doronicum pardalanchis</i> L.		Ir	Eur As
<i>Dysodia papposa</i> Hitch.		Ir	N Am.
<i>Echinops Ritro</i> L.	G.		N Am
<i>Elephantopus tomentosus</i> L.	G. L.	Em	N Am
<i>Elytropappus glandulosus</i> Less.	G.		S. Afr
<i>Eupatorium amarissimum</i>	G.	Ins	
<i>Eupatorium cannabinum</i> L.	G. L.	Diur	Eur N As.
<i>Eurydia moschata</i>	G.		
<i>Flaveria Contrayerba</i> Pers.	G.	Ant	Trop. Am.
<i>Gnephosis eriocarpa</i> Benth.	G.		Austr
<i>Grindelia robusta</i> Nutt	G. L.	Antispas	Cal
<i>Grindelia Tournefortii</i> L.	G.	Er	
<i>Haplocarpha lyrata</i> Hav	G.		S. Afr

NAME OF PLANT	Authority	Properties	Locality
<i>Helianthus autumnalis</i> L.	Ches. G. L.	Ir Nar	N Am
<i>Helianthus tenuifolius</i> Nutt.	Ches. G. L.	Ir Nar	N Am S. States
<i>Helianthus tuberosus</i> L.	C		N Am
<i>Helichrysum apiculatum</i> D. Don	G.	Ant	Austr
<i>Hieracium vicosum</i> Pall.	G.		N As Eur
<i>Ichthyothere Cunaba</i> Mart.	E. G. R.	F P	Brazil
<i>Inula Royleana</i> DC	G		Him. Reg.
<i>Lactuca Scariola</i> L.	G C. M. L.	Nar	E. Eur
<i>Lactuca taraxacifolia</i> Schum.	G.	Nar	Tr Afr
<i>Lactuca tatarica</i> C. A. Mey	G.		W As. In
<i>Lactuca virosa</i> L.	G. M. L.	Nar	Eur N As.
<i>Launaea pinnatifida</i> Cass	G.		As. Trop. Afr
<i>Montanoa floribunda</i> C. Koch.	G.		Mex
<i>Montanoa tomentosa</i> Cerv	G.		Mex
<i>Mutisia viridifolia</i> Cav	G. L.	Sap. Sed.	Chd.
<i>Oldenburgia Arbuscula</i> DC	G	Sap	S Afr
<i>Olearia macrodonta</i> Bak	G.	Sap.	N Zeal.
<i>Olearia moschata</i> Hook.	G		N Zeal.
<i>Onopordon Acarhium</i> L.	G.		Eur N Afr
<i>Othonopsis intermedia</i> Boiss.	G.		Orient
<i>Parthenium Hysterophorus</i> L.	L.		Pernia
<i>Parthenium integrifolium</i> L.	G.	Diur	N A Aus.
<i>Pentzia virgata</i> Less	G.		S Afr
<i>Perezia oxylepis</i> Sch.	G.		Mex
<i>Petasites officinalis</i> Moench.	G.		Eur N As
<i>Picris hieracioides</i> L.	L.	Bitter	Eur Tem As.
<i>Prenanthes albasima</i> L.	G. L.	Bitter	Aus. N Zeal.
<i>Pterigeron adscendens</i> Benth.	G. M.		N A.
<i>Pterocaulon pycnostachyum</i> E. L.	G.	Nar	Aus.
<i>Puhcaria dysinterica</i> Gaertn.	L.	los	N A.
<i>Senecio gaudalensis</i> .	Ches.		Rur
<i>Senecio Grayanus</i> Hemsl.	G		Mex
<i>Senecio Jacobaea</i> L.	G	Abort.	Mex
<i>Senecio toluccarus</i> DC.	G L. Ches.	Anodyne	Eur N As. V.
<i>Senecio vulgaris</i> L.	G		Mex
<i>Senecio vulneraria</i> DC	G L.	Astr	Eur
<i>Silybum Marianum</i> Gaertn.	G.		Mex
<i>Solidago Virgaurea</i> L.	L.	Exp.	Eur
<i>Spilanthes Acmella</i> Murr	G L.	Diur	N Tem. Reg
<i>Spilanthes oleracea</i> L.	G L.	Purg	Cosm. Trop. & Sub-trop.
<i>Tanacetum umbelliferum</i> Boiss.	L.	Purg.	S A.
<i>Tanacetum vulgare</i> L.	G.	Abort.	Afghan.
<i>Vernonia anthelmintica</i> Willd.	G Rusby L.	Abort.	Eur N As.
<i>Vernonia nigritiana</i> Oliv. and Hiern.	G. L.	Ant.	Him Reg.
<i>Xanthoxia texanum</i> DC.	G		Tr Afr
<i>Xanthium canadense</i> Mill.	G.	Sap	Texas
<i>Xanthium spinosum</i> L.	G. Ches.	Mech	N Am.
<i>Xanthium Strumarium</i> L.	G Ches.		Cosm.

CONIFERAE

<i>Cephalotaxus drupacea</i> Sieb and Zucc.	G.		Jap.
<i>Ginkgo biloba</i> L.	G.	Acr	Jap.
<i>Juniperus Sabina</i> L.	C.G. M. L.	Abort.	Eur N As
<i>Juniperus oxycedrus</i> L.	Sm.	Abort	S. Eur Cauc. Reg.

NAME OF PLANT	Authority	Properties	Locality
<i>Juniperus virginiana</i> L.	C. L. G. M.	Abort	E. N. A.
<i>Picea excelsa</i>	White	Ir	Eur Cult. N. Am.
<i>Pinus Laricio</i> Poir	Sm.	Ineb.	S. Eur
<i>Pinus sylvestris</i> L.	Sire	Ineb.	Eur N. Am.
<i>Sequoia sempervirens</i> Endl.	G.		Eur
<i>Taxus baccata</i> L.	M. R. L.	F. P.	W. N. A.
<i>Taxus brevifolia</i> Nutt.		Astr.	E. N. Am.
<i>Taxus canadensis</i> Marsh.	Ches. Wh. L.	Astr.	N. A.
<i>Thuja occidentalis</i> L.	G. L.	Abort.	Jap
<i>Torreya nucifera</i> Sieb. and Zucc.	G.		N. N. Am.
<i>Tsuga canadensis</i>	Wh. L.	Astr.	

CONNARACEAE

<i>Agelaea emetica</i> H. Bu.	G.		
<i>Cnestis corniculata</i> Lam.	G.		Trop. Afr.
<i>Cnestis glabra</i> Lam.	G.		Mal. Is.
<i>Cnestis polyphylla</i> Lam.	G.		Mad. Maurit.
<i>Cnestis glabra</i> Lam.	G.		Trop. Afr.
<i>Connarus africanus</i>	G.		Trop. Afr.
<i>Rourea glabra</i> HBK.	G.	Con.	Venez.

CONVOLVULACEAE

<i>Convolvulus Scammonia</i> L.	C. Sm. L.	Cath.	As. Minor
<i>Convolvulus sepium</i> L.	G.		Eur. N. A.
<i>Convolvulus venenatus</i> West.	G.		St. Croix Is.
<i>Cuscuta americana</i> L.	G.		N. A.
<i>Cuscuta australis</i> R. Br.	G.	A. P.	Aus.
<i>Cuscuta Dorycnium</i> L.	G.		
<i>Cuscuta epithymum</i> Murr.	Sm.	Cath.	Eur. N. Am.
<i>Cuscuta europaea</i> L.	Sm.	Cath.	Eur. Orient. N. Am.
<i>Ipomoea emetica</i> Choisy	G.	Em.	Mex.
<i>Ipomoea Jalapa</i> Cox	G.	Em.	Mex.
<i>Ipomoea pandurata</i> G. F. W. Meyer	Rusby	Pur.	S. E. U. S.
<i>Ipomoea Purga</i> Hayne	Sm.	Pur.	Mex.
<i>Ipomoea sinuata</i> Ortega	G.	Em.	Am. Calif.
<i>Ipomoea tuberosa</i> L.	G.		Trop. Reg.

CORIACEAE

<i>Coriaria myrtifolia</i> L.	G. C. L.		Med. Reg.
<i>Coriaria nepalensis</i> Wall.	G.		Him. China
<i>Coriaria ruscifolia</i> L.	G.		Peru
<i>Coriaria sarmentosa</i> Frost	G. L.		N. Zeal.
<i>Coriaria thymifolia</i> Humb. & Bonpl.	G.		Peru

CORNACEAE

<i>Alangium Lamarckii</i> Thwait	G.		Cosm. Trop.
<i>Amyris balsamifera</i> L.	G.	Ir	Jamaica
<i>Aucuba japonica</i> Hamb.	Sm.	Exp.	Japan
<i>Cornus Amomum</i> Mill.	G. L.	Exp.	E. N. Am.
<i>Cornus circinata</i> L. Her.	L.	Astr.	N. A.
<i>Cornus Mas</i> L.	L.	Astr.	N. A.
<i>Cornus paniculata</i> L. Her.	Panamel	Exp.	E. N. Am.
<i>Garrya Fremontii</i> Torr.	G. L.		W. N. Am.
<i>Maieria Vitiensis</i> Berth.	G.		Austr. Pac. Is.

CRASSULACEAE

NAME OF PLANT	Authority	Properties	Locality
<i>Cotyledon ventricosa</i> Burm.	G.		Afr Austr
<i>Kalanchoe spathulata</i> DC.	G.		
<i>Sedum acre</i> L.	G. Mi. L.	Ir	Trop. As.
<i>Sempervivum montanum</i> L.	G.	Ir	Eur N As N Am Cent Eur

BITTER

<i>Barbarea vulgaris</i> R. Br.	G. L.	Antisc	Eur Temp. As. N Am
<i>Brassica alba</i> Boiss.	Sm.	Ir	Eu As N An
<i>Brassica arvensis</i> L.	C. L.	Ir Em.	Eu As N As
<i>Brassica nigra</i> Koch.	C. L.	Ir Em.	Eu As N As
<i>Capucina Bursa-pastoris</i> Medic	G.	Sap	Temp. Reg
<i>Cardamine amara</i> L.	G.		Eur N As.
<i>Cardamine pratensis</i> L.	G. L.	Antisc	Temp. Reg
<i>Crambe filiformis</i> Jacq.	G.		Pat Ch.
<i>Diploxys erucoides</i> DC.	G. L.	Ir	Med. Reg As. Minor
<i>Diploxys tenuifolia</i> DC.	L.	Ir	Eur
* <i>Eruca sativa</i> Mill.	Wilcox*		Med. Reg W As N Am
<i>Erysimum cheiranthoides</i> L.	Sm. L.	Ir Ant	Eur
<i>Erysimum crepatifolium</i> Reischb.	G.		Eur
<i>Isatis corniculata</i>	G.		
<i>Lepidium Draba</i> L.	G. R.	F P	Eur Orient.
<i>Lepidium latifolium</i> L.	Sm.	Ir	Eur As Orient
<i>Lepidium ceraceum</i> Forst.	G. L.	F P	N Zeal
<i>Lepidium Oahuense</i> Cham. & Schlecht.	G.	F P	Hawai. Is.
<i>Lepidium pycnanthum</i> Forst.	G. R.	F P	Pac Is.
<i>Lepidium sativum</i> L.	G. Wh. L.	Ir	Orient.
<i>Matthiola livida</i> DC.	G.		Egypt, Arab.
<i>Ranunculus Armoracia</i> (L.) Robinson.	Antisc	Ir	Eur As N Am
<i>Raphanus Raphanistrum</i> L.	C. L.	Ir	Eur
<i>Raphanus sativus</i> L.	L.	Antisc	Asia
<i>Senebiera pinnatifida</i> DC.	G.		Cosmop
<i>Sisymbrium Althaea</i> Scop.	C.		Eur Orient
<i>Sisymbrium Sophia</i> L.	G.		Him. Reg.
<i>Sisymbrium officinale</i> (L.) Scop.	L.	Antisc	Temp. Reg.
<i>Sisymbrium toxophyllum</i> G. A. Mey.	G.		Eur N As.
<i>Vesicaria gnaphalodes</i> B. Iss.	G.		N As Persia

CUCURBITACEAE

<i>Bryonia alba</i> L.	Mi. C. L.	Ir	
<i>Bryonia dioica</i> Jacq.	G. Mi. C. L.	Ir	Eur Med Reg
<i>Cucurbita ficifolia</i> Cogn.	G.		Brazi.
<i>Citrullus Colocynthis</i> Schrad.	G. C.	Pur	Afr Cal'd
<i>Cucurbitaria epigaea</i> B. & H.	L.	Ant.	E. Ind
<i>Cucumis africanus</i> Lindl.	G.	Em.	S. Afr
<i>Cucumis dipsaceus</i> Ehrenb.	G.		Arab. Trop As.
<i>Cucumis metuliferus</i> F. Mey.	G.		S. Afr
<i>Cucumis myriocarpus</i> Naud.	L.	Em.	S. Afr
<i>Cucumis prophetarum</i> L.	L.	Pur	Arabia

*A case of probable poisoning of sheep by this weed was reported to Dr. Wilcox from Nebraska.

NAME OF PLANT	Authority	Properties	Locality
<i>Cucumis Sacleuxii</i> Hort.	G.	Sap.	
<i>Cucumis trigonus</i> Roxb.	G. L.	Ant	Ind
<i>Cucurbita maxima</i> Duchesne	G.	Sap.	
<i>Echadum eateium</i> A. Rich.	C. Sm. L.	Pur	Med Reg
<i>Echinocystis fabacea</i> Naud.	G. L.	Sap.	Calif
<i>Echinocystis macrocarpa</i> Greene	G. L.	Sap.	Calif
<i>Fevillea cordifolia</i> L.	G. L.	Em	Ind
<i>Lagenaria vulgaris</i> Ser	G. L.	Sap.	Trop. Reg
<i>Luffa acutangula</i> Roxb.	G.		Trop. old world
<i>Luffa aegyptiaca</i> Mill.	G.		Trop. old world
<i>Luffa purgans</i> Mart.	L.	Cath.	Tr Am
<i>Melothra acrobiculata</i> Cogn.	G.	Taen	Abyss
<i>Momordica Balsamina</i> L.	L.	Em.	Trop. Reg.
<i>Momordica Charantia</i> L.	L.	Ant Em.	Trop. Reg
<i>Momordica cochinchinensis</i> Spreng.	G.		Trop. As
<i>Momordica Cymbalaria</i> Fenzl.	G.		Trop. Afr
<i>Sicydium monospermum</i> Cogn.	G.	Em	Brazil
<i>Telfairia pedata</i> Hook.	G.	Taen	Trop. Afr
<i>Trichosanthes Celebica</i> Cogn.	G.		Celebes
<i>Trichosanthes globosa</i> Bl.	G.	F P	Java
<i>Trichosanthes pubera</i> Blom.	G.		Ind, Malay
<i>Trichosanthes trifoliata</i> Bl.	G.		Malay
<i>Trichospathes walachiana</i> Cogn.	G.		Ind

CYCADACEAE

<i>Catkins</i> sp.	G.		Austr
<i>Cycas circinalis</i> L.	G.		Molucc, I
<i>Cycas media</i> R. Br.	G. M.		Austr
<i>Macrozamia</i> sp.	G. M.		Austr
<i>Zamia nuda</i> Jacq.	G.		W I

CYPERACEAE

<i>Cyperus articulatus</i> L.	G. L.	Ant	Common Trop.
<i>Kyllingia monocephala</i> Rottb.	G.		Trop. old world
<i>Schoenus apogon</i> Roem. and Schult.	M.		Austr N Zest

CHAILLETIACEAE

<i>Chaletia cymosa</i> Hook.	G.		S. Afr
<i>Chaletia toxicaria</i> Don.	G. R.	F P	Trop. Afr
<i>Tapara gwanensis</i> Aubl.	G. R.	F P	Guay

DILLENIACEAE

<i>Davilla rugosa</i> Poir.	G.		Brazil
<i>Dohocarpus Roandri</i> J. F. Gmelin.	G.		Guiana, Brazil
<i>Hibbertia glaberrima</i> F. Muell.	G. M.		Austr
<i>Hibbertia longifolia</i> F. Muell.	G.		Austr
<i>Tetracera alufolia</i> Willd.	G. R.	F P	Tr Afr
<i>Tetracera Asa</i> DC.	E.	F P	Ind Malay

DIOSCOREACEAE

<i>Dioscorea</i> sp.	G.		
<i>Dioscorea alata</i> L.	G.		Tr. As.
<i>Dioscorea bulbifera</i> L.	G.		Tr. As.
<i>Dioscorea deltoidea</i> Wall.	G.		Ind

NAME OF PLANT	Authority	Properties	Locality
<i>Dioscorea hirsuta</i> Mart.	G.		Mex
<i>Dioscorea macrocarpa</i> Harm.	G.		
<i>Dioscorea toxicaria</i> Bojer	G.		Tr Afr
<i>Dioscorea triphylla</i> Schump.	G.		Abyss.
<i>Tamus communis</i> L.	G. C. L.	Ir Diur	Eur Persia Afr

DIPSACEAE

<i>Scabiosa Succisa</i> L.	Sm. L.		Eur Tr N A & S. A
----------------------------	--------	--	----------------------

DIPTEROCARPACEAE

<i>Anastrocladus</i> Vahlb. Arn.	G.		Zeylan.
<i>Dryobalanops can phora</i>	Sm. L.	Del.	Zeylan.

DROSERACEAE

<i>Dionaea muscipula</i> Ellis.	G. Sm. L.	Ir Hyd	N Am
<i>Drosera binata</i> Labil.	G.	Hyd.	Austr. N Zeal.
<i>Drosera communis</i> S. Hil.	G.		Brazil
<i>Drosera intermedia</i> Hayne.	G.	Hyd.	Eur
<i>Drosera peltata</i> Sw.	G. M.		Austr.
<i>Drosera rotundifolia</i> L.	G. Sm. L.	Ir Hyd	N Reg
<i>Drosera Whitakeri</i> Planch.	M.		Austr.
<i>Drosophyllum lusitanicum</i> Link.	G.	Hyd.	Spain Morocco.

EUPHYORBIACEAE

<i>Diospyros acris</i> Hemsl.	G.	Acr	Solomon Is.
<i>Diospyros amara</i>	G.		
<i>Diospyros Ebenum</i> Koen.	G. L. R.	Astr. F. P.	Ind. Malay
<i>Diospyros embryopteris</i> Pers.	L.	Astr.	Tr Afr Philipp.
<i>Diospyros malacarpa</i> A. DC.	G.		Trop. Afr.
<i>Diospyros montana</i> Roxb.	G.		Philipp.
<i>Diospyros multiflora</i> Blanco.	G.	F. P.	N. A.
<i>Diospyros obtusifolia</i> W. L. D.	L.	Astr.	
<i>Diospyros samoensis</i> A. Gr.	G.	F. P.	Samoa Is. Nav.
<i>Diospyros tessellata</i> Poir.	G.	F. P.	Mascari Is.
<i>Diospyros toxicaria</i> Hiern.	G.		Madagascar
<i>Diospyros virginiana</i> L.	G. L.	Ant. Astr.	N. A.

EQUISETACEAE

<i>Equisetum arvense</i> L.	G. M. L.	Diur	Eur N A
	Ches.		
<i>Equisetum hyemale</i> L.	L.	Diur	Eur As. N A
<i>Equisetum palustre</i> L.	Ches.	Diur	Eur

ERICACEAE

<i>Andromeda Polifolia</i> L.	Ches.	Nar	E. N. Am. Eur.
<i>Arbutus Andrachne</i> L.	G.	Nar	As.
<i>Arbutus Unedo</i> L.	G. L.		Med. Reg.
<i>Arbutus variegatus</i> Benth.	G.		Med. Reg. Mex

NAME OF PLANT	Authority	Properties	Locality
<i>Arbutus xalapensis</i> H. B. K.	L.	Diur Nar	Mex. S W U S.
<i>Arctostaphylos polifolia</i> H. B. K.	G.	Nar	Mex.
<i>Arctostaphylos Uva-ursi</i> L.	Sm. L.	Ir Diur	Eur N A.
<i>Epigaea repens</i> L.	Ches.	Nar	E. N Am.
<i>Gaultheria microphylla</i> Hook.	G.		Magellan Reg.
<i>Gaultheria nummularoides</i> D. Don.	G.		Him Reg. Java.
<i>Gaultheria procumbens</i> .	G. L.	Astr	E. N A.
<i>Kalmia angustifolia</i> L.	G. Rusby	Nar Astr	E. N A.
	L. Ches.		
<i>Kalmia glauca</i> A. r.	L.	Nar	E. N A.
<i>Kalmia hirsuta</i> Walt.	L.	Nar	E. N A.
<i>Kalmia latifolia</i> L.	G. Ches.	Nar Astr	E. N. A.
	L. Rusby		
<i>Ledum glandulosum</i> Nutt.	L. Ches.	Nar	N and Arc. Reg.
<i>Ledum latifolium</i> Jacq.	C. Ches.	Nar	N. N A.
	L. Rusby		
<i>Ledum palustre</i> L.	G. C.	Nar	N & Arc. Reg.
	Rusby L.		
<i>Leucothoe Catesbaei</i> A. Gr.	C. L. Ches.	Errhine	N. A. S. States
<i>Leucothoe racemosa</i> A. Gray.	Ches.	Errhine	
<i>Oxydendron arboreum</i> DC.	Whi. L.	Ir	N. A.
<i>Pieris floribunda</i> Benth. & Hook.	Ches.	Inch.	N A.
<i>Pieris mariana</i> Benth. & Hook.	G. Rusby	Inch.	E. N. A.
	L. Ches.		
<i>Pieris nitida</i> Benth. & Hook.	G. L.	Inch.	E. N A.
<i>Pieris ovatifolia</i> D. Don.	G.		Him Reg. Java.
<i>Rhododendron arborescens</i> Torr.	C.	Nar	N A.
<i>Rhododendron arboreum</i> Sm.	Sm. G.	Nar	Ind Him. Reg.
<i>Rhododendron californicum</i> Hook.	Ches.	Nar	N A Calif.
<i>Rhododendron campanulatum</i> D. Don.	G.	Nar	Him Reg.
<i>Rhododendron catawbiense</i> Mx.	L.	Nar	E. N A.
<i>Rhododendron caucasicum</i> Fall.	G.	Nar	Caucas.
<i>Rhododendron chrysanthum</i> Fall.	G. Sm. L.	Nar	Davur.
<i>Rhododendron cuneatum</i> Hook.	G.	Nar	Him. Reg.
<i>Rhododendron dauricum</i> L.	G. R.	Nar	Eur As.
<i>Rhododendron ferrugineum</i> L.	G. C. Sm. L.	Nar	Eur As.
<i>Rhododendron hirsutum</i> L.	G. C.	Nar	Eur.
<i>Rhododendron indicum</i> Sw.	C.	Nar	China.
<i>Rhododendron ledifolium</i> G. Don.	C.	Nar	China.
<i>Rhododendron maximum</i> L.	G. Ches. C.	Nar	E. N A.
	Rusby L.		
<i>Rhododendron nudiflorum</i> Torr.	C.	Nar	N A.
<i>Rhododendron occidentale</i> A. Gray.	Ches.	Nar	W N A.
<i>Rhododendron ponticum</i> L.	G. C. Sm.	Nar	A M.
<i>Rhododendron punctatum</i> Andr.	C.		N A.
<i>Rhododendron sinense</i> Sw.	C.	Nar	Japan.
<i>Rhododendron subanceolatum</i> Mx.	C.	Nar	Japan.
<i>Zenobia speciosa</i> D. Don.	G.		N A.

ERYTHROXYLACEAE

<i>Erythroxylon coca</i> Lam.	Sm. L. G.	Del Stim	Peru
		Anes.	

EUPHORBIACEAE

<i>Acalypha colchica</i> Fisch. & Mey.	G.	Hyd.	Caucas.
<i>Acalypha cordifolia</i> Mueßl.	G. Wat.		Him. Reg.
<i>Acalypha indica</i> L.	G.	Em.	Tr As & Afr.
<i>Acalypha virginica</i> L.	L.	Exp. Diur Ir	N. Am.

NAME OF PLANT	Authority	Properties	Locality
<i>Ecdospermum axillare</i> Bl.	G.		Ind. Malay
<i>Eyeria viscosa</i> Miq.	G. M.		Aus.
<i>Bridelia retusa</i> Spreng.	G.		Ind. Malay
<i>Chrozophora plicata</i> A. Juss.	G.		Eur. Afr. As. Ind.
<i>Chrozophora tinctoria</i> A. Juss.	G.		Eur. Ori. Ind.
<i>Cleistanthus colinus</i> Benth. & Hook.	G.		Afr.
<i>Cnesmone javanica</i> Bl.	G.		Ind.
<i>Croton muricatus</i> Zipp.	G.		Java
<i>Croton Tiglium</i> L.	G. C. Sm. R.	Pur F P	Timor Is.
	L.		Ind. Malay
<i>Croton Verreauxii</i> Baill.	G.		Austr.
<i>Daphniphyllum bancanum</i> Kurz	G.		Java
<i>Elaeococca verrucosa</i>	C.		
<i>Eremocarpus setigerus</i> Benth.	L.	A. P. F. P.	Calif.
<i>Euphorbia aleppica</i> L.	R.	Ir. F. P.	
<i>Euphorbia alsiniferia</i> Baill.	G. M.	Ir.	Austr.
<i>Euphorbia amygdaloides</i> L.	G. R.	Ir. F. P.	Eur. Orient.
<i>Euphorbia antiquorum</i> L.	G.	Ir.	Ind.
<i>Euphorbia arborescens</i>		Ir.	
<i>Euphorbia bicolor</i> Engelm. & Gray.	Ches.	Ir.	N. Am.
<i>Euphorbia candelabrum</i> Trem.		Ir.	
<i>Euphorbia caracasana</i>	R. G.	Ir. F. P.	
<i>Euphorbia cerebrum</i> L.	G.	Ir.	
<i>Euphorbia chamalsyce</i> L.	G.	Ir.	
<i>Euphorbia characias</i> L.	G.	Ir.	Med. Reg.
<i>Euphorbia corollata</i> L.	L.	Ir.	N. A.
<i>Euphorbia cotinoides</i> Miq.	G. R.	Ir. F. P.	Guiana
<i>Euphorbia cotinifolia</i> L.	G. E. R.	Ir. F. P.	Ind.
<i>Euphorbia cyparissias</i> L.	Mi. C.	Ir.	Eur. N. Am.
<i>Euphorbia dendroidea</i> L.	G. R.	Ir. F. P.	Med. Reg.
<i>Euphorbia Drummondii</i> Boiss.	G. M.	Ir.	Austr.
<i>Euphorbia eremophila</i> A. Cunn.	Mi.	Ir.	Austr.
<i>Euphorbia Esula</i> L.	R. Mi. E.	Ir. F. P.	Eur.
<i>Euphorbia Gerardiana</i> Jacq.	Mi. C.	Ir.	Eur.
<i>Euphorbia helioscopia</i> L.	Mi. C.	Ir.	Eur. and N. As.
<i>Euphorbia heptagona</i> L.	G.	Ir.	S. Afr.
<i>Euphorbia heterodoxa</i> Muell.	L.	Afr.	Brazil
<i>Euphorbia hyberna</i> L.	G. R.	Ir. F. P.	Eur.
<i>Euphorbia Ipecacuanhae</i> L.	M. Rusby	Ir.	N. A.
	L.		
<i>Euphorbia Lathyrus</i> L.	C. R.	Ir. F. P.	Eur. N. Am.
	Ches. L.	Ir.	
<i>Euphorbia linearis</i> Retz.	G.	Ir.	W. I.
<i>Euphorbia lingulata</i>	G.	Ir.	
<i>Euphorbia marginata</i> Pursh.	G. Ches.	Ir.	N. Am.
<i>Euphorbia mellifera</i> Ant.	G.	Ir.	Madeira Is.
<i>Euphorbia nerifolia</i> L.	G. R.	Ir. F. P.	Ind. Malay
<i>Euphorbia officinarum</i>	Sm.	Ir.	N. Afr.
<i>Euphorbia palustris</i> L.	G. Mi. C.	Ir.	Eur.
<i>Euphorbia Paralias</i> L.	Mi.	Ir.	Med. Reg.
<i>Euphorbia Peplus</i> L.	Mi. C. Sm.	Ir.	Eur. N. As.
<i>Euphorbia pascuosa</i> Ant.	G. E. R.	Ir. F. P.	Mader Is.
<i>Euphorbia pithyusa</i> L.	Mi.	Ir.	Med. Reg.
<i>Euphorbia platyphyllos</i> L.	G.	Ir.	Eur. Afr.
<i>Euphorbia Preslii</i> Guss.	Sm.	Ir.	N. Am.
<i>Euphorbia prunaeifolia</i> Baker	G.	Ir.	Madagas.
<i>Euphorbia pulcherrima</i> Willd.	G.	Ir.	Mex.
<i>Euphorbia punicea</i> Sw.	G. E.	Ir. F. P.	Jamaica
<i>Euphorbia pungiformis</i> Boiss.	G.	Ir.	

NAME OF PLANT	Authority	Properties	Locality
<i>Euphorbia Regis-Jubae</i> Webb. & Benth.	G.	Ir	Teneriff Is
<i>Euphorbia resinifera</i> Berg.	L.	Ir	Morocc.
<i>Euphorbia Royleana</i> Boiss.	G. R.	Ir F P	Him. Reg.
<i>Euphorbia Subthorpa</i> Boiss.	G. R.	Ir F P	Greece
<i>Euphorbia Sieboldiana</i> Moor & Decne.	G.	Ir	Japan
<i>Euphorbia hymenifolia</i> L.	G.	Ir	Tropics
<i>Euphorbia Tirucalli</i> L.	G. R.	Ir F P	Afr Ind.
<i>Euphorbia venetica</i> Trem.	G.	Ir	Tr Afr
<i>Euphorbia verrucosa</i> Lam.	M. C.	Ir	Eur
<i>Euphorbia Wulfenii</i> Hoppe.	G.	Ir	Dalmatia
<i>Excoecaria Agallocha</i> L.	G. R. M. C.	Acr F P	Tr As. Malay
	L.		Pac. Is.
<i>Excoecaria virgata</i> Zoll and Mor	G.		Java
<i>Fluggea Leucopyrus</i> Willd.	G. R. E.	F P	Ind.
<i>Furges obovata</i> Baill.	G.		Tr Afr
<i>Gymnanthes lucida</i> Sw.	Sm.		W. Ind.
<i>Hippomane Mananilla</i>	L. G. Sm.	Ir	Ind.
<i>Homalanthus crepitans</i> L.	E. & G. C. R.	F P	
<i>Honulanthus Leschenaultianus</i> A. Juss.	G. M.		Ind Malay
			Austr.
<i>Hyoemanche globosa</i>	C.		
<i>Jatropha Curcas</i> L.	G. C. R. L.	F P	Tr Amphig.
<i>Jatropha glandulifera</i> Roxb.	G.		Tr As & Afr
<i>Jatropha macrocarpa</i> Benth.	G.		Mex
<i>Jatropha multifida</i> L.	G.	Ir	Tr Amphig.
<i>Jatropha urens</i> L.	G. Sm.	Ir	Tr Am.
	Ches.		
<i>Joannea princeps</i> Vell.	G. L. E.	Cath F P	Braz
<i>Macaranga</i> sp.	G.		Java
<i>Madotus philippinensis</i> Muell.	Sm. L.	Pur	As. Amr
<i>Maritot Glax</i> sp. Muell.	G.		
<i>Maritot utusuma</i> Pohl.	R. G.	F P Hyd.	Braz.
	Wh. Sm.		
	C. L.		
<i>Mecurialis annua</i>	C. L.	Cath. Ir	Eur
<i>Mecurialis perennis</i> L.	G. C. L.	Ir, Cath.	Eur
<i>Mecurialis tomentosa</i>	C.		
<i>Pedanthus ethymaloides</i> Pont.	G. L.	Ir Em	Mex S. Am.
<i>Petalostigma cordifolia</i> F. M.	G.		
<i>Phyllanthus Conami</i> Sw.	F. G.	F P	Tr Am.
	Sm. M.		
<i>Phyllanthus Emblica</i> L.	G.		Tr As
<i>Phyllanthus epiphyllanthus</i> L.	G.		Cuba
<i>Phyllanthus Gasstrom</i> Muell.	G. M.		Austr
<i>Phyllanthus lacunarius</i> F. M.	G. M.		Austr
<i>Phyllanthus Niruri</i> L.	G.		Tr Reg
<i>Phyllanthus placitum</i> H. B. & K.	G. M. E.		Venezuela
<i>Phyllanthus urinaria</i> L.	G.	F P	Tr Amphig.
<i>Pteranthea trifoliata</i> Baill.	G. M.		Braz.
<i>Platygyne urens</i> Mercier		F P	
<i>Putranjiva Roxburghii</i> Wall.	G.		Ind Burma
<i>Ricinus communis</i> L.	G. Sm. C. L.	F P	Tr Reg.
	Ches.	Pur.	
<i>Sapium indicum</i> Willd.	G. M. E.	F P Nar	Ind.
<i>Sapium insigne</i> Trim.	G.	Ir	Ind
<i>Sebastiania Palmeri</i> Riley.	G.		
<i>Sebastiania Pavoniana</i> Muell. Arg.	G.		
<i>Securinega ramiflora</i> Muell. Arg.	G.	Hyd.	As.
<i>Stillingia oreata</i> Muell.	G.		Ind. Maurt

NAME OF PLANT	Authority	Properties	Locality
<i>Stillingia sylvatica</i> L.	L.	Expect.	U. S.
<i>Toxicodendrum globosum</i>	G.	Con.	
<i>Tragia cordifolia</i> Vahl.	G.	Ves.	Arabia
<i>Tragia involucrata</i> L.	G.	Ves.	Tr. A.
<i>Tragia volubilis</i> L.	G. L.	Ves.	Ind.

FAGACEAE

<i>Castanopsis Tungurrut</i> A. D. C.			Malay
<i>Fagus sylvatica</i> L.	C. M.	Gasir	Eur
<i>Quercus alba</i> L.	L.	Astr. Tymp.	E. N. A.
<i>Quercus Toza</i> Gillet	C.		S. Eur
<i>Quercus lusitanicus</i> Lam.	L.	Astr.	Med. Reg.
<i>Quercus</i> sp.	Ches.		

FICOIDEAE

<i>Aizoon canariense</i> L.	G.		Canar. Is. Afr.
<i>Gueckia pharnacioides</i> L.	G.	Taen.	Orient.
<i>Linum</i> sp.	G.		Taen. Ind.
<i>Mesembryanthemum anatomicum</i> Haw.	G.	Nar.	S. Afr.
<i>Mesembryanthemum crystallinum</i> L.	L.	Idur.	Tr. Reg. & Sub.
<i>Molugo hirta</i> Thunb.	G. M.		Tr. Reg. Afr.
<i>Tetragonia expansa</i> Thunb.	G.	Sap.	Gr. Can. Is. S.
<i>Trianthema</i> sp.	G.	Sap.	Afr. Calif.
<i>Trianthema Portulacastrum</i> L.	G.	Sap. Par.	Austr.
			Japan
			Arab. Afr.
			Austr.
			Tr. An.

FILICALES

<i>Adiantum petatum</i> L.	G.	Astr.	E. As. Jap.
<i>Adiantum peruvianum</i> Kl.	G.		N. A.
<i>Adiantum trapeziforme</i> L.	G.		Peru
<i>Aneides oblongifolia</i> Sw.	G.		Trop. Am.
<i>Angiopteris erecta</i> Hoffm.	G.		Trop. Am.
<i>Aspidium athamanticum</i> Ktze.	G. L.	Ant.	S. Afr.
<i>Aspidium Filix-Mas</i> Sw.	G. L.	Ant.	Eur. N. A.
<i>Aspidium marginale</i> Ktze.	G. L.	Ant.	Eur. N. A.
<i>Aspidium odoratum</i> Willd.	G.		N. Ind. E. As.
<i>Aspidium prestulatum</i> Ten.	G.	An.	
<i>Aspidium rigidum</i> Sw.	G.		
<i>Ceterach officinarum</i> Willd.	G. L.	Abort.	Eur. Ind.
<i>Cheilanthes fragrans</i>	G.		Med. Reg.
			Him. Reg.
<i>Cryptogramme crispa</i> R. Br.	G.		N. Eur. N. W.
<i>Cystopteris alpina</i> Desv.	G.	Hyd.	Am.
<i>Cystopteris bulbifera</i> Bernh.	G.	Hyd.	Eur. Mt. Reg.
<i>Cystopteris fragilis</i> Bernh.	G. L.	Hyd.	N. Am.
			Eur. N. Am.
			Cosm.
<i>Cystopteris montana</i> Bernh.		Hyd.	Cent. Eur. E.
<i>Davallia brassioides</i> Hook.	G.	Hyd.	Malay, Trop.
<i>Davallia elegans</i> Sw.	G.	Hyd.	As. Austr.
			N. A.
			S. A.

NAME OF PLANT	Authority	Properties	Locality
<i>Davallia hirsuta</i> Karf	G.	Hyd.	Japan
<i>Davallia majuscula</i> Lowe	G.	Hyd.	Ceylon
<i>Davallia pentaphylla</i> Don.	G.	Hyd.	Java & ad. Is.
<i>Davallia platyphylla</i> Don.		Hyd.	Ind
<i>Davallia strigosa</i> Sw	G.		N Ind. Japan, Malay
<i>Geiophlebium incarum</i> Sw	G.		
<i>Geichenia flabellata</i> R. Br	G.	Sap	
<i>Gymnogramme cordata</i> Schl.	G.		S. Afr
<i>Lindsaea cultrata</i> Bl	G.		N Ind. Malay
<i>Polypodium laciniatum</i> Bl.	G.		Mad. Austr
<i>Polypodium ligularum</i> Sw	G.		Java
<i>Polypodium perispermum</i> Cav	G.		S. A
<i>Polypodium Phymatodes</i> L.	G.		Malay Ceylon, Austr
<i>Polypodium scandens</i> Forst	G.		N. Z. Austr
<i>Polypodium suspensum</i> L.	G.		Trop. Am
<i>Polypodium vulgare</i> L.	L.	Exp. Dur	Eu N Am
<i>Polystichum spirulosum</i> DC.	G.		N Am
<i>Pteris aquilina</i> L.	G.	Hyd. Ant	Coc Eu N Am
<i>Pteris caudata</i> L.	G.	Ant	

PLAGELLARIACEAE

<i>Sakum artachanthum</i> Bl.	G.		Java
-------------------------------	----	--	------

FRANKENIACEAE

<i>Frankenia ericifolia</i> Chr	Sm. G	(P. P)	Canary Is
<i>Frankenia granatensis</i> Cham. & Schlecht.	L.	(Astr)	Calif

FUMARIACEAE

<i>Corydalis racemosa</i> Pers	G.		Japan
<i>Dicentra canadensis</i> Walp.	L.	Dur	N. A
<i>Fumaria officinalis</i> L.	G. L.	Nar	Temp. old world
<i>Fumaria spicata</i> L.	G.		Laotan.

GENTIANACEAE

<i>Gentiana verna</i> L.	G.		Eur. Cauc. Reg
<i>Menyanthes trifoliata</i> L.	G.		N. Temp. Reg
<i>Tachia guianensis</i> Aubl.	G.		Guian

GERANIACEAE

<i>Erodium cicutarium</i>			Eu. N. Am. S. Am.
<i>Pelargonium peltatum</i> Ant.	G.		S. Afr

GNETACEAE

<i>Ephedra</i> L.	G.		Eur
<i>Ephedra antisiphilitica</i> C. A. Mey	L.	(Astr)	N. A. Mex
<i>Gnetum scandens</i> Roxb.	G.	F. P	Ind. Malay
<i>Gnetum urens</i> Blum.	G.	(U'r)	Guian

GOODENIACEAE

<i>Goodenia grandiflora</i> Sims.	G.		Austr
<i>Scaevola</i> sp.	G.	A. P	Austr
<i>Velleia paradoxa</i> R. Br	G. M		Austr

GRAMINEÆ

NAME OF PLANT	Authority	Properties	Locality
<i>Agrostis alba</i> L.	*Hell.	Ur	N Temp. Reg.
<i>Andropogon annulatus</i> Forsk.	G.		Afr Ind. Austr
<i>Andropogon halepensis</i>		Hyd.	Eur Afr N Am.
<i>Anthisturia prostrata</i> Willd.			Ind.
<i>Anthoxanthum odoratum</i> L.	G.	Con.	Eur As Afr
<i>Arrhenatherum avenaceum</i> Beauv	G.	Sap.	Eur
<i>Avena fatua</i> L.	G.	Mech.	Eur Ori. As.
<i>Briza</i> sp.	G.	Hyd.	S. Am. Eur. N. Am. As.
<i>Bromus catharticus</i> Vahl.	G.		
<i>Bromus molis</i> L.	G.		Eur Afr As.
<i>Bromus secalinus</i> L.	G.		Eur Med Reg. As
<i>Bromus tectorum</i> L.	Pa.	Ir	Eur. N Am.
<i>Catabrosa</i> sp.	G.	Hyd.	
<i>Cenchrus tribuloides</i> L.	G.	Mech.	N Am.
<i>Cortaderia argentea</i> Stopf.	G.		
<i>Cortaderia conspicua</i>	G.	Hyd.	
<i>Cortaderia Kermesina</i>	G.	Hyd.	
<i>Deyeuxia Langsdorffii</i> Kunth.	G.	Sap.	N Am
<i>Ehlorurus</i> sp.	G.		S. Afr. S. Am. Austr
<i>Elymus</i> sp.	G.	Hyd.	Eur N. Am
<i>Festuca</i> sp.	G.	Hyd.	
<i>Festuca quadridentata</i> H. B. & K.	G. C.		Equador
<i>Glyceria</i> sp.	G.	Hyd.	N Am Eur
<i>Glyceria aquatica</i> Wahlb.	G.	Hyd.	N Temp. Reg.
<i>Heteropogon hirtus</i> Pers.			Tr Reg.
<i>Hierochloa australis</i> R. & S.	G.	Con.	Eur
<i>Hierochloa borealis</i> Roem. & Schult.		Con.	N Reg.
<i>Hierochloa ramosa</i> Hook.	G.	Con.	Eur
<i>Holcus</i> sp.		Hyd.	Eur N Am
<i>Lolium Lamarckii</i> L.	G. C.	Hyd. Del.	Eur As
<i>Lolium perenne</i> L.	G. C. Mi.	Hyd.	Eur As.
<i>Lolium temulentum</i> L.	L. Ches.	Nar Intox	Eur As U S.
<i>Melica</i> sp.	G.	Hyd.	
<i>Molinia effusum</i> L.	G.		N Temp. Reg.
<i>Molinia caerulea</i> Moench.	G. C.		Eur As
<i>Panicum</i>	G.	Hyd.	N Temp. Reg.
<i>Panicum juncceum</i> Nees	G.	Hyd.	Braz.
<i>Panicum sanguinale</i> L.	G.	Em	Cosmop
<i>Paspalum scrobiculatum</i> L.	G.	Nar	Tr. and world
<i>Setaria scandens</i> Schrad.	G.		Braz
<i>Sorghum vulgare</i> Pres.	G. C.	Hyd.	Tr Reg.
<i>Spartium juncceum</i>	L.	Em. Cath	Cosmop
<i>Stipa capillata</i> L.	G.		S Am As
<i>Stipa inebrians</i> Hance	G.	Del	China
<i>Stipa palmata</i> Pall.			
<i>Stipa pennata</i> Hohen.	G.	Mech.	Eur As N Am.
<i>Stipa robusta</i> Nash	Ches.	Meb.	N Am. Ry Mts.
<i>Triodia arvensis</i> Brown	G.	Mech.	Austr
<i>Zea Mays</i> L.	G. C. L.	Hyd.	Paraguay

* A case of poisoning by *Agrostis alba* was reported from West Liberty, Ia. by Dr. Hell. Probably forage poisoning.

RUPTAGINACEAE

NAME OF PLANT	Authority	Properties	Locality
<i>Calophyllum Calaba</i> Jacq.	G.	Sap.	Caith. Is.
<i>Calophyllum Inophyllum</i> L.	G.	F P Sap.	Tr. old world
<i>Calophyllum montanum</i> Viekl.	G.		N. Calcd.
<i>Clusia macrocarpa</i> Spreng.	G.		Guiana
<i>Garcinia Cambogia</i> Desr.	Sm.		Ind.
<i>Garcinia Forsteriana</i> Bl.	G.		
<i>Mammea americana</i> L.	G. L.	Ant.	Tr. Am.
<i>Stalagmitis Mangle</i>	G.		

HAEMODORACEAE

<i>Aletris farinosa</i> L.	G.		N. Am.
<i>Gyrotheca major</i> Morong.	Ches.		E. N. Am.
<i>Haemodorum</i> sp.	G.		Austr. Braz.
<i>Ophiopogon</i> sp.	G.		Japan, Ind.
<i>Sansevieria thyrsiflora</i> Thumb.	G.		S. Afr.

HIPPOCASTANACEAE

<i>Aesculus californica</i> Nutt.	Ches.	Sap.	W. N. Am. Calif.
<i>Aesculus flava</i> Ast.	G. R.	Sap. F P	N. Am.
<i>Aesculus glabra</i>	Sm. Ches.	Sap.	N. Am.
<i>Aesculus Hippocastanum</i> L.	G. Ches. L.	Sap.	Turkey
<i>Aesculus Pavia</i> L.	G. R.	Sap. F P	N. Am.
	Ches. L.		

HYDROPHYLLACEAE

<i>Eriodictyon glutinosum</i> Benth.	G. L.	Exp.	Calif.
--------------------------------------	-------	------	--------

HYPERICINEAE

<i>Hypericum crispum</i> L.	G.		Med. Reg.
<i>Hypericum humifusum</i> L.	G.		Eur. As.
<i>Hypericum maculatum</i> Walt.	Ches.	Acr.	E. N. Am.
<i>Hypericum perforatum</i> L.	Ches.	Acr.	Eur. N. Am.

ILICINEAE

<i>Nex Aquifolium</i> L.	G. L.	Em. Cath.	Eur. As.
--------------------------	-------	-----------	----------

ILLECEBRACEAE

<i>Paronychia bonariensis</i> DC.	G.		Argent. Reg.
<i>Paronychia capitata</i> Lam.	G.	Sap.	Eur. Orient.

IRIDACEAE

<i>Belamcanda punctata</i> Moench.	G.		China
<i>Crocus sativus</i> L.	Sm. L.	Ir.	As. Min.
<i>Gladiolus communis</i> L.	C.	Ir.	Eur. Orient.
<i>Gladiolus segetum</i>	C.	Ir.	Med. Reg.
<i>Hemeris collina</i> Vent.	G. M.		S. Afr.
<i>Iris</i>	G.	Ir.	N. Am.
<i>Iris florentina</i> Sair.	Sm. C. L.	Ir.	S. Eur.
<i>Iris foetidissima</i> L.	Sm. C.	Ir.	Eur. Orient.
<i>Iris germanica</i> L.	C. L.	Ir.	Eur. Orient.
<i>Iris hybrida</i> Retz.		Ir.	
<i>Iris neglecta</i> Pari.	Sm.	Ir.	

NAME OF PLANT	Authority	Properties	Locality
<i>Iris prismatica</i> Pursh.	L.	Ir	N A.
<i>Iris Pseudoacorus</i> L.	C. Sm.	It	Eur As Ori.
<i>Iris reticulata</i> Bieb.	Sm.	Ir	As Minor
<i>Iris sibirica</i> L.	C.	Ir	Eur N As.
<i>Iris variegata</i> L.	Wh.	Ir	Eur Orient.
<i>Iris versicolor</i>	L. Rusby	Ir Em. Cath.	N Am
<i>Sisyrinchium angustifolium</i> Mill.	L.	Cath.	N A.

JUGLANDACEAE

<i>Juglans cinerea</i> L.	L.	Cath	N A
<i>Juglans regia</i> L.	C. L.	Ant Astr	Him. Reg.

LABIATAE

<i>Achryospermum</i> G.	G.	Urt	Mad.
<i>Eremostachys superba</i> Royle	G. R.	F P	Him. Reg
<i>Lagochilus inebrians</i> Bunge.	G.	Nar	As.
<i>Leonurus Cardiaca</i> L.	G. L.	Card.	N Tem Reg
<i>Ocimum viride</i> Willd.	Sm.		Trop. Afr
<i>Salvia amarantha</i> Orteg.	G.	Nar	Mex
<i>Salvia officinalis</i> L.	L.	Nar.	Med. Reg
<i>Salvia pratensis</i> L.	G. L.	Nar	Eur Caucas.
<i>Scutellaria galericulata</i> L.	G.	Sed.	N Tem. Reg
<i>Scutellaria lateriflora</i> L.	L.	Sed.	N A
<i>Stachys arvensis</i> L.	G. M. L.	Antisp.	N Tem old wld.
<i>Stachys palustris</i> L.	L.	Antisp	Eur As. N A.
<i>Tetradlea fruticosa</i> Benth.	G.	Pur	Madag
<i>Teucrium Chamaedrys</i> L.	L.		Pur
<i>Teucrium Marum</i> L.	G. C. L.	Diaph	Med. Reg
<i>Teucrium scordium</i>	C. L.	Ant	Eur

LAURACEAE

<i>Hernandia sonora</i> L.	G.	P	Ind
<i>Laurus nobilis</i> L.	Sm. L.		Med Reg.
<i>Lindera Benzoin</i> Musa.	Sm. L.	Ant	E. N Am
<i>Umbellularia californica</i> Nutt.	G. C. L.	Hyd	N Am

LEGUMINOSAE

<i>Abrus precatorius</i> L.	M Sm. L.	Toxal Ira.	India
<i>Abrus pulchellus</i> Wall.	M.		Ind. Afr
<i>Acacia</i> sp.	G. C.	F P	Austr Tr. Reg.
<i>Acacia arabica</i> Willd.	G. L.		Afr As
<i>Acacia catechu</i> Willd.	L.	Astr	Ind Ceylon
<i>Acacia concinna</i> DC.	G.	Astr	Chin.
<i>Acacia Cunninghamii</i> Hook.	G.	Sap.	Austr
<i>Acacia deibrata</i> A.	G.	Sap.	Austr
<i>Acacia falcata</i> Willd.	G.		Austr
<i>Acacia Georginae</i> Bailey	G.		
<i>Acacia Jurema</i> Mart.	L.	Astr.	Brazi.
<i>Acacia pennata</i> Willd.	G.		As. Afr
<i>Acacia penninervis</i> Steber	G.		Austr
<i>Acacia pruriens</i> Kurz	G.	F P	Burm.
<i>Acacia pulchella</i> R. Br.	G.		Austr
<i>Acacia salicina</i> Linal.	G.		Austr
<i>Acacia verniciflua</i> A. Comm.	G. M.		Austr
<i>Acacia verticillata</i> Willd.	G.	Sap.	Austr
<i>Adenanthura pavonina</i> L.	G.		Tr As Malay

NAME OF PLANT	Authority	Properties	Locality
<i>Albizzia anthelmintica</i> Brongn.	G. L.	Sap F P	Abyss
<i>Albizzia Lebbeck</i> Benth.	G.		Tr old world
<i>Albizzia odoratissima</i> Benth.			Tr As. Ind.
<i>Albizzia procera</i> Benth.	G.		Tr As. Austr
<i>Albizzia stipulata</i> Boiv.	G. R. E.	F P	Ind
<i>Anagyris foetida</i>	C. G.		Sm. Cyt
<i>Andira anthe-martica</i> Benth.	G.	Ant	Braz
<i>Andira araroba</i> Aguiar	L.	Ir	Braz
<i>Andira Aubletia</i> Benth.	L.	Ant	Guiana
<i>Andira inermis</i> H. B. K.	L.	Ir	W Ind. Braz.
<i>Andira rosea</i> Mart.	G.	F P	Braz
<i>Andira retusa</i> HBK.	G. L.	Ant	Guiana
<i>Andira spinulosa</i> Mart. & Benth.	L.	Ant.	Braz.
<i>Andira vermifuga</i> Mart. & Benth.	Mills.	Ant	Braz.
<i>Argyrolobium pumilum</i> Eckl. & Zeyh.	G.		Afr Austr
<i>Astragalus baeticus</i> L.	G. L.	Sap.	Sic.
<i>Astragalus Bigelovii</i> A. Gray	Ches.	Loco	W N Am
<i>Astragalus exscapus</i> L.	L.	Dur	Eur
<i>Astragalus galegiformis</i> L.	G.	Sap	Caucas. Reg
<i>Astragalus garbancillo</i> Cav.	G.		Peru
<i>Astragalus glycyphyllos</i> L.	G. L.	Dur	Siber Eur
			Caucas. Reg.
<i>Astragalus hamosus</i> L.	G.		Caucas. Reg.
			Med. Reg. Ind.
<i>Astragalus Horru</i> A. Gray	G. Ches.	Del Loco	N Am.
<i>Astragalus lentiginosus</i> Dougl.	G. L.	Del Loco	N Am.
<i>Astragalus maximus</i> Willd.	G.	Sap	Armen. Caucas
			Reg
<i>Astragalus mollissimus</i> Torr.	G. L. Ches.	Del. Loco.	N Am Ry Mts.
			Col.
<i>Astragalus Mortonii</i> Nutt.	G. L.	Del Loco.	N Am.
<i>Astragalus ochroleucus</i> Cass.	G.	Del	Morocco
<i>Astragalus oocarpus</i> A. Gray	L.	Del. Loco.	W N Am
<i>Astragalus Pattersoni</i> A. Gray.	Ches.	Del Loco	W N. Am
<i>Baptisia</i> sp.	G.	Cyt	N Am
<i>Baptisia tinctoria</i> R. Br.	L.	Ein. Cath	N A
<i>Barbieria maynensis</i> Poepp.	G.		Peruv
<i>Barbieria polyphylla</i> DC.	G.		Porto Rico
<i>Bauhinia coccinea</i> DC.	G.		Cochin China
<i>Bauhinia guianensis</i> Aubl.	G.	F P	Guiana
<i>Bauhinia variegata</i> L.	G. L.	Ant	China, Burma
			Ind
<i>Bowdichia virginoides</i> H. B. K.	R. L.	F P	S. A
<i>Brachysema undulatum</i> Ker Gawl.	M.		Austr
<i>Caesalpinia Bonducella</i> Fleming.	G.		Tr Cosmop.
<i>Caesalpinia coriaria</i> Willd.	L.	Astr	N A
<i>Caesalpinia pulcherrima</i> Sw.	G.		Tr Cosmop
<i>Calandria Houstoni</i> Benth.	G.	Sap.	Mex
<i>Camptosema pinnatum</i> Benth.	G.		Braz
<i>Canavalia obtusifolia</i> DC.	G.		Cos Trop
<i>Cassia acutifolia</i> Delil.	G. L.	Pur	Egypt
<i>Cassia acata</i> L.	G.	F P	Cos. Trop.
<i>Cassia angustifolia</i> Vahl.	L.	Pur	Trop. Afr
<i>Cassia dymobotrya</i> Fres.	G.		Tr Am Abyss.
<i>Cassia hirsuta</i> L. f.	G. R.	F P	Tr Am.
<i>Cassia kituensis</i> Vatke	G.	F P	Tr Afr
<i>Cassia laevigata</i> Willd.	G. M.		Cos Trop.
<i>Cassia marylandica</i> L.	G. L.	Sap.	Tr m.
<i>Cassia occidentalis</i> L.	Sm. L.	Pur	Cosm. Trop.

NAME OF PLANT	Authority	Properties	Locality
<i>Cassia sophora</i> L.	G. M.		Tr. old world
<i>Cassia stipulacea</i> Ait.	G.		China
<i>Cassia Sturtii</i> R. Br.	G. M.		Austr.
<i>Castanospermum australe</i> A. Cunn.	G. M.	Sap. Ir.	Austr.
<i>Centrosema amazonicum</i> Mart.	G. R.	F. P.	Brazil
<i>Centrosema Plumieri</i> Benth.	G. R.	F. P.	S. Am.
<i>Cercis canadensis</i> L.	G.	Sap.	N. Am.
<i>Cercis chinensis</i> Bunge.	G.	Sap.	China
<i>Cladrastis amurensis</i> Benth.	G.		Amur Reg.
<i>Citoria amazonum</i>	R.	F. P.	S. Am.
<i>Citoria arborea</i> Ait.	G. R.	F. P.	Trinidad Is.
<i>Citoria Ternatea</i> L.	L.	Em.	Trop. Reg.
<i>Copaifera Langsdorffii</i> Desf.	L.	Diur.	Braz.
<i>Copaifera Salikorinda</i> Heck.	G.	Cou.	S. Am.
<i>Coronilla Emerus</i> L.	G. C.		S. Eur.
<i>Coronilla juncea</i> L.	G.		Eur.
<i>Coronilla montana</i> Scop.	G.		
<i>Coronilla scorpioides</i> Koch.	G. C. L.	Card.	Per. Reg. Med.
<i>Coronilla varia</i> L.	G. C.		Eur.
<i>Crotalaria alata</i> Hamill.	G. M.		Tr. As.
<i>Crotalaria Mitchelli</i> Benth.	M. G.		Austr.
<i>Crotalaria paniculata</i> Willd.	G.		Tr. As.
<i>Crotalaria sagittalis</i> L.	Ches. L.	Ineb.	N. Am.
<i>Crotalaria striata</i> Schrank.	G.		Braz.
<i>Crotalaria verrucosa</i> L.			Cosm. Trop.
<i>Cytisus Alchingeri</i> Vis.	C.		Dalmatia.
<i>Cytisus austriacus</i> L.	C.		Cauc.
<i>Cytisus biflorus</i> Ker. Gawl.	C.		S. Eur.
<i>Cytisus capitatus</i> Scop.	C.		S. Eur.
<i>Cytisus hirsutus</i> L.	C.		As. Min.
<i>Cytisus nigricans</i> L.	C.		Eur. As.
<i>Cytisus proliferus</i> L.	C.		Teneriffe
<i>Cytisus psaphereus</i> Scop.	C.		Eur. As.
<i>Cytisus scoparius</i> Link.	G. C. L.	Diur.	Eur. Jap. N. Am.
<i>Cytisus sessilifolius</i> L.	C.		S. Eur.
<i>Cytisus supinus</i> L.	C.		Eur.
<i>Dalbergia anacardaria</i> L. f.	G.		Ind.
<i>Dalbergia toxicaria</i> Baill.	G.		Madag.
<i>Derris amoena</i> Benth.	G.		Burm. Malaya
<i>Derris eliptica</i> Benth.	G. L.	F. P. Ina.	Burm. Malaya
<i>Derris guianensis</i> Benth.	G. R.	F. P.	Guiana. Braz.
<i>Derris Negrensis</i> Benth.	G.	F. P.	Braz.
<i>Derris virginica</i> Benth.	L. & G. R. L.	F. P.	Tr. old world
<i>Detarium Senegalense</i> J. F. Gmel.	G.		Tr. Afr.
<i>Entada polystachya</i> DC.	G.		Tr. Am.
<i>Entada scandens</i> Benth.	G. L.	Sap.	Tr. Am.
<i>Enterolobium cyclocarpum</i> Griseb.	G.	Sap.	Jamaica
<i>Enterolobium timbouva</i> Mart.	G.	Sap.	Braz.
<i>Eperua falcata</i> Aubl.	G.	Em.	Guiana
<i>Erythrina Corallodendron</i> L.	G. L.	Diur.	E. Ind. N. Am.
<i>Erythrina Hypaphysae</i> Boerl.	G.		
<i>Erythrina Mulungu</i> Mart.	L.	Hpy.	Braz.
<i>Erythrophleum Comminga</i> Baill.	C.		Mad.
<i>Erythrophleum guineense</i> G. Don.	C. L.	A. P. Nar.	Trop. Afr.
<i>Euchresta Horsfieldii</i> Bena.			Jav.
<i>Flemingia congesta</i> Roxb.		Tacu.	Tr. As.
<i>Galega officinalis</i> L.	G. L.	Sap.	Eur. As.
<i>Galega orientalis</i> Lam.	G.	Sap.	Caucas. Reg.
<i>Gastrolobium bilobum</i> R. Br.	G. M. Sm.	Ineb.	Austr.

NAME OF PLANT	Authority	Properties	Locality
<i>Gastrolobium calycinum</i> Benth.	G. M. Sm.	Inch.	Austr
<i>Gastrolobium Callistachys</i> Meiss.	G. M. Sm.	Inch.	Austr
<i>Gastrolobium grandiflorum</i> F. M.	G.	Inch.	Austr
<i>Gastrolobium obovatum</i> Benth.	G. M. Sm.	Inch.	Austr
<i>Gastrolobium ovalifolium</i> Henry	G.	Inch.	Austr
<i>Gastrolobium oxylobioides</i> Benth.	G. M. Sm.	Inch.	Austr
<i>Gastrolobium spinosum</i> Benth.	G. M. Sm.	Inch.	Austr
<i>Gastrolobium trilobum</i> Benth.	G. M. Sm.	Inch.	Austr
<i>Genista monosperma</i> Lam.	G.	Inch.	Med. Reg
<i>Genista sphaerocarpa</i> Lam.	G.	Inch.	Med. Reg
<i>Genista tinctoria</i> L.	L.	Sur	Eur. Cau. Reg
<i>Geoffrea superba</i>	G.	Ant.	S. Am.
<i>Gleditsia amorphoides</i> Griseb.	G.	Mech.	N. Am.
<i>Gleditsia sinensis</i> Lam.	G.	Sap.	China
<i>Gleditsia triacanthos</i> L.	G.	Mech.	N. A. A.
<i>Glicia maculata</i> HBK.	G.		Guat.
<i>Glycine hirsuta</i> Max.	G.	Fin	Japan
<i>Gompholobium uncinatum</i> A. Cunn.	G. M.	Inch.	Austr
<i>Goodia lotifolia</i> Sausb.	G.	Inch.	Austr
<i>Gymnocladus canadensis</i>	G. C.	Sap.	N. Am.
<i>Gymnocladus chinensis</i> Buell.	G.	Sap.	China
<i>Halimodendron argenteum</i> Fisch.	G.	Sap.	Reg. Caus. Per
			N. Aa.
<i>Hosackia Purshiana</i> Benth.	G.		N. Am.
<i>Hymenaea Courbaril</i> L.	G.	Ant.	S. Am.
<i>Indigofera anil</i> L.	G.		S. Am.
<i>Indigofera australis</i> Willd.	G. M.		Austr
<i>Indigofera gageoides</i> DC.	G.	Hyd.	Malay
<i>Indigofera tinctoria</i> L.	G.	Ins.	Trop. Reg
<i>Isotropis juncea</i> Turcz.	M.		Austr
<i>Laburnum anagyroides</i> Medic.	C. Sm. L.	Depr. Cyt.	Eur.
<i>Lathyrus amoenus</i> Fenzl.	C.		Syria
<i>Lathyrus Aphaca</i> L.	C.	As. Hyd.	Eur. Orient
<i>Lathyrus Cicera</i> L.	C.		Eur. Orient
<i>Lathyrus Clymenum</i> L.	C.		Med. Reg.
<i>Lathyrus odoratus</i> L.	C.		Italy
<i>Lathyrus purpureus</i> Presl.	C.		Sic
<i>Lathyrus sativus</i> L.	P. Sm. C.	Ent.	Eur. N. Afr.
			Orient
<i>Lessertia annularis</i> Burch.	M.		Austr
<i>Laucena glauca</i> Benth.	G.		Trop.
<i>Lonchocarpus densiflorus</i> Benth.	E. G. R.	F. P.	Guiana
<i>Lonchocarpus ichthyocarpus</i> Baill.	G.		Madag.
<i>Lonchocarpus latifolius</i> Kth.	G. R.	F. P.	Tr. Am.
<i>Lonchocarpus Neou</i> DC.	G. R. E.	F. P.	Tr. Afr.
<i>Lonchocarpus ratiflorus</i> Hart.	R.		Braz.
<i>Lonchocarpus violaceus</i> Kunth.	G.	F. P.	Tr. Afr.
<i>Lotus australis</i> Andr.	G. Sm. M.		Austr.
<i>Lotus corniculatus</i> L.	M.		Tem. old world
			Austr.
<i>Lupinus</i> sp.	G.		N. Temp. Reg.
<i>Lupinus albus</i> L.	C. Ches.		S. Eur.
<i>Lupinus angustifolius</i> L.	C.		Med. Reg.
<i>Lupinus densiflorus</i> Benth.	Sm.		Cal.
<i>Lupinus cucophyllus</i> Dougl.	Ches.		W. N. Am.
<i>Lupinus luteus</i> L.	C.		Med. Reg.
<i>Melilotus alba</i> Lam.	L.	Con.	
<i>Melilotus indica</i> Desf.		Con.	Eur. As.
<i>Melilotus officinalis</i> Desr.	L.	Con.	Eur. N. As.

NAME OF PLANT	Authority	Properties	Locality
<i>Milletia caffra</i> Mess.	G.		S. Afr
<i>Milletia ferruginea</i> Hochst.	R.	F P	S. Afr
<i>Milletia pachycarpa</i> Benth.	G.		Malay
<i>Milletia rostrata</i> Meig.	G.		Java
<i>Milletia sericea</i> G. & E.	R. E.	F P	Burm Malay
<i>Milletia splendens</i> Wright & Arn.	G.		Ind.
<i>Mucuna capitata</i> Max.	G.	F P	Him. Reg Malay
<i>Mucuna gigantea</i> DC.	G.		Tr Afr
<i>Mucuna purpurea</i> DC.	Wh.	Ir	Trop
<i>Mucuna urens</i> DC.	L.	Ir	S. Am
<i>Mucuna venenosa</i> A. Murr.	G.		Tr Afr
<i>Muellera moniformis</i> L.	G. E. R.	F P	Trop. Afr
<i>Muellera Telfairi</i> Baker	G.		Mad.
<i>Myroxylon Pereirae</i> Klotzsch.	L.	Sum.	C. A.
<i>Myroxylon peruvianum</i> L.	L.	Sum.	Eq. Peru Braz.
<i>Myroxylon tomentosum</i> HBK.	L.	Con.	Trop. Am.
<i>Nasolia fruticosa</i> Jacq.	G.	F P	Tr Am.
<i>Ormocarpum glabrum</i> Tels. & Binn.	G.		Celebes
<i>Ormosia coccinea</i> Jacks.	G.		Tr Am.
<i>Ougeinia da bergioides</i> Benth.	G.		Ind.
<i>Oxylobium parviflorum</i> Benth.	G. M.		Austr.
<i>Oxytropis foetida</i> DC.	Sm.		Eur.
<i>Oxytropis Lambertii</i> Pursh.	Ches.	Loco	W. N. Am. Ia.
<i>Oxytropis Lapponea</i> Gaud.	G.	Hyd. Sap.	Ry Mts.
<i>Oxytropis sulphurea</i> Fisch.	G.	Hyd.	Eur. N. As.
<i>Pachyrhizus anguatus</i> Rich.	G.		Siber.
<i>Pachyrhizus tuberosa</i> Spreng.	G.	F P	Tr. As.
<i>Parkia africana</i> R. Br.	G.		Ind.
<i>Pentaclethra macrophylla</i> Benth.	G.	Alk.	Tr. Afr.
<i>Phaseolus acutifolius</i> Jacq.	G.	Hyd. Asl.	Ind. Ar.
<i>Phaseolus lunatus</i> L.	Sm.		S. Am. Cosm.
<i>Phaseolus multiflorus</i> Willd.	G.		Cult.
<i>Phaseolus semierectus</i> L.	G. R.	F P	Mex.
<i>Phaseolus vulgaris</i> L.	G. C.		Trop. Reg.
<i>Physosugma venenosum</i> Baff.	G. Sm. L.	Dep. Eser.	N. Am. Cosm.
<i>Piptadenia peregrina</i> Benth.			Cult.
<i>Piscidia Erythrina</i> L. E.	G. Sm. R. L.	Inch. F P	Afr.
<i>Pithecolobium bigeminum</i> Mart.	G.		Braz.
<i>Pithecolobium fasciculatum</i> Benth.	G.	F P	Trop. Am.
<i>Pithecolobium Minabassae</i> Tels. & Binn.	G.		Malay
<i>Pithecolobium montanum</i> Benth.	G.		Malay
<i>Pithecolobium saman</i> Benth.	G.	Sap.	Him. Reg Malay
<i>Pongamia Piscida</i> Steud.	G. E.	F P	Tr. Am.
<i>Prosopis juliflora</i> Benth.	G.		Ind.
<i>Prosopis ruscifolia</i> Griseb.	G.	Sap.	N. Am. Tex.
<i>Psoralea glandulosa</i> L.	G. L.	Em.	Argen.
<i>Psoralea macrostachya</i> DC.	G.	Sap.	Chil. Mex.
<i>Psoralea pentaphylla</i> L.	G.		Cal.
<i>Psoralea tenuiflora</i> Pursh.	G.	Sap.	Mex.
<i>Pterocarpus esculentus</i> Schw.	G.	Sap.	N. Am.
<i>Pterocarpus Marsupium</i> Roxb.	L.	Intox.	Tr. Afr.
<i>Rhynchosia minima</i> DC.	G.	Astr.	Ind.
<i>Robinia Pseudo-acacia</i> L.	G. Ches.	Toxal.	Cosmop. Trop.
<i>Robinia viscosa</i> Vent.	L. Rusby		N. Am. S.
	Rusby	Em.	States
			N. A.
			S. Eur.

NAME OF PLANT	Authority	Properties	Locality
<i>Sabinea florida</i> DC.	G.		S. Eur
<i>Securigera coronilla</i> DC.	G.		N. Am. S.
<i>Sesbania vesicaria</i> Ell.	Ches.		States
<i>Sophora alopecuroides</i> L.	G.		Him. Reg.
<i>Sophora japonica</i> L.	L.	Pur	Japan, China
<i>Sophora molis</i> Grah.	G.		China, Him. Reg.
<i>Sophora secundiflora</i> Lah.	G. Ches. L.	Nar	Mex
<i>Sophora sericea</i> Nutt.	L. Ches.	Nar	N. A.
<i>Sophora tomentosa</i> L.	G. M.	Sap.	Cold Reg.
<i>Stryphnodendron barbatum</i> Mart.	G.	Sap.	Braz
<i>Stryphnodendron polyphyllum</i> Mart.	L.	Astr	Braz
<i>Swainsona coronillaefolia</i> Salisb.	G.	Ineb.	Austr
<i>Swainsona greyana</i> Lindl.	G. M.	Ineb.	Austr
<i>Swainsona ouvea</i> F. Muell.	M.	Ineb.	Austr
<i>Swainsona phacoides</i> Benth.	M.	Ineb.	Austr
<i>Swainsona procumbens</i> F. Muell.	M.		Austr
<i>Swartzia triphylla</i> Willd.	G.		Tr. Am.
<i>Templetonia egens</i> Benth.	G. M.		Austr
<i>Templetonia purpurea</i>	M.		Austr
<i>Templetonia retusa</i> R. Br.	G.		Austr
<i>Tephrosia astragaloides</i> Benth.	G.		Austr
<i>Tephrosia candida</i> DC.	G. R.	F. P.	Tr. As.
<i>Tephrosia cinerea</i> Pers.	G. R.	F. P.	
<i>Tephrosia coronillaefolia</i> DC.	G. R.	F. P.	Borbon Is.
<i>Tephrosia densiflora</i>	G.	F. P.	Tr. Afr.
<i>Tephrosia frutescens</i> DC.	Sm.		Honduras.
<i>Tephrosia macropoda</i> Harv.	G.	F. P.	S. Afr.
<i>Tephrosia nitens</i> Benth.	G.		Tr. Am.
<i>Tephrosia peruviana</i> Baker	G.	F. P.	
<i>Tephrosia purpurea</i> Pers.	G. M. R. L.	F. P.	Austr
<i>Tephrosia tomentosa</i> Pers.	R.	F. P.	Arab.
<i>Tephrosia toxicaria</i> Pers.	G. R. L. E.		Jamaica
<i>Tephrosia virginiana</i> Pers.	G.	A. P.	N. Am.
<i>Tephrosia vogelii</i> Hook.	G. R.		Tr. Afr.
<i>Teramus labialis</i> Spreng.	G.	Sap.	Cosm. Trop.
<i>Tetrapleura caerulea</i> Willd.	G.		Tr. Afr.
<i>Tetrapleura thonnigii</i> Benth.	Ches.	Diur.	W. N. Am.
<i>Thermopsis rhombifolia</i> Rich.	L.	Tymp.	Egypt
<i>Trifolium elegans</i> Sair.	C.	Tymp.	Med. Reg.
<i>Trifolium hybridum</i> L.	C.		S. Eur. As. Min.
<i>Trifolium incarnatum</i> L.	G.	Mec.	S. Eur.
<i>Trifolium nigrescens</i> Viv.	C.		S. Eur.
<i>Trifolium repens</i> L.	C.	Tymp.	N. Temp. Reg.
<i>Trigonella cretica</i> Boiss.	G.		As. Min.
<i>Vicia ervilia</i> Willd.	C.		Eur. N. Afr.
<i>Wistaria chinensis</i> DC.	G.		China
<i>Xylocabrisiformis</i> Benth.	G.	Sap.	Burma, Malay

LENTIBULARIACEAE

<i>Pinguicula vulgaris</i> L.	G.		Eur. Siber. N. Am.
-------------------------------	----	--	-----------------------

LILIACEAE

<i>Agrostocrinum stypanoides</i> Ev. M.	G. M.		Austr.
<i>Allium ascalonicum</i> L.	L.	Ant.	Cult.
<i>Allium canadense</i> L.	L.	Ant.	N. A.
<i>Allium cepa</i> L.	L.	Est.	Per. Bel.
<i>Allium douglasii</i> Hook.	L.	Ant.	Calif.

NAME OF PLANT	Authority	Properties	Locality
<i>Allium Macleanii</i> Baker	L.	Ant.	Afghan.
<i>Allium Moly</i> L.	L.	Ant.	Eur.
<i>Allium Porrum</i> L.	L.	Ant.	Eur.
<i>Allium sativum</i> L.	L.	Ir.	Eur. N. A.
<i>Allium Schoenoprasum</i> L.	L.	Ant.	Eur. As.
<i>Allium tricoccum</i> Soland.	Wh. L.	Ir.	N. A.
<i>Allium ursinum</i> L.	C. L.	Ant.	Eur. N. As.
<i>Allium Victoriana</i> L.	G. L.	Ant.	Eur. Sib.
<i>Allium vineale</i> L.	C.	Ir.	Eur. N. A.
<i>Aloe abyssinica</i> Lam.	L.	Pur.	Austr.
<i>Aloe africana</i> Mill.	L.	Pur.	S. Af.
<i>Aloe arborescens</i> Mill.	L.	Pur.	S. Af.
<i>Aloe chinensis</i> Steud.	L.	Pur.	China.
<i>Aloe ferox</i> Mill.	G. L.	Pur.	S. Afr.
<i>Aloe saponaria</i> Haw.	G.	Pur.	S. Afr.
<i>Aloe spicata</i> L.	L.	Pur.	S. Afr.
<i>Aloe succotrina</i> Lam.	L.	Pur.	S. Afr.
<i>Aloe tenuior</i> Haw.	G.	Pur.	Med. Reg.
<i>Aloe vera</i> L.	G.	Pur.	Med. Reg.
<i>Anemarrhena asphodeloides</i> Bunge.	G.		China.
<i>Asparagus officinalis</i> L.	L.	Ir.	Eur. As. N. A.
<i>Asparagus scaber</i> Brign.	L.	Ir.	Med. Reg.
<i>Astella Banksii</i> A. Conn.	G.		Cauc.
<i>Bulbine bulbosa</i> Haw.	G. M.		N. Z.
<i>Bulbine semibarbata</i> Haw.	G.		Austr.
<i>Chamaecrista caroliniana</i> Willd.	G. L.	Ant.	Austr.
<i>Colchicum autumnale</i> L.	G. C. Sm. L.		E. N. A.
<i>Colchicum neapontanum</i> Tenore.	C.	Ant.	Eur.
<i>Colchicum speciosum</i> Sw.	Sm.	Ant.	Italy.
<i>Colchicum variegatum</i> L.	C.	Ant.	Cauc. Reg.
<i>Convallaria majalis</i> L.	G. Sm. C. L.		S. Eur.
<i>Dianella nemorosa</i> Lam.	Ches. Rusby	Car.	N. Tem. Reg.
<i>Dracaena arborea</i> Hort.	G.		As. Trop.
<i>Drumma ciliaris</i> Jacq.	G.		Austr.
<i>Drumma Cowarii</i> R. d.	G.	Em.	Tr. Afr.
<i>Erythronium albidum</i> Nutt.	L.	Em.	Ag. Austr.
<i>Erythronium americanum</i> Ker. Gawl.	L.	Em.	Madag.
<i>Erythronium Dens-canis</i> L.	G.	Em. Ant.	N. A.
<i>Erythronium grandiflorum</i> Pursh.	G.		N. A.
<i>Erythronium purpurascens</i> S. Wat.	G.	Sap.	Eur. N. As.
<i>Fritillaria Imperialis</i> L.	G. C. M.	Ir.	N. Am.
<i>Fritillaria Meleagris</i> L.		Ir.	Cal.
<i>Fritillaria pinnata</i> Spreng.	L.	Ineb.	Persia. Him.
<i>Gloriosa simplex</i> L.	G.	Nar.	Reg.
<i>Gloriosa superba</i> L.	G.	Em.	Eur. Cauc.
<i>Heliosias frigida</i> Lindl.	G.		Reg.
<i>Hyacinthus orientalis</i> L.	G.		W. N. Am.
<i>Leucocoryne</i>	M.		Tr. Afr.
<i>Leucocrinum montanum</i> Nutt.	Ches.		Trop. As.
<i>Medeola virginica</i> L.	G.	Sap.	Austr.
<i>Meibomia cochinchinense</i> Lour.	G.		N. Am. Ry.
<i>Melanthium virginicum</i> L.	G. L.	Fly Poison	Mts.
<i>Muscaria comosum</i> Mill.	G.	Sap.	N. Am.
			Cochin. China.
			N. Am.
			Med. Reg. Ori.

NAME OF PLANT	Authority	Properties	Locality
Muscaria racemosum Willd.	Sm.	Ir	Med Reg. Cauc
Narthecium ossifragum Huds.	G. C		Eur As. Min. N. Am.
Nothoscordum striatum Kunth.	Ches.		N Am S States
Ornithogalum sp.	Mt.		Eur As Afr
Ornithogalum nutans L.	G.	Em.	Eur As. Min
Ornithogalum thyrsoides Jacq.			S. Afr
Ornithogalum umbellatum L.			N Afr As Min.
Paradisa Liliastrium Bertol.	G.	Card.	Eur
Paris obovata Ledeb	G.		Siber
Paris quadrifolia L.	G. C Mt. Sm. L.	Ir. Nar.	Eur. N. As
Polygonatum tuberosum L.	G.		Mex
Polygonatum biflorum Ell.	G.		N Am
Polygonatum officinale All.	G.		Eur Siber
Polygonatum verticillatum All.	G.		Eur N As
Ruscus aculeatus L.	G. L.	Sap.	Eur Ori
Schoenocaulon officinale A. Gr.	G. Sm. L.	Ast. Par	Mex.
Scilla bifolia L.	Sm.	Ir	Eur As. Min.
Scilla festans Salisb	Sm.	Ir	Eur
Smilax pseudo-syphilitica Kunth.	G.	Sap.	Braz.
Stypandra glauca R. Br	G. M		Austr
Toxicaria calyculata Wahl.	G.	Ant	Eur N Am
Trillium erectum L.	L. Rusby	Em	E. N Am.
Trillium grandiflorum Salisb.	G.	Sap	N Am
Tulipa Genersiana L.	G.		Eur Orient
Tulipa sylvestris L.	G.		Eur
Urginea Scilla Steinh.	G. C Sm. R. L.	Ast F P	S. Afr
Veratrum sp.			N Temp. Reg.
Veratrum album L. E.	G Mt. C Sm. L. E.	Ast. F P	N As.
Veratrum californicum Dur	Ches. L.	Ir	W N Am. Ry Mts.
Veratrum viride Asl.	G. Ches. L. Rusby	Ir Em.	E. N Am.
Xanthorrhoea sp.	G.		Austr
Xerophylla Douglasii S. Wat.		Sap	E. N Am S States.
Xerophylla setifolium Michx.	G.		E. N Am
Yucca filamentosa L.	G.	Sap.	W N Am. Ry Mts.
Yucca glauca (Nutt.) Carr	G.	Sap.	E. N Am S States.
Zygadenus angustifolium S. Wats.	L.		N A
Zygadenus elegans Pursh.	Ches. L.		E. N Am Ry Mts
Zygadenus venenosus S. Wat	Ches. G L.		N W Am

EUPHORBACEAE

Hugonia Mystax L.	G.		Ind
Linum catharticum L.	G L.	Pur	Eur Orient
Linum rigidum Pursh.	Ches.	Pur	N Am. N M Tex

NAME OF PLANT	Authority	Properties	Locality
<i>Linum toxicum</i> Boiss.	G.		Syria
<i>Linum usitatissimum</i> L.	G. L.	Ir	Eur. Orient N. Am.

LOGANIACEAE

<i>Anthracista grandiflora</i> Gilg.		Alk.	
<i>Buddleia brasiliensis</i> Jacq.	G.	F P	Hab.?
<i>Buddleia curviflora</i> Hook. & Arn.	G		Chili, Peru
<i>Buddleia garbosa</i> Hope	G	Sap.	China.
<i>Buddleia Landleyana</i> Fort.	G	F P	Madag
<i>Buddleia madagascariensis</i> Lam.	G	Taen.	Alysa.
<i>Buddleia polystachya</i> Friesen.	R G	Sap. F P	Brazil
<i>Buddleia variabilis</i> Hemsl.		Sap.	
<i>Buddleia verticillata</i> HBK.		F P	Mex
<i>Fagraea eslanica</i> Thumb.	Sm. G	Con.	Zeyl
<i>Gelsemium sempervirens</i> Ant.	G C Sm. L.	Ant.	E N Am. S. Sta.
<i>Nicodemia diversifolia</i> Ten.	G	Sap.	Mascar Is
<i>Spigelia Anthelmia</i> L.	G L.	Ir Nar	Trop. Am
<i>Spigelia glabrata</i> Mart	G.		Brazil
<i>Spigelia marylandica</i> L.	G. Sm. L.	Ir Nar	E N Am
<i>Spigelia pedunculata</i> HBK.	G		N Old World.
<i>Strychnos angustifolia</i> Benth.	G.		China
<i>Strychnos brachyata</i> Rox & Pav	G	Con	Peru
<i>Strychnos Castanea</i> Wedd.	G Sm. L.	Con	India
<i>Strychnos colubrina</i> L.	L.	A. P	Guiana
<i>Strychnos cogens</i> Benth.	L.	A. P	Braz.
<i>Strychnos Crevauxiana</i> Baill.	G L.	A. P	Guiana.
<i>Strychnos gableri</i> Planch.	L.	A. P	Venez.
<i>Strychnos Icaja</i> Baill.	G L.	Ordeal Poi	Trop. Afr
<i>Strychnos Ignatii</i> Berg.	G. Sm. L.	Con.	Philipp. Is
<i>Strychnos innocua</i> Del	G.		Nubia.
<i>Strychnos malaccensis</i> Benth.	L.		Bur Malay
<i>Strychnos Melinoniana</i> Bank.	G		Guiana
<i>Strychnos Nux-Vomica</i> L.	G Sm. R. L.	Con. F P	Burma.
<i>Strychnos paniculata</i> Champ.	G.		China
<i>Strychnos potatorum</i> L.	G		Burma.
<i>Strychnos Pseudo-quina</i> A St Hl.	G.		Braz
<i>Strychnos Santhierana</i> Pierre	G		
<i>Strychnos spinosa</i> Lam.	G.		Madag.
<i>Strychnos Tienta</i> Leach.	G Sm. L.	Con. A P	Java
<i>Strychnos toxifera</i> Schomb.	G. Sm.	Con. A P	Guian
<i>Strychnos yapirensis</i> Planch.	L.	A. P	Ecuador

LORANTHACEAE

<i>Phoradendron flavescens</i> Nutt.	L. Hyams.		N Am S Sta.
<i>Viscum album</i> L.	Sm. L.	Ir.	Eur Temp. As

LYCOPODIACEAE

<i>Lycopodium clavatum</i> L.	Mi. L.	Diur.	N Am. Eur
<i>Lycopodium Selago</i>	Mi. L.	Par	Eur

LYTHRACEAE

<i>Ammannia baccifera</i> L.	G.	Acr	Tr Old World
<i>Cuphea viscosissima</i> Jacq.	G.	Card.	N Am. S. Sta.
<i>Gnora mexicana</i> Lam.	G.	Em.	Mex.

NAME OF PLANT	Authority	Properties	Locality
<i>Lagerstroemia Flos-reginae</i> Retz.	G.	Nar.	Trop. As.
<i>Nesaea verticillata</i> HBK.	G.		N. A. S. Am.
<i>Punica Granatum</i> L.	G. L.	Taen.	S. Eur. Maurit.

MAGNOLIACEAE

<i>Drinys aromatica</i> P. Muell.	G.	Hyd.	Austr.
<i>Drinys Winteri</i> Forst.	G.	Hyd.	S. Am.
<i>Licium anisatum</i> Gaertn.	G. Sm. L.		China.
<i>Licium floridanum</i> Ellis.	G. Ches. L.		N. Am. S. Sta.
<i>Liriodendron chinense</i> Sarg.	G.		China.
<i>Liriodendron tulipifera</i> L.	G.	Hyd. Sap.	E. N. Am.
<i>Magnolia grandiflora</i> L.	G.		N. Am. S. Sta.
<i>Manglietia glauca</i> BL.	G.		Java.
<i>Michelia Champaca</i> L.	G.	Alk.	Malay.
<i>Michelia fuscata</i> Bl.	G.	Alk.	China.
<i>Michelia parviflora</i> Bl.	G.	Alk.	Java.
<i>Talauma macrocarpa</i> Zucc.	G.		Mex.
<i>Talauma ovata</i> A. St. Hil.	G.		Braz.

MALPIGHIACEAE

<i>Byrsonima amazonica</i> Griseb.	G.		Braz.
<i>Byrsonima crassifolia</i> HBK.	G.		Guiana.
<i>Byrsonima spicata</i> Rich.	G.		Trop. Am.
<i>Heteropteria syringaeifolia</i> Griseb.	G.		Braz.
<i>Malpighia oxycocca</i> Griseb.	G.		W. Ind.
<i>Sphedamnocarpus angoliensis</i> Planch.	G.		Trop. Afr.

MALVACEAE

<i>Aithaea rosea</i> L.	G.	Abort.	Ori. Cult. Eur.
			N. Am.
<i>Gossypium herbaceum</i> L.	G. L.	Abort.	Trop. As.
<i>Hubertus diversifolius</i> Jacq.	G.	Abort.	Tr. Old World
<i>Pavonia zeylanica</i>	G.	Ant.	
<i>Sida jamaicensis</i> L.	G.		Trop. Am.
<i>Sida parviflora</i> L.	G.	Nech.	Trop. Am.
<i>Sida urens</i> L.	G.	Ir.	Trop. Am. Afr.

MELASTOMACEAE

<i>Mouriria rhizophoraefolia</i> Gard.	G.		Martinq. Is.
--	----	--	--------------

MELIACEAE

<i>Carpis moluccensis</i> Lam.	G.		Tr. Old World
<i>Dysoxylum arborescens</i> Miq.	G.		Borneo.
<i>Flindersia Schottiana</i> Fy. Muell.	G.		Austr.
<i>Lansium domesticum</i> Jack.	G.	Ant.	Malay.
<i>Melia Azedarach</i> L.	G. C.	Ant.	Himal. Ch.
	Ches. L.		Cult. S. States
<i>Melia birmannica</i> Kurz.	G.		Burma.
<i>Naregamia alata</i> W. & A.	G. L.	Em.	Ind.
<i>Ptaeroxylon obliquum</i> Rolik.	G.		Afr.
<i>Sandoricum indicum</i> Cav.	G.		Trop. As.
<i>Swietenia humilis</i> Zucc.	G.		Mex.
<i>Swietenia Mahagoni</i> Jacq.	L.	Astr.	S.A. Mex. W. I.
<i>Trichilia</i> sp.	G.	Sap.	S. Am.
<i>Trichilia Moschata</i> Sw.	L.	Astr.	Jamaica.
<i>Trichilia trifolia</i> L.	G.	Abort.	Venz.
<i>Walsura piscidia</i> Roxb. G. & E.	R. E.	F. P.	India.

MELIANTHACEAE

NAME OF PLANT	Authority	Properties	Locality
Bersama sp.	G.		Arab. & Abyss.
Melanthus comosus Vahl.	G.		S. Afr.
Melanthus major L.	G.		S. Afr.

MIMNEMACEAE

Anamirta paniculata Colebr.	G. Sm. C R L E	Del. F P	Ind.
Anomospermum japonense Each.	G.		Braz.
Cissampelos Pareira L.	G.		Trop. Reg.
Cocculus Ferrandianus Gaudich.	G.	F P	Hawai Is.
Cocculus glaucescens B.	G.	Sap.	Bar Jav.
Cocculus laurifolius DC.	G.		Him. Reg. Jap.
Cocculus toxiferus Wedd.	G.		
Coscinum Borneanum Mierr.	G.	Sap.	Ind.
Menispermum canadense L.	L.	Diur.	N. A.
Pachygone ovata Mierr.	G. R.	F P	Malay Ind.
Pericampylus incarius Mierr.			Ind. Malay.
Sarcopetalum Harveyanum Fv. M.	G.		Austr.
Stephania discolor Spreng.	G.		Tr. Old World.
Stephania aculeata Walp.	G.		
Tiincora racemosa Colebr.	G.		Ind. Jav.

MONIMIACEAE

Kibara angustifolia	Sm.		
---------------------	-----	--	--

MONOTROPACEAE

Monotropa uniflora L.	G. L.	Antroa	E. N. Am. Jap.
Pterospora andromedea Nutt.	G.		Him. Reg. N. Am.

MORACEAE

Morus alba L.	L.	Taen.	Temp. As.
Morus nigra L.	L.	Taen.	Temp. As.
Morus rubra L.	L.	Taen.	N. A.

MYOPORACEAE

Eremophila maculata Fv. M.	M. G.		Austr.
Myoporum deserti A. Cunn.	G. M.		Austr.
Pholidia maculata Baill.	G.		

MYRICACEAE

Myrica cerifera	L.	Astr. Nar.	N. A.
-----------------	----	------------	-------

MYRISTICACEAE

Myristica gibbosa Hook.	L.	Astr.	India.
Myristica philippensis Lam.	L.	Astr.	Phil. Is.

MYRSINACEAE

Aegiceras majus Gaertn.	G. E. R.	Sap.	Tr. Old World.
Aegiceras minus Gaertn.	G. R.	F. P.	India.
Angularia dioica R. Br.	M.		Aust.
Cavendishia macrocarpa Ruiz. & Pav.	G.		S. Am.

NAME OF PLANT	Authority	Properties	Locality
<i>Embelia micrantha</i> A. DC.	G.	Ant.	Maurit Is.
<i>Embelia Ribes</i> Burm.	G.		Trop. As.
<i>Jacquinia arborea</i> Vahl.	G. R.	F P	Trop. Am.
<i>Jacquinia armillaris</i> Jacq.	G. E. R.	F P	W Ind.
<i>Jacquinia obovata</i> Schrad.	G. R.	F P	
<i>Maesa indica</i> Wall.	G.	F P	Ind Malay
<i>Maesa lanceolata</i> Forsk.	G.		N Afr
<i>Myrsine africana</i> L.	G.	Tact	Hum. Reg. Tr & S. Afr
<i>Theophrasta americana</i> L.	G.		Azores W Ind.

URTACEAE

<i>Euclea frutescens</i> L.	G.	Abort.	Malay China.
<i>Barringtonia acutangula</i> Gaertn.	G.		Malacca Is
<i>Barringtonia alba</i> Blume.	G.		Malacca Is
<i>Barringtonia Butonica</i> Forst.	G. R.	F P	Pac fic Is
<i>Barringtonia Carey</i> Fv. M.	G.		Austr
<i>Barringtonia insignis</i> Miq.	G.		Malay
<i>Barringtonia intermedia</i> Viell.	G.	F P.	N Caled.
<i>Barringtonia neo-caledonica</i> Viell.			N Ca ed
<i>Barringtonia racemosa</i> Roxb.		F P	Many Polyn.
<i>Eucalyptus Globulus</i> Labul.	L.	Astr	Austr
<i>Eucalyptus microtheca</i> Fv. Muell.	G.	F P	Austr.
<i>Eucalyptus rostrata</i> Schlecht.	L.	Astr	Austr
<i>Eugenia Jambos</i> L.	G.		Trop. As.
<i>Lecythis amara</i> Aubl.	G.		Guan.
<i>Lecythis lanceolata</i> Poir.	G.	Nar	Brazi.
<i>Melaleuca leucodendron</i> L.	Sm. L.		Austr.
<i>Napoleona Whitfieldii</i> Van Houtte	G.	Sap.	Trop. Afr
<i>Pimenta acris</i> Kostel.	Whi.	Ir	Ind.
<i>Podium montanum</i> Sw.	G.		W Ind.

URACEAE

<i>Musa sapientum</i> L.	G.	Sap.	Trop. Reg.
<i>Nepenthes gracilis</i> Korth.	G.		Borneo

NYCTAGINACEAE

<i>Boerhaavia erecta</i> L.	G.	Em.	N Am. W Ind.
<i>Boerhaavia hirsuta</i> L.	G.	Em.	N Am. W Ind.
<i>Boerhaavia repens</i> L.	G.	Em.	Cosm. Trop.
<i>Pisonia obtusata</i> Jacq.	G.	Em.	W Ind.
<i>Pisonia tomentosa</i> Casar.	G.	Em.	Braz.

NYMPHAEACEAE

<i>Euryale ferox</i> Salisb.	Sm.		Ind China
<i>Nymphaea acaena</i> Ait.	L.	Astr.	N A.
<i>Nymphaea alba</i> L.	G.		N Temp. Reg.
<i>Nymphaea lutea</i> L.	L.	Astr	Eur

OCHNACEAE

<i>Ochna</i> sp. Schreb.	G.		Afr Ind
--------------------------	----	--	---------

VILLACEAE

<i>Villaresia Moorei</i> F v. M.	G.		Austr
----------------------------------	----	--	-------

COROLLACEAE

NAME OF PLANT	Authority	Properties	Locality
<i>Chionanthus picrophloea</i> Fv Muell.	G.		
<i>Chionanthus virginica</i> L.	G. Sm. L.	Nar	N Am
<i>Forsythia intermedia</i> Zabel.	G.	Sap.	
<i>Forsythia suspensa</i> Vahl.	G.	Sap.	Japan, China.
<i>Forsythia viridissima</i> Lindl.	G.		China
<i>Jasminum floribundum</i> R. Br	G.		Abyss.
<i>Jasminum officinale</i> L.	Sm.	Ast	
<i>Jasminum Sambac</i> Ant.	G.		Trop. As
<i>Ligustrum vulgare</i> L.	Sm. Ches.	Ir	Eur As. N Am
<i>Olea dioica</i> Roxb.	G.	Em.	Ind
<i>Phillyrea media</i> L.	G.	Sap.	Med Reg

ONAGRACEAE

<i>Ludwigia erigata</i> L.	G.		Ind
<i>Montinia caryophyllacea</i> Thunb.			S. Afr
<i>Trapa natans</i> L.	G.		Ori. Ind Eur

ORCHIDACEAE

<i>Agrostis fragans</i> Thou.	G. L.	Ant	Mascar Is
<i>Catasetum</i> sp.	G.		Ind
<i>Cymbidium alofolium</i> Sw	G.	Eu	Ind.
<i>Cypripedium pubescens</i> Willd.	G.	Ir	N Am.
<i>Cypripedium spectabile</i> Salisb.	G.	Ir	N Am.
<i>Dendrobium nobile</i> Lindl.	G.		China
<i>Eria stelata</i> Lindl.	G.		Philipp. Is.
<i>Eulophia viridis</i> Spreng.	G.	Ant.	Ind
<i>Habenaria nigra</i> R. Br	G.		Eur.
<i>Neottia Nidus-avis</i> Rich.	G.	Ant.	Eur N As
<i>Orchis coriophora</i> L.	G.	Con.	Eur Orient
<i>Orchis odoratissima</i> L.	G.	Cou.	Eur
<i>Orchis purpurea</i> Huds.	G.	Cou.	As M n.
<i>Orchis Sima</i> Lam.	G.	Cou.	Eur Ori.
<i>Phajus calceus</i> Lindl.	G.		
<i>Phalaenopsis amabilis</i> Blume	G.		Malay
<i>Phalaenopsis Lucantheana</i> Reichb.	G.		Philipp. Is

OROBANCHACEAE**OXALIDACEAE**

<i>Oxalis Acetosella</i> L.	L.	Vit	Eur As. N
<i>Oxalis amara</i> A. St Hill.	G.		Afr N A
<i>Oxalis Pes-caprae</i> L.	G. L.	Diar	Braz
<i>Oxalis purpurata</i> Jacq			S. Afr
<i>Oxalis Sunthiana</i> Eckl. & Zeyr	G.		S. Afr

PALEACEAE

<i>Areca Catechu</i> L. var. <i>nigra</i> .	G. L.	Nar	Ind Malay
<i>Arenga saccharifera</i> Labill.	G.		Malay
<i>Borassus flabellifera</i> Murr	G.		Ind.
<i>Cocos amara</i> Jacq.	G.		W Ind.
<i>Cocos nucifera</i> L.	L.	Ver	
<i>Corypha umbraculifera</i> L.	G.		Trop. Afr Ind
<i>Hyphaene thebaica</i> Mart.	G.	Taen.	Trop. Afr
<i>Hyphorbe indica</i> Gaertn.	G.		
<i>Phoenix dactylifera</i>	G. L.	Cur	N Afr Arab.

FAMILIAR

NAME OF PLANT	Authority	Properties	Locality
<i>Pandanus odoratissimus</i> L.	G.		Ind. Arab.
<i>Pandanus Thomensis</i> Henrig.	G.		

PAPAVERACEAE

<i>Argemone alba</i> Lestib.	L.	Nar	S U S.
<i>Argemone intermedia</i> Sw	L.	Nat.	Mex.
<i>Argemone mexicana</i> L.	Sm. Ches. L.	Nar	N Am. Tex Mex.
<i>Bocconia cordata</i> Willd.	L.	Ant	China Japan
<i>Bocconia frutescens</i> L.	G. L.	Ant	Mex. Peru
<i>Chelidonium majus</i> L.	C Rusby G Sm. Ches. Ms.	Ir Nar	Eur As. N Am
<i>Eschscholzia californica</i> Cham.	G.		Calif
<i>Glaucium corniculatum</i> Corb.	L.	Puc	Eur.
<i>Glaucium flavum</i> Crantz.	C. Sm. L.	Ir	Eur Med. Reg
<i>Hypocoum procumbens</i> L.	G.		Med. Reg Arabia.
<i>Papaver aculeatum</i> Thumb.	G.		S. Afr Austr
<i>Papaver Rhoeas</i> L.	G Sm. Ches. C L.	Ir	Eur Ori. Afr
<i>Papaver somniferum</i> L.	C Mi. Ches. L.	Nar	Eur As. Ind N Am
<i>Roemeria violacea</i> Medic.	Sm	Ir	Eur N Afr
<i>Sanguinaria canadensis</i> L.	G Rusby L.	Ir Fin	N Am
<i>Stylophorum diphyllum</i> Nutt.	G.		N Am

PASSIFLORACEAE

<i>Andrena lobata</i> Engl.	G.	Ant.	Trop. Afr
<i>Carica Papaya</i> L.	G.		Trop. Amr
<i>Carica quercifolia</i> H. St. Hil.	G.		
<i>Carica spinosa</i> Aub.	G.		Braz Guiana.
<i>Modecca palmata</i> Lam.	G.		Ind
<i>Modecca trilobata</i> Roxb.	G.	Sap	Ind.
<i>Modecca venenata</i> Forsk.	G.		
<i>Ophiocaulon gummifera</i> Han.	G.		Trop. Afr
<i>Passiflora caerulea</i> L.	G.	Eur	Brazil.
<i>Passiflora foetida</i>	Sm. L.	Antisp.	
<i>Passiflora Herbertiana</i> Ker-Gawl.	G.		Austr
<i>Passiflora hispida</i> DC.	L.	Nar	Jamaica
<i>Passiflora incarnata</i> L.	L.	Antisp.	Trop. Amer
<i>Passiflora laurifolia</i> L.	G L.	Hyd.	Trop. Amer
<i>Passiflora quadrangularis</i> L.	G. L.	Nar	Trop. Amer
<i>Passiflora rubra</i> L.	G.	Nar	W. Ind

PHYTOLACCACEAE

<i>Anisomeria drastica</i> Mey	G.		
<i>Gailesia Scorodendron</i>	G.	Ant	
<i>Petiveria alba</i> L.	G. L.	Abort.	W Ind
<i>Petiveria tetrandra</i> Gomez	G.	F P	Braz.
<i>Phytolacca abyssinica</i> Hoffm.	G.	Sap.	Trop. & S. Afr
<i>Phytolacca acinosa</i> Roxb.	G.	Del Sap.	Him Reg Chin.
<i>Phytolacca decandra</i> L.	G. C. L.	Nar Em	N Amer
	Ches. Rusby	Sap.	

NAME OF PLANT	Authority	Properties	Locality
<i>Phytolacca dioica</i>	G.	Sap.	S. Amer
<i>Phytolacca rosandra</i> L.	G.	Sap.	Ind
<i>Pircunia</i> sp. (<i>Phytolacca</i>)	G.	Sap.	S. Am
<i>Rivina humilis</i>	G.		Trop. Am

PIPERACEAE

<i>Piper Carpanya</i> Ruiz & Pav	G. L.		India
<i>Piper darriense</i> C. DC	L.	Sol	Chil & Peru.
<i>Piper methysticum</i> Forst	R. E.	F. P.	Panama
<i>Piper nigrum</i> L.	L.	Ineb.	Pacific Is
<i>Piper Palmeri</i> C. DC.	G.	Sap.	
<i>Piper peltatum</i> L.	L.	Dur	Trop. Am.
<i>Piper umbellatum</i> L.	L.	Dur	Coast Tr. Reg.

PIROLACEAE

<i>Chimaphila maculata</i> Pursh.	G. L.	Astr	N. Am.
<i>Chimaphila umbellata</i> Nutt.	L.	Astr	Eur. N. As.
<i>Monotropa uniflora</i>	Sm.		N. A.
			Him. Reg.
<i>Pyrola chlorantha</i> Sw.	L.	Astr	Japan. N. A.
<i>Pyrola elliptica</i> Nutt.	L.	Astr	Eur. N. A.
<i>Pyrola rotundifolia</i> L.	G. L.	Em.	N. Temp. Reg.
<i>Pyrola minor</i> L.	L.	Astr	N. Temp. Reg.

PITTOSPORACEAE

<i>Bilardiera longiflora</i> Labill.		Sap.	
<i>Pittosporum coriaceum</i> Ait.	G.	Sap.	Madeira Is.
<i>Pittosporum cornifolium</i> A. Cunn.	G.	Sap.	N. Zeal.
<i>Pittosporum crassifolium</i> Soland.	G.	Sap.	N. Zeal.
<i>Pittosporum eugenioides</i> A. Cunn.	G.	Sap.	N. Zeal.
<i>Pittosporum ferrugineum</i> Ait.		Sap.	Malay Austr
<i>Pittosporum floribundum</i> W. & A.	G.		Ind.
<i>Pittosporum huttonianum</i> T. Kirk.	G.	Sap.	N. Zeal.
<i>Pittosporum javanicum</i> Bl.	G.	F. P.	Java
<i>Pittosporum Moorei</i> F. v. Muell.	G.	Sap.	
<i>Pittosporum phyllitacoides</i> DC.	G.	Sap.	Austr
<i>Pittosporum rhombifolium</i> A. Cunn.			
<i>Pittosporum tobira</i> Ait.	G.		Japan. China
<i>Pittosporum undulatum</i> Vent.	G.	Sap.	Austr

PLANTAGINACEAE

PLATANACEAE

<i>Platanus occidentalis</i> L.	G.	Hyd	N. Am.
<i>Platanus orientalis</i> L.	G.	Hyd	S. Eur. Orient

PLUMBAGINACEAE

<i>Armeria elongata</i> Hoffm.	L.	Astr	Eur
<i>Plumbago capensis</i> Thunb.			
<i>Plumbago europaea</i> L.	L.	Em.	S. Eur. Caucas.
<i>Plumbago rosea</i> L.	G. L.	Nar. Acr	Ind
<i>Plumbago scandens</i> L.	G. L.	Pur	Trop. Am
<i>Plumbago toxicaria</i> Bertol.	G.		Trop. Afr
<i>Plumbago zeylanica</i> L.	G.	Abort. Ir	Tr. Old World
<i>Statice pectinata</i> Ait.	G.	F. P.	Canary Is.

POLYTOXIC

NAME OF PLANT	Authority	Properties	Locality
<i>Cantua buxifolia</i> Lam.	G.		Peru
<i>Cantua pyrifolia</i> Juss.	G.		Peru
<i>Cobaea scandens</i> Cav.	G.	Sap.	Mex Ry. Mts. N. Am.
<i>Gilia achilaeifolia</i> Benth.	G.	Sap.	Calif.
<i>Gilia aggregata</i> Spreng.	G.	Sap.	N. A. Ry. Mts.
<i>Gilia laciniata</i> Ruiz. & Pav.	G.	Sap.	Peru. Chili.
<i>Loeselia caerulea</i> G. Don.	G. L.	Em.	Mex
<i>Phlox</i> L.	G.		N. Am.
<i>Polemonium boreale</i> Adams.	G.	Sap.	Siberia
<i>Polemonium caeruleum</i> L.	L.	Sap.	N. Temp. Reg.
<i>Polemonium flavum</i> Greene	G.	Sap.	N. A. N. Mex.
<i>Polemonium gracile</i> Wild.	G.	Sap.	
<i>Polemonium humile</i> Wild.	G.	Sap.	Arct. Reg.
<i>Polemonium paniculatum</i> Ser. Wat.			W. N. Am.
<i>Polemonium reptans</i> L.	G. L.	Sap.	N. Am.

POLYGALACEAE

<i>Krameria triandra</i> R. & Pav.	L.	Ast.	Peru
<i>Monarda</i> sp. R. & P.	G.	Sap.	Trop. Am.
<i>Polygala alba</i> Nutt.	L.	Ir.	N. A.
<i>Polygala amara</i> L.	G.		Eur.
<i>Polygala angulata</i> DC.	G.	Sap.	Brazil.
<i>Polygala nepalensis</i> L.	G.		Braz.
<i>Polygala Rayana</i> Nutt.	L.	Ir.	N. A.
<i>Polygala Cyparissias</i> A. St. Hil.	G.	P.	
<i>Polygala glandulosa</i> HBK.	G.		Mexico
<i>Polygala Javiana</i> DC.	G.	Sap. Ins.	Malaga
<i>Polygala Senega</i> L.	G. Sm. L.	Sap. Ir.	N. Am.
<i>Securidaca longepedunculata</i> Fres.	G.	P.	Abyss.

POLYGONACEAE

<i>Fagopyrum esculentum</i> Moench.	C.	Ir.	Eur. N. As.
<i>Polygonum acre</i> HBK.	R. L. E.	Ir. F. P.	Corn.
<i>Polygonum barbatum</i> L.	G.	F. P.	Tr. Old World.
<i>Polygonum bistorta</i> L.	C. L.	Astr.	N. Reg.
<i>Polygonum convolvulus</i> L.	C.		N. Temp. Reg.
<i>Polygonum flaccidum</i> Meiss.	G.	Ir.	Ind. Malay.
<i>Polygonum hydropiper</i> L.	G. C. L.	Ir.	Temp. Reg. N. & S.
<i>Polygonum hydropiperoides</i> Michx.	G.	Ir.	N. Am.
<i>Polygonum orientale</i> L.	G.	Ir.	Tr. Old World.
<i>Rheum emodin</i> Walb.	L.	Pur.	Him. Reg.
<i>Rheum hybridum</i> Murray	L.	Pur.	Mongolia
<i>Rheum officinale</i> Baill.	L.	Pur.	Tibet
<i>Rheum palmatum</i> L.	Sm.	Pur.	Mong.
<i>Rheum Rhaponticum</i> Delarb.	L.	Pur. Astr.	Siberia.
	Newspaper Death in Iowa		
<i>Rumex abyssinicus</i> Jacq.	G.	Tact.	Abyss.
<i>Rumex acetosa</i> L.	L.	Astr.	Eur. N. As.
<i>Rumex acetosella</i>	C. L.	Oxal. Acid	Eur. N. Am.
<i>Rumex crispus</i>		Oxal. Acid	Eur. N. Am.
<i>Rumex Ecklonianus</i> Meiss.	G.	Tact.	S. Afr.
<i>Rumex hymenosepalus</i> Torr.	G. L.	Astr.	W. & S. W. N. Am.
<i>Rumex obtusifolius</i> L.	L.	Astr.	Eur. As. N. A.

PORTULACACEÆ

NAME OF PLANT	Authority	Properties	Locality
<i>Lewisia rediviva</i> Pursh.	G.		W N Am. Mont.
<i>Talinum polyandrum</i> Ruiz. & Pav.	G.	Acce.	Peru.

PRIMULACEÆ

<i>Anagallis arvensis</i> L.	G. C. Ches.	Sap	Eur Temp As.
	Mt. L.		
<i>Cortusa Matthioli</i> L.	C.	Sap	N. As.
<i>Cyclamen europæum</i> L.	R. C. E. G.	Sap F P	Eur Cauc. Reg.
	L.		
<i>Cyclamen graecum</i> Link.	R.	F P	Greece
<i>Cyclamen hederacifolium</i> Ait.	G.		S. Eur
<i>Cyclamen latifolium</i> Sibth. & Sm.	G.		Greece, A. Min.
<i>Lysimachia Nummularia</i> L.	G. L.	Aste.	Eur
<i>Primula Auricula</i> L.	G.		Eur
<i>Primula obconica</i> Hance.	G. L.	Ir.	China.
<i>Primula Parryi</i> A. Gray.	Pam.		Ry. Mts. N. A.
<i>Primula reticulata</i> Wail.	G.		Him. Reg.
<i>Primula sinensis</i>	Sm.	Ir.	China

PROTEACEÆ

<i>Brabejum</i> L.	G.	Hyd.	S. Afr.
<i>Grevillea ramosoides</i> R. Br.	G.		Austr.
<i>Heucia robusta</i> Wail.	G.	Hyd.	Him. Reg. Bur-
			ma Malay
<i>Knightia excelsa</i> R. Br.	G.	Sap.	N. Zeal.
<i>Macadamia ternstrofia</i> F. Muell.	G.	Hyd.	Austr.
<i>Protea cynaroides</i> L.	G.	Hyd.	S. Afr.
<i>Roupala Pohl</i> Meiss.	G.		Brazil.
<i>Roupala vervaneana</i> Hort.	G.		Austr.
<i>Xylomelum pyriforme</i> Knight.	G.		Austr.

RANUNCULACEÆ

<i>Aconitum Anthora</i> L.	C. L.		N. As.
<i>Aconitum chinense</i> Sieb.	L.		Japan
<i>Aconitum ferox</i> Wall.	C. Sm. L.	F P. Ast.	Him. Reg.
<i>Aconitum columbianum</i> Nutt.	Ches.		W N Am.
			Ry. Mts.
<i>Aconitum Fischeri</i> Reich.	L.		N. As. N. Am.
<i>Aconitum heterophyllum</i> Wall.	C. L.		Him. Reg.
<i>Aconitum japonicum</i> Decne.	L.		Japan.
<i>Aconitum Lycoctonum</i> L.	Mt. C. G. L.		Eur. N. As.
<i>Aconitum Napellus</i> L.	Mt. Rusby	Ast.	N. Temp. Reg.
	C. Sm. Wh.		Eur.
	L. Ches.		
<i>Aconitum noveboracense</i> Gray.	Rusby	Ast.	E. N. Am.
<i>Aconitum uncinatum</i> L.	L.		N. A. Japan.
<i>Actæa alba</i> (L.) Mill.	L. Rusby	Em.	N. A.
	Ches.		
<i>Actæa rubra</i> (Ait.) Willd.	Rusby		N. Temp. Reg.
<i>Actæa spicata</i> L.	Wh. G. Sm.	Asth.	N. A.
<i>Adonis D. L. sp.</i>	Mt.		Eur. As.
<i>Adonis aestivalis</i> L.	G.	Ir.	Eur. Ori.
<i>Adonis amurensis</i> Regel & Radde.	G.	Ir.	Manchuria.
<i>Adonis vernalis</i>	C. Sm. L.	Ir.	Eur. N. Am.
<i>Anemone altaica</i> Fisch.	G.		Siber. Altai.
<i>Anemone apennina</i> L.	Mt.	Ir.	N. Am.

NAME OF PLANT	Authority	Properties	Locality
<i>Anemone coronaria</i>	L.	Ir	Med. Reg Ori.
<i>Anemone nemorosa</i> L.	Sm. C. G. Mi. L.	Ir	Eur N As. N Am.
<i>Anemone parviflora</i> Michx.	G		N Am
<i>Anemone patens</i> L. var <i>Wolfgangiana</i> Koch.	Ches. Rusby	Abort Ir	N Cent St Ry Mts. Br Am.
<i>Anemone pratensis</i> L.	C.		Eur
<i>Anemone Pulsatilla</i> L.	G. C. Mi. Sm.	Abort Ir	Eur
<i>Anemone ranunculoides</i> L.	Mi. L.	Ir	Eur Can Siberia.
<i>Anemone sylvestris</i> L.	L.	Ir	Eur Cauc Reg
<i>Aquilegia caerulea</i> James.	L.	Dior	W N Am
<i>Aquilegia canadensis</i> L.	L.	Dior	N. A.
<i>Aquilegia vulgaris</i> L.	G. C. Mi. L.	Ir	Eur Ori.
<i>Caltha arctica</i> K. Br	Sm.	Eur	Arct Reg
<i>Caltha palustris</i> L.	G. C. Sm.	Acr	N Temp. & Arct Reg
<i>Cnicifuga foetida</i> L.	C. Sm.	Acr	Eur Siberia
<i>Cnicifuga racemosa</i> L.	G. Sm. L. Rusby	Ast	N Am.
<i>Clematis aethusifolia</i> Turcz.	G.	Acr	Mong
<i>Clematis alpina</i> Mill.	G. C. Sm.	Sap.	Eur N Am.
<i>Clematis Bergeroni</i> A. Lavall.	G.	Sap.	
<i>Clematis brachiata</i> Thunb.	G.	Ver	S Afr
<i>Clematis Buchaniana</i> DC	G.	Sap.	Him Reg
<i>Clematis caripensis</i> HBK.	G.	Ves	Mex
<i>Clematis cirrhosa</i> L.	G.	Sap.	Med. Reg
<i>Clematis flammula</i> L.	C. G.	Sap.	Med. Reg
<i>Clematis florida</i> Thunb.	G.	Ir Sap.	Japan
<i>Clematis Fremonti</i> S. War	G.	Sap. Hyd.	W N Am
<i>Clematis Gouiana</i> Roxb.	G.	Ir	Ind. Malay
<i>Clematis integrifolia</i> L.	G. C.	Hyd	S Eur N As
<i>Clematis lanuginosa</i> Lindl.	G.	Ir Hyd	China W N Amer Ry Mts.
<i>Clematis mauritiana</i> L.	G.		Madag Maurit Is.
<i>Clematis orientalis</i> L.	G.	Hyd	Him Reg Ori.
<i>Clematis Pitcheri</i> Torr & Gray	G.	Sap.	E. N Am
<i>Clematis pseudo-flammula</i> Sch.	G.	Sap.	
<i>Clematis reticulata</i> Walt.		Sap.	N Am.
<i>Clematis Vitalba</i> L.	G. Sm. C. Mi. L.	Ir	Eur N Afr Cauc
<i>Clematis Wightiana</i> Wall		Hyd	Ind.
<i>Clematis Williamsii</i> Gray	G. Sm.		Japan.
<i>Delphinium Ajacis</i> L.	M. G. L.	Ir	Eur
<i>Delphinium Brunonianum</i> Royle	G.	Card.	Him. Reg
<i>Delphinium caeruleum</i> Jacq.	G.		Him. Reg N Am
<i>Delphinium Consolida</i> L.	C. Mi. L. Ches.	Ast	Eur N As
<i>Delphinium Geyeri</i> Greene	Ches. L. Ches.	Acr	N A Ry Mts
<i>Delphinium Menziesii</i> DC	G. L. Ches.	Car	W N Am
<i>Delphinium mauritanica</i> Coss.	G.		N Afr
<i>Delphinium peregrinum</i> L.	G.		S. Eur Ori
<i>Delphinium recurvatum</i> Greene	Ches. L. Ches.	Acr	W N Am.
<i>Delphinium Reicheni</i> DC	C.		Med Reg
<i>Delphinium scopolorum</i> A. Gray.	L. Ches.		N Am

NAME OF PLANT	Authority	Properties	Locality
<i>Delphinium Staphisagria</i> L.	G.Sm.W.C. L.	Asi Paras.	Med Reg
<i>Delphinium tricornis</i> Michx.	G. L. Ches.	Car	E N A. S. Sta.
<i>Delphinium consolida</i> Gray	G. Ches.		W N Am
<i>Delphinium uncinatum</i> Hook.	G. L.	Ir	Him Reg.
<i>Eranthis hyemalis</i> Salisb.	Sm. G.	Asi.	Eur
<i>Helleborus foetidus</i> L.	C.G.Sm.L.	Pur	Eur
<i>Helleborus niger</i> L.	G.W.C.Sm. L. M.	Pur	Eur
<i>Helleborus odorus</i> Waldst.	Sm		Hungary
<i>Helleborus orientalis</i> Lam.	C Sm		A Mon. Greece
<i>Helleborus viridis</i> L.	G Ches. L.	Pur	Ea.
<i>Hydrastis canadensis</i> L.	L.	Verb.	N A
<i>Isopyrum fumarioides</i> L.	G	Hyd	Eur N As.
<i>Isopyrum thalictroides</i> L.	G.		S. Eur
<i>Knowltonia vesicatoria</i> Sims.	G L.	Acr Vea.	S Afr Cape Reg.
<i>Nigella damascena</i> L.	L.	Sap	Med. Reg
<i>Nigella arvensis</i> L.	G. L.	Sap.	Med. Reg
<i>Ranunculus abortivus</i> L.		Ir	E. N Am.
<i>Ranunculus acris</i> L.	C.G.Sm.M. L.	Acr	Eur N As.
<i>Ranunculus alpestris</i> L.		Acr	Eur
<i>Ranunculus aquatilis</i> L.	M. Sm.	Ir	Tem. Reg.
<i>Ranunculus arvensis</i> L.	G.Sm.C.L.	Acr	Eur N As.
<i>Ranunculus asiaticus</i> L.	Sm.	Ir	Asa
<i>Ranunculus auricomus</i> L.	M. Sm. L.	Ir	N Temp. Reg.
<i>Ranunculus bulbosus</i> L.	G.C.M.Sm.	Ir	Temp. Reg.
<i>Ranunculus ficaria</i> L.	G.C.Sm.M.	Ir	Eur Caucas.
<i>Ranunculus flammula</i> L.	G.C.Sm.M. Rusby	Ir	N Temp. Reg.
<i>Ranunculus hybridus</i>	G. C. M.	Ir	S Eur
<i>Ranunculus lanuginosus</i> L.	G.		Eur Cau Reg
<i>Ranunculus lappaceus</i>	Sm. M.	P	Austr
<i>Ranunculus lingua</i> L.	C. M.		Eur Siber
<i>Ranunculus polyanthemus</i> L.	M.		Cauc Reg
<i>Ranunculus repens</i> L.	C M.		N Temp Reg
<i>Ranunculus sceleratus</i> L.	G M. Sm.	Acr Ir	Ru. N Am.
<i>Ranunculus septentrionalis</i> Poir.	RusbyChes.		
<i>Ranunculus Thora</i> L.	Ir	Acr	E. N Am
	G M.	Vea.	S. Eur
	Rusby		
<i>Thalictrum flavum</i> L.	C.		Eur N As
<i>Thalictrum foetidum</i> L.	Sm		Eur Siberia
<i>Thalictrum macrocarpum</i> Gren.	C G.		
<i>Trochilus asiaticus</i> L.	G.	Sap.	Siberia
<i>Trochilus europaeus</i> L.	G.	Sap	Eur Caucas.
<i>Trochilus pumilus</i> D. Don.	G	Sap	Him. Reg.
<i>Zanthoxyla</i>	M.		Austr

RESEDACEAE

<i>Reseda Tourn</i> sp.	G.	Hyd.	Eur
<i>Reseda luteola</i> L.	L.	Dmr	Eur

RHAMNACEAE

<i>Ceanothus americanus</i> L.	G. L.	Sap.	E. N Am.
<i>Ceanothus azureus</i> Desf.	G.		Mex.
<i>Ceanothus caeruleus</i> Lag.	L.		Mex

NAME OF PLANT	Authority	Properties	Locality
<i>Ceanothus integerrimus</i> Hook & Arn.	G.	Sap.	N W Am.
<i>Ceanothus ovatus</i> Desf.	G.	Sap.	N Am
<i>Ceanothus thyrsiflorus</i> Eschw.	G.	Sap.	Calif
<i>Ceanothus velutinus</i> Dougl.	G. L.	Sap.	N W Am. Ry Mts
<i>Coilectia spinosa</i> Lam.	G.	Sap.	S. Am
<i>Disarcia serratifolia</i> Benth. & Hook	G.	Sap.	Chi. Pat
<i>Gouania</i> Jacq. sp.	G.		
<i>Karwinskia Humboldtiana</i> Zucc.	G. Coult (Mitchell)		W N Am. Tex
<i>Paliurus aculeatus</i> Lam.	G. L.		S. Eur W As
<i>Rhamnus Alaternus</i> L.	C		Med. Reg. As Minor
<i>Rhamnus californica</i> Esch.	G.	Ir	As Min.
<i>Rhamnus caroliniana</i> Walt.	L.	Lax	N A.
<i>Rhamnus cathartica</i> L.	C. Sm. L.	Ir	Eur As & N Afr
<i>Rhamnus Frangula</i> L.	G. C. L.	Lax	Asia N Afr Eur
<i>Rhamnus Purshiana</i> DC.	L.	Lax	N A
<i>Rhamnus Wightii</i> W. & A.	L.	Lax	Ind
<i>Zizyphus Joazeira</i> Mart.	G.	Sap.	Brazil.
<i>Zizyphus Lotus</i> Lam.	G.		Med Reg
<i>Zizyphus sativa</i> Gaertn.	G.		Med Reg. Temp. As.

RHIZOPHORACEAE

<i>Rhizophora Mangle</i> L.	G Sm L.	Tan.	Tr Shores
-----------------------------	---------	------	-----------

ROSACEAE

<i>Agrimonia eupatoria</i> L.	G. L.	Ant.	N Tem.
<i>Brayera anthelmintica</i> Kunth.	G. L.	Ant	Abyss.
<i>Cercocarpus parviflorus</i> Nutt	G.	Sap.	Ry Mts. W N Am.
<i>Cormus foliosa</i> Planch.	G.	Hyd	
<i>Cotoneaster</i> Rupp.	G.	Hyd	N Temp. Reg
<i>Galenia stipularia</i> Nutt.	G. L.	Em	N Am
<i>Gilenia trifoliata</i> Moench.	L. G.	Em	N A
<i>Kageneckia angustifolia</i> D. Don.	G.	Hyd	Chil.
<i>Kageneckia oblonga</i> Ruiz. & Pav	G.	Hyd	Chil.
<i>Licania hypoleuca</i> Benthm.	G.		New Grar
<i>Nuttallia cerasiformis</i> T & G	G. L.	Amvg Hyd	W N Am.
<i>Osteomeles arbutifolia</i> Lindl.	G.	Hyd.	
<i>Peraphyllum ramosissimum</i> Nutt.	G.	Hyd	N W Am.
<i>Poterium canadense</i> A. Gray	G.	Em	N Am.
<i>Poterium officinale</i> A. Gray.	L.	Astr	Eur N Am N As.
<i>Prunus amara</i>	Sm.	Ast	Asia
<i>Prunus Amygdalus</i> Stokes.	C. G. Sm.	Hyd	N Afr Eur W As. Orient
<i>Prunus Capulin</i> Zucc.	G.	Hyd.	Mex.
<i>Prunus caroliniana</i> Ait.	Ches.	Hyd.	N A
<i>Prunus cerasus</i> D Dietr	Ches.	Hyd	Ry Mts W N Am.
<i>Prunus domestica</i> L.	L.	Lax	Eur Cauc. Reg
<i>Prunus Laurocerasus</i> L.	Sm.Mi.C.L.	Hyd	Orient
<i>Prunus Mahaleb</i> L.	G.	Hyd	S Eur Orient
<i>Prunus persica</i> Stokes	G. C.	Ast Hyd	Temp As

NAME OF PLANT	Authority	Properties	Locality
<i>Prunus serotina</i> Ehrh.	G. Ches.	Hyd.	Temp. Reg.
<i>Prunus undulata</i> Ham.	G.	Hyd.	Eur N Am.
<i>Prunus virginiana</i> L.	G. C. L.	Hyd.	Him. Reg.
<i>Purshia tridentata</i> DC.	G.	Em.	E. N Am.
			Ry Mt. W N Am.
<i>Pygeum africanum</i> Hook.	G.	Hyd.	Tr Afr
<i>Pygeum parviflorum</i> Tejasm. & Binn.	G.	Hyd.	Java
<i>Pyrus</i> L.	G.	Hyd.	N Tem Reg
<i>Pyrus Aucuparia</i> Gaertn.	G.		Eur N As.
<i>Pyrus lanata</i> D Don	G.		Him. Reg.
<i>Pyrus Sorbus</i> Gaertn.			Eur N Afr
<i>Quilla, a saponaria</i> Molina	C. L.	Sap.	Chib.
<i>Rhodotypos hercoides</i> Sieb. and Zucc.	G.	Sap.	Spain.
<i>Rosa canina</i> L.	L.	Astr.	Eur Tem. As.
<i>Rosa galica</i> L.	L.	Astr.	Eur Cauc Reg.
<i>Rubus chamaemorus</i> L.	L.	Dir	N & Arc. Reg.
<i>Rubus cuneifolia</i> E. Merc.	L.	Astr.	Eur
<i>Rubus villosus</i> Ait.	G. L.	Sap. Mech.	E. N Am.
<i>Spiraea Aruncus</i> L.	G.	Sap.	N Tem. Reg.
<i>Spiraea bella</i> Sims.	G.	Sap.	Him. Reg.
<i>Spiraea canescens</i> D. Don	G.	Sap.	Him. Reg.
<i>Spiraea Filipendula</i> L.	G.	Astr.	N. As.
<i>Spiraea Humboldtii</i> Hort.	G.	Sap.	?
<i>Spiraea hypericifolia</i> L.	L.	Astr.	Rus. Tem. As.
<i>Spiraea japonica</i> L.	G.	Sap.	Japan.
<i>Spiraea laevigata</i> L.	G.	Sap.	Siberia
<i>Spiraea palmata</i> Pall.	G.	Sap.	Siber. China
<i>Spiraea tomentosa</i> L.	L.	Astr.	N A

OOIN

<i>Asperula cynanchica</i> L.	G.		Eur As Min.
<i>Asperula odorata</i> L.	G. L.	Cou.	Eur
<i>Basanacantha tetracantha</i> Hook.	G.	F P	Trop Am.
<i>Bothriospora corymbosa</i> Hook.	G.		S A
<i>Cephaelis toxica</i> A. St. Hil.	G.	Dras.	Mex. S. A.
			Galapagos Is.
<i>Cephalanthus occidentalis</i> L.	G.	Sap.	E. N A
<i>Chiococca</i> P. Br sp.	G.	Dras.	Mex S. A
<i>Chiococca</i> P. Br sp.	L.	Dras.	Trop. Am.
<i>Chiococca brachiata</i> Ruiz and Pav.	L.	Dir	Trop. Am.
<i>Chiococca racemosa</i> L.	L.	Dir	Ird.
<i>Cinchona</i> sp.	G. Wh.	Ir F. P	S. A.
<i>Coffea mauritiana</i> Lam.	G.		Mascar Is.
<i>Coffea odorata</i> Forst.	G.		Amic Is.
<i>Coprosoma linatufolia</i> Hook. f	G.		N Zea.
<i>Coptosapeita flavescens</i> North.	G.	A P	Malay
<i>Gahum Aparine</i> L.	L.	Dir	N Tem Reg
			Magellan.
<i>Gahum asprellum</i> Mx.	L.	Cou.	N A.
<i>Gahum triflorum</i> Mx.	G. L.	Cou.	N Tem Reg
			N A. Eur
<i>Geophila macropoda</i> DC.	G.	Em.	Peru.
<i>Geophila reniformis</i> G. Don.	G.		Peru
<i>Hameha patens</i> Jacq.	G.		Tr Am
<i>Mitchella repens</i> L.	G. L.	Sap.	E. N Am. Jap.
<i>Mitragyna speciosa</i> Korth.	G.		Borneo.
<i>Morelia senegalensis</i> A. Rich.	G.	F P	Tr Afr

NAME OF PLANT	Authority	Properties	Locality
<i>Mossacunda frondosa</i> L.	G.	Sap.	Malay Ind
<i>Oenanthe senegalensis</i> Hiern.	G.	Ant	Ind. Tr Afr
<i>Odenandia umbellata</i> L.	L.	Ant	India
<i>Paederia foetida</i>	Sm.		Ind. Malay
<i>Palaucorea rigida</i> HBK			S. Am.
<i>Pavetta reticulata</i> Bl.	G.	Em	Java.
<i>Plectronia discocoe</i> Burck.	G.	Hyd	
<i>Psychotria emetica</i> L.	L.	Em	N Gran.
<i>Psychotria Ipecacuanha</i> Stokes	G Wh. L.	Dep Ein.	Braz
<i>Randia aculeata</i> L.	Sm.		W Ind
<i>Randia dumetorum</i> La.	G. L.	F P	Tr Old World.
<i>Sickungia rubra</i> Mart.	G.	Aik	Braz.
<i>Spermacoce capitata</i> Willd.	G.		Peru
<i>Spermacoce semierecta</i> Roxb.	G.	Coa.	Sumatra
<i>Tricalypsa Sonderiana</i> Hiern.	G.	P	Tr Afr.
<i>Vangueria spinosa</i> Roxb.	G.		Tr As

RUTACEAE

<i>Aceronychia laurifolia</i> Bl.	G.	F P	Tr As.
<i>Caesalpinia edulis</i> L.	G. L.	Nar	Mex.
<i>Chosya ternata</i> HBK.	G.	Sap.	Mex.
<i>Citrus Aurantium</i> L.	C.		Tr As.
<i>Citrus medica</i> L.	G. Wh.	Hyd.	Tr As.
<i>Cuscuta febrifuga</i> H & B.	L.	P P	New Gran
<i>Cuscuta toxicaria</i> Engl.	G.		Braz.
<i>Eyod-a rutacarpa</i> Hook.	L.	Pur	Him. Reg
			China. Japan
<i>Lumnitz philippensis</i> Planch.	G.	A P	Philipp.
<i>Melicope erythrococca</i> Benth.	G.		Annar
<i>Peganum antidysentericum</i> Kostel.	L.	Astr	S Afr
<i>Peganum Harmala</i> L.	G. C. L.	Ant	Cent. As
<i>Pilocarpus</i> sp.	G.		W Ind. S Am.
<i>Pilocarpus officinalis</i> Poehl.	L.	Diaph.	S. A
<i>Pilocarpus pennatifolius</i> Lem.	Sm. Wh.	Dep. Ir	Braz. I.
<i>Pilocarpus racemosus</i> Vahl.	L.	Diaph	W Ind.
<i>Pilocarpus spicatus</i> A. St. Hil.	L.	Diaph	Brazil
<i>Ptelea trifoliata</i> L.	G.	Sap.	E. N Am
<i>Ruta graveolens</i> L.	G.C.Sm.L.	Abort Ir	S. Eur
<i>Ruta montana</i> Mill.	G.	Abort.	Med. Reg. Cau.
<i>Skimmia japonica</i> Thunb.	G.	P.	Japan.
<i>Thevetia Ahouai</i> A DC			Braz.
<i>Thevetia peruviana</i> Juss.			
<i>Zanthoxylum alatum</i> Roxb.	G.	F P	Him. Reg Chin.
<i>Zanthoxylum caribaeum</i> Lam.	G.		New Grau.
<i>Zanthoxylum Naranjillo</i> Griseb.	G. L.	Drur	Argent.
<i>Zanthoxylum scandens</i> Bl.	G.		Java

SALICACEAE

<i>Populus balsamifera</i> L.	Halsted	Tr	E. N Am Asia
-------------------------------	---------	----	--------------

SALVADORACEAE

<i>Salvadora persica</i> L.	G.		Orient. Ind N. Afr
-----------------------------	----	--	--------------------

SAMYDACEAE

NAME OF PLANT	Authority	Properties	Locality
<i>Casearia graveolens</i> Dal.	G.	F P	Ind
<i>Casearia guineensis</i> G. Don.	G.	F P	Tr As.
<i>Casearia tomentosa</i> Roxb.	G.	F P	Tr As Austr

MADAGASCARIACEAE

<i>Exocarpus cypressiformis</i> R. Br.	G. M.		Austr
--	-------	--	-------

SAPINDACEAE

<i>Alectryon excelsum</i> Gaert.	G.	Hyd.	N Zeal
<i>Blighia sapida</i> Kon.			Afr W Ind.
<i>Cardiospermum Halimifolium</i> L.	G. L.	Sap.	Tr Reg N A.
			S. States
<i>Cupania</i> sp.	G.		
<i>Cupania Pseudorhus</i> A. Rich.	G.	F P	Austr
<i>Ditelaia Karak</i> DC.	G.	F P	Malacca
<i>Dodonaea physocarpa</i> Fy. Muell.	G.		Austr
<i>Dodonaea viscosa</i> Jacq.	G. R.	F P	Cosmop Tr
<i>Ganophyllum falcatum</i> Bl.	G.	Sap.	Malacca.
<i>Harpulia arborea</i> Radlk.	G. R.	F P	
<i>Harpulia capensis</i> Roxb.	G. R.	F P	Ind
<i>Harpulia thurifera</i> Bl.	G.		N Cum
<i>Koeleria paniculata</i> Laxm.	G.		China
<i>Magonia glabrata</i> A. St. H.	G. R.	Sap. F P	Braz.
<i>Magonia pubescens</i> St. H.	G. R. E.	Sap. F P.	Braz.
<i>Nephetum lappaceum</i> L.	G.		Malay
<i>Nephetum Longana</i> Camb.	G.		Ind, Burma.
<i>Paulinia costata</i> Sch. recht	R. G.	Sap. F P	Mex
<i>Paulinia Cupania</i> HBK.	G. R. L.	Sap. F P.	Venez.
<i>Paulinia curassavica</i> Jacq.	R.	Sap. F P	W Ind
<i>Paulinia macrophylla</i> Kunth.	G. R.	Sap. F P	N Gran
<i>Paulinia menziesii</i> Juss.	G.	Sap. F P	Braz.
<i>Paulinia thurifera</i> Juss.	G.	Sap. F P.	Trop. Amer
<i>Paulinia trigona</i> Vell.		Sap. F P	Brazil
<i>Sapindus abyssinicus</i> Fres.	G.	Sap. F P	Abyss.
<i>Sapindus aborescens</i> Aubl.	G.	F P	Indian.
<i>Sapindus marginatus</i> Willd.	Sm. L.	Sap.	N Am. S. Sta.
			Tex
<i>Sapindus Mukorossi</i> Gaertn.	G.	F P	Trop. Asia
<i>Sapindus Saponaria</i> L.	G. L. R.	F P	N & S Am.
<i>Sapindus trifolatus</i> L.	G. L.	F P	Trop. As.
<i>Scheuchzeria</i>	G.	Hyd.	
<i>Serjania acuminata</i> Radlk.	G. R.	Nar	Braz.
<i>Serjania curassavica</i> Radlk.	R. E.	Sap. F P	
<i>Serjania cuspidata</i> Cambess.	G.	Sap.	W Ind.
<i>Serjania erecta</i> Radlk.	E & G. R.	Sap. F P	Brazil.
<i>Serjania ichthyocarpa</i> Radlk.	E. R.	Sap. F P	Brazil.
<i>Serjania meibomia</i> Radlk.	E & G. R.	Sap. F P	Costa Rica.
<i>Serjania orthocarpa</i> A. St. H.	E. G. R. L.	Sap. F P	Brazil.
<i>Serjania mexicana</i> Willd.	L.	Sap. F P	Mex.
<i>Serjania nodosa</i> Radlk.	G. R. E.	Sap. F P	W Ind
<i>Serjania piscatoria</i> Radlk.	E & G. R.	Sap. F P	Brazil
<i>Serjania polyphylla</i> Radlk.	G. R.	Sap. F P	Gu an.
<i>Talsia stricta</i> Triana & Planch.	G.	Sap.	New Granad.
<i>Ungradia speciosa</i> Endl.	G. L.	Sap.	Texas.

SAPOTACEAE

NAME OF PLANT	Authority	Properties	Locality
<i>Achras sapota</i> L.	G. L.	Sap.	S Am
<i>Bassia butyracea</i> Roxb.	G.	P. P.	Ind.
<i>Bassia latifolia</i> Roxb.	G.		Ind.
<i>Bassia longifolia</i> L.	L.	Astr.	Ind.
<i>Bassia Mottleyana</i> Miq.	G.	Hyd.	Malay
<i>Ilipe Macclayana</i>	G.		New Guin
<i>Lucuma Bonplandia</i> HBK.	G.		Cuba.
<i>Lucuma deliciosa</i> Planch & Lander.	G.	Hyd.	New Gran
<i>Lucuma glycyphloca</i> Mart.	L.	Astr.	Braz
<i>Lucuma mammosa</i> Gaertn.	G.	Hyd.	S Am.
<i>Lucuma multiflora</i> A. DC.	G.	Hyd.	W In.
<i>Lucuma saucifolia</i> HBK.	L.	Sap.	Mex
<i>Omphalocarpum procerum</i> Beauv.	G.		Tr. As.
<i>Pavonia latifolia</i> Burck.	G.	Sap.	Sunda Is.
<i>Sideroxylon borbonicum</i> A. DC.	G.		Barbon Is.
<i>Sideroxylon dulcificum</i> A. DC.	G.		Tr Afr
<i>Sideroxylon toxiferum</i> Thunb.	G.	A. P.	

SARRACENIACEAE

<i>Sarracenia flava</i> L.	L.	Astr.	N. A.
<i>Sarracenia purpurea</i> L.	G. L.	Astr.	E. N. Am.
<i>Sarracenia variolaria</i> Mx.	L.	Astr.	N. A.

SAURURACEAE

<i>Saururus cernuus</i> L.	G. L.	Sap.	E. N. Am.
----------------------------	-------	------	-----------

SAXIFRAGACEAE

<i>Aphanopetalum resinosum</i> Endl.	G.	Alk.	Austr.
<i>Calceema serratifolia</i> Andr.	G.	Sap.	
<i>Ceratopetalum apetalum</i> D. Don.	G.	Con.	Austr.
<i>Chrysosplenium alternifolium</i> L.	G.	Pot.	N. As. E. N. Am.
<i>Chrysosplenium americanum</i> Schw.		P.	N. Am.
<i>Deutzia staminea</i> R. Br.	G.	Sap.	H. m. Reg.
<i>Dichroa febrifuga</i> Lour.	G.	Em.	H. m. Reg.
<i>Escallonia myrtilloides</i> L. f.	G.		Malay China.
<i>Francoa appendiculata</i> Cav.	G.		New Gran.
<i>Hydrangea arborescens</i> L.	G. L.	Alk.	Chili
<i>Hydrangea Hortensia</i> Sieb.	G.	Sap. Hyd.	N. Am.
<i>Hydrangea involucrata</i> Sieb.	G.	Hyd.	Asia. Or. ent.
<i>Hydrangea Lindleyana</i>	G.	Hyd.	Japan.
<i>Hydrangea Thunbergii</i> Sieb.	G.	Hyd.	Japan.
<i>Jamesia americana</i> Torr. & Gray		Hyd.	Ry. Mt. N. Am.
<i>Philadelphus coronarius</i> L.	G.	Sap.	S. Eur.
<i>Philadelphus grandiflorus</i> Willd.	G.	Sap.	N. Am.
<i>Philadelphus Lemonei</i>	G.	Sap.	N. Am. Ry. Mts.
<i>Philadelphus Lewisii</i> Pursh.	G.	Sap.	
<i>Philadelphus microphyllus</i> A. Gray	G.	Sap.	N. Am. N. Mex.
<i>Ribes aureum</i> Pursh.	G.		Ry. Mts. W. N. Am.
<i>Ribes cereum</i> Dougl.	G. Sm.	Intox.	N. Am. Ry. Mts.
<i>Ribes macrobotrys</i> Ruiz. & Pav.	Sm.		Peru.
<i>Ribes nigrum</i> L.	L.	Dur.	Eur. N. As.
<i>Ribes prostratum</i> L. Her.	Sm.		N. Am.

NAME OF PLANT	Authority	Properties	Locality
<i>Saxifraga Andrewsii</i> Harv.	G.	Sap.	
<i>Saxifraga cortusaefolia</i> Sieb. & Zucc.	G.	Sap.	Japan.
<i>Saxifraga cuneifolia</i> L.	G.	Sap.	Spain.
<i>Saxifraga Sibthorpii</i> Boiss.	G.	Sap.	Greece

SCITAMINEAE

<i>Kaempferia rotunda</i> L.	Kirtikar		Tr As
------------------------------	----------	--	-------

SCROPULARIACEAE

<i>Reyrichia scutellarioides</i> Benth.	G.	Em.	Brazil.
<i>Bonnaya veronicaefolia</i> Spreng.	G.	Ant.	Malay China,
<i>Calceolaria scabiosaefolia</i> Sims.		Em.	Peru.
<i>Capriaria biflora</i> L.	G.		Trop. Am.
<i>Castilleja canescens</i> Benth.	G.	Car	Mex
<i>Chelone glabra</i> L.	G. L.	Ant.	E. N. Am.
<i>Digitaria ambigua</i> Murr	C. L.	Card.	Eur. W As.
<i>Digitaria lutea</i> L.	C.		S. Eur
<i>Digitaria purpurea</i> L.	G. Mi. Sm.	Card. Ast	Eur. W N. Am.
	Ches. L.		
<i>Digitaria Thapsi</i> L.	E. C R C	F P	Spain.
<i>Gerardia tenuifolia</i> Vahl	Ches.		E. N. Am.
<i>Gratiola officinalis</i> L.	M. G. C Sm	tr	Eur
	Ches. L.		
<i>Gratiola peruviana</i> L.	G.		Trop. Am Aus.
			New Zeal.
<i>Limosella aquatica</i> L.	G.	Sap.	N & S Temp.
			Reg.
<i>Limosella Cymbalaria</i> Mill.	C		Eur.
<i>Limosella Elatine</i> Mill.	C		Eur. Orient
<i>Limosella spuria</i> Mill.	C		Eur. N Afr.
<i>Linaria vulgaris</i> Mill.	C. Sm Mi. L.	lr	Eur N. Am.
	Rusby		
<i>Melampyrum arvense</i> L.	C. L.		Caucasus
<i>Melampyrum silvaticum</i> L.	G. L.	Thur.	Eur N As.
<i>Pedicularis palustris</i> L.	G. M. Ches.		N Tem. & Arc.
			Reg
<i>Pedicularis sylvatica</i> L.	Mi.		Eur
<i>Rhinanthus major</i> Ehrh.	C.		Eur
<i>Rhinanthus minor</i> Ehrh.	C. L.	Ins	Eur
<i>Scrophularia aquatica</i> L.	G Sm. C. L.	Card.	Eur. Cauc.
<i>Scrophularia nodosa</i> L.	G Sm. C. L.	Card.	N Temp. Reg.
<i>Striga euphrasioides</i> Benth.	G.	P	India.
<i>Vandellia crustacea</i> Benth.	G.	Card.	Coast. Trop
<i>Vandellia minuta</i> Miq.	G.	Em.	Java.
<i>Verbascum Blattaria</i> L.	G.		Eur N As
			N Am
<i>Verbascum crassifolium</i> Hoffm.	G.		Lusitania.
<i>Verbascum dubium</i> Roem & Schult.	G.		Lusitania
<i>Verbascum orientale</i> Bich.	E. R.	F P	Caucas.
<i>Verbascum phlomoides</i> L.	E. R.	F P	Eur As. Min.
<i>Verbascum pulverulentum</i> Vill.	G.		S. Eur
<i>Verbascum simplex</i> Labill.	E. G. R.	F P	Med. Reg. Ori.
<i>Verbascum sinuatum</i> L.	G.		Syria.
<i>Verbascum thapsoides</i> L.	G.		Eur
<i>Verbascum Thapsus</i> L.	E. G. R. C. L.	F P.	Eur N Am.
			Ori. H. m. Reg.

NAME OF PLANT	Authority	Properties	Locality
<i>Veronica Beccabunga</i> L.	L.	Diur	N Temp. Reg. Eur. As. Min.
<i>Veronica officinalis</i> L.	L.	Diur	N As.
<i>Veronica virginica</i> Nutt.	G. L. Rusby	Car. Em.	E. N. Am.

SELAGINELLACEAE

<i>Globularia Alypum</i> L.	G. L.	Pur	Med. Reg.
-----------------------------	-------	-----	-----------

SIMARUBACEAE

<i>Allanthus glandulosa</i> Desf.	G. Rusby	Tacn.	China
<i>Balanites aegyptiaca</i> Deble.	L. G.	Sap.	N. Afr. Arab. Palest.
<i>Balanites Roxburghii</i> Planch.	G.	Sap.	Ind.
<i>Brucea sumatrana</i> Roxb.	G. L.		Trop. As. Aust.
<i>Cneorum tricocctum</i> L.	C.		S. Eur.
<i>Picrasma excelsa</i> Lindl.	L.	Ins.	W. Ind.
<i>Picrasma quassoides</i> Benn.	G. L.	Ins.	Him. Reg. China
<i>Samadera indica</i> Gaertn.	G. L.		Ind.
<i>Simaba Waldenii</i> Planch.	G.		
<i>Simaruba amara</i> Aubl.	Sm.		Trop. Am.

SOLANACEAE

<i>Aenictus arborens</i> Schlecht.	G.	Nar	Mexico.
<i>Anthocercis</i> Lab. sp.	G.		Austr.
<i>Atropa Belladonna</i> L.	G. C. M. Sm. L. Rusby	Del.	Eur. Ori. Ind.
<i>Brunfelsia Hopeana</i> Benth.	G. L.	A. P.	Brazil.
<i>Capsicum annuum</i> L.	Sm. L.	Ir.	Trop. Reg. Mex. S. Am.
<i>Capsicum frutescens</i> L.	Sm. L.	Ir.	Tex. Mex. Trop. Reg.
<i>Capsicum minimum</i> Blanco.	Wh. L.	Ir.	Philipp. Is.
<i>Cestrum aurantiacum</i> Lindl.	Sm.	Nar	Guatemala.
<i>Cestrum auriculatum</i> L'Her.	G.	Nar	S. Am.
<i>Cestrum macrophyllum</i> Vent.	G.		W. Ind.
<i>Cestrum nocturnum</i> L.	G.		S. Am.
<i>Cestrum paniculatum</i> Lam.	G.		Jamaica.
<i>Cestrum Parqui</i> L'Her.	G.		Trop. Amer.
<i>Cestrum vespertinum</i> L.	G.		W. Ind.
<i>Datura arborea</i> L.	G. C.		S. Amer.
<i>Datura fastuosa</i> L.	G. L.	F. P.	Tr. O. d. World
<i>Datura ferox</i> L.	C.		China
<i>Datura Metel</i> L.	G. C. L.	Nar	Trop. Am.
<i>Datura meteloides</i> DC.	L.	Nar	W. N. A.
<i>Datura sanguinea</i> Riz. & Pav.	G. L.	Nar P.	S. Am.
<i>Datura Stramonium</i> L.	C. G. Rusby Sm. M. Ches. L.	Hyp.	Cosm. Trop.
<i>Datura suaveolens</i>	C.	Hyp.	
<i>Datura Tatula</i> L.	Ches. Rusby	Hyp.	Cosm. Trop.
<i>Duboisia Hopwoodii</i> F. Muell.	Blyth. L.	Hyp.	Austr.
<i>Duboisia myoporoides</i> R. Br.	G. M. L.	Card.	Austr.
<i>Hyoscyamus albus</i> L.	G. L.	Hyp. Del.	Orient.

NAME OF PLANT	Authority	Properties	Locality
<i>Hyoscyamus aureus</i> L.	C.		A. Min. Mesop.
<i>Hyoscyamus Falezler</i> Coss.	C.		W. As.
<i>Hyoscyamus muticus</i> L.	C. L.	Nar	Med. Reg.?
<i>Hyoscyamus niger</i> L.	R Ches Sm. M. Rusby E. G. L.	Hyp. Del.	Eur W. As. Ium Reg.
<i>Hyoscyamus physaloides</i> L.	G.	Hyp. Del.	Siberia.
<i>Hyoscyamus reticulatus</i> L.	G.	Hyp. Del.	As. Min Egypt.
<i>Lathraea venenosa</i> Phil.	G.		Chil.
<i>Lycium barbarum</i> L.	G.		Orient.
<i>Lycopersicon esculentum</i> Mill.	G. M. Sm. C. L.	Sap	S. Am Mex Tex Tem Reg.
<i>Mandragora autumnalis</i> Bertol.	Sm.		Med. Reg.
<i>Mandragora officinarum</i> L.	C. G. L.	Del.	Med. Reg.
<i>Nicotiana physaloides</i> Gaertn.	G.	P. berry	Peru
<i>Nicotiana glauca</i> Link & Otto.	C.		Bras.
<i>Nicotiana glauca</i> Grab.	C.		China.
<i>Nicotiana glauca</i> Grab.	C. G. M.		Arg.
<i>Nicotiana quadrivalvis</i> Pursh.	C.		W. N. Am.
<i>Nicotiana rustica</i> L.	C. L.	Naus	Mexico
<i>Nicotiana suaveolens</i> Lehm.	G. M.		Australia
<i>Nicotiana Tabacum</i> L.	Rusby C. R. Ches. G. L. E.	Naus 1- P	S. Am Temp. Reg.
<i>Nierembergia hippomanica</i> Miers.	G.		Arg. Rep.
<i>Physalis Alkekengi</i> L.	L.	Dur.	Eur Japan
<i>Physalis foetida</i> Pour.	G.	Nar.	Trop. Am.
<i>Physalis virginiana</i> Mill.	G.	Nar	E. N. Am.
<i>Physoclaina orientalis</i> G. Don.	C. G.		Orient.
<i>Physoclaina praecox</i> Miers.	G.	P.	Him. Reg.
<i>Scopolia carniola</i> Jacq.	C. L.	Atropin	Eur
<i>Scopolia japonica</i> Max.	G. L.	Atropin P	Japan
<i>Scopolia lurida</i> Dun.	G.	P.	Him. Reg.
<i>Scopolia physaloides</i> Donl.			Siberia Altai
<i>Solanandra grandiflora</i> Sw.	G.	Alk.	Trop. Amer.
<i>Solanum aculeatum</i> Jacq.	G.		Tr. As. & Am.
<i>Solanum aviculare</i> Forst.	G.		Aus. & N. Zeal.
<i>Solanum caripense</i> Hamb. & Bopl.	G.		Venez.
<i>Solanum carolinense</i> L.	L.	Dur.	N. A.
<i>Solanum crispum</i> Ruiz & Pav.	Sm.	Del.	Chil.
<i>Solanum Dulcamara</i> L.	Ches. Rusby L.	Sol. Nar	Eur. N. Am. Austr.
<i>Solanum ellipticum</i> R. Br.	G.		Austr.
<i>Solanum esuriale</i> Lindl.			Austr.
<i>Solanum eremophilum</i> Fy. M.	G. M.		Austr.
<i>Solanum grandiflorum</i> R. & P.	G.		Peru
<i>Solanum mammosum</i> L.	G.		Trop. Am.
<i>Solanum Melongena</i> L.	C. Sm.		Tr. Old World
<i>Solanum nigrum</i> L.	G. M. Rusby C. M. Ches. L.	Nar	Austr. N. Am. Eur. N. Am. As.
<i>Solanum saponaceum</i> Dun.	G.		Peru.
<i>Solanum sphaerocarpum</i> L.	G. Sm.		Trop. Old World Cald.
<i>Solanum Sturtianum</i> F. & M.	G. M.		Austr.
<i>Solanum stramonifolium</i> Jacq.	G.		Cosm. Trop.
<i>Solanum torum</i> Sw.	G.		Cosm. Trop.

NAME OF PLANT	Authority	Properties	Locality
<i>Solanum triflorum</i> Nutt.	G. Chea.		W N Am. Ry
<i>Solanum tuberosum</i> L.	M. Rusby	Sol.	Mts. Neb.
<i>Solanum villosum</i> Willd.	Chea.		S Am Cult
<i>Solanum Xanthi</i> Gray	Mi.		Temp. Reg
<i>Triguera ambrosiaca</i> Cav.	G.		Eur As. Afr
<i>Vestia lycioides</i> Willd.	G.		Calif
<i>Withania somnifera</i> Dun.	G.	Abort.	Spain.
	G.		Chili
	G.		Med. Reg Ori

STERCULIACEAE

<i>Pterospermum diversifolium</i> Bl.	G.	Ir	India Malay
<i>Sterculia alata</i> Roxb.	G.	Nar	India.

TERNSTROEMIIACEAE

<i>Cameilla japonica</i> L.	G.	F P	Japan, China.
<i>Cameia Sasanqua</i> Thunb.	G.	F P	Japan.
<i>Carapa fasciculata</i> Cambora.	G.		Guiana.
<i>Caryocar glabrum</i> Pers.	G.		
<i>Lianosa Toquian</i> Blanco.	G.	F P	Pacific Isl.
			Guiana.

THYMELAEACEAE

<i>Daphne Genkium</i> L.	L.	Ir	Med Reg
<i>Daphne Laureola</i> L.	L.G.C.M.	Ir F P	Eur N As.
<i>Daphne mezereum</i> L.	L.	Ir	Eur N As.
<i>Daphne striata</i> Tratt	G.		Alp S Eur
<i>Daphnopsis castrifolia</i> Meiss.	G.		N Great
<i>Daphnopsis Cneorum</i> L.	G.R.M.E.	F P	Eur Austr
<i>Daphnopsis Genkium</i> L.	G.R.C.E.	F P Ir	Med Reg.
<i>Daphnopsis oleoides</i> Schreb.	Brandt	Ir	Eur As. Min.
<i>Dicra palustris</i> L.	G. L.	Sap.	E N Am
<i>Edgeworthia Garconeri</i> Meiss.	G	F P	Imm. Reg Chi.
			Japan.
<i>Lasiadenia rupestris</i> Benth.	G	P	Guinea
<i>Lasiophon anthyllioides</i> Vess.	G.	P	S Afr
<i>Lasiophon eriocarpus</i> D. C.	G.	P	India.
<i>Thymelaea Tartaronia</i> All.	G.	Ir	S. Eur
<i>Wikstroemia Chamaedaphne</i> Meiss.	G.	F P	China.
<i>Wikstroemia viridiflora</i> Meiss.	G R	F P	Tr As. Austr

TILIACEAE

<i>Corchorus capsularis</i> L.	G.		Trop. Cosm.
<i>Echinocarpus Sigun</i> Bl.	G.	Hyd	Java
<i>Grewia asiatica</i> L.	G. R.	F P	India.
<i>Grewia bracteata</i> Roth.		F P.	India.
<i>Grewia Maroccoca</i> L.	G. R.	F P	Amicor Is
<i>Grewia orientalis</i> L.	G.		Tr As & Aus
<i>Grewia pilosa</i> Roxb.	G.		Ind Trop. Afr
<i>Grewia piscatorum</i> Hance			China
<i>Prockia theaeformis</i> Willd.	G.	F P	Mascar Is.

TROPAEOLACEAE

<i>Tropaeolum majus</i>	Wh.	Ir	Peru
-------------------------	-----	----	------

TYPHACEAE

<i>Typha latifolia</i> L.	G.	P.	N Temp. Reg.
---------------------------	----	----	--------------

ULMACEAE
UMBELLIFERAE

NAME OF PLANT	Authority	Properties	Locality
<i>Aethusa Cynapium</i> L.	G.C.Mi.Sm.	Ir	Eur Orient.
<i>Ammi Visnaga</i> Lam.	G. L.	Dur	Med. Reg
<i>Anthriscus cerefolium</i> Hoffm.	L.	Dur	Eur N Asia
<i>Anthriscus sylvestris</i> Hoffm.	L.		Eur N As.
			Can Reg
<i>Anthriscus vulgaris</i> Bern.	G. C. L.		Eur Orient
<i>Apium graveolens</i> L.	Mi. L.	Ir	Eur Ori. Ind.
			Calif
<i>Apium leptophyllum</i> F. & M.	G. M.		Austr
<i>Apium nodiflorum</i> Reischb.	Sm.	Ir	
<i>Azorella Glehbaria</i> A. Gray	G.	P	
<i>Carum capense</i> Sond.	G.		S Afr
<i>Carum copticum</i> Benth. & Hooker	G.		Eur N Afr
			N Asia.
<i>Carum Petroelinum</i> Benth & Hooker	C. L.	Ir	Eur.
<i>Caucasia daucoides</i> L.	C.		Eur Temp. As
<i>Chaerophyllum temulum</i> L.	G.		Eur N Afr
			Caucas Reg
<i>Cicuta Bolanderi</i> A. Gray	Ches.	Con.	Calif
<i>Cicuta bulbifera</i> L.	Rusby	Con.	N A
<i>Cicuta californica</i> A. Gray	Rusby	Con.	Calif
<i>Cicuta maculata</i> L.	Rusby	Con.	
	G. Ches. L.		E N Am
<i>Cicuta occidentalis</i> Greene	Ches.	Con.	W N Am Ry.
			Mts.
<i>Cicuta vagans</i> Greene	G. Ches. L.	Con.	W N Am.
<i>Cicuta venenata</i> Nutt	G.	Con.	W N Am
<i>Cicuta virosa</i> L.	G.Sm.C Mi.	Con.	Eur N Am.
	L. Rusby		
<i>Conium maculatum</i> L.	L. Mi.	Dep.	Eur Ori N Am
	Rusby G.		
	Sm Ches.C.		
<i>Conopodium denudatum</i> Koch.	G.		Dur
<i>Coriandrum sativum</i> L.	G. L.		S Eur Ori
<i>Crithmum maritimum</i> L.	G.	Ant	Eur
<i>Daucus Carota</i> L.	C. L.	Dur	Eur Ori N Am.
<i>Elaeagnum ascepium</i> Boiss	Sm.	Ir	S Eur
<i>Elaeagnum foetidum</i> Boiss	Sm.	Ir	S. Eur N Afr
<i>Ferula communis</i> L.	G.		Med Reg
<i>Ferula foetida</i> Regel	L.	Antisp.	Turkest
<i>Ferula gabariflua</i> Boiss. & Buhse	Wh. L.	Ir	Persia
<i>Heraeleum lanatum</i> Michx.	G. L.	Acr	Eur N Am
			W As.
<i>Heraeleum Spondylium</i> L.	C. L.	Acr	Eur N As.
<i>Heteromorpha arborescens</i> Cham. & Schl.	G.		S Afr
<i>Hydrocotyle asiatica</i>	L.	Nar	
<i>Hydrocotyle javanica</i> Thunb.	G.	F P	Java
<i>Hydrocotyle umbellata</i> L.	G.	Ent	Trop Reg
<i>Hydrocotyle vulgaris</i>	Sm. Mi.		Eur
<i>Lichstenstemma Beinhana</i> Eckl. & Zeyh.	G.		S Afr
<i>Lebsten cimia pyrethifolia</i> Cham. & Schlecht	G.		S Afr
<i>Molopospermum cicutarium</i> DC	G.	Nar	Eur
<i>Oenanthe crocata</i> L.	G. C. L.	Ir	Eur
<i>Oenanthe fistulosa</i>	C. Mi.	Ir	Eur Casp. Reg.
<i>Oenanthe Lachenalia</i> G. Gmel.	G.		Eur Casp. Reg.

NAME OF PLANT	Authority	Properties	Locality
<i>Oenanthe Phellandrium</i> Lam.	G. Sm. C. L. Mt.	Ir	Eur N. As.
<i>Oxypolis rigidus</i> Britton.	Ches.		E. N. A.
<i>Pastinaca sativa</i> L.	C. L.	Diur	Eur N. A.
<i>Peucedanum ambiguum</i> Nutt.	G.		W. N. A.
<i>Peucedanum foeniculaceum</i> Nutt.	L.		W. N. A.
<i>Peucedanum Ostruthium</i> Koch.	L.	Acc	Eur
<i>Pimpinella Saxifraga</i> L.	L.	Naus.	Eur
<i>Sanicula marilandica</i> L.	G. L.	Astr	E. N. A.
<i>Sium cicutaeformis</i> Schrank.	L. Ches.		N. Temp. Reg.
<i>Sium erectum</i> Huds.	G. Sm. Mt.	Ir	Eur. Austr. Reg.
<i>Sium latifolium</i> L.	Mt. C. Ches.		Eur N. A.
<i>Thapsia garganica</i> L.	G. L.	Ir	Med. Reg.
<i>Thapsia villosa</i> L.	G.	Ir	S. Eur
<i>Trachymene australis</i> Benth.	G. M.		Austr.

URTICACEAE

<i>Antiaris toxicaria</i> Lesch.	Sm. L.	Ast. A. P.	Ind. Malay
<i>Artocarpus calophylla</i> Z. & M.		F. P.	Tennasserim
<i>Cannabis sativa</i> Lam.	Mt. Sm. L.	Def.	Cent. As. N.
	Rushy		Ham. Eur. N. Am.
<i>Ficus a. ussima</i> Bl.	G.		Tr. Asia, Malay
<i>Ficus Carica</i> L.	G. L.	Lax.	S. Eur. Orient
			Afr. N. & S.
<i>Ficus hispida</i> L.	G.	P.	Am. Austr.
			Trop. As. & Austr.
<i>Ficus hypogaea</i> .	G.		
<i>Ficus leucartatomia</i> Poit.	G.		Malaya
<i>Ficus Roxburghii</i> Wall.	Sm.	Ast.	Sim. Burma
<i>Humulus Lupulus</i> L.	Mt. L.	Sed.	
<i>Laportea canadensis</i> Gaud.	Wh. L.	Ir.	N. Am.
<i>Laportea crenulata</i> Gaud.	L.	Ir.	Trop. As.
<i>Laportea Gaudichaudiana</i> Webb.	L.	Ir.	Phil. Is.
<i>Laportea stimulans</i> Mtg.	G. L.	Ir.	Java
<i>Trema aspera</i> R. Br.	M.		Austr.
<i>Urtica dioica</i> L.	Sm. Mt. L.	Ir.	N. Temp. Reg.
	Ches.		
<i>Urtica gracilis</i> Ast.	Ches. L.	Ir.	E. N. Am. Ry
			Mta.
<i>Urtica holosericea</i> Nutt.	Pam.	Ir.	Ry Mta. W. N.
			Am.
<i>Urtica pilulifera</i> L.	Sm. L.	Ir.	Orient. Med.
			Reg.
<i>Urtica urens</i> L.	Mt. L.	Ir.	Tem. old world

VALERIANACEAE

<i>Valeriana dioica</i> L.	C.		Eur.
<i>Valeriana officinalis</i> L.	Sm. C. L.	Ir.	Eur. N. As.

VERBENACEAE

<i>Callicarpa americana</i> Lam.	G.		N. Am. Tex.
<i>Callicarpa cana</i> L.	G.		Trop. As. Austr.
<i>Callicarpa longifolia</i> Lam.	G.	F. P.	Malay, Austr.
<i>Callicarpa tomentosa</i> Willd.	G.		China
<i>Clerodendron infortunatum</i> Gärt.	G.		Malay

NAME OF PLANT	Authority	Properties	Locality
<i>Clerodendron Siphonanthus</i> R. Br.	G.		Ind & Malay
<i>Duranta Plumieri</i> Jacq.	G.	Sap.	Trop. Am.
<i>Lantana salicariifolia</i> Jacq.	G.		S. Afr
<i>Stachytarpheta indica</i> Vahl.	G. L.	Abort.	Cosm Trop.
<i>Verbena hastata</i> L.	L.	Ir	N A.
<i>Verbena officinalis</i> L.	Sm. L.	Ir	Amphig. temp.
<i>Verbena venosa</i> Gilb. & Hook	Sm.	Ir	Arg. Reg.
<i>Vitex</i> sp.	G.		S. Am. Afr As.
			Austr
<i>Vitex pteropoda</i> Miq.	G.		Malay
<i>Vitex vestita</i> Wall.	G.		Malay

VIOLACEAE

<i>Anchetea salutaris</i> St. Hil.	G. L.	Em.	Brazil
<i>Ionidium Ipecacuanha</i> Vent.	L.	Em.	Brazil
<i>Noisetia</i> HBK.	G.	Em.	S. Am.
<i>Viola</i>	G.	Em.	N Temp. Reg.
<i>Viola odorata</i> L.	C. L.	Em.	N Afr & As.
<i>Viola pedata</i> L.	L.	Em.	N A.
<i>Viola sepiicola</i> Jord.	C		Eur

VITACEAE

<i>Cissua nivea</i> Hochst.	G.		Abyss.
<i>Cissua pruriens</i> Planch.	G.		
<i>Vitis hederacea</i> Ehr.	L.	Ir	E. N. Am. Ry.
(<i>Ampelopsis quinquefolia</i> Michx.)			Mta.
<i>Pseodera quinquefolia</i> (L.) Greene)	Hala.		
<i>Vitis elongata</i> Wall.	G.		Ind.
<i>Vitis Mnabau</i>	G.		
<i>Vitis Saponaria</i> Seem.	G.		Austr. Pac. Is.
<i>Vitis sessilifolia</i> Baker	B.		Brazil

XYRIDACEAE

<i>Xyris communis</i> Kunth			Trop. Am.
-----------------------------	--	--	-----------

ZINGIBERACEAE

<i>Glozza Beaumetzii</i> Heck.	G.	Tacn.	
<i>Hedychium longecornutum</i>	G.	Ant.	
<i>Thaumatococcus Danieli</i> Benth.	G.		Trop. Afr.
<i>Zingiber officinale</i> Rose.	Sm.	Ir	India

ZYGOPHYLLACEAE

<i>Guaiacum officinale</i> L.	G. L.	Sap.	India
<i>Larrea mexicana</i> Moric.	G. L.	A. P.	Mex. S.W. U.S.
<i>Nitraria tridentata</i> Desf.	G.	Inch.	Syria, N. Afr. & Tropics
<i>Tribulus cistoides</i> L.	G.		Cosm. Trop.
<i>Tribulus maximus</i> L.	G.	Sap.	N. Am. Guiana
<i>Zygophyllum coccineum</i> L.	G.		N. Am. Arab.
<i>Zygophyllum Fabago</i> L.	G.		Spain, N. Afr.
			W. Asia
<i>Zygophyllum yodocarpum</i> F. v. M.	G.		Austr.

**EUTHALLOPHYTA SCHIZOPHYTA SCHIZOMYCETES.
BACTERIA.**

NAME OF PLANT	Authority	Properties	Locality
<i>Bacillus botulinus</i> von Erm.	Jordan		N Am. Eur
<i>Bacillus piscidus</i> Sieber (1) Schum.	G.		Eur

For other toxic species see Buchanan, Part I of this Manual.

**EUTHALLOPHYTA. EUPHYCEAE. ALGAE.
RHODOMELACEAE**

<i>Alsidium Helminthochortos</i> Kütz.	G.		Atlantic Ocean
<i>Chondria vermicularis</i> Hook.	G.		Atlantic Ocean
<i>Hypnea musciformis</i>	G.		Atlantic Ocean
<i>Rhodomenia palmata</i> Gr.	G.		Atlantic Ocean

See Parts I and II of this Manual.

EUTHALLOPHYTA. PHYCOMYCETES.

<i>Saprolegnia</i> sp.	G.		Cosm.
------------------------	----	--	-------

ASCOMYCETES

Helvellaceae

<i>Gyromitra esculenta</i> Fr.	H.		Eur N Am.
--------------------------------	----	--	-----------

Hypocreaceae

<i>Claviceps purpurea</i> Tul.	G. Mi.	(Abort)	Eur N Am. As. Austr.
--------------------------------	--------	---------	-------------------------

BASIDIOMYCETES

Agaricaceae

<i>Amanita cothurnata</i> Atk.	A.		E. N Am.
<i>Amanita floerkecephala</i> Atk.	A.		E. N Am.
<i>Amanita Frostiana</i> Pk.	Cl. A. H. Pk.		E. N Am.
<i>Amanita magnivelaris</i> Pk.	H.		E. N Am.
<i>Amanita mappa</i> Fr.	H.		E. N Am.
<i>Amanita muscaria</i> L.	Cl. G. H. Earl. A. Pk. Fr.		Eur. N Am. As.
	Ches. Ph.		
<i>Amanita phalloides</i> Fr.	Cl. Pk. H. Earl.		Eur N Am.
<i>Amanita recutita</i> Fr.	A. Mi. Bull.		Eur N Am.
<i>Amanita rubescens</i> Fr.	Cl. A. Fr. Ph. Pk.		Eur N Am.
<i>Amanita solitaria</i> Bull.	Bull. Cl. A. Fr. Ph. Pk.		Eur N Am.
<i>Amanita spreta</i> Pk.	A. Pk. H.		N Am.
<i>Amanita strobiliformis</i> Fr.	H. Fr. A.		Eur N Am.
<i>Amanita verna</i> Bull.	H. A.		Eur N Am.
<i>Amanita virosa</i> Fr.	Fr. H. A.		Eur N Am.
<i>Cantharellus aurantiacus</i> Fr.	G. H. A. Ph.		Eur N Am.
<i>Clitocybe ludens</i> Schw.	H. A. Cl.		N Am.
<i>Coprinus narcoticus</i> Batsch.	G. Ph.		Eur
<i>Entoloma clypeatum</i> (L.) Miquel.	H. Cl.		Eur N Am. Fr.
<i>Entoloma graveolens</i> Pk.	Cl.		N Am.
<i>Hebeloma crustuliniforme</i> Bull.	Cl.		N Am.

NAME OF PLANT	Authority	Properties	Locality
<i>Hebeloma fastibile</i> Fr	H. Cl		Eur N Am
<i>Inocybe scaber</i> Müll.	H		Eur N Am.
<i>Lactarius scris</i> Bolt.	G. H.		Eur N Am.
<i>Lactarius atrovirens</i> Pk.	Pk. H.		N Am.
<i>Lactarius camphoratus</i> B.	H.		Eur N Am.
<i>Lactarium chrysorrheus</i> Fr	Fr H		Eur N Am.
<i>Lactarius insularis</i> Fr	Fr H	Acr	Eur N Am.
<i>Lactarius pergamentus</i> Fr.	Fr. H	Acr	Eur N Am.
<i>Lactarius rubus</i> Scop.	G.	Acr.	Eur N Am.
<i>Lactarius scrobiculatus</i> Fr	Fr H	Acr	Eur N Am.
<i>Lactarius subdulcis</i> Fr	H. Fr	Acr	Eur N Am.
<i>Lactarius torminosus</i> Fr	G.	Acr	Eur N Am.
<i>Lactarius trivialis</i> Fr	H. Fr	Acr	Eur N Am.
<i>Lactarius turpis</i> Fr	Fr H	Acr	Eur N Am.
<i>Lactarius vellereus</i> Fr	Fr H	Acr	Eur N Am.
<i>Lactarius zonarius</i> B.	G.	Acr	Eur
<i>Lentinus suavissimus</i> Fr	G. Fr		Eur
<i>Lentinus stypticus</i>	G.		Eur
<i>Lepiota Morgani</i> Pk	Pk. Cl. H.		N Am
<i>Marasmius oreades</i> Bolt.	G.		Eur N Am.
<i>Marasmius urens</i> Bolt.	G. H.		Eur N Am.
<i>Panæolus papilionaceus</i> Fr	H.		N Am.
<i>Pholiotæ caperata</i>	Cl.		N Am.
<i>Pholiotæ racemosa</i> B.	G.	Acr	Eur N Am.
<i>Russula Barlea</i> L.	G.		Eur N Am.
<i>Russula emetica</i> Sch.	H. G. A. Cl.	Em.	Eur N Am.
<i>Russula foetens</i> Pers.	G. Cl.		Eur N Am.
<i>Russula fragilis</i>	H.		Eur N Am.
<i>Russula rubra</i> Fr	G. H. Fr		Eur N Am.
<i>Stropharia aeruginosa</i> Curt.	H.		N Am.
<i>Tricholoma album</i> Schæff.	H.	Acr	Eur N Am.
<i>Tricholoma saponaceum</i> Fr.	Fr. Cl.	Acr	Eur N Am.
<i>Tricholoma sulphureum</i>	Cl.	Acr	Eur N Am.
<i>Volvaria volvacea</i>	Cl.		N Am.

HYDNACEAE

<i>Hydnum amarescens</i> L.	G.		Eur
<i>Hydnum graveolens</i>	G.		Eur

LYCOPERDACEAE

<i>Lycoperdon Bovista</i> Pers.	G. Mi.		Eur N Am.
<i>Lycoperdon gemmatum</i>	Mi.		Eur N Am.
<i>Lycoperdon giganteum</i> Fr	Fr. G.		Eur N Am.
<i>Lycoperdon saururus</i> Lam.	G.		Eur
<i>Scleroderma vulgare</i>	G.		Eur N Am.
<i>Secotium acuminatum</i>	Cl.		N Am.

PHALLACEAE

<i>Ithyphallus impudicus</i>	A. Cl.		N Am. Eur
<i>Lasurus borealis</i> Burt.	H.		N Am.
<i>Mutinus caninus</i> Fr	Fr. H. Cl.		Eur N Am.
<i>Mutinus elegans</i> Mont.	H.		N Am.
<i>Phallus duplicatus</i> Bosc.	H. Cl.		Eur N Am.
<i>Phallus Ravenelii</i> (B&C) E. Fisch.	H.		Eur N Am.

RHYTHMIDACEAE

<i>Crucibulum vulgare</i>	Cl.		Eur N Am.
<i>Cyathus striatus</i>	Cl.		Eur N Am.
<i>Cyathus vernicosus</i>	Cl.		Eur N Am.

POISONOUS FUNGI

NAME OF PLANT	Authority	Properties	Locality
<i>Boletus alveolatus</i> B. & C.	H.		N. Am.
<i>Boletus felinus</i> Bull.	A. H.		Eur. N. Am.
<i>Boletus Frostii</i> Russell	H.		N. Am.
<i>Boletus luridus</i> Schaeff.	M. H.		Eur. N. Am.
<i>Boletus satanas</i> Lenz	H. Cl.		Eur. N. Am.
<i>Polyporus anthelminticus</i> Berk.	G.		Eur.
<i>Polyporus hispidus</i> Bull.	G.		Eur.
<i>Polyporus squamosus</i> Fr.	Fr. G.		Eur.

UREDINEAE

<i>Colectosporium Sonchis-arvensis</i> (Schw.) Thunb.	Ches.	Ir.	Eur. N. Am.
<i>Puccinia coronata</i> Cda.	Panniel		Austr. Cosmp.
<i>Puccinia graminis</i> Pers.	Virchow		Eur. N. Am.
<i>Puccinia rubigovera</i> DC.	Panniel		Eur. N. Am.
<i>Puccinia xylostrizae</i>	Nelson		N. Am. Ry. Mts.
<i>Uromyces trifolii</i> (Hedw.) Lev.	Mohler		Eur. N. Am.

USTILAGINACEAE

<i>Ustilago avenae</i> (Pers.) Jens.	Mt.	Ir.	Eur. N. Am.
<i>Ustilago Hordei</i> (Pers.) Well. & Swingle	Mt.	Ir.	Cosmp.
<i>Ustilago neglecta</i> Nicot.	Panniel	Ir.	Eur. N. Am.
<i>Ustilago ruga</i> (Jensen) Kell. & Swingle	Mt.	Ir.	Eur. N. Am.
<i>Ustilago Triticis</i> (Pers.) Jens.	Mt.	Ir.	Eur. N. Am.
<i>Ustilago utriculosa</i> Tul.	Panniel	Ir.	Eur. N. Am.

TILLETIACEAE

<i>Tilletia foetida</i> (B. & C.) Trel.	Schaffner	Ir.	N. Am. Eur.
<i>Tilletia Tritici</i> (Berk.) Went.	Schaffner	Ir.	N. Am. Eur.

FUNGI IMPERFECTI

<i>Cladosporium</i> sp.	G.	Ir.	Eur. N. Am.
<i>Dipicoidia zeae</i> Lev.	E. F. Smith	Peilagra	Eur. N. Am.
<i>Fusarium</i> sp.	G.	Ir.	Eur. N. Am.
<i>Fusarium roseum</i> Lk.	Panniel	Ir.	Eur. N. Am.
<i>Gibberella</i> sp.	G.	Ir.	Eur. N. Am.
<i>Macrosporium Brassicae</i> B.	Panniel	Ir.	Eur. N. Am.
<i>Phragmotus eximius</i> Kuhn.	Mohler	Ir.	Eur. N. Am.
<i>Polythrincium trifolii</i> Kuntze	Panniel	Ir.	Eur. N. Am.

See Part II of this Manual for other species

LICHENES

<i>Cetraria juniperina</i> Ach.	G.		Eur. N. Am.
<i>Cetraria pinastri</i> Ach.	G.		Eur.
<i>Evernia vulpina</i> Ach.	G.		Eur. N. Am.
<i>Parmelia vulpina</i> Ach.	G.		Eur.
<i>Peltigera horizontalis</i> L.	G.		Eur. N. Am.
<i>Varicaria amara</i> Ach.	G.		Eur.

BIBLIOGRAPHY OF POISONOUS PLANTS

By HARRIETTE S. KILLOGG

In this bibliography we have included such modern works bearing upon the subject of poisonous plants as are more easily accessible to the student but to make this list more complete many titles of books by older writers have been added. These works are of interest not only from an historical point of view but also from an artistic standpoint because many of them are handsomely illustrated. This is particularly true of the works of Brand and Ratzeburg, Berge and Riecke, Golet, Bullhard, Dietrich, Henry Hegetschweiler, Otto, and Miguel which were published many years ago.

The bibliography also contains the more important articles published by the various experiment stations and by the United States Department of Agriculture besides a long list of rather recent papers appearing in technical chemical journals and in professional journals of medicine, veterinary medicine, and pharmacy. However this bibliography is by no means complete so far as these technical journals are concerned, but from the papers indicated in the bibliography the student can easily find further literature.

Miss Kellogg has attempted to make a subject catalogue covering a great many different topics such as the sale of poisons, general treatises on poisonous plants, vegetable toxicology, poisons from Abrus, vegetable alkaloids, hydrocyanic acid, immunity, luminous, ricinus, poisonous seeds, geographical papers, poisons from a legal standpoint, pellagra antidotes, etc.

While it would be impossible to indicate a reference to every species, the catalogue of the poisonous plants of the world indicates, to some extent, where the species was discussed. We have had to omit from the catalogue references to many popular treatises such as Lehmann's "Giftpflanzen mit besonderer Berücksichtigung der wirksamen Stoffe," and the works of Golet "Viral Hens und Poisonous Plants in Field and Garden" and the "Giftpflanzenbuch" by Berge and Riecke. The later as well as the earlier treatises, frequently refer to the literature where the species is mentioned as being poisonous. An example of this is found in *Helicoborus niger*. Berge and Riecke gave seventeen references where the plant is mentioned as poisonous, eleven references refer in regard to Oleander, and five upon *Teucoma radicans* showing that many of these plants have passed as poisonous for a considerable length of time.

Of peculiar and local popular treatises Germany has contributed more than any other country. The treatment of the poisonous plants in books such as Goepfert's "Ueber die Giftige Pflanzen Schlesiens," and Krause's "Studies of Poisonous Plants, in the German Colonies," aid materially in determining the distribution of certain plants. The United States has done something along this line especially through its Department of Agriculture. The publications of Coville and Chesnut, are excellent illustrations of this. Popular accounts in such treatises as that by Miss Huntington are helpful. The subject of poisoning probably has been treated more exhaustively than any other subject indicated. Dr. Warren has recently published a long list of papers on this subject, and he shows especially how popular impressions get into current literature.

without having much foundation in fact. On the subject of the anatomy very little has been published. Attention may be called to the excellent treatise of Collin, "Traite de Toxicologie Vegetale, Application du Microscope a la Recherche des Poisons Vegetaux." 1907. The German work by Mitlacher, "Toxicologische oder forensische wichtige Pflanzen und vegetabilische Drogen, mit besonderer Berücksichtigung ihrer mikroskopischen verhältnisse", also has an admirable discussion of the subject. There are, however, many pharmacopeal treatises like the work of Flückiger (English translation "Pharmacographia"), "Pharmacognosy" by Powers, Kraemer's "Botany and Pharmacognosy", the "Organic Materia Medica" by Sayre, "Foods and Drugs" by Greenish, Tschirch and Oesterle's "Anatomischer atlas, der Pharmakognosie und Nahrungsmittelkunde," and "Microscopy of Vegetable Foods" by Winton, and the English translation of works of Moeller, which will indicate the sources from which information of this kind can be secured.

We have added some "second hand" references which have been taken from what we believe to be reliable sources. It was quite impossible to obtain all of the original papers referred to in the bibliography. Many of them, however, have been seen in the original, either by Miss Kellogg or myself.

We wish to express our thanks to Dr. Wm. Trelease of the Missouri Botanical Garden, Dr. B. L. Robinson and Dr. W. G. Farlow of Harvard University, Dr. C. S. Sargent of the Arnold Arboretum, Mr. C. G. Lloyd of the Lloyd Library, Cincinnati, to the Surgeon General of the United States Army, and Mr. Johnson Brigham of the Iowa State Library, Des Moines, for courtesies in the loan of books and papers.

L. H. PAMMEL.

PARTIAL CLASSIFICATION OF THE BIBLIOGRAPHY

The numbers following the topics correspond with the numbers in the Bibliography.

Annus, 158, 508a, 550
 Aconitum, 94
 Aconite, { 186, 263, 302, 366, 474,
 Aconitum, { 503a, 529, 923, 947, 1078
 Acorn, 2
 Actinomyces, 88, 123, 138a, 463,
 809, 832, 859a, 1072
 Aethusa, 147, 302, 1078
 Agaricus, 934
 Alexoparmica, 679
 Alfa fa, 414, 628, 1036
 Algae, 33, 34, 70, 74, 126, 285, 286,
 676, 728, 742, 991
 Alkaloids, Vegetable, 121, 171, 187,
 397, 484, 779, 779a, 780, 781, 808,
 817, 873a, 938, 1082
 Almonds, 688
 Amanita, 307, 510, 512, 633a, 660, 797
 Anatomy, 236, 237, 763, 887, 997
 Andromeda, {
 Andromedotoxin, { 695, 799, 801
 Anemone, 687, 708
 Anhalonium, 193, 593, 599
 Anise, see Star anise
 Anthoceros, 670

Antium, { 662, 916, 917
 Antiaris, {
 Antidotes, 157, 307, 319, 376, 410, 478,
 507a, 508a, 522, 804, 932, 973
 Antioxina, 561, 718
 Arbutus toxaria, 981
 Arecia, 183, 302
 Argemone, 885
 Aroideae, { 754, 897
 Arum, {
 Arrow Poisons, 43, 50, 133, 156, 314,
 329, 394, 598, 898, 916, 917
 Aspergillus, { 95, 154a, 318, 375, 527,
 Aspergillus, { 618, 939
 Astragalus, 332, 640, 725
 Atractylis, 602
 Atropa, 887
 Atropin, 151a, 302, 944, 1078
 Auto intoxication, 986
 BACILLUS,
 anthracis, 51, 561
 oedematis, 135, 361
 suppuris, 711
 tetani, 292, 561
 virgula, 1011

Bacterial Poisons, 126a, 144, 321, 544, 1007
 Bacterium,
 diphtheriae, 561, 944
 tuberculosis, 38, 217, 463, 561, 826
 Baneberry, 41
 Barium, 209, 482, 487, 984
 Beans, 115, 256, 260, 388, 393, 394, 420, 855
 Beech, 316
 Belladonna, 302, 559, 621
 Berberis, 302, 862
 Bibliography 3a, 477 502a, 561, 579a, 727a, 823, 895, 1025, 1030
 Black Tongue, 352
 Blastomycosis, 449a, 526a, 776b, 987a
 Borage Plants, } 367, 368
 Boraginaceae, }
 Botulism } 420, 463, 561, 633
 Botulism }
 Boxwood, 334

Cactaceae, 193, 444, 599
 Caabar Bean, 302, 311, 412
 Calandrina, 720
 Camas, 253
 Camphor, 856
 Cannabis, 284, 816
 Capsella, 532
 Cardiac Poisons, 493
 Cascar, 290
 Cashew, 1063
 Cassava, 19, 224, 657, 990, 1058
 Castor Oil, See Ricinus
 Catapa, 1024
 Celastrus scandens, 78
 Cerebritis, 128, 247, 247a, 466, 655, 749, 1061a
 Cherry, 250, 681
 Chickens, See Ornithogalum
 Cholera, 135
 Cicuta maruata, 96, 219, 442, 474, 803, 929, 1094
 virosa, 96, 302, 531, 736, 1078
 Cicutoxin, 96. Also see references under Cicuta
 Claviceps See Ergot
 Cleistanthus, 480
 Clover, 202, 414
 Coca, 921
 Cocaine, 35a, 52a, 579a, 921 1031a, 1078
 Cocos, 271
 Coffee, 953
 Cochicum, 6, 6a, 129, 302, 689, 832a, 1078
 Conium, 501
 Conium maculatum, 336, 501, 538, 1078, 1086
 Corvalaria, 474
 Convulsives, 845
 Copafera, 302, 440
 Copas, 481

Corn, 86, 86b, 247a
 Broom, 427
 Kaffir, 767
 Cornstalk Disease, 12, 207, 819
 Corvidae, 642
 Cotoneaster, 460
 Coumarin, 440
 Cowbane, 732
 Creosote, 213
 Crotalaria, } 331, 945
 Crotalism, }
 Croton, 518a, 561
 Croton, 257 302, 553, 645
 Cryptogams, 75, 714, 778, 805
 Curare, 50, 788, 789
 Cyanogenesis, 258, 262, 264, 583
 Cynoglossum, 246, 883. See Borage
 Cypridium, 474, 513, 625, 626, 699, 700, 963
 Cytisin, } 189, 302
 Cytisus, }

DALBERGIA, 199
 Damiana, 72
 Dandel, 384
 Datura, 159, 239, 302, 337, 835, 887, 1053
 Dermal Mycosis, } 659, 675, 690, 713,
 Dermatitis, } 741, 783, 784,
 Dermatomycosis, } 1043
 Derris, 727, 763
 Decuria, 35
 Digitalis, 270, 302, 563, 669
 Dioscorea, 903
 Dioscorin, 905
 Diphtherial Poison, See Bacterium
 diphtheriae
 Dipodomys, 836a, 930
 Duboisia, 1078

ECHITES, 410
 Economic Plants, General, 238, 287, 696, 911
 America, 228, 433
 Mexico, 911
 United States
 California, 160
 Montana, 90
 Puerto Rico, 185, 761
 Vermont, 518
 Asia and Adjacent Islands,
 Japan, 60
 Australia, 282, 636
 Miscellaneous,
 Cuba, 655b
 Guam, 870a
 Entoloma, 464
 Equisetoids, } 842, 843, 843a, 961
 Equisetum, }
 Ergot, } 1002
 Ergotism, } 52, 231, 423, 530, 632,
 Ericaceae, 799

- Erigeron, 243, 574
 Eriobotrya, 458
 Erythrophloeum, 596
 Escholtzia, 297
 Eupatorium, 474
 Euphorbiaceae, 474, 925
 Exargin, 816
 FAVUS 94, 191
 Ferns, 296, 683
 Fish Poisons, 277, 278, 279, 371, 480,
 543, 727, 746, 829
 Flax, 259
 Floras,
 General, 47a, 274, 523
 North America 117, 363
 United States, 116, 117, 155, 359
 Alabama, 673a
 California, 1030a
 Connecticut, 355a
 Iowa, 364a
 Maryland, 924
 Minnesota, 631
 Washington, 786a
 See also Bulletins from the
 various Experiment Stations
 South America, See Medicinal and
 Poisonous Plants
 Asia, See Economic, Medicinal and
 Poisonous Plants
 Australia 645a
 Europe,
 England, 70a
 Germany, 554
 Switzerland, 554
 Flowers, Poisonous, 22, 23, 853
 Foods, 140, 421, 616, 1064
 Forage Poisoning, 125, 291, 465, 469,
 749, 1061
 Fungi, Poisonous, 73, 94, 95, 503,
 509, 610, 611, 629, 903, 1091
 Also see Mushrooms
 GALEGA, 682
 Gastro-enteritis, 542
 Gautheria 399, 407, 474, 514, 786,
 820, 859
 Geomium 273, 474, 875, 926a, 1078
 Gemata, 227, 242
 Germeides, 1009
 Gift-apple, 232
 Ginger, 417, 499
 Gleditschia, 745
 Glucosides, 306, 308, 349, 392, 458,
 520, 938
 Golden Rod, 910. See Hay Fever
 Grasses, 729, 731, 739, 740
 Darnel, 384
 Foxtail, 696
 Johnson, 214
 Porcupine, 82
 Red Top, 423
 Sleepy, 213
 Sweet, 276
 HAY FEVER, 6b, 338, 661, 814, 814a,
 852, 879a, 976
 Hava Poison, 596
 Heavea, 1062
 Hemlock, 865
 Hemlock Water, 501, 572, 573
 Heracleum, 335
 Hippomane, 411, 846
 Honey, Poisonous, 529, 800
 Horsetail. See Equisetum
 Hydrocyanic Acid in Plants, 80, 105,
 125, 196, 224, 261, 349, 369, 372, 385,
 386, 387, 388, 389, 390, 391, 392, 393,
 394, 435, 458, 459, 520, 630, 992,
 1014, 1078
 Hyocyamus, 887
 IMMUNITY, 229, 252, 306, 502, 561
 Ipoh, 327, 916, 917
 Ivy, Ground, 890
 Ivy Poison. See Rhus
 JAMURANDI, 419
 Janipha, 409
 Jatropha, 409, 553, 1058
 Jequirity, 661
 KALMIA, 204, 212, 533, 695, 864
 Kenyah, Dart Poison. See Ipoh
 Kumanga. See Komanga
 Kirodro Poison, 199
 Kistoumpa. See Menabea
 Kokra. See Coena
 Komanga, 439, 439
 Ksopo, 765
 Lac, Japanese, 948, 998
 Larkspur, 74, 211, 302, 339, 1047
 Lathyrus, 267, 396, 900
 Laurel. See Kalmia
 Lauro-cerasus, 1006
 Leguminosae, 465, 557, 684, 761, 857,
 889
 Lemon, Salts of, 111
 Lepiota, 161
 Leucomaines, 484, 1008
 Lichens, 553b
 Linamarin, 520
 Loco Weeds, { 89, 209, 210, 233, 234,
 Locomism. { 235, 268, 356, 472, 482,
 487, 533, 648, 649, 655a, 705, 721,
 728a, 743, 8.0, 876, 946
 Loganiaceae, 549
 Lotus, 265
 Lupine {
 Lupinosis { 141, 565, 889
 Lychius Githago, 148, 554
 Lymphangitis, 727a, 727d, 749a, 827a
 MACLEURA, 404
 Macrosporium, 822
 Macrozama, 591
 Manchinal Tree. See Hippomane

- Mangels, 665
 Materia Medica, 9, 39, 55a, 57, 248, 350, 365, 592, 1066, 1074
 Mechanical Injuries, 82, 87, 537, 696, 729, 731, 735
 Medicinal Plants, General Distribution, 71, 169, 170a, 302, 604, 612, 620, 834, 943, 1090
 Africa,
 Soudan, 575
 America, 84
 United States, 57, 172, 281, 830
 California, 333
 Massachusetts, 1060
 Minnesota, 1034
 Missouri, 733
 North Carolina, 497
 Tennessee, 324
 Asia and Adjacent Islands,
 Afghanistan, 26
 Japan, 60
 Malay, 338
 Philippines, 44
 Medical Jurisprudence, 65, 155a, 169a, 304a, 586, 804, 828, 837, 967, 968, 1078
 Menages, 437, 438, 765
 Meningitis, 291, 561, 749
 Meningo-encephalitis. See Cerebritis
 Mesquite, 1063
 Millet, 219a, 467, 468
 Mistletoe, 580
 Morphine, 92, 299, 302, 516, 562, 877, 886, 967, 1045, 1074, 1078
 Mucor, 354
 Mucormycoses, 35
 Mulberry, 103, 367
 Mushrooms, 36, 173, 175, 201, 203, 205, 304a, 381, 382, 569, 629, 654, 751, 815, 971. See Fungi
 Mycosis, 145a, 673, 874
 Nicotina, 93, 302, 758a
 Nuts,
 Areca, 183
 Betel, 302
 Nutmeg, 1052
 Nymphaea, 873
 Nuxvomica, 241
 OAK. See Acorn, Quercus
 Oedema. See Bacillus, oedematis
 Oenanthe crocata, 61, 1078
 Oils,
 Castor, 230, 302. See Ricinus
 Croton. See Croton
 Gaultheria, 820
 Pennyroyal, 1029
 Peppermint, 417
 Savin, 1029
 Tansy, 1029
 Vervain, 104
 Oleander } 892, 1021, 1067
 Oleogrin }
 Opium, 229, 302, 639. See Morphine
 Orange, 107
 Orange, False, 106
 Ordeal Poisons, 311, 438, 694, 1033, 1057
 Ornithogalum, 495
 Oxalic Acid, 561, 1078

 PAPAVERACEAE, 747
 Parasitic Diseases, 703
 Parsnip, 119, 473, 730
 Peas, 507
 Pellagra, 29, 86a, 150, 154a, 351a, 581, 661, 756, 825, 836a, 930, 987
 Pelletaria, 443
 Peppermint, 417
 Pharmacognosy } 301, 302, 558, 877,
 Pharmacographia } 1046, 1079
 Pharmacopoeia, 302, 450
 American, 182
 United States, 774, 776a
 British, 115a
 French, 176a
 German, 244a
 Italian, 288
 Swedish, 775
 Phaseolus. See Beans
 Physostigma, } 272, 412
 Physostigmin }
 Phyto-bezoars, 202, 425, 991a
 Phytolacca, 474, 1042
 Pink-root. See Spigelia
 Poison Lore, 4, 5a, 92a, 587, 978, 1018
 Poison Tree. See Copas
 Poisons, Detection of, 92, 93, 118, 154, 157, 429, 446, 552, 933
 Poisons in General, 4, 92, 97, 130, 310, 380, 552, 614, 723, 846, 940
 Mineral Poisons, 723, 963
 Vegetable Poisons, 30, 31, 197, 475, 490, 764, 827, 919, 920, 1000, 1016
 Poisonous Plants, General Treatises,
 28, 42, 75, 145, 188, 194, 197, 248, 249, 252, 299, 341, 343, 403, 455, 457, 521, 523, 556, 578, 588, 668, 709, 714, 744, 773, 812, 824, 834, 853, 858, 884, 902, 915, 933, 982, 1010, 1018, 1032a, 1048, 1077
 Geographical Distribution
 Africa, 839, 851
 Soudan, 575
 South Africa, 374, 1021b, 1059
 America, North, 23
 Mexico, 911
 United States, 27, 162, 163, 166, 167, 340, 500, 555, 805
 California, 67, 533, 894, 1005
 Colorado, 339
 Idaho, 927
 Indiana, 195
 Iowa, 734, 738
 Massachusetts, 922

- Minnesota, 175
 Missouri, 733
 Montana, 84, 90, 168, 1049
 Nebraska, 82, 83
 Nevada, 534, 536
 New Jersey, 405, 406
 New York, 751, 988
 New York City, 863
 North Carolina, 497, 626
 North Dakota, 1004
 Ohio, 878, 913, 914
 Pennsylvania,
 Eric, 395
 Rhode Island, 47
 Vermont, 518
 America, South,
 Brazil, 143, 571
 Chili, 787
 Guiana, 747, 898
 Surinam, 747
 West Indies,
 Cuba, 180
 Porto Rico, 185, 761
 Asia, and Adjacent Islands
 Afghanistan, 26
 China, 218
 India, 888, 947
 Bombay, 543
 Japan, 948, 998
 Malay, 69, 225
 Philippines, 43
 Australia, 46, 282, 634, 635, 636,
 684, 996
 Victoria, 282
 Europe, 445
 Austria, 564
 Belgium, 139
 Central Europe, 445
 Denmark, 377
 England (also Great Britain),
 21, 457, 515
 France, 131, 344, 973, 1015
 Germany, 112, 245, 342, 353,
 373, 383, 402, 449, 454, 538,
 556, 778, 792, 881, 904, 1073a,
 1087
 Colomes, 562
 Duisberg, 99
 Thuringia, 726
 Schwarzburg-Rudolstadt, 926
 Hungary, 393, 566
 Netherlands, North, 666
 Poland, 73
 Saxony, 275
 Silesia, 941
 Switzerland, 446, 447, 470, 1015
 Russia, 869
 Potato, 617, 627
 Primula, 701, 702, 831
 Prunthogatum. See Ornithogalum
 Prulaurocerasin, 459, 460
 Prunus, 459, 474, 1014. Also see
 Cherry
 Ptomaines, 121, 484, 1008
 Quercus, 4, 474, 1021a
 Ranunculus, 149, 424, 580, 759
 Rape, 49, 469
 Rattle Weed. See Crotalaria
 Raphanus, 606
 Rhizopus, 192
 Rhododendron, 800, 1071
 Rhus, 3, 11, 15, 30, 53, 76, 124, 134,
 138, 146, 181, 223a, 422, 426, 432,
 474, 476, 485, 491, 517, 525, 531,
 539, 613, 637, 639a, 678, 741, 758,
 771, 772, 783, 784, 790, 868, 907,
 949, 951, 955, 965, 969, 972, 974,
 1025, 1038, 1044, 1056.
 Bibliography on Rhus Poisoning,
 1025, 1026, 1027
 Rhus and Bacterial Infection, 136
 Ricin, 508a, 561
 Ricinus, 85, 220, 229, 683
 Robinia, 811
 Rosta, Poisonous, 836
 Rosaceae, 396
 Rumex, 1078
 Sakalava, 438, 439, 765
 Sale of Poisons, 93, 280, 448
 Salikounda, 440
 Sambucus, 474
 Sanguinaria, 474
 Santonin, 345
 Saponin, 45, 372, 896
 Savanna Flower. See Echites
 Sawdust, Poisonous, 271
 Scarlet Fever Poison, 120
 Seeds, Poisonous:
 Beans, 115
 Colchicum, 302
 Cotton, 215
 Croton, 302
 Cynoglossum, 883
 Eriobotrya, 458
 Garcia, 152
 Larkspur, 79
 Leguminosae, 889
 Lupine, 141
 Lychnis, 589
 Nux vomica, 302
 Omphalea, 152
 Rape, 49
 Ricinus, 85, 302
 Sabadilla, 302
 Samarubaceae, 185
 Sinapis, 302
 Vicia, 80
 Wormwood, 100
 Septicaemia, 320
 Shepherd's Purse, 532
 Sicyos, 404
 Silage, 208
 Sinarubaceae, 199

- Sinapis, 860
 Sium, 610a, 777
 Sneezeweed, 641
 Snowdrop, 601
 Solanaceae, 674
 Solanum, 283, 474
 Sophora, 190, 798
 Sorghum, 24, 48, 767, 1022
 Spargelia, 295, 315, 846, 954, 979
 Sporotrichosis, 3a, 90a, 135a, 145a, 536b, 727a, 727c, 859a, 993a
 Spruce, 752
 Spurge, 568
 Staphisagria. See Larkspur
 Star-anise, 222, 242, 494
 Sterilaria media, 150
 Stomatitis, 294, 671
 Strawberry, 814b
 Streptococcus, 174, 176, 561
 Strophantus, 311, 329
 Strychnin, 283, 376, 658, 1045
 Strychnos, 133, 241, 436
 Stylophorum, 886
 Sugar Beets, 665
 Suicide, 689
 Sumach, 491, 951, 1050. Also see Rhus
 Symptoms from Poisoning, 37, 37a, 39, 92, 93, 118, 376, 382, 552, 932. Also see General Treatises upon Poisoning
 TACHIN, 311, 437, 438
 Tartaric Acid, 1078
 Taxus, 150a, 461, 901, 959
 Tea, 953
 Temperature affecting Toxicity of Poisons, 418, 872, 1088, 1089
 Templetonia, 190
 Tetanus. See Bacillus tetani
 Thalictrum, 505, 506
 Therapeutics, 39, 55a, 288, 783, 893, 953, 1070
 Thrush, 379, 607, 608, 609
 Tobacco. See Nicotin
 Tonka Bean, 440
 Toad Poison, 605
 Toxicology, General, 155a, 551, 553a, 567, 597, 651, 724, 796, 828, 918, 937
 Vegetable, 173, 177, 197, 553, 561, 831, 107
 Veterinary, 592, 706, 715
 Toxins, 718
 Treatment for Poisons, 39, 595, 646, 689, 693, 722, 804, 933, 956, 962, 1068
 Trees, Malignant Effect of, 838
 Trichophyton, 401, 870, 975
 Tropaeolum, 389
 Tulip Grass, 139
 Tunas, 415
 Turpentine, 958
 UMBELLIFERAE, 547, 603, 803
 Upas, 241
 Ustilago, 276
 VERATRUM, 58, 114, 221, 302, 313, 760
 Verbascum, 669
 Verhena, 104
 Veronal, 328
 WATER BLOOM, 650
 Weeds, Poisonous, 734
 Yew. See Taxus

BIBLIOGRAPHY OF POISONOUS PLANTS

BY HARRIETTE S. KELLOGG

- 1 Abbott, A. C. Principles of Bacteriology, a Practical Manual for Students and Physicians. 641 Philadelphia and New York. Ed 8.
- 2 Acorn Poisoning Vet Rec. 13.369. London.
- 3 Acree, S. F. and Syme, W. A. Some Constituents of the Poison Ivy Plant. Am. Chem. Jour. 36.301-341.
- 3a Adamson, H. G. Sporotrichosis, a Resume of the Literature Relating to Sporotrichal Infection of the Skin. Brit. Jour. of Derm. 20.296.
- 4 Adoynis, Santes de Liber de venenis. 101, Venice, 1492.
- 5 Aggio, C. Two Interesting Outbreaks of Vegetable Poisoning. Am. Vet. Rev. 33.368.
- 5a Ayar N. S. Certain Facts Regarding the Poison-lore of the Hindus. Trans. India Med Cong. 1894.307-310. Calcutta.
- 6 Albert. On Colchicum Poisoning. See Zeitsch f Staatsarznei 64.202. 1862.
- 6a Albertoni and Casali On Colchicum Poisoning. See Boll.dac.med. d. Bologna. 1890. 9 ser. 1.38.
- 8b Alberts Het Pollen-asthma. Amsterdam. 1903.
- 7 Alexandersen See Sarr.
- 8 Alkaloids in Plants Botanical Magazine 16. (Written in Japanese.)
- 9 Allen, Timothy F. Allen Encyclopedia. The Encyclopedia of Materia Medica. A Record of the Positive Effects of Drugs, etc. 10 vols. Philadelphia. 1877-1879.
- 10 Alpers W. C. The Antidote Cabinet. Phar. Era. 22.136-138.
- 10a Alsberg C. L. See Black, O. F.
- 11 Aumbaugh, W. E. On Death from Rhus Poisoning. See Med. World. 21.176.
- 12 Alway, F. J. and Peters, A. T. Losses from Cornstalk Disease in Custer Co., Nebraska. during the Winter of 1906-1907. Press Bull. Neb. Agr. Exp. Sta. 27.4.
- 13 Anderson, F. W. Poisonous Plants and the Symptoms they Produce. Botanical Gazette. 14.180-181.
- 14 Andrews, F. M. The Effect of Alkaloids and other Vegetable Poisons upon Protoplasm. Ind. Acad. Sci. 1905.195.
- 14a Andrews, L. See Graves, C. B.
- 15 Anide, John Chemical Observations on Rhus Toxicodendron. Therapeutic Gazette. 13.676.
- 16 Anonymous. Acorn Poisoning. Vet. Rec. London. 13.359.
- 17 Anon. Alkaloids in Plants. Bot. Mag. 16.
- 18 Anon. Are Birds Affected by Eating Poisonous Food? Scientific American 79.424.
- 19 Anon. Cassava Poisoning. Agr. News. Barbadoes. 1.5.
- 20 Anon. On Death Ensuing from Rhus Poisoning. See Boston Med. and Sur Jour. 137.302.

21. Anon. English Poisonous Plants. Gentleman's Magazine. No. 1755.
22. Anon. Flowers that are Poisonous. Current Literature. **23** 331-333.
23. Anon. Giftigkeit der Amerikabluthen. Pharmaceutische Centralhalle. **1903** 136.
24. Anon. Green Sorghum Poisoning. West Indian Bulletin **3** 326-333.
25. Anon. Die Immunität von Pflanzen gegen ihr eigenes Gift. Zeitsch.fang. Mikr **13** 60-62. Leipzig. 1907
26. Anon. Medical Plants in Afghanistan. Jour Applied Science **1860** 142
27. Anon. Native Poisonous Plants. Harper's Bazaar. **34** 941
- 27a. Anon. Poisoning by Water Hemlock. See Med. Times and Gazette **1855** 51
28. Anon. Our Poisonous Plants. Pathfinder **8** 7.
29. Anon. Pellagra. Boston Evening Transcript. Aug. 18. **1898**.
30. Anon. Plant Poisons. Sci Am **98** 361
31. Anon. Plant Poisons. Gardener's Chronicle. **45** 111, 366.
32. Are Birds Affected by Eating Poisonous Food? Sci. Am. **79**:424
33. Arthur, J. C. Some Algae of Minnesota Supposed to be Poisonous. Bull. Minn. Acad. Nat. Sci. **2**:4. (Appendix) **3**:97 Second Report in Ann. Rept Minn. Univ **4** Supplement 1. Also Report U. S. Dept. Agr. **1897** 97
34. A Supposed Poisonous Seaweed in the Lakes of Minnesota. Proc. Am. Assoc. Adv. Sci. **22** 305. (Abs.)
35. Asahina, Y Ueber die Alkaloide von *Dicentra pusilla* Sieb. et Zucc. Arch. Pharm. **247**:201. 1909.
- 35a. Ashworth. On Cocain. See Lancet **1889**:273.
36. Atkinson, G. F Mushrooms, Edible, Poisonous, etc. 322. Ithaca, N. Y. 1901. Ed. 2.
37. Atkinson, V. T Action of Poisons. Special Report, Diseases of Cattle and on Cattle Feeding. U. S. Dept. Agr. Bur. An. Ind. **1896** 63-76
- 37a Poisons and Poisoning. In Special Report, Diseases of Cattle, U. S. Dept. Agr. **1908** 53-69. Revised by Leonard Pearson.
38. Auclair, J. Les poisons du bacille-tuberculeux humain et la degeneration caseuse. Revue de la Tuberculose **1898**.
39. Aurand, S. Herbert Botanical Materia Medica and Pharmacology; giving the Botany, Pharmacology, Physiological Action, Therapeutic Range, Treatment of Poisons and Antidotes. 406. Chicago. 1899.
40. Avery, Samuel See Peters, A. T.
41. Bacon, Alice E. An Experiment with the Fruit of Red Baneberry. Rhodora **5** 77-79.
42. Bacon, C. W Handbook to Accompany Chart of Poisonous Plants. 30. London. 1899.
43. Bacon, R. F. Philippine Arrow Poisons. Philippine Jour. Sci. **3-A** 41-44.
44. The Physiologically Active Constituents of Certain Philippine Medical Plants. Phil. Jour. Sci. **1**:1007-1036.

45. Bacon, R. F. and Marshall, H. I. . . . The Toxic Action of Saponin. *Phil. Jour Sci.* **1**:1037-1042.
46. Bailey, F. M. and Gordon, P. R. . . . Plants Reputed to be Poisonous and Injurious to Stock. Gov't Printer, Brisbane. 4:112. 1887.
47. Bailey, W. W. . . . Rhode Island's Poisonous Plants. *Trans. R. I. Med. Soc.* **8**:64-72.
- 47a. Bailon, H. . . . Histoire des Plantes. 13 vols. Paris. 1868-1896.
48. Ball, C. R. . . . Saccharine Sorghums for Forage. *Far Bull U. S. Dept. Agr.* 246-37.
- 48a. Ballardini . . . On Aconite Poisoning. See Withaus and Becker's Medical Jurisprudence 4:648. Abs. of Article in *Ann. Univ. d. Med. Milan* **8**: 635. 1840.
49. Bang, B. . . . Cases of Poisoning from Rape-Seed Cakes. *Ugeskr. Landw.* **43**: 589-591, 604-605. 1897. Abs. in *Exp. Sta. Rec.* **9**:954.
50. Barbosa, R. J. . . . L'uraery du curare. 180. 7pt. Brussels. 1903.
51. Bare, C. L. . . . The Ravages of Bacillus Anthracis in California. *South California Practitioner* **1894**: 121-153.
52. Barger, G. and Carr, F. H. Note on Ergot Alkaloids. *Phar Jour* **23**: 257.
- 52a. Barker . . . On Cocain. See *Am. Jour. Oph.* **10**:342. 1893.
53. Barrows, W. B. . . . On the Non-poisonous Quality of the Ripe Fruit of *Rhus Toxicodendron*. See *Bull. Div. Orn. Mam. U. S. Dept. Agr.* **6**:85.
54. Barth, Herman . . . Studien über den mikrochemischer Nachweis von Alkaloiden in pharmaceutisch verwendeten Drogen. *Botanischer Centralblatt* **75**: 275-231, 261-267, 292-300, 326-644; 369-378, 401-409.
55. Barthelat, C. J. . . . Mucormyces, Spontanées, Historique. I, Les Mucormyces chez l'homme. *Arch. de Parasitologie.* **1903**.
- 55a. Barthelow, R. . . . A Practical Treatise on Materia Medica and Therapeutics. Ed. 7. New York. 1889.
56. Barton, W. C. . . . A Flora of North America. Philadelphia. 1820.
57. . . . Vegetable Materia Medica of the United States, or Medical Botany. 2 vols. 1817-1818.
58. Bastin, E. S. . . . Structure of *Veratrum viride*. *American Jour. of Pharmacy.* **77**:196-203.
59. Bastin, E. S. and Tremble. *Coniferae*
60. Batchelor, J. and Miyabe, K. . . . Japanese Economic Plants. List of those used Medicinally by the Aines. *Trans. Asiatic Society of Japan.* **21**. Abs. in *Proc. Am. Phar. Assoc.* **44**:501-505.
61. Baume, M. . . . Poisoning by *Oenanthe crocata*. *Archiv des Mal. Mentales.* **1881** No. 3. Abs. in *Ed. Vet. Rev.* **4**:704.
62. Baumgarten, P. . . . Lehrbuch der pathologischen Mycologie. 1 Hfte. 220. Braunschweig. 1836.
63. . . . Sur les microorganismes morbigenes d'origine végétale. Traduit par Steinen. *Jour. de médecine de chirurgie et de pharmacologie.* **1886**. Mai.

64. Baumgarten, P. and Muller, R. Versuche über accommodative Zuchtung von Schimmelpilzen. Berliner klin. Woch. 1882.
65. Beal, J. H. The Legal Responsibility of the Pharmacist in the Sale of Poisons and Dangerous Substances. Pharm. Rev. 10:292-298, 338-344.
66. Becker, T. C. See Witthaus, R. A.
67. Behr, H. M. On the Poisonous Plants indigenous to California. Repr.
68. Beilstein, F. Handbuch der organischen Chemie. 4 vols. Hamburg and Leipzig. 1895-1899.
69. Benedicente, A. Recherches pharmacologiques sur quelques poisons par les negritos de l'archipel Malais. Archives Italiennes de Biologie. Fasc. 1. 1899.
70. Bennett, A. W. On Vegetable Growths as Evidence of the Purity or Impurity of Water. St. Thomas Hospital Report. 20:8.
- 70a. Bentham, George Illustrated Handbook of the British Flora 1: 503. 2:505-1076.
71. Bentley, Robert and Tri-men, Henry Medicinal Plants. Description and original figures of the Principal Plants employed in Medicine. 4 vols. London. 1890.
72. Bentley, W. H. Poisoning by Damiana—Recovery. Therap. Gaz. N. S. 4:57. 1883.
73. Berdan, T. Tablica grzybon trujacych w kraju. (Table of poisonous Fungi in Poland.) Warsaw. 1888.
74. Berr, Ueber die Schwinnnen bei Kindern. Bremen, 1842. (Clinique des hôpitaux des enfants de Paris, 1842.)
74. Berge, Fr. and Riecke, V. A. Giftpflanzenbuch, oder Allgemeine und besonders Naturgeschichte sämtlicher inländischen so wie der wichtigsten ausländischen phanerogamischen und kryptogamischen Giftgewächse, mit treuen Abbildungen sämtlicher inländischen und vieler ausländischen Gattungen. 11-329. 72pt. Stuttgart. 1845.
76. Berenger G. M. Rhus Poisoning. Am. Jour. Phar. 68:18-20.
77. Berلمان, M. See Sabanijeff.
78. Bernhard, H. Celastrus scandens, L. Amer. Jour. Phar. 4. 12.
79. Bernon, Empoisonnement par les semences de Staphysaigre. Jour de Phar. et de Chimie. 1882. Aug.
80. Bertrand, G. La vicianine, nouveau glucoside cyanhydrique contenu dans les graines de vesce. Bull. Soc. chim. France. 4 ser. 1. 151-154.
81. Bertuch, F. See Guenther, J.
- 81a. Beslev, F. W. See Shreve, F.
82. Bessey, C. E. Injuriousness of Porcupine Grass. Am. Nat. 18:929.
- 82a. Notes on Poisonous Plants of Nebraska. Proc. Soc. Prom. Agr. Sci. 23:34.
83. Poisonous Weeds. Rept. Neb. Agr. Exp. Sta. 16:14-62.
- 83a. Besta, C. See Cenn, C.

84. Bigelow, Jacob American Medical Botany A Collection of the Native Medicinal Plants of the United States. 3 vols. Boston. 1817-1820.
85. Bishpam, W. N. Report of Cases of Poisoning by Fruit of *Ricinus communis*. *Am. Jour. Med. Sci.* **124**:319.
- 85a. Bissell, C. H. See Graves, C. B.
86. Bitting, A. W. The Effects of Eating Mouldy Corn. See *Rept. Ind. Agr. Exp. Sta.* **1899**:44-45.
- 86a. Berregard, A. P. See Hare, R. F.
- 86b. Black, O. F. and Alsberg, C. L. The Determination of the Deterioration of Maize, with incidental References to Pellagra. *Bull. U. S. Dept. Agr. Bur. Pl. Ind.* **199**:32.
87. Blanchard, Raphael Accidents causés par une graminée américaine. *Archives de Parasitologie* **10**:187-194.
88. Quelques cas anciens d'actinomyose. *Arch. de Parasit.* **2**:362-376. 1899.
89. Blankinship, J. W. The Loco and other Poisonous Plants of Montana. *Bull. Mon. Agr. Exp. Sta.* **42**:75-104.
90. Native Economic Plants of Montana. *Bull. Mon. Agr. Exp. Sta.* **56**:36.
- 90a. Bloch, Bruno. Die Sporotrichose. *Mediz. Klinik* **1909**.
91. Block Beitrag zur Kenntnis der Prolidung u. der Geweben des thierischen Organismus. Thèse. Stuttgart. 1870.
- 91a. Bodgett, F. H. See Shreve, F.
92. Blyth, A. W. Poisons. Their Effects and Detection, A Manual for the Use of Analytical Chemists and Experts. 712. London, 1894.
- 92a. Old and Modern Poison Lore.
93. Blyth, A. W. and Blyth, M. W. Fourth Edition of "Poisons." Revised, Enlarged and Rewritten. 772. London. 1906.
94. Bodin, E. Sur le champignon du foyon de la souris *Achorion Quinckeianum*. *Archiv. de Parasit.* **5**:5.
95. Bodin, E. and Gauthier, L. The Toxins Produced by *Aspergillus fumigatus*. *Annales de l'Institut Pasteur* **20**:209-224. *Abstr. in Exp. Sta. Rec.* **18**:8.
96. Boehm Cicutoxin. See *Archiv. f. exp. Path. u. Pharmacol.* **5**:279-310.
97. Böhm, R., Naunyn, B., and von Boeck, H. Handbuch der Intoxicationem. Ed. 15 of the German Ed. of Ziemssen's Cyclopaedia.
98. Boehmer, G. R. Commentatio de plantis segetis infectis ac cedit commentatio de plantis auctoritate publica extirpandis, custodiendis, et a foro proscribendis. 198. Wittenberg. 1785.
- 98a. Cyano segetum nuper imputatum virus limitatur. 12. Wittenberg. 1787.
99. Boeninger, T. K. De plantis venenatis et speciatim de plantis venenatis agri Duisburgensis. 133. Duisburg. 1790.
100. Bond, A. R. Poisoning by Wormwood Seed. *Phar. Jour. Ser. 4.* No. 1483.
101. Boote, L. E. See Dunstan, W. R.
102. Bose, C. L. See Evans, J. F.
103. Bouchard, Charles Upon the Role of Poisons of Microbic Origin in Infectious Maladies. *Gaz. hebdom. de Med. et de Chim.* **1889**:120-122.

104. Bourdier, L. . . . Sur la "verbenaline", glucoside nouveau re-
ture de la verveine (*Verbena officinalis*, L.)
C R. Soc. Biol. Paris. Nov. 1. 1907
- 104a. Bourgoing, Ed. . . . See Rabuteau, A.
105. Bourquelot, E. and Danyon, E. . . . Sur la presence d'un glucoside cyanhydrique
dans les feuilles du sureau. C R. Acad. Sc.
Paris. 1905. July 3
106. . . . Sur un empoisonnement par le fausse orange,
survenu à Bois-le-Roi, Seine-et-Marne, le 6
septembre 1898. Bull. de la Soc. myc. de
France. 12:146-150.
107. Boutet, and Clement Empoisonnement par la fleur Orange. Bull. de
la Soc. des Nat. Sci. de l'Ouest de la France.
1901:45.
108. Boutmy, E. . . . See Brouard, P.
109. Bowhill Tulip-grass Poisoning. Vet. Rec. London, 13:
229.
110. Bower, F. O., Translator See De Bary, A.
111. Brashwate, P. F. . . . Suicidal Poisoning with Salts of Lemon Per-
foration of the Stomach. Brit. Med. Jour
1:183. (1905)
112. Brandt, J. F. and Ratze-
burg. . . . Deutschlands phanerogamische Giftgewächse
in Abbildungen u. Beschreibungen. Ed. 2.
30. 48 pl. Berlin. 1838.
113. Brebeck. See Fischer, A.
114. Bredeman, G. . . . Alkaloids of *Veratrum viride* and their De-
termination. Phar. Jour 22:283.
115. Bretin, Yvesque and Ver-
dier Haricots toxiques dits de Hongrie. Jour
Phar. et Chim. 26:348.
- 115a. British Pharmacopoeia 1898. (Ph. Brit.)
116. Britton, N. L. . . . Manual of the Flora of the Northern States
and Canada. 1860. New York. 1901
117. Britton, N. L. and Brown,
A. Illustrated Flora of the Northern United
States and the British Possessions from
Newfoundland to the Parallel of the South-
ern Boundary of Virginia, and from the At-
lantic Ocean westward to the 102d Meri-
dian. Vol. I:612 f. 1-1425, II: 643 f.
1426-2892, III:588 f. 2893-4081. New York.
1896-1898.
118. Brodie, E. C. . . . Further Experiments and Observations on the
Action of Poisons on the Animal System.
Phil. Trans. 102:218.
119. Brodie, D. A. . . . Poison Parsnip in Western Washington. Bull.
Wash. Agr. Exp. Sta. 45:12.
120. Brooke, J. . . . Activity of the Scarlet Fever Poison After a
Year. Med. Rec. 2:126.
121. Brouard, F. and Boutmy, E. . . . Ptomaines et alcaloides végétaux. Bull. de l'
Acad. de Med. 1881:19.
122. Brown, A. See Britton, N. L.
123. Brown, E. L. . . . A Case of Actinomycosis. Chicago Medical
Record. 1894:251.
124. Brown, S. A. . . . On Rhus Poisoning. See Med. Rec. 30:221.
125. Brunnich, J. C. . . . Hydrocyanic Acid in Fodder Plants. Jour.
Chem. Soc. 1903:788-796.
126. Buchanan, R. E. . . . Notes on the Algae of Iowa. Proc. Ia. Acad.
Sci. 14:47-64.

- 126a. Buchner, H. Ueber Bacteriengifte und Gegengifte Munchener Medicinisch Wochenschrift. **1893**: 449-452, 480-483.
127. Buchner, Max Ueber pathogenetische Wirkungen der Polkeimkörper Jahrb. med. Wissensch. 1868.
- 127a. Buckley, J. S. See Mohler, J. S.
128. Buckley, S. S. A Special Investigation of the So-called "New Horse Disease" in Maryland. Bull. Md. Agr. Exp. 53 110-114.
129. Budd, Charles Probable Poisoning by Colchicum Lancet No. 3001
130. Buechner, J. G. De venenis Halle. 1746.
131. Bulliard Histoire des plantes veneneuses et suspectes de la France 398. Paris 1798. Ed 2
133. Bureau, Ed Sur les Strychnos africains et les plantes servant à empoisonner les armes en Afrique. Bull. Mus. Hist. nat. Paris. **1901** 418-425.
134. Burgess, T. J. W. The Beneficial and Toxical Effects of the various Species of Rhus. Canadian Jour. Med. Sci. (Phar. Jour. and Trans. **1881**)
135. Burke, R. W. The Bacillus of the Malignant Oedema Vet Jour **1890** 246-247
- 135a. Burling, J. M. Subcutaneous Abscesses Caused by the Sporothrix Schrenckii Southern California Pract. **24**:1
136. Burrill, T. J. On Bacterial Infection and Rhus Poisoning See Am. Mo. Mic Jour **3**:192, Proc. Am. Assoc. Adv. Sci. **31**:515; Am. Nat. **17**:3, 9.
- 137 Rhus Poisoning. Garden and Forest **9**:386-369
138. Bussey, S. C. On Recurrence of Attacks of Rhus Poisoning without Re-exposure Am. Jour. Med. Sci. **66**:436.
- 138a. Byron, J. M. A Case of Actinomyces in Man. N. Y. Med. Jour **2**:216, (1889)
139. Caels, T. P. De Belgis plantis quasitate quam hominibus cacterisve animalibus nociva seu venenata praeditis symptomatibus ab earum uso productis nec non antidotis adhibenda. 66. Brussels. 1774
140. Cahn, A. On Toxins in Foods. See Strasburger-Medizin. Zeit. Jan. **1906** 27
141. Callsen, J. Ueber die Alkaloide der Samen von Lupinus angustifolius und von Lupinus perennis, var. polyphyllus. Archiv der Phar **237**:566-595
142. Cambier, Jacob See Power, F. B.
143. Caminhos, J. M. Das plantas toxicas do Brazil. Rio de Janeiro 1871. Trans. into French by F. Rey. (Catalogue des plantes toxiques de Bresil. Paris, 1880.)
144. Campbell, H. P. Bacterial Poisoning through Medicine Am. Jour. Phar. **1890**:113-118.
145. Card, F. W. Poisonous Plants. Gar. and For. **9**:8.
- 145a. Carougeau, M. Sur une nouvelle mycose sous-cutanée des équides. Madagascar 1906. Jour. de Med. veter. et de Zootechnie **80**:8-22, 75-90, 148-153. Abs. in Centrbl. f. Bakt. Abt. I. **45**:610-612
146. Carpenter, C. R. Rhus Toxicodendron. Ther. Gaz. **14**:93.
147. Carruthers, Wm. Aethusa Cynapium. Jour. Roy. Agr. Soc. of Eng. **63**:280.
148. Lychnis Githago. id. **63**:279

149. *Ranunculus bulbosus*. id 62:279
 150. *Stellaria media*. Is it Poisonous? id 64:308.
- 150a. Carruthers, Wm. and Others Yew Poisoning. id Ser III 3:698
151. Carter, Marion H The Medical Mystery of Today (Pellagra) McClure's 34:94-104.
- 151a. Cartwright, C B. A Case of Atropin Poisoning N Y Med Jour Sept 5, 1903.
- 151b. Casali See Albertoni.
152. Cash, J T The Characters and Actions of the Seeds of *Omphalea megacarpa* (diandra), *Omphalea triandra* and *Garcia nutans* (from Trinidad) Phar. Jour, 81:351-352
153. Cassava Poisoning Agr News. Barbadoes. 1-5.
154. Cattel, H W Post-Mortem Pathology A Manual of Post-Mortem Examinations and the Interpretations to be Drawn therefrom A Practical Treatise for Students and Practitioners. 551 6 pl. 183f. Ed. 2. 1905
- 154a. Ceni, Carl and Resta, Carlo Ueber die Toxine von *Aspergillus fumigatus* und *flavescens* und deren Beziehung zur Pellagra Centrbl f Allgem. Path u Path. Anat. Jena. 13:930-941. 1902
155. Chapman, A. W. Flora of the Southern United States. 698. Ed. 2. New York. 1894
- 155a. Chapman, H. C. A Manual of Medical Jurisprudence and Toxicology Philadelphia. 1892
156. Charteris, F On the Active Principles of Poisoned Spears Obtained from the Metch or Mouch. Country Phar Jour London 1901 Ser 4. 13:43.
157. Chaussier, H Contre-poisons, ou moyens reconnus le plus efficaces pour combattre l'effet des diverses especes de poisons, suivis de l'indication des secours à donner aux personnes mordues ou piquées par des animaux ou des insectes vénéneux, aux noyés, aux asphyxiés, aux enfants nouveaux-nés, etc. Paris. 1818.
158. Chestnut, V. K. Abrus Poisoning. Asa Gray Bulletin 5:35-36.
159. Datura Poisoning. id 5:87
160. Plants Used by the Indians of Mendocino County, California. Cont. L. S. Nat. Herb. 7:295-408.
161. Poisonous Spores of the Green-spored *Lepiota* Asa Gray Bull. 5:87-93.
162. Preliminary Catalogue of Plants Poisonous to Stock. Ann. Rept. Bur. An. Ind. U. S. Dept. Agr 15:387-420
163. Principal Poisonous Plants of the United States. Bull. L. S. Dept. Agr. Div. Bot. 20:60. See also Repr. II. St. Bd. Health 60. 1899.
164. Problems in the Chemistry and Toxicology of Plant Substances. Science (N. S.) 15. 1016-1028.
165. Some Common Poisonous Plants. Year Book U. S. Dept. Agr. 1898 137-146.
166. Some Poisonous Plants of Northern Stock Ranges. Yr Bk U. S. Dept. Agr. 1900: 305-324.

- 167 Thirty Poisonous Plants of the U. S. Farmer's Bulletin, U. S. Dept. Agr. 86.32
168. Chesnut, V. K. and Wilcox, E. V. The Stock Poisoning Plants of Montana Bull. U. S. Dept. Agr. Div. Bot. 26.150.
169. Chevalier, J. Considerations sur les causes qui peuvent influencer la teneur en principe actifs des plantes médicinales. Bull. Sc. Pharmac. 18:390.
- 169a. Chevers, N. A Manual of Medical Jurisprudence for India. Ed. 3. Calcutta. 1870.
170. Chrysler, M. A. See Shreve, F.
- 170a. Church, H. J. Medical Botany. London. 1827.
171. Ciaatjian, G. Nature et signification des alcaloïdes végétaux. Rec. Inst. Bot. Bruxelles. 5.
172. Clapp, A. A Synopsis, or Systematic Catalogue of the Indigenous and Naturalized Medicinal Plants of the United States. Trans. Am. Med. Assoc. 1852.
173. Clarke, J. F. Chemistry and Toxicology of Mushrooms. See Chapter 12, pp. 288-298. Atkinson's Mushrooms, Edible, Poisonous, etc.
174. Claude, H. Mycètes aigus par toxines strepto-staphylococciques. Compte. Rend. d. l. Soc. de Biol. 1894:547-550.
175. Clements, F. E. Minnesota Mushrooms. Geol. and Nat. Hist. Survey of Minn. Minn. Plant Studies No. 4. 169.
176. Coats, J. An Inflammatory Lesion of the Kidneys of a Ram with Streptococcus in the Glomeruli and Tubules. Glasgow Medical Journal 1890. 103-112.
- 176a. Codex medicamentarius Galicus Pharmacopée Française. 1908. (Ph. Fr. 1908)
- 177a. Coulon, Eug. Traité de Toxicologie végétale. 210. Paris 1917.
178. Collins, G. W. See Cook, O. F.
179. Colville, Y. E. Post mortem Indications of Death from Poisoning. Oklahoma Medical News 15. 193-196.
180. Combs, Robert Some Cuban Medical Plants. Phar. Rev. 12: 87-91, 109-112. 136. Also Cont. Bot. Dept. Ia. St. Coll. Agr. and Mech. Arts. 5.20.
181. Connor, J. J. On Poisoning from the Internal Administration of Tincture of Rhus Toxicodendron. See Am. Jour. Der. Gen. Ur. Dis. 11.368.
182. Connor, J. T. American Homeopathic Pharmacopoeia. Philadelphia. 1893. Ed. 2.
183. Cook, Ernest H. A Case of Poisoning by the Areca Nut. Chem. News No. 1934. 1895.
184. Cook, O. F. Origin of the Hindi Cotton. Cir. Bur. Pl. Ind. U. S. Dept. Agr. 12.12.
185. Cook, O. F. and Collins, G. N. Economic Plants of Porto Rico. Cont. U. S. Nat. Herb. 8.269.
186. Cooke, N. C. Oriental Aconite. Phar. Jour. III. 8.563-565. 1873.
187. Cordier, Action des alcaloïdes sur les végétaux. Assoc. franc. pour l'Avance des Sci. 1908.388-391.
188. Cornevan, Ch. Les plantes vénéneuses et ses empoisonnements qu'elles déterminent. 524. Paris. 1893.

189. Sur l'empoisonnement par quelques espèces de *Cytisus*. *Compte Rend. des seances de l'Acad. des Sci. de Paris*. **1886**, 13.
190. Venenosité de quelques légumineuses exotiques appartenant aux genres *Templetonia* et *Sophora*. *Compte Rend. hebdom. de la Soc. de Biol. à Paris*. 1893. 29 Avril.
191. Constantin, J. Remarques de la Favus de la Poule. *Bull. de la Soc. de Myc. de France*. **1893** 166.
192. Constantin, J. and Lucet, A. Sur un *Rhizopus* pathogène. *Bull. de la Soc. de Myc. de France*. **19**, 200.
193. Coulter, J. M. Preliminary Revision of the North American Species of *Anhalonium* and *Lophophora*. *Cont. U. S. Nat. Herb.* 8:91-132.
194. Coulter, S. M. Plants Poisonous when Taken into the System. *Ann. Rept. Dept. Geol. and Nat. Resources of Indiana*. **1899** 553-1074.
195. The Poisonous Plants of Indiana. *Proc. Ind. Acad. Nat. Sci.* **1904** 51-63.
196. Couperot, Sur quelques végétaux à acide cyanhydrique. *Jour. Phar. et Chim.* **27** 542.
197. Coupin, H. Contributions à l'étude des substances toxiques pour les plantes. *C. R. Assoc. franc. Acad. Sci.* **30**:414-415. 1902.
198. See Israel, O.
199. Courchet, M. L. On Kironiro Poison not from *Dalbergia toxicaria*, as stated by Baillon, but from one of the *Simarubaceae*. See *Ann. de l'Inst. Colon. de Marseille*. **1905** 195-247. (Rev. in *Phar. Jour.* **1906**:463.)
200. Court, G. See Pictet, A.
201. Courtet, A. Notes sur divers cas d'empoisonnement par les champignons à Pontarlier. *Bull. Soc. myc. France*. **24**:133-137.
202. Coville, F. V. Crimson Clover Hair Balls. *Cir. U. S. Dept. Agr. Div. Bot.* 8 4.
203. Observations on Cases of Mushroom Poisoning in the District of Columbia. *Cir. U. S. Dept. Agr. Div. Bot.* 13-24.
204. Poisoning by Broad-leaved Laurel (*Kalmia latifolia*). *Proc. Am. Assoc. Adv. Sci.* **44**:189. (Abs.).
205. Coville, F. V. and Funston, F. The Botany of Yakutat Bay, Alaska. *Cont. U. S. Nat. Herb.* 3 325-353.
206. Cowley, A. H. See Paul, B. H.
207. Craig, R. A. Cornstalk Disease. *Cir. Pur. Univ. Agr. Exp. Sta.* 3 10.
208. Silage Poisoning. *Supp. to Bull. Pur. Univ. Agr. Exp. Sta.* 147-628.
209. Crawford, Albert C. Barium, a Cause of the Loco Weed Disease. *Bull. Pur. Pl. Ind.* 129-87.
210. Laboratory Work on Loco-weed Investigation. *Bull. U. S. Dept. Agr. Bur. Pl. Ind.* 121 39-48.
211. Larkspurs as Poisonous Plants. *Bull. U. S. Dept. Agr. Bur. Pl. Ind.* 111-1 12.
212. Mountain Laurel, a Poisonous Plant. *Bull. U. S. Dept. Agr. Bur. Pl. Ind.* 121 21 35.
213. Notes, mainly bibliographical, on two American Plants—Sleepy Grass and Creosote Bush. *Phar. Rev.* **26**:230-235. Repr.

214. . The Poisonous Action of Johnson Grass. Bull. U. S. Dept. Agr. Bur. Pl. Ind. 90:31-34.
215. . A Poisonous Principle in certain Cottonseed Meals. Jour. Phar. and Exp. Ther. 1:519-548.
216. Crook, E. and Powell, R. D. . See Powell, R. D.
217. Crookshank, E. M. . On the Morphology, Cultivation and Toxic Products of the Tubercle Bacillus. Lancet. 1:296-300. 1891.
218. Crow, W. E. . Criminal Poisoning in Hongkong. Lecture before Odd Volumes in Hongkong. Rev. in Pharm. Rev. 14:164.
219. Crozier, A. A. . Another Death from Eating Cicuta maculata. Bot. Gaz. 14:17.
- 219a. . Millet. Bull. Mich. Agr. Exp. Sta. 17:64.
220. Cushny, R. . Ueber das Ricinusgift. Archiv für experimentelle Pathologie et Pharmacologie. 41. 1898.
221. Czapek, Dr. F. . . Biochemie der Pflanzen. I 584, II:1026. Jena 1905.
222. Da Gama, J. A. . . Poisoning by Star-anise. Trans. Grant Coll. Med. Soc. 1898, (Bombay 1899):13. Discussion, p. 15.
223. Dakin, R. . . On Rhus Toxicodendron. See Am. Jour. Med. Sci. 4:98.
- 223a. Damen, Carl. . . Die Gesundheitspflege der landwirtschaftlichen Hausausgetiere. Ed. 3. 873. Berlin 1902.
224. Daniell, F. W. . On the Production of Hydrocyanic Acid from Bitter Cassava Root. Phar. Jour. II. 6:312-314.
225. Daniels, C. W. . . A Plea for the Scientific Study of Native Drugs and Poisons. Jour. Malay. Br. Med. Assoc. Singapore. 1895. N. S. Nos. 2, 3, 5.
226. Darjon, Em. . . See Bourquelot, Em.
227. Darlington, Wm. . . Agricultural Botany. Useful Plants and Weeds. Philadelphia 1847.
228. . . American Weeds and Useful Plants, Being a Second Edition of the Above. New York. 1859.
- 228a. Dassonville, C. . . See Matruchot, L.
229. Daugherty, C. M. . The Castor-oil Industry. Yr. Bk. U. S. Dept. Agr. 1904:287-298.
230. Davenport, C. B. and Neal, H. V. . On the Acclimatization of Organisms to Poisonous Chemical Substances. Arch. f. Entwicklungsmech. d. Organ., Leipz. 2:564-583. 1895-1896.
231. Davidson, . . Fatal Case of Poisoning by Ergot of Rye. Lancet No. 2083. (1882).
232. Davies, D. H. . The Gift-apple of South Africa. Chem. and Drug. London. 64:696.
233. Day, M. G. . Experimental Demonstrations of the Toxicity of the "Loco Weed." N. Y. Med. Jour. 49:237.
234. . Loco Weed. Ref. Book of Prac. Ther. by F. P. Foster. 1:587.
235. . Separation of the Poison of the "Loco Weed." N. Y. Med. Jour. 50:604.

236. De Bary, A. Comparative Anatomy of the Vegetative Organs of the Phanerogams and Ferns. Translated and Annotated by F. O. Bower and D. H. Scott. 659. Oxford, 1884.
237. Comparative Morphology and Biology of the Fungi, Mycetozoa and Bacteria. Trans. by H. E. F. Garnsey. Rev. by I. B. Balfour. 525. Oxford, 1887.
238. De Candolle, Alphonse. Origin of Cultivated Plants. English Trans. 468. New York. 1892.
239. Deffernez, Edmond. Triple empoisonnement par le *Datura Stramonium*. Extr. du Bull. med. de Charleroi. 6. 1895.
240. Delacroix, G., See Priehaux, E.
241. Delile, A. Raffineau., Dissertation sur les effets d'un poison de Java appelé Upas tueur, et sur la noix vomique, la fève de St Ignace, le *Strychnos potatorum* et la pomme de Vontac qui sont du même genre de plantes, que l'Upas tueur. 48. Paris. 1809.
242. De lotte Intoxications aiguës produites par l'avis étoilé et les fleurs de genêt. Limousin med., Limoges. 25:36-39. 1901.
243. De Puy, C. E. An Enquiry into the Botanical History, Chemical Properties, and Medicinal Qualities of the *Erigeron Canadense*. 1815.
244. Desaint See Moussu.
- 244a. Deutsches Arzneibuch Ed. 4. 1900. (Ph. Germ. IV.)
245. Dietrich, D. V. F. Deutschlands Giftpflanzen, nach natürlichen Familien aufgestellt mit Abbildungen. 64. 24 col. pl. Jena, 1826.
246. Dieudonné, Jakob., *Cynoglossum officinale*. St. Petersburg 1868. See Eckhard's Beiträge zur Anatomie u. Physiologie.
- 246a. Dinmore, S. C., See Kennedy, P. B.
247. Dinwiddie, R. R. On the Toxic Properties of Molds. Bull. Ark. Agr. Exp. Sta. 40:35-36.
- 247a. Unsound Corn and Forage as a Cause of Disease in Live Stock. Bull. Ark. Agr. Exp. Sta. 25:1-9.
248. Dioscorides, Pedanius Anazarbeus De materia medica libri quinqué. Ejusdem de venenis libri duo. Interprete Jano Antonio Sarraceno. 767. 1598.
249. Doetz, J. C. Neue Versuche und Erfahrung der einige Pflanzengifte. Herausgegeben von J. C. G. Ackermann. 53. Nuremberg. 1892.
250. Dohme, A. R. L. and Engelhardt, H. Further Experiments with Wild Cherry Bark. Phar. Rev. 14:13-15. See also Phar. Rund. 13:260.
251. Doten, S. B. See Kennedy, P. B.
252. Dragendorff, G. Die gerichtlich-chemische Ermittlung von Giften in Nahrungsmitteln, Luftgemischen, Speiseresten, Körpertheilen, etc. St. Petersburg. 1868. Ed. 3. Göttingen. 1888.
- 252a. Die Heilpflanzen der verschiedenen Völker und Zeiten; ihre Anwendung wesentliche Bestandtheile und Geschichte. 1896.
- 252b. Dungen, E. V. Beiträge zur Immunitätslehre. Munch. med. Woch. 1900. Nos. 20 and 28.

253. Dunghison, Robley A Dictionary of Medical Science, New Edition by R. J. Dunghison. Canas Poisoning, 820-829. Philadelphia. 1874.
254. New Remedies. Ed. 4. Philadelphia. 1874
- 254a. Dunghison, R. J. See Dunghison, R.
255. Dunning, H. A. B. See Hynson, H. P.
256. Dunstan, W. R. A Report on the Chemical Examination of the Beans of *Phaseolus lunatus*. Agricultural Ledger. 1905. No. 2 11-16.
257. Dunstan, W. R. and Boole, L. E. Croton Oil. Abs in Proc. Am. Phar. Assoc. 44:641-642.
258. Dunstan, W. R. and Henry, T. A. The Chemical Aspects of Cyanogenesis. Rept. Brit. Assoc. Adv. Sci. 1906:145-157
259. Le glucoside cyanogénétique du lin. Bull. de l'Acad. roy. de Belgique. (Classe des Sciences). No. 7 790-793
260. The Poisonous Properties of the Beans of *Phaseolus lunatus*. Jour. Board of Agr. Great Britain and Ireland. XIV 12:722-731
261. Sur la formation de l'acide cyanhydrique dans les végétaux. Ann. Chim. et de Physique 8 ser 10:118-128.
262. Dunstan, W. R., Henry, T. A. and Aud, S. J. M. Cyanogenesis in Plants. Proc. Royal Soc. London. 78.
263. Dunstan, W. R. and Umney, C. I. Contributions to our Knowledge of the Aconite Alkaloids. II. The Alkaloids of the true *Aconitum Napellus*. III. The Formation and Properties of Aconine and its Conversion into Aconine. Jour. of the Chem. Soc. 51: 1892.
264. Dunstan, W. R., Wynneham, R., and Henry, T. A. Cyanogenesis in Plants. Pt. II. Proc. Roy. Soc. 70 153-154. Part III (by Dunstan and Henry) Proc. Roy. Soc. 72 285-294.
265. The Nature and Origin of the Poison of *Lotus arabicus*. Philos. Trans. Roy. Soc. London. 1943 515-533.
266. Dupony. See De Nabaa.
267. Duvernoy, G. D. De Lathuri quadam venenata specie in comitata Monhelgardensi culta. 19 Basle. 1770.
- 267a. Eames, E. H. See Graves, C. B.
268. Eastwood, A. The Loco-weeds. Zoe 2:53-58.
269. Eccles, R. G. What is a Poison? Am. Med. Philadelphia. 10-986; 1007.
270. Eden, R. T. Digitalis and its Cumulative Action. Ther. Gaz. 6:89.
271. Elborne, Wm. Cocos (or Kokra) wood, Irritant Properties of the Sawdust. Chem. and Drug. 189:85-86. Abs. in Proc. Am. Phar. Assoc. 44:642.
272. Elphinstock, Edward Case of Physostigmine Poisoning. Am. Vet. Rev. 24:50.
273. Elwert, C. P. Gelsemium sempervirens. Proc. Cal. Phar. Soc. 1881. (abs.).
- 273a. Engelhardt, H. See Dohme, A. R. L.

290. Fenn, C. M. Untoward Effects of Cascara Sagrada, *Ther Gaz.* 4:522.
291. Ferguson, A. T. Cerebro-spinal Meningitis or Forage Poisoning. *Am. Vet. Rev.* 34:383.
292. Fermi, C. and Pernosi, L. Ueber das Tetanugift Vergleichende Studien mit Berücksichtigung anderer Gifte und der Enzym. *Zeitsch. f. Hygien.* 10:385-444.
293. Fernald, M. L. See Gray's Manual.
294. Freydel, Paul Toxicité de l'acide filicique 75. Toulouse. 1897.
295. Fischer, A. Vorlesungen über Bakterien. Ed. 2 374 69f. 1903.
296. Fischer, A. and Brebeck. Zur Morphologie, Biologie and Systematik der Kahnpilze, der *Monilia candida* und der Soorerregers. Jena. 1894.
297. Fischer, R. The Alkaloids of *Echoltzia californica*. *Proc. Am. Phar. Assoc.* 49:443-453.
298. Fischer, T. Ein Fall von Stomatitis. *Bacterium stomatog foetidum*, ein aerober Faulnisserreger. *Zeitsch. f. Hyg.* 49:329.
299. Flandin, C. De l'empoisonnement par les végétaux ou leurs principes immédiats ou général, par l'opium ou la morphine en particulier. *Compt. Rend. Acad. Sci. Paris.* 25:150 154. 1847.
300. Fleming, Geo. See Neumann, J. G.
301. Flückiger, F. A. *Pharmakognosie des Pflanzenreiche.* 2 Aufl. Berlin. 1881-1882.
302. Flückiger, F. A. and Ham-
bury David. *Pharmacographia. A History of the Principle Drugs of Vegetable Origin, Met with in Great Britain and British India.* London, 1879.
303. Flugge, C. *Fermenta und Mikroparasiten.* 308. Leipzig. 1893.
- 303a. Die Mikroorganismen mit besonderer Berücksichtigung der Aetiologie der Infektionskrankheiten. 692. Leipzig. 1886.
304. Foelléré, F. E. *Voyage aux alpes maritimes, ou Histoire naturelle agricole, civile et médicale du Comté de Nice et pays limitrophes.* 2 v. 376-426. Paris. 1821.
- 304a. *Traité de médecine légale.* 6v. Paris. 1813.
305. Fogel, Estelle D. See Pammel, L. H.
306. Ford, W. W. Artificial Immunity to Glucosides. *Proc. Am. Assoc. Adv. Sci.* 1908. Abs. in *Science.* 37:652-657.
307. The Toxins and Antitoxins of Poisonous Mushrooms, *Amanita phalloides*. *Jour. Inf. Dis.* 3:191.
308. Antibodies to Glucosides with especial Reference to *Rhus Toxicodendron*. *Jour. Inf. Dis.* 4:541-541.
- 308a. The Distribution of Poisons in Mushrooms. *Science.* 20:97-108.
309. Forrest On Hydrocyanic Poisoning from Kidney Bean. See *Glasgow Med. Jour.* 18:68. 1882.
310. Forster, E. L. Poisons their Source, Use and Antidotes. *Brit. Jour. Nursing.* London 30:64, 185, 225, 246, 309, 353, 397, 429, 469, 508. 31:25, 64, 104, 144, 185.

311. Fraussenet, J. F. F. Poisons d'épreuve tanguin, M'Boundou, strophantus, fève de Caabar Bordeaux. 1896.
312. Francis George Poisonous Australian Lake Nature 18 1.
313. Frankforter, G. B. The Alkaloids of Veratrum. Minn Bot. Studies. 1897:1024-1043
314. Frazer, T. B. and Tilhe, Joseph Wa-Nyika Arrow Poisons. Phar Jour 1896:76-77
315. Free, I. E. Pink-root Poisoning. Med and Sur Reporter 70:633
- 315a. Freeman On Celsium Poisoning. See Lancet 1873:473.
- 315b. Fricberg, L. H. Translator See Kobert, R.
316. Friedberger, F. and Frohner E. Veterinary Pathology Translated by M. H. Hayes With Notes on Bacteriology by R. H. Hewlett Ed 6. 1731, 2702
- 316a. Frohner Eugen . Lehrbuch der Toxicologie für Tierärzte, Ed. 3. 391. Stuttgart. 1910.
317. Frohner E. See Friedberger, F.
- 317a. Frothingham, L. See Page, C. G.
318. Fuchs, F. A. Keratom, Aspergill. Wiener Klin. Wochenschr
319. Leber Keratom als Ursache der Hypopyonkeratis. Graefes Arch. 25 abt 2 p. 285. 1879
320. Gaffky, George Experimentell erzeugte septikämie mit Rücksicht auf progressive Virulenz u akkommodative Zuchtung Mittb d Kaim. Ges. Amt 1881. 126.
321. Gamaleia, N. The Bacterial Poisons. Trans. by E. P. Hurtd. 136. Detroit. 1893
322. Gardiner, W. W. Poisoning from Eating Locust-tree Bark. Am. Vet. Rev. 27:599-600.
323. Garnsey, H. E. F. See De Bary, A.
324. Gattinger, A. The Medicinal Plants of Tennessee. 128. Nashville. 1894
325. Gautier A. Les toxines microbiennes et animales Paris. Soc d'edit. scientif. 1896.
326. Gautier L. and Bodin, E. See Bodin, E.
327. Geiger, Paul Beitrag zur Kenntniss der Ipoh-Pfeilgifte. Mit einem Anhang. Pharmakognostische Mitteilungen über einige zur Heerstellung von Ipohverwendete Giftpflanzen. 103 Bsp.
328. German, John. On Vernal Poisoning. See Jour Am. Med. Assoc. 1906: (No. 26) 1999.
329. Gerrard, A. W. "Wangka," a new African Arrow-poison its Composition and Properties (Strophanthus). Phar Jour. III. 11:833-834 1889. See also pp. 849-850.
330. Gibbs, H. Actinomyces bovis. New Am. Pract 1890-337-339. Chicago.
331. Gibbons, H. Poisonous Effects of Crota aria-Vulgo Rattle Weed, Loco Weed. Pacific Med and Sur Jour 21:496.
332. Poisonous Species of Astragalus. Am. Journ. Phar 51:237-240.
333. Gibbons, W. P. The Medicinal Plants of California Proc. Am. Phar Assoc. 19:297-308. (abs)

- 334 Gibson, R. J. H... . On the Occurrence of a poisonous Alkaloid in "West African Boxwood." *Quart Jour Inst Comm. Research Topics*, **1**:32-40. Liverpool.
- 335 Gilot, X. . Note sur l'action vésicante de *Heracleum*. *Ann. Soc. Bot. Lyons*, **26** 133-150.
- 336 Galman, J. F. . *Conium maculatum*. *Jour belge d'homéopathie*, **6**:254-261 (1899)
- 337 Gimlette, J. D... . On *Datura* Poisoning. See *Brit. Med Jour* **1903** 1137
338. . Several Species in the Malay Peninsula Used for Asthma. *Brit. Med. Jour* **1903** 1137
- 339 Glover, G. H. . Larkspur and other Poisonous Plants. *Bull. Col. Agr. Exp. Sta* 113:24
340. . The Poison Weed Problem of the Arid West. *Am. Vet. Rev.*, **34**:493-502
341. . Poisonous Weeds. *Bull. Col. Agr. Exp. Sta* 137 3-8.
342. Grmelin, J. F. . Abhandlung von den giftigen Gewachsen, welche in Deutschland und vornehmlich in Schwaben wild wachsen. 228. Ulm, Stuttgart. 1775.
343. Allgemeine Geschichte der Pflanzengifte 525. Nuremberg. 1775.
344. Godet, C. . Les Plantes vénéneuses du Canton de Neuchâtel, avec 26 planches coloriées 59 Neuchâtel. 1864
- 345 Godfrey, B. . Santonine Poisoning. (Of two Dogs) *The Veterinarian* **73**:150-151.
346. Godfrey, George. See Michael, L.
- 347 Godlee, R. S. and Powell, R. D. See Powell, R. D.
348. Goeppert, H. R. . Ueber die giftige Pflanzen Schlesiens. Breslau. 1832
349. Gola, G. . L'auido cannabico e i glucosido nel regno vegetale. *Suppl. Encicl. Chimica*, **23**:43-79
350. Good, Peter. A *Materia Medica Botany*. New York 1846.
- 351 Gordon, P. R. See Bailey, F. M.
- 351a Gono, B. . Ricerche Batteriologiche e Chimiche sulle Altecrazione del Man. Contributo all'etiologia della Pellaagra (Memoria 2a) *Rivista d'Igiene e Sanità Pubblica*, **7**:825. 1896.
352. Gotthel. Black Tongue its Etiology. *Archives of Pediatrics*, April **1889**.
353. Goze .. . Unsere Giftpflanzen
- 354 Graham, H. .. *Mucor corymbifer* in the External Auditory Meatus. *Lancet* **2**:1379 1890
355. Grawitz, Paul Beitrage zur systematischen Botanik der pflanzlichen Parasiten. *Virsch. Archiv* **1887**:546.
- 355a Graves, C. B. et al. Catalogue of the Flowering Plants and Ferns of Connecticut growing without Cultivation. 50 Hartford, 1910.
- 356 Gray, Asa . The Animal Poison of the Far West—"Loco or Crazy Weed." *Am Agr* **1878**:380-381
357. . The Botanical Text-Book, an Introduction to Scientific Botany. New York, 1850.
358. A Contribution to the Flora of North America. American Academy of Arts and Sciences. 1883
359. Genera of the Plants of the U. S. 2 vols. New York. 1848.

360. Manual and Lessons in Botany and Vegetable Physiology New York 1874.
361. A Manual of the Botany of the Northern United States. 710. Boston. 1848.
362. Ed. 7 of Gray's Manual. A Handbook of the Flowering Plants and Ferns of the Central and Northeastern United States and Adjacent Canada. Rearranged and Extensively Revised by B. L. Robinson and M. L. Fernald. 926. New York. 1908.
363. A Synoptical Flora of North America. 2 vols. New York. 1878-1884.
364. Green, J Reynolds . . . The Soluble Ferments and Fermentations. 512. Cambridge. 1901.
- 364a. Green, Wasey . . . Plants of Iowa. 264. Bull. St. Hort. Soc of Iowa. 1907.
365. Greenish, H G. . . . An Introduction to the Study of Materia Medica, being a Short Account of the more Important Crude Drugs of Vegetable and Animal Origin. 571. London, 1899.
366. Greenleaf, R. W . . . A Case of Acute Poisoning. Rept. Boston Med. Surg. Jour.
367. Greimer, Karl . . . Ueber giftige Boragineen Alkaloide. Phar Zeitung. 43: No. 20.
368. Ueber giftig wirkende Alkaloide einiger Boragineen. 1900.
369. Greshoff, M. . . . The Distribution of Prussic Acid in the Vegetable kingdom. Rept. Colonial Museum Haarlem, Holland. 1906. (Reprint).
370. Rept. Brit Assoc Adv Sci. 1906:138-144.
371. Monographie der giftigen en betwermende plantem bij de vischvangst in gebruik. 2 vols. completed, v 3 incomplete. 1893-1910.
372. Phytochemical Investigations at Kew. Bull. Misc. Inf. Roy Bot. Gar. Kew. 1908:397-418.
373. Gressler, F G. L. . . . Deutschlands Giftpflanzen mit zahlreichen Illustrationen und 8 Tafeln in Farbendruck. 16 Aufl. 40. Langensalza. 1897.
374. Grey George . . . Herb-poisoning at the Cape of Good Hope. Phar Jour III 5:248-250. (1879).
375. Grigorescu, G . . . Action des substances toxiques sur l'excitabilité des nerfs et des muscles périphériques: un antidote de la strychnine. Arch de physiol norm. et Path. Paris. 1894. ser 66:32-39.
376. Grohe . . . Experiments über die Injektion der Pilzsporen von *Aspergillus glaucus* u. *Penicillium glaucum*. Berlin Klin. Wochenschr. 1870.
377. Grönlund Chr . . . Danske Giftplanter. Kjøbenhavn 1874.
378. Gross, Herman . . . Locust Bark Poisoning Horses. Breeder's Gazette. 55:455.
379. Gruby . . . Recherches anatomique sur une plante cryptogame qui constitue le vrai muguet des enfants. Compte Rend. des Séan de l'Acad de Paris. 14. 1842.
380. Guardia, J . . . Poisonous Plants and their Poisons. Bul of Phar 7:542-548. 1893.
381. Gueguen, J. . . . Les champignons vénéneux et leurs caractères. Revue scientifique 1908:321-328.
382. Toxicologie des champignons. Symptômes, causes et traitement des empoisonnements. Rev sci. 1908:361-366.

- 383 Guenther, J and Bartuch, F Pinakothek der deutschen Giftgewächse. 16. Jena. 1840.
384. Guerin, P The Probable Cause of the Poisonous Effect of the Darnel, *Lolium temulentum* Bot Gaz. 28:136-137
- 385 Guignard, L. Le haricot à acide cyanhydrique (*Phaseolus lunatus*, L.) Rev. Vit. 1906.
386. Nouveaux exemples de Rosacées à acide cyanhydrique. C. R. Acad. Sci. Paris. 143: 461
- 387 Recherches physiologiques sur la greffe des plantes à acide cyanhydrique Ann se Nat ser 9. Bot. 6:261-305
388. A Study of Beans yielding Hydrocyanic Acid. Rev Vit. 28:5-9, pl. 1, 33-37, 341-350 f.p. 543-547, 573-576, 626-631; 663-687, 689-694, 715-721
389. Sur la localisation des principes actifs chez les Tropeolées. Compt Rend des Séan de l'Acad. des sci. de Paris. 117:587-590 Abs. in Beihefte zum Bot. Centrbl. 4:220-221
390. Sur l'existence, dans le sureau noir d'un composé fournissant de l'acide cyanhydrique. Bull. Sci. Pharmacol. 12 and 13
- 391 Sur la greffe des plantes à acide cyanhydrique. C. R. Acad. Sc. Paris. 30 Dec. 1907
392. Sur la métamorphose des glucosides cyanhydriques pendant la germination. C. R. Acad. Sc. Paris. 147:1023
393. Sur la prétendue toxicité des Haricots de Hongrie. C. R. Acad. Sc. Paris. 9 Dec 1907
394. Sur les quantités d'acide cyanhydrique fournir par le *Phaseolus lunatus*. L., cultivé sous le climat de Paris. Bull. Sc. Phar. 14:565.
395. Guttenberg, Gustave. Poisonous Plants which Grow in and around Erie. (Read before the Nat. Hist. Soc. Sept. 5, 1870. 21-25
396. Hager, Hermann Ueber die giftige Wirkung einiger Lathyrus-Arten. Monatliche Mittheilungen aus dem Gesamt-Gebiete der Naturwissenschaften 6:153.
- 397 Haines, W. S. Alkaloidal and other Organic Poisons. In Hamilton and Godkins System of Legal Medicine. 1:415-473.
398. See Peterson.
- 399 Halderman On Gaultheria Poisoning. See Lancet. 1886.
- 400 Hale, E. M. The special Symptomology of the New Remedies. Philadelphia. 1877
- 401 Halvand, F. Etude sur les trichophytes de la barbe Arch. de Parasit 8:590.
- 402 Hale, J. S. Die deutsche Giftpflanzen. 2 v 119+126. 24 col. pl. 1784-1795.
- 403 Halsted, B. D. Notes on Poisonous Plants. Gar and For 8: 172.
- 404 Other Poisonous Plants. (*Maclura aurantiaca* and *Sicyos angulata*.) Bot. Gaz. 19:200
405. The Poisonous Plants of New Jersey Bull. N. J. Agr. Exp. Sta. 135:28

406. Preliminary Report of the Above. Rept Bot Dept. N. J. Agr. Exp. Sta. **1894**:401-419 1895:351-355.
407. Hamilton On Recovery from Gaultheria Poisoning. See N. Y. Med. Jour **1875**.
408. Hamilton, A. M. and Godkin
A System of Legal Medicine 1894. Articles on Poisons by C. E. Pelieu, W. S. Haimes, and V. C. Vaughan.
409. Hamilton, W. On the Genus *Jatropha* and the *Jan pha* Manihot and *J. Loeflingii*. Phar Jour **5**:27-31 31-33.
410. On the Noxious Properties of the *Echites suberecta* or Savanna flower and their Antidotes. Phar. Jour **8**:25-28. (1846.)
411. On the Properties of the Hippomane *Manchella* or Manchinel Tree. Phar Jour **5** 408-412.
412. Hanbury, David Note on the Ordeal Bean of Calabar (*Physosigna venenosum* Balf.) Phar Jour, **11**. 4 559-561. (1863.)
413. See Flückiger, F. A.
414. Hansen, N. E. The Wild Alfalfas and Clovers of Siberia, with a Perspective View of the Alfalfas of the World. Bull. Bur Pl. Ind. 156-31.
415. Hare, R. F., Mitchell, S. R. and Berregard, A. P. Denatured Alcohol from Tunas and other Sources. Bull. N. M. Agr. Exp. Sta. 72 57.
- 415a. Harger, E. B. See Graves, C. B.
416. Harknis, W. D. and Swan, R. E. The Chronic Arsenical Poisoning of Herbivorous Animals. Jour Am Chem Soc. **30**: 928.
417. Harlan, Herbert Blindness and Death from Drinking Essences of Ginger, Peppermint, etc., due to Methyl Alcohol. Phar. Rev. **19**:51-53.
418. Harnack, E. and Hochheim, W. Ueber die temperatur erniedrigende Wirkung kramperregender Gifte. Ztschr. kun. Med. Berlin **25** 16-45.
419. Harnack, E. and Meyer, H. Untersuchungen uher die Alkaloide der Jakobandblätter. Lie. Ann. d. Chem. **204**: 67-84.
420. Harrington, C. On Poisoning by Canned Beans, Caused by Toxins rather than by Living Bacteria. Botulism suggested. Rev. in Am. Jour Med. Sci. **133**:811. From Article by Roly. Münch. Med. Wochenschr. **1906**. No 37 1798.
421. On Poisoning from Foods. Rev. in Am Jour Med. Sci. **133**:810. From Art. by Dr. W. G. Willoughby in Pub. Health. **1906**:626.
422. Harrison, W. H. The Poison Ivy. Gar and For **8**:68.
423. Harshberger, J. W. Ergotism, (Account of an Outbreak at Scranton, Pa., due to Eating Red Top. *Agrostis alba vulgaris* with *Claviceps microcephala*. (Wallr.) Tul.) Rept Pa. Dept. Agr. **1902** 160.
424. *Ranunculus acris*, L. An Additional Poisonous Plant. Bot. Gaz **19**:159. Gar and For **7** 170.

425. A Review of our Knowledge of Phytobezoars.
Jour. of Com. Med. and Vet. Arch **19**:
143.
426. When is *Rhus Toxicodendron* most Active?
Gar. and For **8**:239.
427. Hartley, C. P. . . . Broom Corn. Far. Bull. U. S. Dept. Agr.
174-30.
428. Hartman, G. . . . Mechanische Vergütungen. Berlin 1896.
429. Harvey, P. F. Observations on the Toxic Effects of some
Common Drugs. N. Y. Med. Jour **78**
148-150.
430. Hartzel, Stella . . . See Michael, L. G.
431. Harz, C. D. Landwirthschaftliche Samenkunde **2** 1315.
- 431a. Haslam, Thos. P. . . . Meningo-Encephalitis. Bull. Kan. Agr. Exp.
Sta. 173-233-251 (Contains Bibliography.)
432. Hatch, W. G. On Loss of Lamb from *Rhus* Poisoning. See
Cal. State Med. Jour **1**:465.
433. Havard, V. Drink Plants of the North American Indians.
Bull. Torr. Bot. Club. **23**:33-46.
434. Heald, E. D. See Israel, O.
435. Hebert, A. Recherches sur la presence de l'acide cyan-
hydrique chez diverses plantes. Bull. de la
Soc. chim. de Paris, 1896, No. 74:310.
436. Sur les principes actifs des fruits d'un *Strych-*
nos africain. Jour. Phar. et Chem. **27** 151.
437. Heckel, E. Nouvelles sur le Tanghin du Ménabé (*Mena-*
bea venenata, Baill.) et sur sa racine tox-
ique et medicamenteuse. C. R. Acad. Sc.
Paris, **134**:441-443.
438. Sur la *Menabea venenata* Baillon, qui four-
nit, par les racines, le Tanghin de Ménabé
ou des Sakalaves (poison d'épreuve) en-
core denommé Kistoumpa ou Kimanga, à
Madagascar, sa patrie. C. R. Acad. Sc.
Paris, **134**:364-366.
439. Sur un autre poison des Sakalaves appelé
Komanga ou Kimanga et fourni par l'*Ery-*
throphloeum Couroungou Baillon. Rev. des
cult. Coll. **11**:129-134.
440. Heckel, E. and Schlagden-
hauffen, Fr. Sur la *Copaifera Salikounda* Heckel de l'
Afrique tropicale et sur ses graines à cou-
marine (*Salikounda* des peuples Sousous,
au point de vue botanique et chimique. Com-
paraison avec la fève de Tonka. Ann. de
la fac. des sci. de Marseille III Sep-abd.
1-18.
441. Hedges, Florence . . . See Smith, E. F.
442. Hedrick, U. P. Cicuta, a Plant that Poisons Cattle. Bul. Ore
Agr. Exp. Sta. 46:12.
443. Heffter, A. Pellotina, a new Hypnotic. Phar. Rev. **114**:
153-154.
444. Ueber Cacteen Alkaloide. Ber. deutsch. chem.
Gesell. **34**:3004.
445. Heft, J. J. Ein Beitrag zur Kenntniss der speziell in
central Europa. Giftpflanzen und Pflanzen-
gifte. Schwanden, 1889.

446. Hegetschweiler, J . Flore des plantes vénéneuses de la Suisse contenant leur description l'époque de leur floraison, les lieux, l'indication de celles qui sont employées en médecine, les symptômes qu'elles produisent et les premiers soins à donner dans les empoisonnements. Destinée à l'usage compagne. 117 23 pl. 1849.
447. Hegetschweiler, J. and Labram, J. D. Die Giftpflanzen der Schweiz. 46. Zurich.
448. Hehr, P. The Sale of Poisons in India. Ind. Med. Chir. Rev. 3:351-360. Bombay.
449. Hehn, Heinrich. Deutschlands Giftpflanzen. Eine kurze Beschreibung der giftigen und verdächtigen einheimischen Pflanzen nebst Hinweis auf deren Verwendung in der Pharmacie, zugleich als Text für die vom Verwässer, herausgegebenen Giftpflanzenherbarien für Lehrer, Schüler, und zum Hausgebrauche. 74. Hamburg. 1880.
- 449a. Hektoen, Ludwig. Systematische Blastomycose und coccidioidales Granulom. Chari-Festschr. herausgeg. von Prof. Paul Dittich in Prag. Wien u. Leip. 1908 116-142.
450. Hempel, C. J. . New Homoeopathic Pharmacopoeia of Jahr, Buchner, and Gruner. New York. 1856.
451. Henkel, Alice. . American Medicinal Barks. Bull. Bur. Pl. Ind. U. S. Dept. Agr. 139 49.
452. . Weeds Used in Medicine. Far. Bull. U. S. Dept. Agr. 188 45.
453. . Wild Medicinal Barks of the United States. Bull. Bur. Pl. Ind. U. S. Dept. Agr. 89 76.
454. Henry, Arne. . Die Giftpflanzen Deutschlands zum Schulgebrauch und Selbstunterricht durch Abbildungen und Beschreibungen erläutert. Bonn. 48. 32 col. pl. 1836.
455. Henry, J. . . Les principales plantes vénéneuses. Brussels.
456. Henry, T. A. . . See Dunstan, W. R.
457. Henslow, G. . . Poisonous Plants in Field and Garden. 189. London. 1897.
458. Herissey, H. . . Ueber das Blausäure liefernde Glykosid der Samen von Eriobotrya japonica. Arch. der Phar. 245:469.
459. . Ueber das Prulaurocerasin, das Blausäure liefernde Glukosid der Blätter von Prunus lauro-cerasus. Arch. der Phar. 245:463-469.
460. . . Ueber das Vorkommen des Prulaurocerasins in Cotoneaster micropaylla, Wal. Arch. der Phar. 245:473.
461. Herissey, H. and Lefebvre, Ch. Ueber das Vorkommen der Raffinose in Taxus baccata. Arch. der Phar. 245:481 486.
462. Herman, L. . Untersuchungen zur Physiologie der Muskeln und Nerven. Heft 3.
463. Herzog, M. Disease Producing Microorganisms. 644 New York. 1910.
464. Hetier, Fr. Empoisonnement par l'Entoloma hydnium. Bull. de la Soc. myc. de France. 18 127 129.
465. Heyer, F. Ueber die Eigenthümliche giftige Wirkung einer zu Leguminosen gehörigen Futterpflanze. Landw. Post. Berlin. 1887. April. 26.

466. Hickman, R. W. Epizootic Cerebro-spinal Meningitis of Horses. Rept. Bur. Ani. Ind. U. S. Dept. Agr. **1906**, 165-172.
- 466a. Hulger, A. See Husemann, A.
467. Husemann, T. D. Feeding of Millet to Horses. Bul. N. D. Agr. Exp. Sta. 26:89-105.
468. Rheumatism in Horses. Bull. N. D. Agr. Exp. Sta. 7:15.
469. Hitchcock, A. S. Rape as a Forage Crop. Far. Bull. U. S. Dept. Agr. 164:16.
470. Hochsetter, C. F.. Die Giftgewächse Deutschlands und der Schweiz. Esslingen, 1884.
471. Hoffman, E. R. Lehrbuch der gerichtlichen Medizin. Ed. 5. Vienna, 1890-1891.
472. Hoffman, Fr. Loco Weeds. Phar. Rund. **1889**, 168.
473. Hoffman, Ralph. Virulence of the Wild Parson. Rhodora. 4:188.
474. Holm, Theo. Medicinal Plants of the United States. Merck's Report 1907-1910 (not yet complete).
475. Holmes, E. M. Agricultural and Horticultural Poisons. Chem. and Drug. London, **63**:247.
476. Rhus Toxicodendron. Phar. Jour. **81**:231-232.
477. Some Books of Medicinal and Toxicological Interest. Phar. Jour. Ser. IV. **18**:5-7.
478. Vegetable Poisons and their Antidotes. Phar. Jour. III. 4:769-770.
479. Homen, E. A. De l'action du streptocoque et de ses toxines sur les nerfs, les ganglions apicaux et la moelle épinière. C. R. le la Soc. de Biol. **1896**:518-523.
480. Hooper, D. The Bark of the Cleistanthus albus as a Fish Poison. Phar. Jour. IV. 1465.
481. Horsfield, Thomas. An Essay on the Copra or Poison Tree of Java, Addressed to the Hon. Stamford Raffles, Lieut. Gov.
482. Howard, C. D. Occurrence of Barium in the Ohio Valley Brines and its Relation to Stock Poisoning. Bull. W. Va. Agr. Exp. Sta. 103:281-295.
483. See Morse, F. W.
484. Huard, Louis. Les alcaloïdes, les leucomaines, et les ptomaines. Ann. de Phar. **1899**. No. 5.
485. Hubbard, R. On Poisoning from Internal Administration of Rhus Preparations. See Med. Rec. **30**:601.
486. Hugemeyer Ueber Abschwächung pathogener Schimmelpilze. Bonn, 1888.
487. Hughes, D. A.. Loco-weed Poisoning and Barium Salts. Am. Vet. Rev. **33**:613-617.
488. Poisonous Plants Destructive to Live Stock with Special Reference to a Recent Book on the Subject. Am. Vet. Rev. **86**:672-679.
489. Hughes, Richard. A Manual of Pharmacodynamics. Ed. 3. London, 1876.
490. Hunt, R. and Jones, W. Non-alkaloidal Organic Poisons. In Text-book Leg. Medicine (Peterson and Haines) 2:532-643. Philadelphia, 1904.
491. Huntington, Annie O. Poison Ivy and Swamp Sumach. 58 p. Jamaica Plain, Mass. 1908.

492. Huseman, A., Hilger, A.
and Huseman, Th. Die Pflanzenstoffe in chemischer, physiologischer und toxicologischer Hinsicht 320.
Berlin, 1883.
493. Huseman, Th. The Distribution of Cardiac Poisons in the Vegetable Kingdom. (abs.) Phar. Jour III 7:795-796, 834-835.
494. Poisonous Property of "Star Anise" Phar. Jour III 11:453-454. (1880) From Phar Zeit
495. Hutcheon, D. Poisoning of Horses by *Printhogalum thyrsoides* or "Chuckerinchee" (*Ornithogalum thyrsoides*) Agr Jour Cape of Good Hope 28:165-172
496. Hutyra, Franz, and Marek, Josef Specielle Pathologie und Therapie der Haus-thiere
497. Hyams, C W Medicinal Plants which have been Collected and Used in North Carolina Bul N C Agr. Exp. Sta. 150 329-409.
498. Hyde, J V See Montgomery, J H
499. Hynson, H. P and Dunning H. A B Examination of "Jamaica Ginger" that Caused Blindness. Phar Rev 19:54-61
500. Illinois State Board of Health Principal Poisonous Plants of the United States. Springfield, 1899. (Repr. of Chestnut's Principal Poisonous Plants, etc.)
501. Imbert-Gourtheyre For the Identity of Hemlock and the Athenian Poison, see "De la mort de Socrate" Paris, 1875.
502. Immunität (Die) von Pflanzen gegen ihr eigenes Gift. Zeitsch f ang Mikr 13:60-62. Leipzig, 1907
- 502a. Index Catalogue (Of the Surgeon General's Office, U S Army Ser 2. 13 594-603. Bibliography on Poisons and Allied Subjects). See also Corresponding Volume of Ser 1 There are many cross references to other volumes that are helpful.
503. Inokn, Y Zur Kenntniss der Pilzvergiftung Fortschritte der Medicin 1893:444-449
- 503a. Irving On Aconite See Med. Times and Gaz, 1:63. 1882
504. Israel, O., Klingman, Tb., Kahlenberg, L., Trine, R H., Head, E. D. and Conpin, H Action of Poisonous Plants by Stronger and Weaker Solutions. Centrbl f Agr 28: 8.-188.
505. Itaire, L. van Sur l'existence dans le *Thalictrum aquilegifolium* d'un composé fournissant de l'acide cyanhydrique Jour de Phar et de Chimie Oct. 1906.
506. *Thalictrum aquilegifolium*, eine Blausäure liefernde Pflanze Arch Phar 243:553-554. Abs. in Bot. Centrbl. 102 5.7
507. Jackson, J R Poisonous Indian Peas Phar Jour 76:521-522.
- 507a. Jacob, J Poisons et Contrepoids Paris 1877
508. Jacot, J Definition des Wortes Gift Deutsch. med. Wochenschr 26:2-4.

- 508a. Jacoby, M. Antiscin, antiabrin, antocrocin. See von Kraus and Levaditi's *Handbuch der Technik*, etc 2:255-262.
509. Jahn, E. Ueber Giftpilze und Pilzgifte. *Apotheker-Zeitung*, 13:No. 93.
510. Jeanmire De la nocivité relative et temporaire de l'*Amanita junquilles*. *Bull. Soc. myc. de France* 24:176-180.
511. Jelliffe, S. M. See Wilcox, R. W.
512. Jennings, O. E. A Case of Poisoning by *Amanita phalloides*. *Jour. Myc.* 13:187.
513. Jesup, H. G. Is *Cypripedium spectabile* Poisonous to the Touch? *Bot. Gaz.* 18:142.
514. Jewett On *Gaultheria* Poisoning. See N. Y. *Med. Gaz.* 1867:68.
515. Johnson, Charles British Poisonous Plants. 59. London. 1856.
516. Johnson, Lawrence A Manual of the Medicinal Botany of North America. 292. New York. 1884.
- 517 On Toxicity of Tincture of *Rhus Toxicodendron* which had been Kept Thirty Years. See *Med. Bot.* 119.
518. Jones, L. R. Venomous Shrubs and Woody Vines. *Bull. Vt. Agr. Exp. Sta.* 145:56-199.
- 518a See Rich, F. A.
519. Jones, W. See Hunt, R.
520. Jørgensen, A. La linamarine glucoside générateur d'acide cyanhydrique. Réponse à la Note de MM. Dunstan et Henry. (*Bull. de l'Acad. Roy. de Belgique—Classe de Sci.* 7:773-778. *Abstr. in Bot. Centralbl.* 107:242-243).
521. Juch, K. W. Die Giftpflanzen. Zur Belehrung für Jedermann. Augsburg. 1817.
522. Judkins, F. H. Antidotes to Poison. *N. Albany Med. Herald* 16:270.
523. Knäsch, W. Das Pflanzenleben der Erde. Eine Pflanzengeographie für Laien und Naturforscher. Nach dem Tode des Verfassers mit einem Vorwort versehen von H. A. Berlepsch. 642. Hanover. 1855.
524. Kahlenberg, L. See Israel, O.
525. Kalm, P. On Poisoning from *Rhus Toxicodendron*. See *Travels* 1:79. 1770.
526. Karsten, Georg See Strasburger, E.
- 526a. Kartulis, S. Ueber Blastomycosis glutealis fistulosa. *Zeitschr. f. Hyg. u. Infektkr.* 84:285.
527. Kaufmann, Paul Nouvelle expériences sur l'injection le spores de *Aspergillus glaucus*. *Lyon Med.* No. 1832. No. 10.
528. Kaupp, B. F. Acute Poisoning. *Am. Vet. Rev.* 30:454-456.
529. Kebler, L. F. Poisonous Holes. *Proc. Am. Phar. Assoc.* 4:167-173.
530. Keeler, D. D. Poisoning by Ergot. *Am. Vet. Rev.* 38:251-252.
531. Kellerman, W. A. Poison Ivy and Ivy Poisoning. *Ohio Naturalist* 14:2:227.
- 532 Poisoning by Shepherd's Purse. *Bot. Gaz.* 20:325-326.
533. Kellogg, A. California and Loco Poisoning. *Proc. Cal. Acad. Sci.* 8:3.
- 533a. Kennedy, G. W. *Kalmia latifolia*. *Am. Jour. Phar.* 47:5.
534. Kennedy, P. B. Summer Ranges of Eastern Nevada Sheep. *Bull. Nev. Agr. Exp. Sta.* 55:53.

535. Kennedy, P. B. and Dinmore, S. C. Digestion Experiments on the Range. *Bull. Nev. Agr. Exp. Sta.* 71:20-32
536. Kennedy, P. B. and Doten, S. R. A Preliminary Report of the Summer Ranges of Nevada Sheep. *Bull. Nev. Agr. Exp. Sta.* 51:56.
537. Kerner, Anton von and Oliver, F. F. Poison from Stinging Hairs. See *The Natural History of Plants*. 1:441
538. Kerner, J. D. Deutschlands Giftpflanzen. Hannover 1798.
539. Kerner, J. On Rhus Toxicodendron. See *Abh. Am. Jour. Phar.* 30:542
540. King, John American Dispensatory Cincinnati 1877-1879.
541. King, John, and Lloyd, J. L. Supplement to the American Dispensatory Cincinnati. 1880
542. Kingsley, A. T. Gastro-enteritis in Pigs. *Am. Vet. Rev.*
- 542a. Kinnebrew, C. See Michael, L. G.
543. Kirtikar, K. R. The Poisonous Plants of Bombay. *Jour. Bombay Nat. Hist. Soc.* 1892:61-76, 312, 317, 447-493
544. Kitt-Thou Bakterienkunde. 1899.
545. Klees Handbuch der pathologische Anatomie 1869.
546. Klingman, Th. See Israel, O
547. Knoke, Plantae venenatae in belliferae. *Levinsk.* 177.
548. Knowles, M. F. Loco Poisoning. *Becce's Gazette* 89:372
549. Knowlton, P. H. The Loganiaceae. *Mexico's Report* 10:363.
550. Kober, A. Ueber Arum pectoratum L. Sitzungsber. der Naturf. Gesell. bei der Univ. Dorpat 8:114-117
551. Kober, E. R. Compendium der praktischen Toxikologie 1903.
552. Lehrbuch der Intoxicationen. 816. Stuttgart. 1893
553. On the Toxic Constituents of Jatropha Curcas and Croton Tiglium. *Bull. of Phar.* 7:200 (1893)
- 553a. Practical Toxicology for Physicians and Students. Translated and Edited by L. H. Friedburg. 201. New York. 1897
- 553b. Ueber Giftstoffe der Flechten. *Sep. Abh. aus Sitzungsber. der Dorpat Naturf. Gesell.* 1892:156-186
554. Koch, D. G. D. J. Synopsis Florae Germanicae et Helveticae, etc. Ed. 3. 875. Leipzig 1877
555. Koeng, A. The Principal Poisonous Plants of the United States. *Penn. Med. Jour.* 2:260-262
556. Kohlhaus, J. J. Giftpflanzen auf Stein abgedruckt mit Beschreibungen. Regensburg. 1805
557. Kohlschreier, F. Sur les legumes toxiques. *Rev. internat. d. falsific.* 20:25. Paris. 1907
558. Kraemer, Henry A Textbook of Botany and Pharmacognosy. 850. Ed. 3. Philadelphia. 1905
559. Some of the distinguishing Morphological Characters of Belladonna Scopolia. *Proc. Am. Phar. Assoc.* 56:819-824. f. 77-79
560. Krapf, K. von Experimenta de Ranunculorum venerea qualitate. Vienna. 1838.
561. Kraus, R. V. and Levaditi, C. Handbuch der Technik und Methodik der Immunitätsforschung. 2 v. 1138+1219. Jena. 1908 (Contains complete Bibliographies)

- 561a Krause, A. See Mueller, G.
- 562 Krause, M. Tier-und Pflanzengifte in den deutschen Kolonien. Arch. b. Schiffs- u. Tropen Hyg. **11** 650-656. See also Verhandl. d. Gesellsch. deutsche Naturf. u. Aerzte **1907** Leipzig 1908. 2 Teil. Heft 349.
- 563a Vergleichende Untersuchungen über Pflanzengifte. Glykoside und andere Glykoside der Digitalis-Gruppe mit Hilfe des Brechungsexponenten und der Dispersion. Zeitsch. f. experim. Pathologie u. Therapie. **1**. 1905.
- 563b. Kren and Schramek Ueber Sporotrichose. Wiener klin. Wochenschr. **1909**. No. 44.
564. Kreutzer, K. J. Oesterreichs Giftgewächse. Vienna 1838.
565. Krocke Zur Lupinenkrankheit der Schafe. Landw. Jahrb. **9**:27-35.
566. Kubinyi, A. Plantae venenosae Hungariae. Merges novenyik, etc. Buda. 1842.
567. Kunkel, A. J. Handbuch der Toxicologie. 1117. Jena 1901.
568. Kye, H. M. On the Action of the Spurge on Salmonoid Fishes. Proc. Roy. Soc. **70**:48-66.
569. Labasse Un Champignon vénéneux servant à la fabrication des poisons chez les peuplades de l'Afrique Centrale. Anjou Med. Angers. **10**:132-135. 1903.
570. Labram, J. D. See Hegetschweiler, S.
571. Lacerda, J. B. de Experiencias fisiologicas con algunas plantas toxicas do Brazil. Rio de Janeiro. 1890.
572. Laud, E. F. Water Hemlock Poisoning. Bul. N. D. Agr. Exp. Sta. **44** 563-569.
573. See also Am. Chem. Jour. **20**:262.
574. Lahitte, Henry Erigeron canadense. Phar. Post. **1887**:802-803.
575. La Glesch, Louis, and Vuillet, Jean Plantes médicales et toxiques du Soudan français. Schweiz. Wochenschr. Chem. Pharm. **40**:546-548.
576. Lambert, L. See Noel, C.
577. Lang, W. H., Translator See Strasberger, E.
578. Langguth, G. A. Programma de plantarum venenatarum ar-cendo scelere. 12. Wittenberg 1770.
579. Larsen, C. See Michael, L. G.
- 579a. Latte For Bibliography of earlier cases of Cocaine Poisoning see Diss. Berlin 1888.
580. Laurent, Emile Sur l'existence d'un principe toxique pour le Pourrier, dans les baies, les graines et les plantules du Gui. C. R. Acad. Sci. Paris. 1901.
581. Lavender, C. H. Pellagra à Précis. Pub. Treas. Dept. Marine Hospital Service U. S. A. **22** 1 pl. Wash-ington, 1902.
582. Law, James Text-book of Veterinary Medicine. **2**:601.
- 582a. Lazell, F. J. Gathering Mushrooms. Cedar Rapids Repub. **1910**. Sept. 25.
583. Leather, J. W. Cyanogenesis in Plants. Agric. Jour. India. **1**:220-225.
584. Leclerc, N. De l'empoisonnement et des moyens que l'on doit employer pour constater. Strasburg 1803.
585. Lefebvre, G. See Herissey, Ch.

586. Legrand du Saulle, H.,
Berryer, G., and Pouchet,
G. *Traité de médecine légale, de jurisprudence
médicale et de toxicologie, etc.* Ed. 2
Paris, 1886.
587. Legue, G. *Medicins et empoisonneurs au XVII^e siècle*
Paris, 1896.
588. Lehmann, Otto *Giftpflanzen mit besonderer Berücksichtigung
der wirksamen Stoffe und zahlreiche in den
Text gedruckten Illustrationen, 126.* Ham-
burg, 1892.
589. Lehmann, K. B., and Mori,
R. *Ueber die Giftigkeit und Entgiftung der Sa-
men von Agrostemma Githago.* *Arch v f.*
Hyg. 9:257. (*Bot. Gaz.* 14:233,
590. L'Engle, E. M. See Price, T. W.
591. Leuterer, Dr. *Investigations of Poisonous Characters of*
Macrozamia. *Chem. and Drug.* 1896:171.
- 591a. Levaditi, A. See Kraus, R. V.
592. Lev G. *Materia Medica pharmacologica e toxicologia*
italiana de veterinaria 388. Milan 1899.
- 592a. Lewandowski, F. *Ueber Impetigo contagiosa s. vulgaris (nebst*
Beiträgen zur Kenntnis der Staphylo- und
Streptokokken bei Hautkrankheiten) *Arch.*
dermat. u. Syphilis 94. Heft 2 u. 3.
Anhalonium Lewini. *Ther. Gaz.* 4:231.
593. Lewis, I. *Arrow Poisons. Review of Different kinds*
and their Sources. *Phar. Centrid.* 1895:
509-510. See also *Naturw. Rund.* 1895:
86. Also in *Proc. Am. Phar. Assoc.* 44:
508-509.
595. *Erste Hilfeleistung bei Vergiftungen. Cong.*
internat. d'hyg. et de démog. C. r. 1894,
Budapest, 1896. 3:286-294 pt. 5.
596. *The Flaya Poison and Erythrophlaene.* *Ther.*
Gaz. 4:145.
597. *Lehrbuch der Toxicologie.* Ed. 2, x + 509. 7f
7 pl.
598. *Die Pflanzengifte. Historische und experimentelle*
Untersuchungen. 152. Berlin 1894.
599. *Ueber Anhalonium Lewini und andere giftige*
Cacteen. *Ber. der Deutsch. bot. Gesell.*
1894:283-290.
600. *Was ist Gift?* *Deutsch. Med. Wochenschr.*
32:1748.
601. Liastard, A. *Poisoning of Horses by Snow Drop.* *Am.*
Vet. Rev. 30:298.
602. Lignozat *Empoisonnement par la racine d'aconit*
*(Aconitum gummifera, el heddad des Ara-
bes)* *Arch. de med. et phar. m.*, Paris
37:164-168. 1901.
603. Linderle d., Fr. *Dissertatio circa Phytotoxicon agnam rechinam,*
*plantas venenatas Cechiae indigenas Umbelli-
feras exhibens.* 108. Prague 1834.
- 603a. Linderberg, Adolph *Un nouveau mycetome.* *Arch. de Parasit.*
13. No. 2.
604. Linley, John *Flora Medica. A Botanical Account of all*
the more Important Poisonous Plants Used
in Medicine in different Parts of the World
606. London, 1838.
605. Linsley, W. I. *The Tool-poison of New Zealand.* *Phar.*
four. U. 5:371-373. 1864. (*Proc. Brit.*
Assoc.)

606. Linné, Carl v. De Raphanis. 21 Upsala. 1763
607. Lenoissier G. and Roux, G. Recherches morphologiques sur le champignon du muguet. Arch. exp. med. et d' anat. pathol. **1890**:63-87
608. Sur la morphologie et la biologie du Champignon du muguet. Compte Rend Acad. Sci. **1889**.
609. Sur la mycose expérimentale due au champignon du muguet. Lyon medical. **1889**.
610. Last, A. Untersuchungen über die in und auf dem Körper des gesunden Schafes vorkommenden niederen Pilze. Dissertation. Leipzig. 1885
- 610a. Lattle. On Poisoning by Sam sp. Cinic Cincinnati. **7**:49. 1874.
611. Lloyd, C. G. Mycological Notes. Cincinnati Series of Papers covering many of the important fleshy fungi.
- 611a. Lloyd, J. U. See King, John.
612. Lloyd, J. U. and C. G. Drugs and Medicines of North America. I. Ranunculaceae 304 Cincinnati 1884-1885
613. Lodeman, E. G. On Recurrence of Attacks of Rhus Poisoning without Reexposure. See Gar and For **8**:368.
614. Loew O. Ein natürliches System der Gift-Wirkungen. Munch. 1893.
615. Loudon, J. C. An Encyclopedia of Plants 1574 London, 1846.
616. Lubeman. On Poisoning from Foods. See Centrbl. f. Bakt. **1906**. I. Abt 433.
617. Lucas E. De solano tuberoso ejusque principio narcotico **32**. Hirschberg 1846.
618. Lucet A. De l'aspergillus fumigatus chez les animaux domestiques et dans les œufs en incubation (étude clinique et expérimentale) 108 Paris. 1899
619. Also see Constantin, J
620. Luerksen, Ch. Medicinisch-pharmaceutisch Botanik. Leipzig
- 620a. Lutembacher. See Pattrier
621. Lyons, A. B. Assay of Third Extract of Belladonna Leaves. Phar. Rev. **26**:22-24
622. Plant Names. Scientific and Popular. Ed 2. 630. Detroit. 1907
623. McCarthy, Gerald. The Poisonous Plants of North Carolina. Bull. N. C. Agr. Exp. Sta. (St. Bd. Agr.) **22**:48.
624. McChntock, C. F. See Vaughan, V. C.
625. MacDougal, D. T. On the Poisonous Influence of *Cypripedium spectabile* and *C. pubescens*. Minn. Bot. Studies. **1894**:32-36.
626. Poisonous Influence of various Species of *Cypripedium*, Minn. Bot. Studies. **1895**. 455-451
627. McFaydean, J. Poisoning Horses by Spoiled Potatoes. Jour. Roy. Soc. England. **1897**. Ser III (No. 29). 115-135. Aba. in Exp. Sta. Rec. **9**:389
628. McGinnis, J. W. Intoxication from Alfalfa Hay. Am. Vet. Rev. **38**:238-239.
629. McIlvaine, C. M. Fungi, Edible and Poisonous. Am. Jour. Phar. **68**:651
630. McKelway, F. L. Three Cases of Poisoning by Potassium Cyanide. Am. Jour. Med. Sci. **129**:684
631. MacMurtrei, Conway. Metaspermae of the Minnesota Valley. Minn. Nat. Hist. and Geol. Sur. Bot. Ser. **1**:826.

- 632 McNeill, J. H. and Pammel, L. H. The Danger of Feeding Hay that Contains Ergot. Press Bull. la Agr. Exp. Sta. 1909.
- 632a MacQueen, James See Neumann.
- 632b Madsen, B. A. See Michael, L. G.
- 633 Madsen, Thorwald Botulinismustoxin. Chapter in von Kraus and Levitt's Handbuch der Tech. u. Meth. etc 1:137-143. (Contains Bibliography.)
- 633a Maguin, L. Les cas d'empoisonnement par l'Amantia muscaria. Bull. de la Soc. myc. de France. 19:173-175.
- 634 Maider, J. H. Native Plants Poisonous to Stock. Agr. Gaz. 6:57-59. New South Wales.
- 635 Notes on Some American and Australian Plants Injurious to Stock. Agr. Gaz. 4:679.
- 636 Plants Reported to be Poisonous to Stock in Australia. N. S. W. Dept. Agr. Misc. Pub. No. 477. 1901.
- 637 Poison Ivy. Roy. Soc. N. S. Wales. 1903:3.
- 638 Lethal Native Plants of Australia including Tasmania. 693. London. 1890.
- 639 Marsh, J. M. California Opium. Am. Jour. Phar. 46:106.
- 639a On Rhus Toxicodendron. See Proc. Am. Phar. Assoc. 13:166.
- 640 Poisonous Species of Astragalus. Am. Jour. Phar. 51:239.
- 641 Sueszweed. Proc. Am. Phar. Assoc. 20:231-236.
- 642 Makoni Leber Alkaloide der chinesischen Corydalis Knollen. Arch. der Phar. 246:381.
- 643 Mann, E. A. Examination of the Western Australian Poison Plants. Jour. Dept. Agr. W. Austr. 12:550-561.
- 643a Mann, J. D. Forensic Medicine and Toxicology. London and Philadelphia. 1893.
- 644 Mantheim On Cocain. See Zeitsch. f. Klin. Med. 1890. 30.
- 645 Marchand, L. Du Croton Tigium recherches botaniques et therapeutiques. 94. 2 pl. Paris. 1861.
- 645a See Numan, A.
- 646 Marcus, H. D. Poisons and their Treatment. Internat. Clinie 4:48. Philadelphia. 1898.
- 647 Marek, J. See Huttya, Fr.
- 648 Marsh, C. W. The Loco Weed Disease. Far. Bul. U. S. Dept. Agr. 390:16.
- 649 Results of Loco-weed Investigation in the Field. Bull. Bur. Pl. Ind. U. S. Dept. Agr. 121:37-38.
- 650 Marsh, D. C. Water Bloom. Bull. Wis. Geol. and Nat. Hist. Sur. 12:36.
- 650a Marshall, H. T. See Bacon, R. T.
- 651 Maschwa, J. v. Handbuch der gerichtlichen Medicin 4 v. Tullingen, 1831-1832. (The Toxicology is contained in Vol. 2, and is by B. Schachardt, M. Seidel, Th. Husemann, and A. Schauerstein.)
- 652 Mason, P. De la rapidité d'absorption des poisons par l'organisme. Arch. Internat. de pharmacod. 11:465-481. Brussels and Paris. 1903.
- 653 Massey, W. F. Medicinal Plants which have been Collected and Used in North Carolina. Bul. N. C. Agr. Exp. Sta. 150:331-409.

- 653a. Matruchot, L. and Dassonville, C. On Dermatomycozosis of Fowls. *Rev. Gen. Bot.* **11**:429-444. *Abstr. of Exp. Sta. Rec.* **12**:94.
654. Maurizio, Adam Les Maladies causes aux poisons et aux oeufs par les champignons. *Revue Myc.* **1897** (No. 75): 79-85. *pl.* 170-172.
655. Mayo, N. S. Erizootic Cerebritis, or "Staggers" of Horses. *Bull. Kan. Agr. Exp. Sta.* **24** 107-116.
- 655a. Same Observations upon Loco. *Bull. Kan. Agr. Exp. Sta.* **35** 113-119.
- 655b. Maza, M. G. de la Flora Habanera. *Fanerogams.* **1**:268, **2**:269-597.
656. Mead, Richard A Mechanical Account of Poisons. London. 174.
657. Meo, George Poisonous Principle of the Bitter Cassava Root. *Phar. Jour.* **11**. 322-333.
658. Meltzer, S. J. On the Absorption of Strychnine and Hydrocyanic Acid from the Mucous Membrane of the Stomach, an Experimental Study on Rabbits. *Trans. Assoc. Am. Phys.* **11**:221-228. Philadelphia. 1896.
659. Melvin, A. D. and Mobler, J. R. Dermal Mycosis Associated with Sarcopic Mange in Horses. *Proc. Am. Vet. Med. Assoc.* **1907**:175. *2 pl.* See also *Rept. Bur. An. Ind.* **24** 259-275.
660. Meiner Empoisonnement par l'Amanita phalloides à Noir-Montier (Vendée). *Bol. Soc. Mys. de France* **24**:68-71.
661. Merck, E. Annual Report of Recent Advances in Pharmaceutical Chemistry and Therapeutics. **23**. See also earlier Volumes. See Merck's Weekly Report.
662. Merrill, E. D. The Occurrence of Antiaris in the Philippines. *Phil. Jour. Sci. Sect. C* **2**:111.
663. Meyer, Arthur Beiträge zur Kenntnis pharmaceutisch wichtiger Gewächse. *Archiv d. Phar.* See Harnack, E.
664. Meyer, H.
665. Michael, L. G., with the Collaboration of R. T. Robbins, Carl Larsen, R. A. Madson, Stella Hartzell, Claude Kinnebrew, and George Godfrey The Influence of Feeding Sugar Beets and Mangels to Breeding Animals with Reference to the Formation of Renal and Urinary Calculi. *Bull. Ia. Agr. Exp. Sta.* **112** 35-77.
666. Miquel, F. A. W. De Noord-Nederlandsche vergiftige Gewassen. **220**. Amsterdam. 1838.
667. Millspough, C. F. American Medical Plants. An Illustrated and Descriptive Guide to the American Plants Used as Homeopathic Remedies, their History, Preparation, Chemistry, and Physiological Effects. **1**:99 *pl.* 1-99, **2**:100-190. *pl.* 100-180.
- 667a. Mitchell, S. R. See Hare, R. F.
668. Mülbacher, W. Toxicologisch oder forensisch wichtige Pflanzen und vegetabilische Drogen, mit besonderer Berücksichtigung ihrer mikroskopischen Verhältnisse. Berlin and Vienna. 1904.
669. Moeller, J. Digitalis and Verbascum. *Phar. Post* **37**: 677-680.

- 669a Moellier, Josef. See Winton, A. L.
- 670 Moffit, J. E. A Case of Vegetable Poisoning (Anthracosis luteola) Med. Gaz. **16** 155. Australia.
- 671 Mohler, J. R. Mycotic Stomatitis in Cattle. Cir. Bur. An. Ind. U. S. Dept. Agr. 5136.
- 672 Mohler, J. R. See Melvin, A. D.
- 673 Mohler, J. R. and Buckley, J. S. Pulmonary Mycosis of Birds with Report of a Case in a Flamingo. Rept. Bur. An. Ind. U. S. Dept. Agr. **20** 122-138, pl. 6-10.
- 673a Mohr, Ch. Plant Life of Alabama. 921. Geol. Sur. of Alabama. 1901.
- 674 Mole, Ph. La localisation des alcaloïdes dans es Solanacees. Bull. de la Société belge de microscopie **21** 28-20. 1894-1895.
- 675 Montgomery, F. H. and Lyde, J. N. A Practical Treatise on Diseases of the Skin. Ed. 7. 1904.
- 676 Moore, G. T. The Contagion of Public Water Supplies by Algae. Year Book U. S. Dept. Agr. **1902**: 175-186.
- 677 Mori, R. See Lehmann, K. B.
- 678 Moorman, J. W. On Poisoning from Internal Administration of Rhus Preparations. See Am. Jour. Med. Sci. **51** 570.
- 679 Morris, Daniel On the Use of Certain Plants as Alex. vari- eties or Snake-bite Antidotes. Annals of Botany. **1**:153-161.
- 680 Morrow, P. A. Drug Eruptions. 165. New York. 1887.
- 681 Morse, F. W. and Howard, C. L. Poisonous Properties of Wild Cherry Leaves. Bull. N. H. Agr. Exp. Sta. **56**:113-123.
- 682 Moussau and Desaint. Poisoning of Sheep by Galega officinalis. Rev. Med. Vet. **64** 562-569. Abs. in Exp. S. a. Rec. **19** 785.
- 683 Moutet. Un nouveau cas d'intoxication par l'association de la fougère mâle et de l'huile de ricin. Bull. med. de l'Algérie. **1** 263. ser. 2. Algiers. 1903.
- 684 Mueller, Ferdinand Von Notes on Leguminous Poisonous Plants of Southwest Australia. Ann. Chem. and Drug. **1884**.
685. See also Trans. Roy. Soc. Victoria. **6** 47.
- 685a Also Jour. of Bot. **3** 325.
686. Second Systematic Census of Australian Plants. Pt. 1. 224. Melbourne, 1889.
686. Select Extra-Tropical Plants readily Engage for Industrial Culture or Naturalization. With Indications of their Native Countries and Some of their Uses. 449. Detroit. 1884.
- 687 Mueller, Georg and Krause, C. Ueber die Giftwirkung der Anemone nemorosa. Archiv. für wissenschaftliche u. praktische Tierheilkunde. **23**:326-335.
688. Mueller, O. Selbstmord durch Vergiftung mit bitteren Mandeln. Ztschr. f. Med.-Beamt., Berlin. 1907. **20**:37.
- 689 Mueller, Karl. Colchicum autumnale. Die Natur. **44** 550.
690. Mueller, R. See Baumgarten, P.
691. Murrel, W. What to do in Cases of Poisoning. Ed. 10. London. 1907.

692. Water, John An Introduction to Pharmaceutical and Medical Chemistry. I 397, II 216. Philadelphia, 1820
693. Mutual Life Insurance Company of New York Poisons Remedies 45 New York 903.
694. De Nabias and Dupont Sur un poison d'épreuve du Haut Oubanghi. Jour de med. de Bordeaux **33**:273 1903.
695. Nagelvoort, J. B. Notes on Andromedotoxin in Leaves of *Kalmia latifolia*. Drug. Bull. **1890** 77
- 695a. Neat, H. V. See Davenport, C. B.
696. Nelson, A. Squirrel-tail Grass (Fox-tail), one of the Stock Poats of Wyoming. Bull Wyo Agr Exp. Sta. 19 73-79
697. Nelson, S. B. Feeding of Wild Plants to Sheep. Rept U S Dept Agr Bur An. Ind. **10**:425
698. Feeding Wild Plants to Sheep. Bul Wash Agr Exp Sta 73-64
699. Nestler, A. Das Hauptgift der Cypripeden. Weiser-Festsch **1908** 200-206.
700. Das Sekret der Drüsenhaare der Gattung *Cypripedium* mit besonderer Berücksichtigung seiner hautreizenden Wirkung. Ber der Deutsch. Bot Gesell. **28**:554-567 1 pl
701. Die hautreizende Wirkung der *Primula Obconica* Flance and *Primula sinensis* Lindl. Sep. Ber. der Deutsch. Bot Gesell. **18**:189-202 pl 7-8.
702. Hautreizende Primeln. Berlin, 1904
703. Neumann, L. G. A Treatise on the Parasites and Parasitic Diseases of the Domesticated Animals. Trans by Fleming. Ed. 2. Rev and Rev. by Jas MacQuern. 697 New York 1907
- 703a. Nichols, H. J. See Phasen, J. M.
704. Nibernurt See Widal.
705. Nocke de C. Poisoning by Loco Weed. Am Vet Rev **20** 570.
706. Noe, J. Action de divers poisons sur les animaux herbivores (herissons); variabilité et spécificité des effets des substances toxiques. Archives internat de pharmacod **12** 153-158. Brussels and Paris. 1903
707. Noel, Charles Contributions à l'étude médicale de l'Anémone Pulsatilla. 81 Nancy 1897
708. Nor, Ch. and Laubert L. Recherches expérimentales sur l'Anémone Pulsatilla. Arch inter de phar **1897**:Fasc 1-2
709. N. Frey F. Plantes vénéreuses croissant dans les prairies et dans les artificiels. Agriculture nationale **1898**. No. 15
710. Nou, Fritz See Strassberger, F.
711. Novy F. G. The Toxic Effects of the Bactera of Hog Cholera. Med. News **2** 23 237
712. See also McClintock, C. F.
713. Nicknolds, Coleman Vesicant Dermatits. Am Vet Rev **25** 635-659.
714. Neman, A. and Marchand, I. Sur les propriétés nuisibles que les fourrages peuvent acquérir pour différents animaux domestiques par les productions cryptogamiques. 115. 5 pl. Groningen 1830
715. Neen Joshua Veterinary Toxicology. New York. 1901 1907
716. Oesterle See Tschirsch, A.
717. Olver, F. W. See Kettner, Anton von.

- 718 Oppenheim, Carl Toxines and Antitoxines. Trans. by C. A. Mitchell. London. 1906.
- 719 Opwyrd, R. J. See van Praag, L.
- 720 Orcutt, C. R. The Calandrina Plant. West American Scientist 7:190.
- 721 The Loco Weed. do 7
- 722 Orfila, M. J. A Popular Treatise on the Remedies to be Employed in Cases of Poisoning and Apparent Death. Trans. by W. Price. Philadelphia. 1818.
- 723 Traité des poisons tirés des règnes minéral, végétal, et animal ou Toxicologie générale. Paris. 1813.
- 724 Traité de Toxicologie. Ed. 5. Reviewed, Corrected and Considerably Enlarged. 2 v. Paris. 1852.
- 725 Ot, J. Physiological Action of Astragalus mollissimus. New Remedies 2:227.
- 726 Otto, K. Die vorzüglichsten in Thüringen wildwachsenden Giftpflanzen mit besonderer Rücksicht auf ihren Standpunkt in Fürstenthum Schwarzburg-Rudolstadt. 28. 15 pl. Rudolstadt. 1834. Ed. 2. 1842.
- 727 Patlach, H. Ueber eine Tuba-Wurzel (Derris elliptica, Benth.) Ein Beitrag zur Kenntniss der indischen Fuchsigifte. Phar. Centralhalle. 26:697-706.
- 727a Page, C. G. Sporothrix and Epizootic Lymphangitis. Jour. Inf. Dis. 23:137-149. (Contains Bibliography)
- 727b. Page, J. B. See Page, C. G.
- 727c. Patriet and Lutembacher Sub-cuti-réaction positive obtenue chez deux sporotrichosiques par l'injection sous-cutanée de cultures jeunes de sporotrichose, broyées, diluées dans du sérum et stérilisées. Compt. Rend. Soc. de Biol. 97. No. 24.
- 727d. Palan, W. A. A Treatise on Epizootic Lymphangitis. London. 1904.
728. Pammel, L. H. Regginton alba and the Dying of Fish in Iowa. Aba. in Proc. Ia. Acad. Sci. 1889:90-91. (Vol. 1.)
- 728a. Loco Weeds. Vis Medicatrix 1:40.
729. Mechanical Injuries and Obstructions from Grasses. See Bull. Ia. Geol. Sur. 1 176-182.
730. Is Parsnip Poisonous? Car and For 8:228.
731. Poisonous Effects of Grasses. See Bull. Ia. Geol. Sur. 1 171-176.
732. Poisoning from Cowhane. Bull. Ia. Agr. Exp. Sta. 28 215-228.
733. Poisonous and Medical Plants of Missouri. Bull. Mo. St. Bd. Hort. 14 46.
734. Poisonous Weeds. See pp. 423-448 in "Some Weeds of Iowa." Bull. Ia. Agr. Exp. Sta. 70.
735. Squirrel-tail Grass or Wild Barley. Bull. Ia. Agr. Exp. Sta. 30 302-319.
736. Vergiftung durch Wasserschierling (Cicuta virosa, var. maculata.) Phar. Rund. 13:102-103. 4 f.
737. Also see McNeil, J. H.
738. Pammel, L. H. and Fogel, E. D. A Catalogue of the Poisonous Plants of Iowa. Proc. Ia. Acad. Sci. 14 147-172.

739. Pammet, L. H., Hall, C. R.,
and Scribner F. L. Grasses of Iowa Pt. 2. 436. Bull. Ia. Geol.
Sur. Suppl. Rept. 1904
740. Pammet, L. H., Weems, J.
B., and Scribner, F. L. Grasses of Iowa. Bull. Ia. Geol. Sur. 1 525
741. Park, R. *Dermatitis venenata*, or *Rhus Toxicodendron*
and its Action. 1879.
742. Parker, G. H. Report on the Organisms excepting Bacteria,
found in the Waters of the State. July 1887-
June 1889. Report St. Bd. Health of Mass.
1890-581-620
743. Parker, W. T. Loco-Weed. Science 23 101
744. Patek, Johann. Die Giftpflanzen. Prague. 1866-1867
745. Paul, B. H. and Cownley,
A. H. *Gleditsia triacanthos*. Phar. Jour 1887
746. Pauwels, W. M. I. B. Bijdrage tot kennis der Surinaamsche Viscer-
giften. 87. Leyden. 1903.
747. Pavesi V. Studi comparativi su tre specie di papaveri
nostrali. Atti dell'ist. bot. di Pavia. 9:45.
748. Payne, G. T. On Death from *Rhus* Poisoning. See Proc.
Am. Pharm. Assoc. 42 135.
749. Pearson, L. A Preliminary Report on Forage Poisoning in
Horses (So-called Cerebro-spinal Meningi-
tis). Jour. Comp. Med. and Vet. Arch.
21:654-657
- 749a. Epizootic Lymphangitis in Horses and Mules.
Cir. Penn. St. Livestock Sanitary Bd. 8.
750. See Atkinson, V. T.
751. Peck, Charles H. Annual Report of the State Botanist of New
York. 43 colored Plates. 1 Booke and Pro-
sonymous Fungi with Full Descriptive Text.
241. Albany. 1897
752. The Black Spruce. Albany Institute. 1875
753. Peckolt, Th. The Medicinal Plants of Brazil. Phar. Arch.
1898 No. 9
754. Peuler, A. and Warbler H. On the Nature of the Toxic Principle of the
Aroniaceae. Jour. Asiatic Soc. of Bengal, Cal-
cutta. 1890 106-117
755. Pellagra. See Boston Evening Transcript. "The Choke"
Aug. 18. 1900
756. Notes on Pellagra. McClure's 34:104-107
- 756a. Pellew, C. E. See Hamilton, A. M.
757. Pellon, H. H. See Wilcox, R. W.
758. Penhallow, D. P. *Rhus* Poisoning. Gar. and Hor. 8:359
- 758a. Penn, W. A. The Sovereign Herb. A History of Tobacco.
326. London. 1901
759. Percival, C. H. *Ranunculus acris* as an Irritant. Jour. Bot.
Brit. and Foreign. 32:184
760. Percy, S. R. An Inquiry into the Properties of *Veratrum*
viride. Trans. Am. Med. Assoc.
761. Perkins, J. The Leguminosae of Porto Rico. Cont. U. S.
Nat. Her. 10:220.
762. Pernossi, L. See Ferni, C.
763. Perredez, P. E. F. The Anatomy of the Stem of *Derris alagnosa*,
Benth. Bull. Wellcome Chem. Res. Lab.
23 10.
764. Perret, A. H. Recherches des poisons pruritiques dans les vé-
gétaux. C. R. Soc. de Biol. Paris 59:602-
604.
765. Perrot, Em. Sur la Ksopo, poison des Sakalaves (*Menabea*
venenata). Revue des Cultures Colon. 10:
See also C. R. Acad. Sci. Paris 134:303-
306.

766. Peters, A. T. See Alway, F. J
767. Peters, A. T., Slade, H. B.,
and Avery, S. Poisoning of Cattle by Common Sorghum and
Kaffir Corn. Bull. Neb Agr Exp. Sta. 77
16.
768. Peterson, F. and Haines,
A. F. American Text-Book of Legal Medicine and
Toxicology 815. 1903.
769. Peterson, Maude G. How to Know Wild Fruits. A Guide to
Plants when not in Flower, by Means of
Fruit and Leaf 340 1905.
770. Peyre, A. See Rabuteau.
771. Pfaff, Franz Ivy Poisoning and its Treatment. Bull. Roy.
Bot. Gar. Kew 1903 13-16.
772. On the Active Principle of Rhus Toxicoden-
dron and Rhus venenata. Jour. Exp. Med.
2 181-195 1897.
773. Poisonous Plants. Phar. Rund 9:8. 1891.
774. Pharmacopoeia of the United States. 6th Re-
vision New York 1882.
775. Pharmacopoeia Svedica. Ed. 7, 290. Stock-
holm. 1899. Review.
776. Pfaff, Fr. and Orr, S. S. On Rhus Toxicodendron. See Science n. s.
1.119.
- 776a Pharmacopoeia of the United States of Amer-
ica 8th. Dec. Rev 1905. (U. S. P. VIII).
- 776b Phalen, J. M., and Nichols,
H. J. Blastomycosis of the Skin in the Philippines.
The Military Surgeon. 242 1900.
- 777 Phillips On Case of Poisoning by Sium aquaticum.
See Trans. Me. Med. Assoc. 10:147 1889.
778. Phoebus, Philipp Deutschlands kryptogamische Giftgewächse in
Abbildungen und Beschreibungen. 114 p
col. pl. Berlin. 1838.
779. Pictet, A. La constitution chimique des alcaloïdes végé-
taux 1897.
- 779a Quelques considérations sur la genèse des alcaloïdes
dans les plantes. Archiv. des Sci.
phys. et nat. 18 329-352.
780. The Vegetable Alkaloids with particular Ref-
erence to their Chemical Composition. Lon-
don. 1894.
781. Pictet, A. and Court, G. Sur quelques nouveaux alcaloïdes végétaux.
Bull. Soc. Chim. France. 1V. 1:1001-106.
782. Pictet, A. and Wolfenstein,
R. See Wolfenstein, R.
783. Piffard, H. G. Materia Medica and Therapeutics of the Sk-
97 New York. 1885.
784. Practical Treatise on Diseases of the Skin.
1891.
785. Pilzmerkblatt Berlin. 1906.
786. Pinkham On Gaultheria Poisoning. See Trans. Mass.
Med.-Leg. Soc. 1878:27 Boston.
- 786a Piper, C. V. Flora of Washington. Cont. U. S. Nat. Herb.
11:637.
787. Pischek, A. Die Giftpflanzen in der Umgebung von Cilli.
Program der K. R. Staats Gymnasiums
Cilli. 1884 513.
788. Planchon, Gustave Plantes qui fournissent Curare Jour de Phar.
1880. Paris.
789. Sur les plantes qui servent de base aux di-
vers Curares. Compte Rend. 90 133. Paris.

790. Planchon, Louis Accidents causes par le contact du *Rhus Toxicodendron*. Montpellier Medical. 1887 Ju. and Sept.
791. Plant Poisons Sci. Am. **93**:361
792. Plato, C. G., Deutschlands Giftpflanzen zum Gebrauche für Schulen fasslich beschrieben. Leipsic, 1829-1840
793. Beitrag zur systematischen Stellung des Sootpilzes in der Botanik. Leipsic 1885.
794. Das organisirte Contagium der Schafpocken und die Mitigation desselben nach Toissaint's Manier. Leipsic, 1882.
795. Die Hyphenpilze oder Eumyceten in W. Kofe u. A. Wasserman Handbuch der pathogenen Mikroorganismen. **1**:526-660 f 55 Jena
796. Pirneck, J. J. Toxicologia, seu doctrina de venenis et antidotis. Vienna 1785.
797. Plowright, C. B. Six Fatal Cases of Poisoning by *Amanita phalloides* (Vaill.) at Ipswich in September 1907. Trans. Brit. Myc. Soc. **1908**:25-26.
798. Plugge, P. C. Das Alkaloid von *Sophora lomentosa*, L. Archiv. der Pharm. **229**:561-165
799. Andromedotoxin haltige Ericaceen. Arch. der Pharm. **229**:552-554
800. Giffiger Honig von *Rhododendron ponticum*. Arch. der Pharm. **229**:554-556.
801. Ueber Andromedotoxin den giftigen Bestandtheil der *Andromeda japonica*. Phar. Zeitsch. f. Russland. **1883**:609-613, 625-631
802. Die Wichtigsten Heilmittel in ihrer Wechselnden chemischen Zusammensetzung und Pharmakodynamischen Wirkung. Jena, 1866.
803. Poh, J. E. De plantis venenatis umbelliferis. 4. Leipsic 1771
- 803a .. See also Arch. f. Exp. Path. u. Ph. **34**:265 1894
804. Poison Register, Universal Containing also Poison Acts of the various States, Lists of Poisons in Groups, with Antidote Treatment, Maximum Doses, etc. Chicago. 1898
805. Porcher, F. P. The Medical, Poisonous and Dietic Properties of the Cryptogamic Plants of the United States. 126. New York. 1854. From the Trans. Am. Med. Assoc. **7**
806. Report on the Indigenous Medicinal Plants of South Carolina. Trans. Am. Med. Assoc. **1849**
807. Resources of the Southern Fields and Forests. Richmond. 1863.
808. Pouchet, G. Importance de l'action exercée par certains albuminoïdes sur la variation de toxicité des alcaloïdes et des principes actifs. Nouv. remèdes. **15**:289-293. Paris. 1899.
809. Powell, R. D., Godlee, R. S. -
and Taylor, H. H. With Report by E. Crook on a Case of Actinomyces hominis. Med. chir. London. **1889**:175-210
810. Power, F. B. Notes on the So-called Loco-weeds. Phar. Rund. **8**:29-38

- 811 Power, F. B. and Cambier, Jacob
On the Chemical Constituents and Poisonous Principle of the Bark of *Rolania pseudacacia*. *Phar Rund.* 8:29-38.
- 812 Praag, L. van, and Opwyda, R. J.
Leerboek voor practische gifteer 3 Pts. Utrecht 1871
- 813 Praschl, W. W.
Plantae venenatae in terra nostra Vindobonense sponte crescentes. 41. Vienna. 1840
- 814 Prusnitz, C.
Die Heisterberggifte von Kraus and Levaditi's Handbuch der Technik und Meth. etc. 1. 317-329 (Contains a Bibliography)
- 814a
814b Prentiss, A. N.
He is heer antioxin, id. 2:262-277
Is the Strawberry Poisonous? *Bot Gaz* 13:19
- 815 Prentiss, D. W.
Five Cases of Mushroom Poisoning, three of which Proved Fatal. Treatment of the Poisoning. *Sep Phil Med Jour* Sept. 24. 1898 16:51
- 816
Report of Cases of Poisoning from *hualgine*, *Cannabis indica*, *Arsenic*, and *Camphor*. *Theor Gaz* 1892:103-106
- 817 Prescott, A. B.
Alkaloidal Poisons. In Text book, *Legal Medicine* (Peterson and Haines) 2:447-531
- 818
Laboratory Notes. A Partial Analysis of the *Oxytropis Lambertii*. *Am Jour Phar* 50:564.
- 819 Price, T. M.
Enzymes in Corn Stalks and their Relation to Cornstalk Disease. *Cat Bar An. Ind. U. S.* Dept. Agr 84:75
- 820 Price, T. W. and L. Cogle E. M.
A Fatal Case of Poisoning with the Oil of *Gaultheria*. *Am Jour Med Sci* 127:265. 1904.
- 821 Price, W., Translator
See Orfila, M. J. B.
- 822 Prillieux, E. and Delacroix, G.
Maladie de l'oeil produite par le Macrosporium parasiticum, *Thum*. *Bul de la Soc Myc de France* 1893:201
- 823 Pritzel, G. A.
Tesaurus literaturae botanicae omnium gentium inde a cernit botanicarum pulvis ad nostra usque tempora, quiddecim milia operum recensens. 547. Leipzig. 1851
- 824 Procter
Poisonous Plants. *Pract Phar*
- 825 Procupin, G.
La Pellagre. Paris 1903
- 826 Prudden, T. M.
On the Poisonous Products of the Tubercle *Bacillus*. *New York Med. Jour* 2:281-284
- 827 Puhb, J. G.
Materia veneraria regni vegetabilis 196. Leipzig 1785
- 827a Quain, E.
Report of six Cases of Tubercular Ulcers and Tubercular Lymphangitis of the Upper Extremity. *St Paul Med.* 1904. (Aug)
- 828 Rahuteau, A.
Elements de Toxicologie et de Medicine Legale, appliquees à l'empoisonnement. Paris 1873. Ed. 2. Edited by E. Bourgoing 1888
- 829 Radtkofer, L.
Leber fischvergiftende Pflanzen. *Sep Abd. Bayerisch. Akad. der Wissenschaften in München* 16:379-416.
- 830 Rafinesque, C. S.
Medical Flora or Manual of the Medical Botany of the United States of North America. 1:268; 2:276. 48 pl.

- 831 Rambusch, W. T. Are the Leaves of the *Primula obconica* Poisonous? *Am. Gar* **17**:215
832. Ransom, W. B. A Case of Actinomyces of the Orbit, with a Summary of Seven other Cases of Actinomyces. *Brit. Med. Jour* **1898** 1353-1555.
- 832a. Ratti. On Colchicum Poisoning. See *Phar Jour* and *Trans.* III. 6:47
833. Ratzeberg, J. F. C. See Brandt, J. F.
834. Rawton, Oliver de. Vegetales que curan y vegetales que atan. **348**. Paris. 1888
835. Read, Alex. On the Use of *Datura Stramonium* in Certain Cases. 1816.
836. Reed H. S. An Alleged Excretion of Toxic Substances by Plant Roots. *Nature*. **78**:541 London. 1908.
- 836a. The Fungus, *Diplodia*, as a Possible Factor in the Aetiology of Pellagra. *Repr. N. Y. Med Jour.* Jan. 22 **1910**:18.
837. Reese, J. J. Text-book of Medical Jurisprudence and Toxicology. Ed. 2 Phila. 1889
838. The Malignant Effect of certain Trees upon Surrounding Plants. *Plant World* **10**:279.
839. Renner. Native Poisons West Africa. *Jour Afr Soc.* **1904**:109-111
840. Ribbert M. H. V. Ueber den Untergang pathogen Schimmelpilzener organismus. 59. *Vers D Naturf* **1886**.
841. Rice, W. S. Suspicious Character of the Wounds. *Outing* **40**:551-554
842. Equisetum Poisoning. *Am. Vet Rev* **36**:944-951
843. Rich, F. A. Poisonous Equisetum. *Proc Am. Vet Assoc* **1902** 178.
- 843a. Rich, F. A. and Jones, L. R. A Poisonous Plant the Common Horsetail. *Bull. Vt. Agr. Exp. Sta* **95**:187-192.
844. Richardson. An Epidemic of Acute Poisoning in Horses. *Am. Vet. Rev.* **23**:566-571
845. Pichet, C. Les poisons convulsivants. *Arch. internat. de pharmacod.* **4**:293-309 Paris. 1897-1898.
846. Ricard-Madonna, J. B. Recherches et expériences sur les poisons d'Amérique, tirés des trois règnes de la nature, etc. (Du Brinvilliers, *Spigelia anthelmintica*, L., II Du Mancenilier, *Hippomane Mancinella*, L.) 169. Bordeaux. 1826.
847. Riecke, V. A. See Berge, F. E.
848. Robin, C. Histoire naturelle des végétaux parasites qui croissent sur l'homme et sur les animaux vivants. Paris. 1853.
849. Robinson, B. L. See Gray's Manual
850. Robbins, E. L. See Michael, L. G.
851. Rochebrune, A. T. de Toxicologie africaine. Etude botanique, historique, ethnographique, chimique, physiologique, thérapeutique, pharmacologique, posologique, etc., sur les végétaux toxiques et suspects propres au continent africain et aux îles adjacentes. Avec figures dans le texte par Charles Ricard. *Bull. de la Soc. de Paris.* Fas. 1-5. 768. 1896-1898. (Not completed.)
852. Rochussen, F. Phytochemistry and Hay Fever. *Phar Rev* **26** 167-176.

853. Rodigas, Em Fleurs vénéneuses. Bull. d'arboricult. et de floricult. potagère. 1889:281
854. Rodin, H Les plantes médicinales et usuelles, des Champs Jardins Forêts. Paris, 1882
855. Roily On Poisoning by Canned Beans. Month Med. Wochenschr. 1906. No. 37 1798.
856. Roretz, A. von Camphor Production in Japan. Proc. Am. Phar. Assoc. (abst.) 24:129
857. Rosenthal, D. A Ueber Giftpflanzen aus der Klasse der Leguminosen. Schles. Gesellsch. 1884:57
858. Rossi, P De nonnullis plantis, quae pro vénéneis habentur, observationes et experimenta a Petro Rossi Florentinae instituta. 66. Pisa. 1762
859. Rotenbiller von On Gaultheria Poisoning. See Klinisch. Ther. Wochenschr. May 1900.
- 859a. Rothe, L Ueber die Agglutination des Sporotrichon des Beurmann durch Serum von Aktinomykose. Deutsch. med. Wochenschr. 1910:30
860. Roub, J. F Poisoning by Feeding upon *Sinapis nigra*, Am. Vet. Rev. 26:437-439
861. Roux, G. See Linossier, G
862. Rudel, C Beiträge zur Kenntniss der Alkaloide von *Berberis aquifolium* und *Berberis vulgaris*. Arch. der Pharm. 229:631-649, 649-660. Abh. in Bot. Centrbl. 2:294.
863. Rusby H. H The Poisonous Plants of the Vicinity of New York City. Repe. Lecture before the Alumni of the College of Pharmacy of the City of New York. 1895. 18. Also Med. and Sur. Jour. 72:839-844.
864. Poisonous Properties of the Mountain Laurel. Drug. Cir. 46:27
865. Ruschhaupt, Fr. and L. I Poisoning by a Hemlock Species. Rept. Wis. St. Bd. Health. 18:99
866. Russell, W Sur les migrations des glucosides chez les végétaux. Compte Rend. Acad. Sci. Paris. 139:1230-1232.
- 867 Sur la sécrétion de quelques principes actifs des végétaux pendant le repos hivernal. Revue Gen. de Bot. 15:160.
868. Ryan, Angie M Poison Ivy and its Effects. Amer. Bot. 4:27-28.
869. Sabanaieff, N. and Berlmann, M Die Giftpflanzen des Gouvernements Jaroslavl, mit Angabe ihres Schadens für die Haus-thiere. Jaroslavl 8-16. 1886
870. Sabouraud Les Trichophytes Humaines. 1894
- 870a. Safford, W. E. The Useful Plants of the Island of Guam. Con. U. S. Nat. Herb. 9:371
871. Sagot, M. P Recherches des plantes très vénéneuses par l'essai sur les tetrads des Batraciens. Sep. Bull. Soc. bot. de France. 25:1
872. Saint Hilaire Influence de la température organique sur l'action de quelques substances toxiques. In Richet, C. Physiol. Trav. du lab. 1:390-430. Paris. 1893.
873. Saunders, C. F. Poisonous Effects of a California Shrub. (Nympha Parryi) Plant World. 6:245-246.
- 873a. Savoure, Camille Etude sur les alcaloïdes d'origine microbienne. 97. Rennes. 1896.

874. Recherches experimentales sur les mycoses internes et leurs parasites. Archiv Parasit **10** 5.
875. Sayre, L. E. A further Study in the Alkaloids of Geisium. Proc. Am. Phar. Assoc. **26**.
876. Loco Weed. Am. Vet. Rev. **11** 555. Also see Proc. Am. Phar. Assoc. **33** 107-108.
877. A Manual of Organic Materia Medica and Pharmacognosy. Ed. 2. 684. Philadelphia.
878. Schaffner, J. H. Poisonous and other Injurious Plants of Ohio. Ohio Naturalist **4** 16-19, 32-35, 69-73.
879. Schenck, H. See Strasburger, E.
- 879a. Scherer. On Hay Fever. See Jour. Mich. St. Med. Soc. 1904. No. 5.
880. Schimper, A. F. W. See Strasburger, E.
881. Schimpfky, Richard. Deutschlands wichtigste Giftgewächse in Wort und Bild, nebst einer Abhandlung über Pflanzengift. Für den Schulgebrauch und zum Selbstunterrichte. Gera-Unterhaus. Verlag von Fr. Eugen Köhler. 1893.
882. Schnaggenhaußen, Fr. . . . See Heckel, E.
883. Schnaggenhaußen, Fr., and Kieb. Note sur la racine et semences de cyanglossé. Jour. der Phar. f. Elsass-Lothringen **18** 285. See also **1892** 61.
884. Schützberger, S. Die Gift und Heilpflanzen. Eine Anleitung, die giftige und heilkräftigen Wirkungen kennen zu lernen und Anweisung, wie man sie gegen allerlei Krankheiten benutzen kann. Leipzig. 1899.
885. Schlotterbeck, J. O. Does Argemone mexicana contain Morphine? Phar. Rev. **19** 458-461.
886. Schlotterbeck, J. O., and Watkins, H. C. Contribution to the Chemistry of Stylophorum diphyllum. Phar. Rev. **19** 453-455.
887. Schlotterbeck, J. O., and Zwa.uwenberg, A. von. Comparative Structure of the leaves of Datura stramonium, Atropa Belladonna, and Hyoscyamus niger. Pharm. Arch. f. **1898**. No. 1. Abs. in Just's Jahresb. **1898**. Abt. 2 50.
888. Schmeiz, J. D. F. Zur Kenntnis indischer Gifte. Internat. Arch. f. Ethnog. **15** 54. Leyden. 1902.
889. Schmidt, Ernst. Ueber die Alkaloide der Lupinosenen. Archiv der Pharmacie **235** Heft. 3.
890. Schwedberg, O. Grund. Art. 167. 1887.
891. On Digitalis. See Archiv. Exp. Path. **18** 149.
892. On Oleandrin. See Ber. Chem. Gesell. **18** 233.
893. Schneidemühl, Georg. Specielle Pathologie u. Therapie der Haus-thiere für Thierärzte, Aerzte und Studierende. Berlin. 1908. 799 f. 53.
894. Scutellier, A. Native and Introduced Poisonous Plants of California. Pacific Phar. **1907**.
895. Literature on Medicinal Plants and Drug Plant Culture. Phar. Rev. **26** 20 24, 236-245.
896. Schneider, Max. Ueber Saponine. Zeitsch. des allgem. österreichischen Apotheker-Vereins. **43** 893-898. 917-21.
897. Schniegans, M. Der Giftstoff von Arum maculatum. J. d. Oh. f. Elsass-Lothr. **62**.

898. Schomburgk, R. H. . . . The Urari or Arrow Poison of the Indians of Guiana. *Phar Jour* 18:500-507 (1857)
889. Schorlemmer, C. . . . A Manual of the Chemistry of the Carbon Compounds or Organic Chemistry. London. 1874.
- 889a. Schramek See Kren.
900. Schribaux, E. . . . The Poisoning of Cattle by Eating the Green Plants of *Lathyrus Cymenium*. *Jour Agr Prat* 59: No. 28. 32-55
901. Schuetz, K. E. . . . De Taxo baccata ejusque veneno. 4). Vrat-slavia. 1840.
902. Schulze, J. E. F. . . . Toxicologia veterum plantas venenosas exhibens Theophrasti, Galeni Dioscoridi, P. m., aliorum que auctoritate ad cetera venena relatas. Loca ex veterum monumentis eruta perpetuo commentario ornari. varia experimenta et observata adjecta 78. Halle 1788
903. Schummel, T. E. . . . Ueber die giftige Pilze mit besonderer Rücksicht auf Schlesien. 28. Breslau 1840
904. Schunemann, H. . . . Die Pflanzen-Vergiftungen. Ihre Erscheinungen und das Vorzunehmende Heilverfahren geschildert an den Deutschen heimischen Giftpflanzen. 88. Braunschweig 1891.
905. Schutte, H. W. . . . Onderzoekingen over Dioscorine, het giftige Alkaloid uit de Knollen van *Dioscorea bifurcata* Bl. *Nederlandsch Tijdschrift voor Pharmacie, Chemie en Toxicologie* 9 Ma 1897
906. Schutz, Josef . . . Ueber das Eindringen von Pilzporen in die dadurch bedingten Erkrankungen der Augen etc. *Mith. a. d. Kais. Gesundheitswarte* 1894 208.
907. Schwalbe, K. . . . On *Rhus Toxicodendron*. See *Munch med. Woch.* 49:1, 616; *Med. Rec.* 63:855.
908. Schwarz, A. . . . Giftpflanzen, Heilpflanzen, und Nährpflanzen. Bilder aus der heimischen Pflanzenwelt mit erläuternden Text. 277. Forth. 1899.
909. Scott, D. H., Translator . . . See De Bary, A.
910. Scott, J. L. . . . Golden-rod Killing Horses. *Gar and For* 8: 477-478.
911. Segura, Jose C. . . . El Maguey. Memoria sobre el cultivo y beneficio de sus productos. 407. Mexico. 1901
913. Selby, A. D. . . . A First Ohio Weed Manual. *Bull. Ohio Agr. Exp. Sta.* 81 249-500.
914. . . . A Second Ohio Weed Manual. *Bull. Ohio Agr. Exp. Sta.* 175 291-383.
915. . . . Poisonous Plants. *Jour. Columbus Hort. Soc.* 1886 119-128
916. Seligman, C. G. . . . On the Kenyah Dart Poison and its Active Principle. *Antiarin*. See *Brit Med Jour* 1903 1131
917. . . . On the Physiological Action of the Kenyah Dart Poison "Ipon" and its Active Principle, *Antiarin*. *Jour. Phys.* 1903 39-58. London.
918. Selmi, F. . . . Studi di Tossicologia Chimica. Bologna 1871
919. Sempie, R. H. . . . Lecture on Vegetable Poisons. Delivered Nov. 30. 1842. *Phar Jour* 2:429-442
920. . . . Lecture on Vegetable Poisons. Delivered at the Establishment of the Pharmaceutical Society, June 22, 1842. *Phar Jour* 2: 73.
- 920a. Seymour On Gelsemium. See *Boston Med. and Sur Jour* 105:590.

- 921 Sharp, G. Coca and Cocaine Studied Historically. *Phar Jour.* 32:28-30; 117-118, 184-186.
- 922 Shenstone, J. C. Poisonous Plants of Essex. *Jour Essex Tech.*
- 923 Shionoyana, J. Chemical Researches on the Aconitum and Other Plants. *Bot. Mag.* 7:441
- 924 Shreve, F. and Others. The Plant Life of Maryland. Maryland Weather Service 2:333
- 925 Segel, A. Ueber die Giftstoffe zweier Euphorbiaceae. *Inaug. Dissert. Dorpat.* 55. 1895.
- 925a Simon, J. F. See Sobernheim.
- 926 Simpson, William. The Names of Medical Plants of Commercial Value that are Gathered in North Carolina, their Value and the Relative Amount Sold in this Country and Exported. *Bull Phar* 8:448-450.
- 926a Sinkler. On Gelsermin. See *Phil Med Times* 8: 150. 1877-1878.
- 927 Slade, H. B. Some Conditions of Stock Poisoning in Idaho. *Bull. Id. Agr. Exp. Sta.* 37 159-190.
928. Twentieth Century Practice.
- 929 Smith, R. F. Death from Eating *Cicuta maculata*. *Bo. Gaz.* 13:128-129
- 930 Smith, E. P. and Hedges, Florence. Diplodia Disease of Maize (A Suspected Cause of Pellagra). *Science* 30:60-61
- 931 Smith, Frederick. Manual of Veterinary Hygiene. Ed. 5. 1035 New York.
- 932 Smith, J. B. Antidotes to Alkaloidal Poisons. *Med Times and Hosp. Gaz. London.* 5:22-24
- 933 Smith, S. Bernhard. Poisonous Plants of all Countries. 88. Bristol. 1905.
- 934 Smith, W. G. Poisoning by *Agaricus dealbatus*. *Gar Chron.* No. 644 556.
- 935 Smith, W. G. as Translator See Tubeuf
- 936 Smithsonian Institution. See various Publications, especially No. 258.
- 937 Sobernheim, J. F. and Simon, J. F. Handbuch der praktischen Toxicologie. Berlin. 1838.
- 938 Sohn, Charles E. Dictionary of the Active Principles of Plants. Alkaloids; Bitter Principles, Glucosides, their Sources, Nature, and Chemical Characteristics, with Tabular Summary, Classification of Reactions, and full Botanical and General Indexes. 194. London. 1894.
- 939 Sour and Alexandresen. Sur la toxicité d'un type d'*Aspergillus fumigatus* isolé du maïs avarié. *Compte. Rend Soc. Biol.* 64:288-289
- 940 Sommerfeld, T. Entwurf einer Liste der gewerblichen Gifte, ert. Aufträge der internationalen Vereinigung für gesetzlichen Arbeiterschutz, unter Mitwirkung, von Thomas Oliver und Felix Putzeys. Jena. 1908
- 941 Spieler, A. J. T. De plantis venenatis Silesiae. 42. Vratislavia. 184
- 942 Spratt, G. A Compendium of Toxicology of the Principal Indigenous and Exotic Plants. London.
943. . . . Flora Medica. A Description of the Medicinal Plants Admitted into London, Edinburgh, and Dublin Pharmacopoeia. 2 vols. London. 1829.

944. Sprouch, C. H. E. Le poison diphtérique considéré principalement au point de vue de son action sur le rein. *Compte Rend. d l'Acad. des Sci de Paris* **109**:260-262
944. Stalberg, S. A Case of Poisoning by an Unknown Quantity of Atropine *Recovery The Wm. Jones Med. Soc. **127** 517
945. Stalker, M. Crotalism: A New Disease among Horses, Rept Ia. Agr Coll. Dept of Bot **1884**: 114-115
946. The Loco Plant and its Effect upon Animals. Rept Bur An. Ind. U. S. Dept Agr **3** 272
947. Stapf, O. The Aconites of India, a Monograph Ann. Roy. Bot. Gar Calcutta **10** 115-179 1905
948. Stevens, A. B. Japanese Lac Doctoral Thesis 1906
949. On Rhus Poisoning See Proc Mich State Phar Assoc **1908**:61-64
950. See Tachurch, A
951. Stevens, A. B. and Warren, L. E. Poison Sumac (*Rhus venenata*) Proc. Am. Phar. Assoc. **55**, 423-443
952. Also See Am Phar Jour **79**:518.
953. Stewart, F. E. The Therapeutics of Tea and Coffee Ther. Gaz. (N. S.) **2**, 202-203
954. Stuckberger, W. W. The Drug known as Pink-root (*Spigelia marilandica*). Bull. Bur. Pl. Ind. U. S. Dept. Agr **100** 41-44
955. Stokes, J. S. On Poisoning by the Internal Administration of Tincture of Rhus Toxicodendron See Med. Sur. Rept **17** 373
956. Stowe, W. A Toxicological Chart, Exhibiting at one View the Symptoms, Treatment, and Modes of Detecting the various Poisons, Mineral, Vegetable, and Animal, to which are Added Concise Directions for the Treatment of Suspended Animation. Ed. R. L. Wa. Chart. London.
954. Strasburger, E., Voll, F., Schenck, H., and Karsten, G. A Text-book of Botany. Third English Edition. Revised by W. H. Lang. 746. London 1908.
958. Stratward, R. A Case of Poisoning with Spirits of Turpentine Abs in Am. Vet. Rev **34**:58.
959. Strickland, C. W. Cattle Poisoning by the Common Yew Gar. Chron. **18** 160
960. Strong, A. B. The American Flora 2 vols New York 1848.
961. Sturdevant, L. B. and Peters, A. T. The Poisoning of Horses by the Common Horse-tail Weed. (*Louissetum arvense*) Rept. Neb. Agr. Exp. Sta. **19** 111-115.
962. Sutton, R. L. The Emergency Treatment in Cases of Acute Poisoning. Kansas City Med. Index
963. Swain, R. E. and Harkins, W. J. Arsenic in Vegetation Exposed to Smelt or Smoke. Jour. Am. Chem. Soc. **30**:915.
964. Also see Harkins.
965. Syme, R. L. Some Constituents of the Poison Ivy Plant (*Rhus Toxicodendron*). 37 Thesis Johns Hopkins University 1906.
966. .. Also see Acree, S. F.

- 967 Taylor, A. S. On Poisons in Relation to Medical Jurisprudence and Medicine. Third Am. and Eng. Ed. Philadelphia. 1875
968. Principles and Practice of Medical Jurisprudence. 3 vols. London. 1873
- 969 On Rhus Toxicodendron. See Jour Cut and Ven. Dis. 1.434.
970. Taylor, H. H. See Powell, R. D.
- 971 Taylor, Thomas Edible and Poisonous Mushrooms
- 972 Ten Bosch, J Does Poison Ivy Discriminate? Gar. and For. 9.239
973. Tesnier, T. J Essai sur les poisons vegetaux la plus commun en France et leurs antidotes. 19. Montpellier. 1809.
974. Thacher, J On Loss of Hair and Nails from Rhus Poisoning. See Am. New Disp. 334. 1817
975. Thur, A. Experimental Researches concerning Trichophyton tonsurans. Brit. Med Jour 1469 397-399
976. Thompson, R. H. A Contribution to the Causes of the So-called Hay Fever, Nasal Asthma, and Allied Affections Considered from a Chemical Standpoint. Med. Rec. 1887 305-306
978. Thompson, C. J. S. Poison Romance and Poison Mysteries. London. 1899
979. Thompson, Hedge An Experimental Dissertation on the Spigelia Marilandica or India Pink Philadelphia 1802
980. Thomson, Thomas Chemistry of Organic Bodies. 183. London
981. Thunberg, C. P. Arbor toxicaria Macassaricensis. Upsala 1788
- 982 De planus venticatus. 15. Upsala 1788.
- 983 Thurber Geo Poisoning by Cypripedium. Bull. Torr Bot Club. 9.15
- 984 Tidy, C. H. On Poisoning by Nitrate of Baryta. Med Pres and Cir. 9.448.
985. Tillie, T. B. See Frazer T. R.
986. Tirelli, V. Auto intossicazione secondarie all'azione dei caustici e del calore ricerche tossicologiche Ann. di freniat. Turin 9.253-273.
- 987 Tizzoni, Guido Neue bakteriologische Untersuchungen über die Pellagra. Centrbl. f. Bakt u Parasit 46 (Abt. 1) 310-311
- 987a Tokishige, H Ueber pathogene Blastomyceten. Centrbl. f. Bakt. Abt. 1 19.105-113. 1896.
988. Torrey, John Natural History of the State of New York Pt. 2. Botany. 2 vols. 1843.
989. Tovey, J. R. See Ewart, A. J.
990. Tracy, S. M. Cassava. Par Bull. U. S. Dept. Agr. 167 31
- 991 Trelease, Wm. The Working of the Madison Lakes. Wisc. Acad. Sci. Arts and Letters 7.121-129
- 991a An Unusual Phyto-bezoar. Trans. St. Louis Acad. Sci. 7.493-497. 1 pt.
992. Treub, M. Nouvelles recherches sur le rôle de l'acide cyanhydrique dans les plantes vertes. Ann. Jardin bot. Buitzenborg. Ser. I. 6.77-106.
993. Trimble See Bastin, E. S.
- 993a Trimble, W. K. A Case of Sporotrichal Infection. Jour. Kan. Med. Soc. 9.305.
994. Trimen, H. See Bentley, R.
995. True, R. H. See Israel, O.
996. Tryon, H. Plants Poisonous to Stock. Brisbane Courier May 24, 1887

- 997 Tschirch, A. and Oesterle. Anatomischer Atlas der Pharmakognosie und Nahrungsmittelkunde. 334. Leipzig. 1900.
- 998 Tschirch, A. and Stevens, A. B. On Japanese Lac, the milk sap of *Rhus vernicifera*. Arch. Phar. **243** 504. Also Am. Phar Jour **78** 53.
- 999 Tuseuf, Carl, and Smith, W. G. Diseases of Plants. 598. New York.
- 1000 Tuzcek, F. Behandlung der Vergiftungen mit Pflanzenstoffen (excl. der chron. Morphinismus und Cocainismus des Ergotismus, der Pellagra, und der Lathyrismus). Handb. d. Spec. Therap. innerer Krankh. **2** 234-334. Jena. 1894.
- 1001 Uriney, C. See Dunstan, W. R.
- 1002 Vanlen, E. Ergot. See Arch. f. exp. Path. u. Pharmak. **55**:131, also Chem. Centrbl. **1906**:2, 690. (Abs. in Phar. Rev. **25** 323-324).
- 1003 Varentin, G. C. Ein Fall von Scur des Mittelohres. Arch. f. Ohrenheilkunde. **28**.
- 1004 Van Es, L. and Waldron, L. R. Some Stock Poisoning Plants of North Dakota. Bull. N. D. Agr. Exp. Sta. **58** 321-352.
- 1005 Vance, George. Plants Poisonous to Cattle in California. Rept. Com. Agr. **1874** 159.
- 1006 Vater, A. De *Laurocerasus indole venenata*. 32. Wittenberg. 1737.
- 1007 Vaughan, V. C. Some New Bacterial Poisons, their Causal Relation to Disease and the Changes in our Theories Suggested by their Action. Med. News. **2** 150-162.
- 1007a See Hamilton, A. M.
- 1008 Vaughan, V. C. and Novy, I. C. Pitomanes and Leucomanes.
- 1009 Vaughan, V. C., Novy, I. C., and McClintock, C. T. The Germicidal Properties of Nucleins. Med. News. **1893**:536-538.
- 1010 de Vaublanc, A. Les plantes qui tuent. Jour. d. Hyg. **28** 153-169. Paris. 1900.
- 1011 Vergerau, L. Le toxine du *Bacillus virgula*. Compt. Rend. Soc. Biol. Paris. **1903**:803-835.
- 1012 Verdier. See Breton.
- 1013 Verschaffelt, F. Bepaling der werking van vergiften op planten. Versl. d. -wis-en-natuurk. Afd. d. k. Akad. v. Wetensch. Amst. **12** 1893-855-859. 1903-1904.
- 1014 Sur la presence de l'acide cyanhydrique dans les bourgeons de *Prunus* en voie d'eclosion. Arch. Neerland. Sc. exact. nat. Ser. **27** 497-509. **1902**.
- 1015 Vicat, P. R. Histoire des plantes veneneuses de la Suisse contenant leur description, leurs mauvais effets sur les hommes et sur les animaux avec leurs antidotes. redigee, d'apres ce qu'on a de mieux sur cette matiere, et surtout d'apres l'Histoire des plantes helvetiques de M. le baron de Haller, mise à la portee de tout le monde avec le lieu natal de chaque plante pour la France, les figures necessaires, et plusieurs observations nouvelles. 29+392; 112+8. Versailles. 1776.
- 1016 Vlier, A. Recherches des poisons vegetaux et animaux. 132. Paris. 1882.

- 1017 Volkens, G Faser-Flecht and Banderstoffe. Notizbl. der Königl. bot. Gar. u. Mus. 24 Berlin-Dahlem 5:42-64. Ap 22
- 1018 Wattwright, J W Secret Poisons and their Use in Ancient Times. Med. Sci 44:286-290. New York
- 1019 Waldron, C A Poisoning from Locust Bark. Am Vet Rev 3:436-439
- 1020 Waldron, L R See Van Es, L.
- 1021 Waldron, L R Poisoning from Oleander. Deut. med. Wochenschr. 17:801. Abs. Am. Jour. Med. Sci 120:725.
- 1021a Wallace, Farmer Clipping on Killing Horses by Oak leaf gird (Based on Rept. For. Ser. U. S. Dept. Agr. 1909. Sept. 3).
- 1021b Walsh, L. H South African Poisonous Plants. Being Notes on South African Plants Poisonous to Stock, with Particulars of Symptoms and Treatment. 56. Cape Town, 1909
- 1022 Warburton, C W The Non-saccharine Sorghums. For. Bul. U. S. Dept. Agr. 26:21
- 1023 Warner, H See Pedler, A.
- 1024 Warder, J. A. The Western Catalpa—a Memoir of the Shav-
anon. Jour. Am. Agr. Assoc. 1881.
- 1025 Warren, L. E. Contributions to the Bibliography of the Poi-
sonous Species of Rhus. Mid. Drug and
Phar Rev 44:149-150, 218-219, 288-290.
- 1026 The Poisonous Species of Rhus. Phar Jour
Oct and Nov 1909. Repr
- 1027 Rhus Michauxii. A Nonpoisonous Plant
Repr Am. Jour. Phar 1910:499-500.
- 1028 See Stevens, A. H.
- 1029 Watkins, W. H Case of Poisoning by the Essential Oils of
Savine, Pennyroyal, and Tansy. New Or-
leans Med. and Sur Jour 5:230.
- 1030 Watson, Serrão Bibliographic Index to North American Bota-
ny. Polypetalae. Washington 1878
- 1030a Botany of California. Geol. Sur. of Cal 2:55
- 1030b Contributions to American Botany. Proc.
Am. Soc. Arts and Sci. 21
- 1031 Wey On Cocain. See Med. News Phil. 1:496.
1887
- 1032 Weatherby, C. A See Graves, C. H.
- 1032a Webb, Paul C Plants Injurious to Animals. Annual Report
Assoc. of Phil. Coll. of Phar 47:27
- 1033 Weddell, E. C The African Poison Test as Observed in the
Portuguese Colony of Angola. 1906.
- 1034 Weschke, Carl The Enumeration of the Medicinal Plants of
Minnesota. Phar Rund. 8:155-157
- 1035 Westbrook, F. F Contributions à l'étude des toxines du choléra.
Ann. de l'Institut Pasteur 1894:318-337
- 1036 Westgate, J. M Alfalfa. For. Bul. U. S. Dept. Agr. 339-48
(Alfalfa Blat p. 37)
- 1037 Wettstein, R. von Handbuch der systematischen Botanik 1:201
Leipzig and Vienna. 1901
- 1038 Wemyss, H. M Rhus Toxicodendron. Bull. of Phar 8:552-
553
- 1039 Whipple, G. C Microscopy of Drinking Water. 119. New
York. Ed. 2.
- 1040 White, E. A A Preliminary Report on the Hymenales of
Connecticut. Bull. Geol. and Nat. Hist. Sur.
of Conn. 3:80.

- 1041 Second Report of the Hymeniales of Connecticut. Bull. Geol. and Nat. Hist. Sur. of Conn. 15:70
- 1042 White, G. R. Phytolacca Poisoning in Cattle. Am. Vet. Rev. 26:339-340.
- 1043 White, J. C. Dermatitis venenata: an Account of the Action of External Irritants on the Skin. 216. Boston. 1887
- 1044 Death from Rhus Poisoning. N. Y. Med. Jour. 17:223
- 1045 Wignall, F. and Nobecourt. Recherches sur l'action antitoxiques des centres nerveux pour la strychnine et la morphine. Bull. et mem. Soc. med. d. hôp. de Paris. Ser. 3. 15:182-184
- 1046 Wignall, F. and Nobecourt. Lehrbuch der Pharmacognosie. Berlin. 1879
- 1047 Wilcox, R. W. Larkspur Poisoning of Sheep. Bul. Mon. Agr. Exp. Sta. 15:37-51. Abs. in Ann. rept. Bur. Ind. U. S. Dept. Agr. 15:473-479
- 1048 Plant Poisoning of Stock. Ann. Rept. Bur. An. Ind. U. S. Dept. Agr. 17:91-121
- 1049 Poisonous Plants of Montana. Rept. Mon. Agr. Exp. Sta. 16:99-51
- 1050 Poisonous Sumach once more. Gar. and For. 8:478.
- 1051 Wilson, E. V. See Chesnut, V. K.
- 1052 Wilcox, R. W. On Nutmeg Poisoning. See Lancet 1895: 50. Rev. in Am. Jour. Med. Sci. 110:90.
- 1053 Wilcox, R. W. and Jeffle, S. V. On Datura Poisoning. See Brit. Med. Jour. 1903 1137. (Article by Dr. J. D. Gamble), Rev. in Am. Jour. Med. Sci. 126: 348.
- 1054 On the Kenyah Dart Poison and its Active Principle, Antiarin. (Article by Dr. C. G. Seligmann in Brit. Med. Jour. 1903 1131. Rev. in Am. Jour. Med. Sci. 126:347
- 1055 Wilcox, R. W. and Pellon, H. H. Review of Article by John Germain on Veronal Poison in Jour. Am. Med. Assoc. 1906 (No. 26) 1999. Am. Jour. Med. Sci. 133: 795
- 1056 Wilcox, T. P. On Recurring Attacks of Rhus Poisoning without Re-exposure. See Gar. and For. 8:478.
- 1057 Winceman, F. de. Poisons d'épreuve de l'Afrique occidentale. Assoc. franc. Avanc. Sci. 1903 736-743
- 1058 Wiley, H. W. Sweet Cassava. Jatropha Manihot. Bot. Gaz. 14:71-76.
- 1059 Williams, A. J. Poisonous Plants in South Africa. Vet. Rec. London. 16:421
- 1060 Williams, S. W. Report on the Indigenous Medical Botany of Massachusetts. Trans. Am. Med. Assoc. 1848.
- 1061 Williams, T. A. A Report on the Grasses and Forage Plants and Forage Conditions of the Eastern Rocky Mountain Region. Bul. Div. Agron. U. S. Dept. Agr. 12:78
- 1062 Williams, W. L. Enzootic Cerebrospinal Meningitis of Horses in Idaho. Rept. U. S. Dept. Agr. Bur. An. Ind. 1897 179-187
- 1062 Williams, W. T. The Etiology of Heaves. Am. Vet. Rev. 26: 955-957

1063. Williams Cashew (Mesquite) Poisoning. Jour Jamaica Agr Soc. 1:319-321 1897
1064. Willis, G. S. V. A Manual of Vegetable Materia Medica 480. London, 1889. Ed. 11
1065. Witloughby, W. G. . . . On Poisoning from Foods. See Public Health 1908:626
1066. Warner, B. . . . Observations on the Poisonous Vegetables which are either Indigenous in Great Britain or Cultivated for Ornament. 103 London. 1781
1067. Wilson, F. W. Oleander Poisoning of Live Stock. Bol. Ariz. Agr. Exp. Sta. 59:381-397 67
1068. Wilson, James New Remedy for Sheep Poisoning on the Stock Ranges. Yearbook U. S. Depr. Agr 1902:22.
1069. Wilson, James and Coville, F. V. . . . See Coville, F. V.
1070. Wilson, J. C. An American Textbook of Applied Therapeutics for the Use of Practitioners and Students 1326. Philadelphia, 1898.
1071. Wislizen, P. Rhododendron Poisoning in a Goat. The Veterinarian 70:639-640.
1072. Wingate, J. B. Actinomycosis. Jour Comp. Med. 17:276-281
1073. Winkler, L. Sammtliche Giftgewächse Deutschlands naturgetreu dargestellt und allgemein sprachlich beschreiben. Mit einer Vorrede von Friedrich Schwalgricher 119 96 col p. Berlin 1831
1074. Wislizen, K. Veterinary Materia Medica and Therapeutics 755 New York 1902 Ed. 2
1075. Winton, A. L. and Moeller, J. Microscopy of Vegetable Foods. 701 1906.
1076. Wuthaus, R. A. Medical Chemistry 1881
1077. Toxicology (Vol 4 of next book named) Vegetable poisons. 600-834
1078. Wuthaus, R. A. and Becker, T. C. Medical Jurisprudence, Forensic Medicine and Toxicology. 4:492, 35f New York 1896.
1079. Wittstein, G. C. Handwörterbuch der Pharmakognosie des Pflanzenreichs. 464 Breslau 1882 1st Hälfte)
1080. The Organic Constituents of Plants and Vegetable Substances and their Chemical Analysis. Melbourne. 1878.
1082. Wolfenstein, R. U. and Pietet, A. Pflanzenalkaloide 134. Ed. 3. 1900
- 1082a. Wood, Alphonso Class Book of Botany 829 New York 1867
1083. Wood, H. C. Text book of Therapeutics Ed. 1.
1084. Woodville, William Medical Botany Ed. 2 4 vols London 1810
1085. Wright, J. S. Botanical Products of the United States Pharmacopoeia. 1890. Proc. Ind. Acad. Sci. 1894 108-119
1086. Wright, R. and Farr, E. H. The Pharmacy of Conium maculatum. Chem and Drug. 49.
1087. Wunchmann, F. Deutschlands Giftpflanzen. Berlin. 1833
1088. Zehl, L. Die Beeinflussung der Giftwirkung durch die Temperatur sowie durch das Zusammenwirken von zwei Giften. Zeitsch. f. allgem. Physiol. 8:140-190. Abs in Bot. Centrbl. 108:328-329.

- 1089 Zoethout, W. D. On some Analogies between the Physiological Effects of High Temperature, Lack of Oxygen, and Certain Poisons. *Am Jour Physiol.* **2**:220-242. 1899. Abs. in *Science* **8**, 776-778. (n. s.)
- 1090 Zoon, J. *Icones plantarum medicarum. Centuria I. Abbildungen von Arzneypflanzen.* Nuremberg. 1784.
- 1091 Zopf, Wilhelm. *Die Pilze in morphologischer, physiologischer, biologischer, und systematischer Beziehung.* 500. Breslau. 1890.
- 1092 Die Polsthere. 1890.
- 1093 Zwaluwenberg, A. van. See Schlotterbeck, J. O.
- 1094 Gienk, Robert. *Cicuta maculata*, Linne. *Am. Jour. Phar.* **63**:328-332.
- 1095 Marsh, J. M. Notes on some North American plants. *Am. Jour. Phar.* **63**:321-328.
- 1096 Mearns, E. A. Two new species of poisonous sumachs from the states of Rhode Island and Florida. *Proc. Biol. Soc. Washington.* **15**:147-149.
- 1097 Nelson, A., and Others. Woody Aster. *Bull. Wyo. Agr. Exp. Sta.* **28**:20.

- Baptitoxin, 530
 Belladonna, 386, 715
 Benzoylecognin, 575
 Berberin, 472
 Berberin, 88, 90, 108, 112, 146, 468, 470, 472, 473, 478, 483
 Bikkacotin, 450
 Brucin, 145, 146, 147, 586
 Bryonin, 751
 Caffein, 628
 Caffein, 144, 146, 606, 614, 628, 742
 Calabar, 148, 528, 531
 Calycanthin, 113, 476
 Carpain, 627
 Cathin, 614
 Celastrolin, 614
 Cephaelin, 742
 Chelerythrin, 113, 480, 485
 Chelidonium, 480, 485
 Chonosidin, 479
 Chrysantheremin, 789
 Clestin, 656
 Cinchonidin, 145
 Cinchonin, 143, 742
 Cinnamylcoralin, 575
 Cocain, 6, 73, 77, 79, 44, 147, 575, 576, 614
 Codamin, 147, 481
 Codein, 39, 77, 112, 113, 146, 47, 479, 481
 Colchic, 77, 78, 79, 146, 148, 375, 479
 Columbin, 473
 Concein, 631
 Coniferin, 582
 Conin, 1, 73, 77, 79, 126, 23, 144, 147, 648, 651, 652, 653, 688, 726, 728, 747
 Conydrin, 631
 Cornutin, 28, 77, 78, 277, 278, 279
 Corybutin, 147, 480
 Corydalin, 147, 480
 Cryptopin, 481
 Curarin, 73, 77, 487
 Curin, 148, 687
 Cusanyrin, 575
 Cusparin, 582
 Cynopin, 126, 639
 Cynoglossin, 706
 Cythrin, 77, 78, 79, 90, 118, 148, 180, 537, 542, 551
 Damascen, 446
 Delphinin, 108, 446, 464
 Delphinoidin, 108, 464
 Delphinin, 108, 464
 Didochanin, 742
 Ditamin, 692
 Dramin, 588
 Dubotain, 716
 Echolin, 149, 277
 Echitamin, 692
 Echitemin, 692
 Ergotin, 277, 278
 Ergotin, 277
 Ergotoxin, 278
 Erythrophobocin, 48, 534
 Escrin, 82, 531, 729
 Esmerdin, 528
 Eupatorin, 138, 755, 772
 Fumarin, 483
 Gelsamin, 77, 80, 129, 147, 689
 Gelsaminin, 147, 689
 Gnoscocin, 481
 Gracilin, 734
 Guvacin, 378
 Heuotropin, 705
 Hydrastin, 108, 146, 468
 Hydrastin, 146
 Hydrococcotin, 481
 Hydroquinon, 585, 742
 Hyxin, 147, 575
 Hyoscin, 148, 716, 726, 727, 730
 Hyoscin-pseudocynin, 148
 Hyoscinin, 3, 60, 61, 77, 133, 148, 716, 726, 727, 730, 732, 736
 Imperin, 375
 Inasin, 692
 Indacotin, 450
 Isopilocarpin, 148
 Isopyrin, 446
 Jaborin, 148, 587
 Japocotin, 149, 450
 Jervin, 103, 148, 381
 Jurechin, 715
 Kassin, 386
 Lanthopin, 481
 Lappachonin, 450
 Lanthanin, 147, 480
 Lanthanin, 481
 Lobelin, 78, 752
 Lophopharin, 657
 Lycopin, 530
 Lycocin, 386
 Maasin, 715
 Mandragotin, 715
 Maltin, 530
 Melonidin, 481
 Menapernin, 112, 148, 473
 Mesalin, 637
 Methylechin, 126, 651
 Morphin, 3, 39, 73, 78, 80, 82, 85, 90, 117, 118, 143, 144, 146, 147, 479, 480
 Moschalin, 140, 787
 Muscarin, 10, 32, 73, 74, 77, 78, 80, 82, 148, 150, 157, 238, 242
 Muscarin, 751
 Narcein, 146, 481
 Narcein, 146, 147, 481
 Narcein, 237
 Narcein, 7, 73, 77, 78, 80, 85, 87, 133, 144, 147, 551, 716, 728, 729
 Nirelin, 446
 Oxycotin, 112, 472
 Pafarin, 450
 Papaveramin, 112, 147
 Papavarin, 59, 112, 146, 147, 481
 Pectenin, 637
 Pectatin, 637
 Pelosin, 413, 478
 Pelosin, 147
 Phytolaccin, 107, 435
 Physostigmin, 146, 148, 587
 Picolin, 728
 Pilocarpidin, 582
 Pilocarpin, 146, 148, 528, 531
 Pilocercin, 637
 Piperin, 396
 Piturin, 5
 Piturin, 147, 716
 Protocurin, 148, 687
 Protopin, 113, 480, 481, 485
 Protopernitridin, 381
 Pseudohyoscyamin, 148, 715, 726
 Pseudojervin, 148, 381
 Pseudomorphin, 481
 Pyrethrin, 754
 Pyridin, 145, 146, 728
 Quebrachin, 692
 Quebrachinamin, 692
 Quebrachin, 147
 Quercitrin, 506, 510, 584, 666
 Quinamin, 742
 Quinin, 77, 78, 87, 145, 147, 741, 742
 Rhoeodin, 481
 Rubijervin, 381
 Sambucin, 747
 Sanguinarin, 113, 480, 484, 485
 Sarracenin, 14, 497
 Scopolamin, 60, 77, 78, 148, 716, 737
 Scrocin, 278
 Septentrionin, 450
 Solanin, 718
 Solanin, 132, 148, 718
 Solanin, 60, 83, 132, 33, 144, 148, 149, 718, 72, 722, 724
 Spathin, 530, 543
 Spathin, 147, 551
 Spigelin, 688
 Staphagrin, 108, 464
 Styracin, 1, 72, 73, 82, 79, 144, 145, 147, 628, 646, 688, 728, 729, 736
 Styacin, 480
 Syntactonin, 480
 Tassin, 101, 148, 328
 Thalictrin, 446
 Thobin, 146, 47, 481
 Thobin, 146, 629, 742
 Theobromin, 144, 146, 627, 628
 Theophyllin, 146, 628
 Trigonidin, 531, 648
 Tropaeocin, 575
 Truxillin, 575
 Tubocurarin, 687
 Tubocurarin, 148
 Uramin, 530
 Uramin, 120
 Xanthopuccin, 108, 468
 Xanthopuccin group, 148
 Volatile, 344
Albana hirtella, 704
 Alkanet, 704
Affamonda cathartica, 806
Alibegum Theropala, 314, 519
 Allen, Dr., 440
 Allen, H. M., 608
 Allen, Timothy P., 809
 Alligator Pear, 478
Allium, 104, 377, 383
ascotum, 375, 383, 384, 833
canadense, 104, 383, 834
Cepa, 383, 833
Douglasii, 833
Macleanii, 834
Moly, 383, 834
Perrini, 373, 383, 834
saricatum, 375, 383, 834
Schoenoprasum, 375, 383, 834
tricolorum, 104, 383, 834
unifolium, 104
versutum, 834
viridifolium, 834
cyneale, 834
Aspice, 639
Atarigo, 576
Almond, 505, 863
Bitter, 88
Oil, 315

- [illegible]

Athanasia amara 8 5
Athyrocarpus pericarpinifol-
ium, 814
 Atlain, 148, 450
 Group, 450
 Atkinson, G. F., 96, 234, 237,
 238, 240, 803, 870
 Atkinson V. T., 870
 Attractylin, 83
Atractylis, 863
gummifera, 83
Atriplex 426
canescens, f. 423
hastata 107, 813
horrida 107, 4-4 & 3
locynata, 107, 813
nutans, 07 813
pauciflora, 813
torreyana, 813
vestita, 8 3
Atriplicum 407, 426
Atropa 863
Belladonna, 4 60 133 48,
 150, 715 f 716 853
Atropamin, 715
Atropin J. 60, 61, 82, 148
 49 238, 584, 715, 715
 726, 728, 730, 731, 73
 863
 Poisoning by, 61 77 78
 Treatment for poisoning,
 79
Atropa Phosphorus, 2
 Atropin of Rose, 699
Aucuba, 870
Aucuba japonica, 664, 817
Aucubol 664
 Aucub. R. J. M., 881
 Aucub. S. H., 870
Aurantia, 78
A. antiochiensis 583
Auricularia, f. 20
Aurora, 8, Economic plants
 of, 864
 Pines of 865
 Poisonous plants of, 867
Aurora, 8, Gastrolobium,
 533
 Indigo, 534
 Salt-hubb, 107, 425
 Wattles, 523
Austria Poisonous plants of
 867
Austrian, Pine, 529
 Turpentine, 329
AUTOBASIDOMYCETES
 220, 233-247
 Auto-intoxication, 10 863
 Auto-poisoning, 73
 Group 339 338 339
Autum 102, 215, f 338, 339,
 826
Autum, 102, 215, 159, f
 360
 Avery S., 96, 348, 870, 904
 Avocado, 478
 Oil of, 478
 Awned Bromo Grass, 339-361,
 f 361
 Axwort, f 523
Azalea, 666 667
 California, f 668, 668-669
 torreyana, 666
Azolla, 315
Azorella glebaria, 856

Babcock, H. H. 393
Baccharis, 75
Baccharis cordifolia, 138, 755,
 815
holimifolia 138

Bachelor's Button, 756
BACILLARIALS, 153, 168-
 189
Bacillus, 171, 863
aceticus, 163
acid-lacticus, 230
alkali, 174
amyloferus, 163
anthracis, 97, f. 160, 775,
 863
anthracis & *symptomatici*,
 175
botulinus, f. 174, 174, 857
cloacalis f. 161
coli, 172
diphtheriae 97
enteridis, 172-173
lactis, 175
marisnigricans, 176
oedematis, 175 863
oedematis v. *maligni*, f. 175
parvulus agilis, 172, 859
prodigiosus, 163
putrificans, 176
Sorghii, f. 163
rubripes, f. 160
subtilis, 171 172, 863
terrestris, 174, f. 174, 863
typhi, f. 161
typhosus, 173, f. 173
virgula, 863
Bacomyces roseus f. 302
Bacon, Alice R., 870
Bacon, C. W., 870
Bacon, H. P. 52, 62, 870, 871
Bacteremia, 170, 173, 176
Bacteria, 10-11, 97, 160, f
 161, 161-163, 775, 899
 Biology of, 161 163
 Nutrient, 163, f. 163
 Poisonous properties, 18,
 163-165
 Saprophytic, 10, 97
 Soil, 163
Bacteriactae, 163, 171 182
Bacterial poisons, 864
 Classified, 10 11
Bacteriolysin, 165, 177 182
Bacterium, 864
anthracis, 172-179, f. 174, f
 179
asthenes, 177
atium, 177
baylisianum, 177
canaliculatus, 176
cholerae, 177
coli commune, 177
diphtheriae, 179, 864
leptum, 180
mallei, 180
necrophorum, 179
pestis, 181
pneumoniae, 76
putrificans, 177
suicida, 177
tuberculosis, 180, f. 181,
 864
Bacteria festuensis, 839
 Bailey F. N., 871
 Bailey W. W., 871
 Bailon, H., 871
Balanites, 89
egyptiaca, 853
Roxburghii, 853
BALANOPHYTES, 416
Balanops, 399
BALANOPHYTES, 399
BALANOPSIDALES, 146,
 399
 Balfour, A. 346
Baliocarpum acillare, 823
 Ball C. R. 345, 871 903
 Ballardini, 871

Balloon Vine, 604
Balm, Sweet, 710
Balm-of-Gilead, 397, f. 399
Balmoney, 134
Balsam Apple, 751
 Calaba, 627
 Family 618-619
 of Copaliba, 78, 529, f. 330
 of Tolu, 528
BALANINACEAE, 123, 607, 618-
 620, 811
Bamboo, 538
Banana, 391, f. 391
Banberry 467-468, 864
 Red, 168 467-468
 White, 108
Bang, B. 871
Banyan Tree, 405-406
Baptisia, 118, 530, 531 834,
 535, 539-542
alba, 90
australis, 90, 540
benecata, 117, f. 540, 541,
 542
leucantha, 90, 117, 541, f.
 541
perfoliata, 90
stictica, 90, 540-541, 829
versicolor, 90
Baptisia, 533, 540
Baptismin, 530
Barbarea vulgaris, 818
Barbatoria, 573
Barberry, 468, f. 469
 Family, 112 469-472
Barbieria maynensis 829
polyphylla, 829
Barbosa, R. J., 871
Barr, C. L., 871
Barger Geo., 278, 871
Barilla, 424
Barium, 864
 in Loco Weeds 40, 566, 567
Bark, *Acacia* 533
Angustura, 582
Calaya, 741, 742
Canella, 627
Cascarilla, 588
Cherry, Wild, 518
Cinchona, 78, 87, 145, 146,
 741, 742
 Clove, 478
 Cotton Root 624-626
Cupress, 742
Dios, 691-692
Dogwood, 664
Eunifera, 642
Hemlock, 327
Logwood, 528
Maple, 613
Mulberry, Paper, 406
Oak, 403
 Paper 642
Persea, 147
Peruvian, 144, 741, 742
Quillaja, 78
St Ignatius, 688
Simaruba 583
 Soap, 505
 Walnut, 401
Barker 248 871
Barigera cristata 692
Lupinus 692
Pronota, 698
Barley, 366-367
 2-rowed, 366
 4-rowed, 366
 Little, 367, f. 368
 Smut, 98, 217
 Wild, 66, f. 67, 103, 338, f
 368

- Po Tash*, 486
Rapa, 490
Brayera anthelmintica 503,
 f. 505, 595, 847
Brazil Nut, 638, f. 638
 Poisonous plants of, 667
Brazilian Ipsecac, 63,
Magonia, 64, 407
stinging Nut, 596
Bread fruit, 405
Breadfruit Tree, 405
Bready G. W., 630
Breckin C 320, 874, 883
Bredeman G. 874
Breeder's Gazette, 561
Breifeid, O., 213, 256
Bremsia Lachrym. 205
Bretin, 874, 884, 914
Bridelia ovata 53
reissig, 822
Brewer, W. H., 564
Brier Green 104
Brigham Johnson 863
Brimhall, 25
Brielle Fern 313
Bristly Puntan, f. 330
Brixton, N. L., 874
Briod, 90, 836
Broad Bean, f. 422, 523
Brodie, J. A., 874
Brodie, E. C., 874
Broken-back, 575
Brome Grass, 66
 Awned, 359-360, f. 361
Bromelia, 89
Pinguin, 378, 811
Бромелия, 89, 372, 811
Bromelia, 273
Bromin 72
Bromus, 226, 340, 359-360
catharticus, 846
malis 826
secalinus, 826
Septoria on 286
tectorum 66, 359-360, f. 361
Branchitis, 170, 171
 Membranous, 263
 Purulent, 170
Brachomyces, 264
Brachopneumomycosis, 261
Brake, J. 874
Broom, 429
 Common, 147
 Lord 144, 864
 Rare, 698
 Scotch, 110, 45
Broom-weed, 821
Bruckmann Alcasimus, 405
Brusard, J. 874
Brown A. 874
Brown, Crum, 146
Brown, E. L. 874
Brown, H. J. f. 662
Brown S. A. 874
Brown Mustard, 490
Brucella simulans, 853
Brice 145, 146, 147, 686
Brinjal 715, 853
latifolia 715
Brimbach, J. C. 34, 874
Brimston, T. L. 518
Brussels Sprouts, 486
Bryonia alba, 751, 818
diversa 135, 751, f. 790, 818
Bryonia n. 751
oryzoides, 751
oryzoides f. 749, 751
Black, 149
 BRYOPHYTES, 155, 308, 312
 Bryophytes, 310
Bryum, 310
Bubonic Plague, 181
Buchanan, R. E., 24, 243,
 720, 859, 874
Buchanania, 805
Buchner, H., 230, 302, 875
Buchner, M., 875
Buchu, 87, 583
Buckhead, 129, 690-691, f.
 691
Buckeye, 123, 616-617
 California, 123, 617-618
 Family, 123, 616-618
 Ohio, 123, 617, f. 619
 Red, 123, 617, f. 619
Buckley, J. S., 25, 263, 265,
 875
Buckley, S. S., 875
Buckthorn, 123, 124, 229, 628
 Family, 620-621
 Sea, f. 640, 641
 Southern, 629
Buckwheat, 12, 78, 106, 419
 419, 421
 Family, 418-421
 Flour, poisonous, 419
Rudd, Charles, 875
Budding, 52, 89
brasilensis, 836
brasilensis, 836
globosa, 836
ladleyana, 836
madagascariensis, 836
pulcherrima, 836
variegata, 836
verticillata, 836
Buechner, J. G., 875
Buffalo Berry f. 641
 Har. 718, f. 722, 722-724
Buglossa, Bristles of, f. 71
Viper's, 131
Bulb-bearing Hemlock, 126
Bulbine bulbosa, 376, 834
semibarbata, 834
Bulbine, 237
Bulbus Crowfoot, 110, 459
 Water Hemlock, f. 458, 657
Bull Nettle, 72, 122, 196, 724,
 725
 Thistle, 141, 798, f. 80
Bullard Pierre, J. 803, 863,
 875
Bully Tree, Black, 679
Bumelia lutea, 679
Bumelia, 679
Burch flower, 101, 379
 Common, 379-380, f. 380
 Root, 719
 Sinking, f. 319, 219-220
 Wheat, f. 319
Buphara disticha, 386, 805
Bur Clover 552
Cucurbit, 135
Marigold, 779
Oak, 403
Bur-reed, 312, f. 312
Burckhardt, 298
Burdock, 47, f. 48, 141, 756
 Root, 755
 Source of drug, 87
Bureau, E., 875
Burgess, T. J. W. 875
Burgundy Pitch, 328
Burke, R. W. 875
Burrow, J. B. 875
Burns, Bean, 574
BURMANNIACEAE, 811
Burnet, 16
Burning Bush, 123
Burrill, T. J. 180, 288, 875
Bursaria ramosa, 575
BURSERACEAE, 75, 811
Roschke 304
Buxey, S. C., 875
Bush, Burnag, 123
 Calen, 669
 Cautic, 696
 Honeysuckle, 35
Busse, Otto, 249, 304
Buxey, B. 486
Butter Tail 292
Butter Vegetable, 681
Butter and Eggs, 134, 735,
 Yellow, 78
Buttercup, 77
 Creeping, 110
 Crownfoot, 457
 Family, 89
 Fall, 110, f. 490
 Tufted, 110
Butterfly Weed, 695
Butternut, 401
Butterprint, 124
Butter-crown, 680
Butterwort, 698
Butter-yellow, 78
Button Bachelor, 756
Bush, 744
 Sinkeroot, 648, 692, 755
Butyrospermum Parhi, 680
B. SACAR, 122, 604, 811
Buxus hainanica 65, 811
temperata 122, 604, f.
 811, 811
Hyron J. M., 875
Buxus amoenica, 875
crassifolia, 837
speciosa, 817
 C
- Cabbage*, 486
 Chinese, 486
 Skull, 371
Cacalia ciliolata, 417
Cacao, 621, 624
 Tree, f. 621
Cactaceae, 121, 634-637, 811,
 864
Cacti Spines cause phytoto-
xicity, 68
Cactin, 637
Cactus, 128, 637
 Family, 634-637
 Giant, 635
 Night blooming, 730
 Old Man, 635
Cadaba indica, 812
Cadaverin 10, 149, 172
Cadinene, 648
Calea, T. P. 875
Cacalia, 89
Bandurcia, 829
comata 533, 829
pulcherrima, 829
Canna, 221
Caffeidin, 628, 742
Caffeine, 144, 46, 606, 64,
 628, 742
Cala A., 875
Carnia, 679
Carpenter 396
Carpinus indicus, 521
Carpus, 639
Carpus, 639
Carpus, 124
Cassia Balsam, 637
Calabar Bean, 72, 85, 148,
 528, 864
Calabar 148, 528, 531
Calamagrostis canadensis, 102,
 276
Caladium bicolor, 808
Calamander Wood, 681
Calceola 105, 369
Calandrina, 864

- Cactelaria*, 733
cercatibora, 733
Labiataefolia, 851
Cactiripin, 464
Calcull, Rena, 107 424-425
Calceola oblongata, 756
Calendula, 756
Calicium, f. 308
Callisema serratifolia, 851
Caio Bush, 669
California
Azalea 668-669, f. 668
Buckeye, 23 617-618
Economic Plants of, 864
Flora of, 865
Hellebore, 31 104, 383
Laurel, 112, 478
Lily, 375
Medicina' Plants of, 866
Poison Ivy, 123. 609-613, f.
609
Poisonous Plants of, 866
Poppy, 479, f. 480
Rhododendron, 137, 667
Walnut, 401
Calla palustris, 371
Calliandra, 89
Callitoni, 829
Callitropa americana, 131,
708, 857
cane, 857
longifolia, 857
tomentosa, 857
Callitrope involucrata, 624
triangulata, 624
Callistophus horreensis, 756
Callitria, 819
quadricolor, 328
Callum, Jacob, 875
Colchocorus, 375
Calluna 666
pulchra, 665
Col quetanid herb 666
Camellia, 78
Sweet, 371
Vegetable, 371
Calophyllum, 627
calaba, 827
Inophilum, 827
montanum, 827
Calistrotia gigantea, 809
procera, 65, 695, 809
Collins, 447 448-449
arctica, 843
palustris, 4, 75, 381 418,
449, 845
Caurip, 120
CALYCANTHACEAE, 312, 448,
449, 475-476, 811
Calycanthin, 112 476
Calycanthus 475
decidua, 112. 47 476, d.
873
glauca, 811
sp. 112
Colyptostoma Goepfertiana,
2, 2
Campas, 377 379, 864
Death, 103 375, f. 376, 377,
378, 466
Poison, 378
Swamp, 103
Cambier Jacob, 875
Comefina, 486, 493
saliva, 493 f. 494
Camellia, 89, 629
Hongkongensis, 629
japonica, 855
sasanqua, 855
Cammaria latifolia, 807
Camphora, J. M., 875
CAMPHORACEAE, 52 136, 811,
812
Campbell, H. P., 875
Campbell, J. R., 379
Campbor, 72, 77 87 477, 861
Camphorin, 478
Lamson, 439
Bladder, 437
Starry, 437
Camptotoma, 52
pendulum, 829
Canada Moonseed, 112
Pitch, 327
Thistle, 141, 798, f. 799
Violet, 631
Canadian Cocklebur, 763
Canadin, 468
Canagrace, 418
Canarium commune, 611
sp., 211
Cantharus ensiforme, 521
obtusifolia, 521, 829
Candlecut Tree, 587
CANOLLACEAE, 748
Candolle, A. de, 375, 732
Candytail, 486
Cana, Fronted, 366
Canella alba, 627 812
black, 627
CANELLACEAE, 627 812
Canna edulis, 391
Becerra, 391, f. 391
indica, 391
Cannabin, 106, 411
Cannabim hybrid, 411
Cannabimol, 411
Cannabar, 72, 410, 864
indica, 411 445, 637
sativa, 74, 106, 410-411 /
no. 857
CANNABEE, 391
Cano gum, 405
Cantharellus cantharellus 859
Cantharella, 2 6, 7
Canthardin, 72
Cantua 89
du. etc. etc. 842
pyralis 843
Canuteho, 488
Lane Bean, 574
Caneberry, 715
Jasmine, 747
Myrtle, 637
Saffron, f. 87
Caper, 495
Family, 111 495-497
Spurge, 121 f. 495, 602
Capillaria, 160
CAPRARIACEAE, 77 114, 479,
486, 495-497, 812
Captos (*nephallifera*)
812
feruginea, 812
glabrata, 812
oblongata, 812
spinosa, 495
Caprya bifida, 629, 852
CAPRIPIACEAE, 53, 115, 741,
744 748, 812-813
Capriotin, 726
Capriota, 726
Capsella, 487 493, 864
Burso pastoris f. 492, 493,
818
Capitata, 727 725-726
ovoides, 77, 395, 714, 725,
726, f. 725, 853
Irregularis, 714, 725, 853
minima, 853
Caragana, 527 534
Corona fruticulata, 855
Carapa, Oil of, 575
procera, 575
Carreen-Cactus, A. 402
Caraway, Oil of, 648
Source of drug, 87
Carbonic acid, 6, 77 78, 79
Carbolism, 78
Carbon disulfide, 72
in alkalis, 143
monoxid, 72, 738
Carbonic acid, Poisoning by,
77
Card, F. W., 875
Cardamine anara, 818
pratensis, 818
Cardamon, Bastard, 391
Bengal, 391
Java, 391
Malabar, 391
Round, 391
Cardiac Poisons, 73, 74-78,
148, 692, 803, 864
non-poisonous to insects,
7
Sediment, 637
Cardinal-flower, 136
Cardiospermum Halimolobos,
606, 850
Cardol, 78, 395, 608
Cardoon, 756
Cardopodium corymbosum,
815
Carduus Flodmannii, 801
natalis, 65
Careya, 89
Carex 368, 369
peruviana 368, f. 369
hurta f. 369
Carya, 89
Papaya, 627 841
quercifolia, 627 841
afinosa, 84
Cae CACTA, 627
Cacuin, 627
Caring cinia, 807
Xyliferum, 807
Carlton, M. A. 227, 230
Carlina dracula, 815
Carnation, 836
Carrubra, 148
Cash Tree f. 64, 377
Caroba, 134, 739
Carobin, 134
Carol na Blue-Ray 389
Jasmine, 147
Larkspur, 108, f. 40
Moonseed, 472
Caragea M. 875
Carya moluccensis, 857
Carcin, 627
Carpenter C. B., 874
Cerr F. H., 228, 871
Carragen, 194
Cotton Flower, 64
Carna f. 68 137 647 663
Family, 647-664
Wild, 663-664
Carruthers, Wm., 875 876
Carter Marion H. 876
Cartamus speciosa, 756
Cartwright, C. B., 876
Carum cophaea, 856
Curri, 648
Copicum, 856
Petroelinum, 648, 856
Caryacol, 709
Cavet Geo. W. 272
Catvul, 648, 709
Carya torreyana, 402
glabra, 402
ilinoensis, 401
jacinate, 401

Chancre. Soft 176
 Chaneroid 176
 Chantemesse, A. 262, 263
 Chapman A. W., 876
 Chapman, H. C., 876
 Chara, f. 92, 194
 CHARALES, 154, 193-194
Chardina seranthesmora, 53
 Charlock, 114
 English, 488, f. 489
 Charrus, 411
 Charteris, P. 876
 Chausser, H. 876
 Chavicol, 648
 Chayote, 750
 Ches, 86
 Pentagon on, 283
 Shut f. 118
 Cheeca, 624
Chenanthus, 90
 fragrans, 824
Chenanthus, 78, 48
Chenanthus, 148
 Chen, 486
 Chenanth, 649
 Chelerythra, 113, 485
 Chelidonium, 480, 483
Chelidonium, 480, 483
 maius, 113, 479, 485, f. 483, 841
Chelone glabra, 134, 733, 853
 Cheney M., 484, 485, 503, 602
 CHEKOTOPACAE, 52, 89, 107
 423, 424-430, 813-814
 Poison in reduced by heat, 83
 Chenopodium, 429
Chenopodium, 52, 89, 107
 427-429
 album, f. 424, 429
 ambrosioides, 107, 424, 425, f. 424, 629, 813
 ambrosioides v. anthelminticum, 107, 428, 429
 anthelminticum, 814
 Bonni-Henricus, 427, f. 427, 814
 Hairy, 428
 coifuratum, 814
 capitatum, 424
 hybridum, 428, 814
 mollisimum, 107, 814
 polypernum, 814
 Quinn, 427
 rubrum, 814
 Tulwar, 814
 Chenqua, 640
 Cherry 505, 514-519, 854
 Rich, 404
 Choke, 40, 116, 303, 514, 515
 English, 505
 Ground, 715, f. 715
 Laure, 85, 117, 503, 504
 Leaves, 57, 629
 Powdery Mildew on, 269, f. 269
 Sand, 116, 503
 Sour 505, 514
 Sweet 514
 Wild, 50
 black, 64, 117, 503, 505, f. 515, 516-516
 red, 116, f. 504, 505, f. 516, 516-519
 Western, 515
 Chestnut, K. 5, 11, 44, 64, 65, 84, 96, 107, 113, 113, 121, 128, 130, 135, 241, 328, 378, 379, 381, 383, 385, 386, 388, 394, 403, 430, 434, 440, 443,

453, 463, 463, 466, 467, 476, 479, 517, 518, 519, 537, 543, 548, 549, 561, 564, 565, 568, 581, 593, 594, 612, 617, 636, 638, 630, 651, 653, 656, 667, 669, 677, 684, 694, 720, 722, 730, 733, 762, 769, 775, 782, 784, 793, 803, 804, 876, 877, 892, 916
 Chestnut, 185
 American, f. 411, 403
 Rus, 105
 European, 403
 Oak, 403
 Spanish, 403
 Water, 640
 Chevalier J., 877
 Chevers, N., 877
 Chibin, 575
 Chick Pea, f. 119, 120, 521, f. 523, 573-574
 Chickasaw Plum, 505, 514
 Chickenshoe, 377, 864
 Chickweed, 107, 442, 444
 Common, 442, 444, f. 441
 Chicoria, 137, 759
 Chiecy, 157, 755, 756, 758, 759
 Root, 759
 Chilian Strawberry, 585
 Chili, Poisonous plants of, 867
Chimaphila maculata, 842
 umbellata, 645, 842
 Chimaphila, 663
 China, Poisonous plants of, 867
 Chingona, 848
 Red, 741
 Chinese later 736
 Cabbage, 486
 Ginseng, 647
 Indigo, 421
 Jin che, 641
 Sumac, 583, f. 581
 Tea, 614
 Umbrella Tree, 821
 Yam, 699
 Wood, 141 of, 582
 Yeast, 198
 Chiquipin, Water, 444
Chiococca brachyate, 848
 racemosa, 848
 Chionanthus, 683
 Chionanthus, 89
 Microphloia, 840
 virginica, 89, 120, 683, 840
 Chios Turpentine, 608
 Chirala, 689
 Chiratin, 689
 Chives, 375, 383
Chiodanthus dichotoma, f. 146
 CHLAMYDOMYCEAE, 165, f. 146, 183-184
 Chlamydospora, f. 17
 Chloral, 75
 Chloral hydrate, 6
Chlorocodon illicium, 809
Chlorophora excelsa, 408
 Chloroform, 6, 23
Chlorogalum, 89
 pomelidionum, 63, 104, 389
 CHLOROPHYCEAE, 98, 154, 190, 193
 Binlogy, 190, 191
Chouya ternata, 849
 Choke Cherry, 40, 116, 303, 514, 515
 Choking Distemper, 21
 Chlorocystis, 172
 Cholera, 864
 Chicken, 177

Blag, 8, 171
 Asianic, 181, 182
 Organism of, 182
 Cholenterin, 62, 506
 Cholin, 106, 237, 238, 648
 As a remedy for ptomaine poisoning, 10
Chondria vermicularis, 859
Chondrodendrum tomentosum, 473
Chondromyces pedunculata, f. 169
Chondrus crispus, 194
 CHORIPETALAE, 136
 Choriza, 624
 Christian, Robert, 746
Choristigma Staphylinum, 809
 Christian, A. H., 222
 CHROMACEAE, 184, 185
Chroococcus viridula, 83
Chroococcus viridula, 824
 fructuosa, 822
 Chrysanthemum, 789
Chrysanthemum, 91, 756, 758, 767, 789
 Ratanita, 815
 cinerariifolium, 789, 813
 coccineum, 789
 coronarium, 736
 frutescens, 736
 indicum, 736
 leucanthemum, 789
 Marshall, 789
 Parthenium, 756, 789, 813
 plumifolium, 813
 simplex, 736
 Summer, 736
 Chrysler M. A., 877
Chrysobalanus Icaco, 373
Chrysomela Camphora, 515
Chrysomela Rhabdodendron, 223
 Chrysophan, 518, 528
 Chrysophanic acid, 48
Chrysophanum Camphora, 679, 581
Chrysophanum alternifolium, 83
 ambrosioides, 851
 apocynifolium, 115
 telandra, f. 8
 Chrysanthemum, 277
 Churchill, J., 877
 CHYTRACEAE, 304
 Cicer, 335, 373-374
 astratum, f. 119, 120, 521, f. 523, 573-574
 CICHORIACEAE, 748, 753, 762
 CICHORIACEAE, 757
 Cichorium, 757, 758, 759
 Endivia, 756, 758
 Intybus, 137, 733, 756, 758, 759
 Cila, 44, 50, 651-659, 740
 Bolanderi, 126, 658, 856
 bulbifera, 49, 126, f. 658, 659, 856
 californica, 856
 Douglasii, f. 644, 656
 maculosa, 49, 79, 83, 126, 648, 651, f. 652, 652, 656, 856, 864
 occidentalis, 6, f. 48, 126, 856
 ragosa, 49, 126, 665-668, f. 657, 856
 tomentosa, 856
 terrestris, 49, 83, 126, 652, f. 657, 656, 864
 Cicuto, 656
 Cicutoxia, 50, 77, 79, 126, 149, 435, 475, 648, 652, 653, 656, 864

- Cocklebur 67 137, 767 770
 Canadian, 763, f 769
 Cock-scomby, Gangrene of, 278
 Cocoa 742
 Cocoon not P. m. 370
 Cocoon Antidote. 731
 Cocos amara, 840
 nucifera, 370, 840
 Oil, 627
 Codamin, 147, 48.
 Codein, 59, 77 113, 113, 146, 479 48.
 Codinene, 330
 Coffea arabica, 629, 744, f 74.
 mooriana, 848
 odorata, 848
 Coffee, 629, 741 743 744, 861
 Adulterant of 759
 Bean, f. 116, 117
 Mocha, 744
 Plant, 742 f. 743, 744
 Seeds, 629
 Substitutes for, 523, 525, 527 814
 Tree, Kentucky f 116, 117
 Wild, 745
 Cohagen, S. L., 352
 Cohosh, 75
 Black, 446
 Blue, 112 469
 Coix Lacryma, 339
 Coin acuminata, 621 742
 Habit, 621
 Nut, 621
 Colchicin, 375
 Colchicum, 77, 78, 79, 146, 149, 378, 479
 Colchicum, 6, 72 83, 85, 148, 375 864
 autumnale, 4, 148, f 176, 834
 neapolitanum, 834
 Seeds, 867
 speciosum, 834
 variegatum, 834
 Coleman 61
 Coleosphaerium, 185
 Colopospermum, 223
 solidogonum 98 222, 775
 Sporichium, 86.
 Colin Root, 104
 Colletia, 124
 spinosa, 621 847
 Collins, Eng 863 877
 Collins, G. W. 877
 Collomia grandis, 698
 Coliaturin, 64
 Colostoma antiquarium, 808
 gigantea, 53, 808
 viride, 808
 Colocynthis, 749, f. 750
 Colocynthis, 78, 749, 750
 Colombia Disease, 279
 Color of animals, 386
 Colorado, Poisonous plants of 866
 Coltsfoot, 753
 Colubrina formosa, 621
 Columba Root, 473
 False, 473
 Columbaria, 473
 Columba Aconite, 46
 Columbine Eastern, 446
 European 446, f 447
 Rocky Mountain, 446
 Columbo, American, 689
 Columella, 160
 Colutea, 534
 arborescens, 534
 orientalis, 551
 alyi, f. G. W. 877
 Colza, Indica, 490
 Coma, 375
 Comandra umbellata, 416
 COMBRETACEAE, 53, 648, 814
 Combretum druceanum, 814
 contractum, 53
 erythrophyllum, 814
 phanerophyllum, 814
 racematum, 814
 trifoliatum, 814
 Combs, R., 396, 587, 877
 Comfrey, 704
 Wild, 131, 786
 Comma bacillus, 149
 Commelina diffusa, 814
 nudiflora, 814
 tuberosa, 814
 COMMUNICACEAE, 89 373, 814
 Commiphora abyssinica, 575
 Comocladia, 608
 globosa, 805
 COMPOSITAE, 52, 53, 89, 93, 137-141, 222, 692, 748, 753-802 814-816
 Composite Family, 137
 Concord Grape, 620
 Condango, 696
 Cone-flower, 139, 779
 Tall, 779
 Western, 779
 Conferus, 198
 CONFRUATILES, 198
 Conhydrin, 147
 Conkain, 651
 Conidiobolus utriculatus, f 21
 Conidiophore of Aspergillus, f 21
 CONIFERAE, 155 126, 327-332, 651, 816-817
 Coniferin, 392
 Crusts, 161, 327
 Conicium, 651
 Conin, 3, 75, 77, 79, 126, 133, 144, 147, 648, 651, 652, 653, 726, 728, 747
 Conydrin, 651
 Conium, 2, 6, 44, 650, 651
 Alkaloids of, 147
 maculatum, 4, f 30, 31, 75, 126, 648, 651 652, 655, 864
 CONJUGATAE, 154, 189 190
 Con-unctivitis, 77, 169, 170
 CONNABACEAE, 498, 817
 Connarus africana, 817
 Connexticut, Flora of, 865
 Connor, J. T. 877
 Conopodium demidatum, 856
 Consolida, 706
 Consumption, 180
 CONTORTAE, 150 681-697
 Convolvulin, 104 148, 381
 Convolvulin, 78, 377 384-385, 864
 majalis, 64 104, f 384, 385, 814
 Convolvulin 104, 148, 385
 Convolvulin, 572
 CONVULSACEAE, 53, 89, 130, 698-702 817
 Convolvulin, 130, 699, 702
 Convolvulin, 701 703
 overana, 702
 Family, 130, 698-702
 Aridus, 699
 Scammonia, 699, 817
 scoparium, 699
 sepium, 130, 701-702, f 702, 817
 venosus 817
 CURATIVE PLANTS, 803, 864
 Conydrin, 120, 147, 651
 Cook, Dr., 607
 Cook, E. H. 877
 Cook, O. F., 877
 Cooke, N. C. 877
 Copahu, Balsam of 78, 539, f 530
 Capaic acid, 529
 Capaifera Langsdorffii, 830, 854
 affinis, 529, f 530
 Santalum, 830
 Copas, 854
 Copernicia cerifera, 370
 Copper sulphate as an astringent, 94-95
 Coprinus 234
 comatus, 235
 maritimus, 859
 Coprosma Hirsutifolia, 848
 Copostella flavescens, 848
 Corah Bean, 517 542, 543
 Pungi, f 218
 Tree, 527
 Coral-berry 135, 748
 Corallorhiza innocens, 815
 Corchorus capillaris, 621 f 622, 855
 Cord Grass, Fresh-water, f 60
 CORDAITALES, 326
 Cordia alba, 704
 Cordier, 877
 Cordyrops, 273
 cinerea, f 273
 militaris, f 273
 aphlognathoides, f 273
 Ratanhi, 273
 Taylori, f 273
 Cordone terminalis, 375
 Coran Poxia, Millet, f 149
 Corchorus, 736, 757
 durandii, 139
 cinerea, 256
 CORIACEAE, 607, 817
 Coriaryrin, 607
 Coriander 648, f 640
 Coriandrol, 648
 Coriandrum sativum, 648, f 640, 856
 Cornaria, 607
 myrsinitis, 607 817
 nauphaensis, 817
 russifolia, 607 817
 sermentosa, 149, 607 817
 thymifolia, 607, 817
 Tutu, 607
 Cork 403
 Tree 581
 Wood, f 40, f 717
 Corkwood 621 f 717
 Cormus solida, 303, 847
 Corn, 102, f 337 338 864
 Broom, 345, 864
 Chamomile, 440
 Corkle, 62 107 108, 439, f 440
 Meal, 441
 Seeds, 440
 Crop, Statistics of, 342
 Dene, 341
 Diplodia on, f 287
 Fusarium on, 287
 Flint, 34
 Kaffir 345, 346, 347, 348, 864
 Leaf-browning of, 281, f 282
 Oil, 342
 Pod, 341
 Pop, 341
 Poppy, 112 483
 Rust, f 23

- Cucurbita foetidissima*, 75.
maxima, 750, 819
moschata, 731
Pepo, 750, 751
CUCURBITACEAE, 89, 135, 740, 749-751, 818-819
 Culture, affects toxicity of plant, 87
 Culver's physic, 733, 736
 Culver's-root, 134, 733, 734
 Cumarin, 90, 533, 551, 552, 773, 803
 Cumin Oil, 648
 Cuminum Cuminum, 648
 cuminum, 648
 Cumolinol, 583
 Cumin Seed, 648
 Cupress, 33
Pseudotsuga, 250
Cupress, 637
 ципресс, 638
 ципрессина, 638
Cupressus Lawsoniana, 317
horizontalis, 327
thyoides, 327
 CUPRESS, 72, 687, 706, 788, 789, 803, 864
 A. kaloides, 48
 Plants, 148, f. 683
 Currie, See Curran
 Curarin, 73, 77, 687
 Curcuma Oil, 70
 Curcuma, 593
Curculigo scorodoneifolia, 805
Curcuma turcorum, 391
longa, 391
Zedaira, 393
 Curc, 48, 687
 Curled Jack, 419, f. 400
 Curly Grass, *Hemil.*, 313
 Maple, 613
 Cuscuta Black, 184, 498
Cuscuta, 500
 Indian, 744, 748
 Mianm., 498, f. 300
 Red, 498
 Septaria on, 264
 Spot Blotch on, 281
 Tomato, 715
 Curry Powder, 392, 527
 Cutared Crowfoot, 77, 110, 437, f. 438
Cuscutaria, 505
Cuscuta, 344
Cuscuta americana, 87
ortensis, 70
pal. collis, 817
Urgentium, 87
Epilinum, 701
Epilinum, 130, 701, 817
Epilinum, 87
Cuscuta, 701
 Cuslaw, 751
 Cuslaw, R., 443, 879
 Cuslaw, W. C., 465
 Cuslawgrass, 375
Cusparia febrifuga, 382, 849
viridula, 582-583, 849
 Cusparin, 582
 Custard Apple, 110
 Family, 476-477
 Cut-leaved Water Parsnip, 76
 Cutch, 530
Cyatopsis tetragonoloba, 526
 Cyanide of Potassium, 6
 Cyanides, 72, 73
 Cyanogercis, 53, 348, 864
 Cyanogenetic species, 315
Cyanocephalus, 307
 Cyanosis, 78, 24
CYATHACEAE, 313
Cyathus striatus, f. 245, 247, 860
terrestris, 860
CYTHACEAE, 819
CYCADALES, 323
Cycas circinnalis, 323, 324, 819
media, 325, 819
Cyclamen, 52, 89, 128, 676
europaeum, 4, 128, 844
 European, 676
gracile, 844
hederacolum, 844
latifolium, 844
persicum, 128
 Persian, f. 657
 Cyclamin, 128, 676, 677
Cyclopoma atriplicifolium, 424
CYCLOSPORAE, 194
 Cyclic acid, 533
 Cynin, 133
Cylindropuntia, 636
Cymbidium aloifolium, 840
Cymene, 648, 709
 Cymol, 730
Cymodactylus aculeatus, 809
sepioides, 809
Cymodactylus, 4
 Cynapin, 124, 659
Cynara cardunculus, 756, 815
CYNARFEE, 757, 758
 Cynoglossin, 706
Cynoglossum, f. 48, 705, 706, 864
officinale, 131, 705-706, f. 706, 811
 Seeds, 867
cynoglossum, 706
cynoglossum, 131
Cynomandra betula, 716
CYNOMANDRA, 356, 367, 369, 819
Cypripus, 368, 369
aristata, 819
Papyrus, 369
 Cypress Plant, 130
 Spruce, 121, 600, f. 600
 Scimus, 107
 Vine, 679
Cypripedium, 105, 392, 395, 864
acule, 395
Colletium, f. 391, 395
conditum, 105, 391, 395
hercynicum, 392-395, f. 394
macranthum, 395
montanum, 395
puberulum or *pubescens*, 392-393, f. 393
pubescens, 105, 395, 840
regina, f. 397
spectabile, 105, f. 393, 840
Cystosperma, 53
latifolius, 53
Mertensii, 888
 Cyathus, 122
Cyathopsis, 315, 319-320
alpina, 824
bulbifera, 320, 824
fragilis, 315, 320, 824
montana, 824
Cyclops, 206
Cylinus, 77, 78, 79, 90, 119, 148, 530, 537, 551, 803, 864
Cylinus, 535, 551, 864
Adonis, 90
alpinus, 90
Alchingeri, 90, 838
Antleus, 90
antleus, 830
biflorus, 90, 830
candicans, 90
capitatus, 830
formosissimus, 90
hirsutus, 90, 830
Laburnum, 63, 85, 90, 148
monophthalmus, 90
nigriflorus, 90, 830
polystichus, 90
proliferus, 90, 534, 830
Ruthenicus, 90
scoparius, 90, 527, 551, 830
seasilifolius, 830
surinus, 830
valdensis, 90
Cytomyces, 264
 Czapke, F., 443, 879
 D
DACRYOMYCETINAE, 233
Dactylis glomerata, 229
Dactylis quercina, f. 339
Dactylis Draca, 370
 Daffodil, 386
 Da Camp, J. A., 879
 Dahha, 756
varialis, 756
 Daisy, 776
 Garden, 736
 On-eye, 789
Delbergia, 864
lanceolata, 830
racemosa, 830
 Dakin, R., 879
 Dais, H., 278
Damaecena, 446
 Damaen, C., 879
 Damaena, 864
 Damaen, 627
 Dandelion, 753, 756, f. 750
 Root, 753
 Damaen, F. W., 879
 Dandelion, C. W., 879
 Dandelion, F. W., 747, 874, 879
Daphne, 52, 642-643
genkwa, 642
genkwa, 642
genkwa, 855
laureola, 853
metastemum, 4, 21, 144, 642, 643, f. 642, 853
racemosa, 64
reticulata, 853
 Daphnetin, 125, 642
 Daphnetin, 125, 642
Daphniphyllum *hanceanum*, 821
Daphniphyllum *restrifolia*, 853
Cuscuta, 853
Cuscuta, 853
oleoides, 853
 Darlington, Wm., 879
Darlingtonia californica, 497, f. 498
 Darnel, 4, 72, 74, 103, 36, 364, 864
 Poison, 361-364, f. 361, 363
 Meal, 363
Dario croutense, 16
Darmstadt, 713
 Darnville, Ch. 13, 5, 352, 879
 Date Palm, 370
Datura cannabina, 628
glomerata, 125, 628
DATISCEAE, 125, 628
Datisce, 628, 769
Datura, 88, 148, 717, 727, 729-733, 864
alba, 88, 133, 729
arborescens, 853
atropa, 729
fastuosa, 729, 863

- [illegible]

- Climbing, 315
 Flamy f. 309, 313
 Flowering, f. 327
 Maiden Hair 101, 313, f. 316, 316-317, f. 317
 Male Shield, 149, f. 300, 315
 Extract of, 77
 Ostrich, 321
 Rays f. 309, 313, 322
 Sennu ve, 320-321
 Shield, f. 320
 Sweet, 399
 Tree, 313
 Fertility, Toxicity of plant affected by, 83
Ferula, 650
communis, 856
foetida, 856
galbaniflua, 648, 856
Narthex, 648
Sumbul, 648
Festuca quadridentata, 826
syriaca, 239
Field Marigold, 139, 140, 784-785, f. 785
Fetterbush, 673-674
 Mountain 674, 675
Fever Relapsing, 182
 Rheumatic, 171
 Scur et. 71
 Spiraea, Bacillus of f. 173
 Typhoid, 173
 Bacillus of, f. 173
 Feverwort, 138, 744, 745
Ficaria verna, 751, 810
 Fydel Pau 883
Fiber Cels, 370
 Cotton 623
 Lute, 421
 Plants, 345, 386, 580-581, 621, 622, 623
 Among Leguminosae, 624
 Ramie, 406
 Raphia, 370
 S. 386
Ficaria, 89
FICOIDAE, 89, 874
Ficus, 89
altissima, 406
benghalensis, 406
Carica, 403, f. 407, 857
castanea, 405
Aspidia, 857
Aspidia, 857
leucosticta, 857
religiosa, 406
Roxburghii, 857
Synmorus, 405
Field Larkspur, 108, 460
 Marigold, f. 139, 140
 Mushroom, 235
 Pea, 521
 Penny Cress, 495, f. 405
 Sow Thistle, 759, 760
Fig, 405
Sicyonia, 405
Sycamore, 405
 Tree, 407
Figwort Family, 134, 733-738
 Fibert, European, 403
FILICALES, 312, 313-322, 824
FILICIA, 89, 90, 417
 Folic acid, 77, 319
 Flucin, 319
Filipendula filiformis, 115, 629
 Fily Fern, f. 309, 313
 Fine-leaved Sneezeweed, f. 754
Fir Balsam, 327
 Black, 327
 Douglas, 327
Fischer, A., 874, 883
Fischer R., 302
Fischer, E., 146
Fischer R., 883
Fischer T., 883
Fisetin, 612
Fish, Poisoning from, 77, 78
 Poison, 1, 2, 52, 53, 54, 120, 121, 131, 472, 606, 621, 804, 865
 California, 125
 Soapwort, 593
Fistulina hepatica, 234
FLACOURTIACEAE, 627
Flag, Blue, f. 388
 Carolina, 389
 Exotic, 389
 Large, 105
 Sweet, 105
FLACOURTIACEAE, 625
FLAGELLATAE, 153, 188
Flandin, C., 883
Plavaria Contrayerva, 815
Flax, 120, 580-581, 865
 Annual Blue-flowered, 582
 Dodder, 701
 False, 493, f. 493
 Family, 120, 580, 581
 Fiber, 580-581
 Large-flowered Yellow, 120, 581
 Red Garden, 580
 Seed, 120, 581
Flenbane, 776-777, 778-779
Flerning, George, 13, 277, 823
Flerning, Luke, 746
Flemingia congesta, 810
Flickinger, Dr. P. W., 720
Flinderna Schottiana, 837
Flora, See page 863
FLORIDEAE, 194
Flower-de-luce, 389
Flowering Almond, 505, 514
 Dogwood, 664
 Fern, f. 316, 322
 Plants, 101
 Spruce, 121, 122, 598, f. 308
 Tobacco, 133, 715
Flowers, Poisoning from, 64
 Poisonous, 865
Flora, 681
Flückiger, P. A., 59, 86, 440, 481, 482, 504, 718, 863, 883, 888
Flügge, C., 883
Flugge Leucopyrus, 823
 obocata, 823
Fly Agaric, 14, 31, 74, 99, 148, f. 236, 236-238
 Fungus, 195, f. 203, 204
 Killer, 34
 Poison, 104, 131, 176, 726
Podder Pea, 45
Podere, P. E., 3, 5, 885
Poinsettia vulgaris, 648
Fogel, Estelle D., 96, 883, 903
Fog-Fruit, 708
Foliaceous Lichens, 307, f. 309
Folkulius, 16
Fomes applanatus, 234
Fooda, Poisonous, 865
Foofa Parsley, 76, 126, 659, f. 660
Foot and Mouth Disease, 277
Foot-rot, of cattle and sheep, 179
 of orange and lemon trees, 179
Forage Plants, 538, 525-526
 Poisoning, 20-24, 80, 287, 343, 865
Ford, W. W., Prof., 31, 35, 36, 235, 238, 241, 242, 243, 613, 883
Forget-me-not, 704
Formalin, 78
Formic Acid, 80, 106, 412
Forrest, 883
Forster, E. L., 883
Forstia, 128
intermedia, 840
surpensa, 128, 683, 840
viridissima, 128, 840
Foss, B., 347
Foster, 687
Foul Brand of Bees, 163
 American, 175
 European, 174, 75
Pouquiera splendens, 627
Powder, See *Powder*
Powder, 16
Powd. Meadow Grass, *Powd.*
 ary Meadow on, 273
Poxglove, 75, 736-737
 Common, 385
 Purple, 134, 736, 737, f. 737
Poxia, Carcan, f. 340, f. 145
 Green, f. 307
 Smut, 27, f. 318
Pragma, 507, 509-510
chilensis, 115, 505, 509
indica, 109
ovata, 115, 505, 509-510, f. 509, f. 309, 629
virginiana, 111, 509, f. 509, 629
virginiana var. *floridissima*, 505
Fragrant Golden-rod, 774
 Violet, 631
Fraser, J. F. F., 884
France, Poisonous plants of, 863
Francia, Dr., 25, 54
Francis, Geo., 91, 97, 884
Francia appendiculata, 851
Frangula, 630
Frank, A. B., 288, 296
Frankia purifolia, 823
grandifolia, 825
FRANKIACEAE, 825
Frankforter, C. R., 884
Fraxea carolinensis, 689
speciosa, 689
Fraser's Honeyuckle, 744
Fraser, 146
Fraser T. B., 884, 913
Fraxinus, 629
americana, 683
nigra, f. 682, 683
ovata, 683
peninsularis v. *lanceolata*, 683
Frederick, Dr. H. J., 176
Free T. E., 884
Freeman, 884
Freeman, E. M., 364
Freezia, 389
refracta, 389
French Honeyuckle, 525
Mulberry, 708
Pharmacoceia, 866
Sorra, 419
Turpentine, 330
"Franching" of cotton, 287
Fresh-water Cord-grass, f. 69
Eel Grass, 332
Friedberg, L. H., 4, 884, 894

- Friedberger, P. 20, 46, 324, 328, 330, 377, 383, 403, 441, 449, 483, 548, 551, 560, 574, 581, 589, 676, 642, 649, 696, 707, 724, 729, 751, 884
- Fries, 803
- Frijolillo, 343
- Fringe Tree, 129, 683
- Fringed Gentian, 690
- Frullaria*, 377
- Frullaria* 4, 834
- Frullaria*, 834
- Frullaria*, 834
- Frühner E. See Friedberger
- Prothingham, L., 884
- Frullaria, 310
- Fruticose Ligustrum, 307
- Furcraea, 94
- Fuchs, P. A. 263, 884
- Fuchsia, 644
- Fuchsia* *variegata*, f. sp. 1, 194, 194
- Fuligo, 160
- Füller's Tequila, 74
- Fumaria officinalis*, 113, 845
- spicata*, 825
- Fumaria* *acuta*, 113, 834
- Fumaria*, 483
- Fumaria*, 113
- Fumaria*, 113
- Fumaria*, 310
- hygrometrica*, f. sp. 1, 194, 194
- Fumaria*, 53, 90, 98, 195-206
- Fumaria*, 293
- Fumaria* and *poisonous*, Dil. ference between, 239-240
- Fumaria*, 220
- Fumaria*, 28
- Fumaria* by, 31-38, 78
- Fumaria*, 2, 5, 72, 881
- Fumaria*, 295, 300
- Fumaria* Imperfect, 100, 154, 281-306, 861
- Fumaria* of disease, 12
- Fumaria*, Bled's nest, f. sp. 1, 273
- Fumaria*, 273
- Fumaria* *laga*, f. sp. 1, 248
- Fumaria* *cap*, 14
- Fumaria* *rot*, f. sp. 1, 238
- Fumaria*, 498
- Fumaria*, 195, 204
- Fumaria*, 293
- Fumaria*, f. sp. 1, 238
- Fumaria*, 160, 183
- Fumaria* *ultra*, 642
- Führer, R. 99
- Furcraea, 89, 386
- Furcraea*, 803
- Furcraea*, See *Furcraea*
- Furcraea*, 260, 287, 294, f. sp. 1, 861
- Furcraea*, 290
- Furcraea*, 100, f. sp. 1, 293
- Furcraea*, 288, f. sp. 1, 287
- Furcraea*, 287
- Furcraea*, 290
- Furcraea*, 100, 273, 288, 364, 861
- Furcraea*, 287
- Furcraea*, 291, 293
- Furcraea* *destructiva*, 284
- Fustic, 406
- G**
- Gaffky, George, 884
- Gadard, A., 237
- Gastardia puchella*, 756
- Galangal*, 392
- Galanthus nivalis*, 386, 805
- Galea*, 534, 865
- adornalis*, 334, 533, 830
- orientalis*, 830
- GALEGEAE, 504
- Calen, 276
- Galeopsis tetrahit*, 131
- Galium*, 90, 741
- Aporosa*, 848
- asprillum*, 848
- triflorum*, 744, 848
- Galium* *Scorodendron*, 841
- Gallic acid*, 503, 608, 612
- Gallic acid*, 533
- Galium* B. T. 565
- Galium*, Aleppo, 403
- Galium*, K., 884
- Galium*, 742
- Galium* J. 565, 567
- Galium*, 627
- Galium*, 629
- Galium*, J., 213, 261, 290
- Galium*, 63
- GAMOPETALAE, 157
- Langrene, Dry, 28, 100
- Erysipelatous*, 688, 601
- Gaophyllum*, 89
- falcatum*, 450
- Garcinia Cambogia*, 827
- Forsteriana*, 827
- Handbury*, 627
- Mongostoma*, 627
- Garden Cress, 113
- Leek*, 303
- Leek*, 756, 768, 762
- Pea*, 521, f. sp. 1, 507
- Poppy*, f. sp. 1, 111, f. sp. 1, 481-482
- Pussley*, 423
- Raspberry*, 505
- Strawberry*, 505
- Thyme*, 709
- Gardenia jasminoides*, 742
- Gardner* W. W. 884
- Gargel*, 434
- Gargel*, 375, 383
- Field*, 383
- Golden*, 383
- Wild*, 383
- Garner* W. W. 88
- Garnsey*, H. E. 884
- Garrya Fremontii*, 127, 817
- Gas* (illuminating) poisoning by, 6
- Gas Plant*, 581
- GASTEROMYCETES, f. sp. 1, 246
- Gastroenteritis*, 10, 20, 217, 328, 330, 371, 375, 377, 386, 421, 581, 643, 652, 694, 725, 726, 804, 865
- Gastrolobic acid*, 332
- Gastrolobin*, 531
- Gastrolobium*, Australian, 533
- bilabium*, 531, 830
- calycinum*, 533, 831
- calistachys*, 831
- grandiflorum*, 533, 831
- obovatum*, 831
- ovalifolium*, 831
- oxylobioides*, 831
- Poisonous to Stock*, 37, 533
- polystachyum*, 533
- spinatum*, 831
- trilobum*, 533, 831
- Gatzinger* A. 884
- Gaultheria*, 666, 669, 865
- microphylla*, 821
- mammosideroides*, 821
- Oil of*, 665, 866
- procumbens*, 404, 665, 821
- Shallon*, 665
- Gaura*, 644-645
- biennis*, 644-645
- coccinea*, 645
- purpurea*, 645
- Scarlet*, 645
- Gautier*, L. 265, 873, 884
- Gautier* *morchellaformis*, f. sp. 1, 848
- Gaylussacia bacata*, 655
- Gaster*, 248
- Gaster*, f. sp. 1, 246
- Geiger*, P. L., 3, 884
- Geissospermum Villosum*, 147, 807
- Gelsomum*, 77, 80, 129, 147, 689
- Gelsomum*, 147, 689
- Gelsomum*, 688-689, 865
- semperflorens*, 129, 147, f. sp. 1, 685, 689, 836
- Gentianella incana*, 825
- Gentia*, 865
- aphrodites*, 90
- arida*, 90
- germanica*, 90
- monosperma*, 90, 831
- romosissima*, 90
- sphaerocarpa*, 531, 831
- spicata*, 90
- elactoria*, 90, 324-325, f. sp. 1, 831
- GENISTEAE, 530, 151
- Gentian*, Cloned, 690
- Family*, 129, 689-691
- Fringed*, 690
- Horse*, 153
- White*, 690, f. sp. 1, 690
- Yel* *owish*, 690
- Gentiana Andreana*, 689, 690
- arida*, 690
- arida*, 690, f. sp. 1, 690
- puberula*, 689
- turno*, 825
- GENTIACEAE, 120, 683, 689, 691, 825
- Gentianella*, 689
- Gentianic acid*, 689
- Gentiana* *superba*, 831
- Gentiana* *macropoda*, 848
- uniflorus*, 848
- GERANIACEAE, 120, 373, 377, 578, 825
- GERANIALKS, 117, 574-604
- Geraniol*, 577, 709
- Geranium*, Cultivated, f. sp. 1, 577
- Family*, 57, 120
- Oil*, 506
- Robertianum*, 178
- sp.*, 150
- Geranium*, 88
- Gerardia*, 734, 737, 738
- grandiflora*, 134, 738
- purpurea*, 134
- tenuifolia*, 134, 738, 852
- Slender*, 134, 738
- Cerhardt*, 145
- Gerlach*, M., 16, 295, 301
- German*, John, 884
- German* *Chamomile*, 754
- Millet*, 102, f. sp. 1, 350, 350
- Pharmacopoeia*, 866
- Velch*, 572
- Germander*, 709
- Germany*, Poisonous plants of, 867
- Germinides*, 865
- Gerard* A. W., 884
- GERANIACEAE, 698
- Gerum nitrat*, 503
- urbanum*, 505
- Geyer*, C. A., 801
- Gherkins*, West Indian, 750

- Giant Cactus, 635
 Puff ball, 99
Gibberina cerealis 767 f 263
Gibberella, 773 861
Sasibemata 273, f 188 282
 Gibbs, H 884
 Gibbon, E. 884
 Gibbon, W. P. 884
 Gibson, R. J. H. 885
 Gid 569
 Giesecke, J.
 Gift Apple, 885
 Gilchrist W. D., 556
Gilia achillaeifolia, 843
 aggregata 130, 608, 843
 lanceolata 843
Gilman stipularia 103 847
 stipularia 3
 trifoliata 113, 847
 Gilot, Victor, 244, 245
 Gilot, X., 885
 Gilman, J. P. 885
 Gilray E., 284
 Ginter Ward, 344
 Gihette J. D., 729 885
 Gio Juniper used in the
 manufacture of 330
 Ginger, 77, f 300, 391 865
 Wild, f 415, 416
 Gingerol, 391
Ginkgo biloba, 328, 815
 Seed, 393
 GINSENG, 328
 GINKGOAL, 328
Ginseng mexicana, 836
 Ginseng, f. 8th, 647
 Chinese, 847
 Family, 125 646-647
 Gipsies, Prison used by 2
Glechhia pharadioides, 824
 Gihagin, 440
 Githagium, 441
Glaucifera, 389
 communis, 827
 segetum, 827
 Glandera, 180
 Bacillus of, f. 181
 Percy 180
 Glauc, n, 1, 3
Glaucium corniculatum 841
 Aurum 841
Gleditsia 61 89 865
 amorphoides 831
 sinensis 831
 triflorata 507 537 831
Glechima, 313
 lobellata, 825
 GLICHENIACEAE, 313
 Glensk, R., 9 8
Glencapra, 184 187 f 309
Gliricidia maculata, 831
Globba Bedumetia 858
Globularia Alatum, 853
Gloiostrichia, 93, 86-87
 Pisum, 187
Gloiostrichia coliformis, 194
Gloriosa simplex, 834
 superba, 376, 834
 Glover G. H., 96, 346, 347,
 467 885
 Glozima, 698
 Gluco-alkaloids, 149
 Achillein, 149
 Solania, 49
 Glucose 489 505
 Guacoides, 865
 Absinthin, 140, 755
 Achi ein, 140, 149, 756
 Adonidin, 148, 446
 Aesculetin, 6 7
 Aesculin, 123, 617 642, 683
 Anagidatin, 7 54, 83, 116,
 117 144, 303, 584, 585,
 506, 512, 517 518, 519
 Aniamin, 148, 486
 Apocynin, 692
 Apocynin, 139, 148, 692
 Arbutin 505, 506, 665, 666
 Asclepiadin, 696
 Asclepioside 130, 697
 Aurubin, 664
 Baptisin, 533, 340
 Bryonin, 751
 Carcin, 627
 Cathalin, 739
 Cephaanthin, 135, 744
 Cheiranthin, 78, 148
 Cheiratin, 689
 Chocorin, 127 759
 Colocynthin, 77, 749 750
 Convallamarin, 104, 148,
 385
 Convallarin, 104, 148, 385
 Convolutin, 120, 699, 702
 Coronillin, 148
 Crocin, 742
 Cyclamin, 52 89, 178, 676
 Daphnetin, 125 642
 Daphnin, 125, 642
 Dharin 54, 348
 Digitatin, 78 88, 138, 114,
 144, 148, 385, 441 334,
 692 736, 737
 Digitonin, 148
 Digitonin, 134 148, 736
 Digitonin, 134, 377 736
 Diocimin, 583
 Elaletin, 749
 Fricolin, 127 665 666, 669
 Erythrocentaurin, 689
 Erythrophlorin, 418 584
 Euonymin, 148, 615
 Euparin, 722
 Eupatorin, 138 753 772
 Frangulin, 620
 Gastrophin, 531
 Gentiopterin, 689
 Glyceretin, 531
 Glycerethan, 527, 531
 Gentioin, 734
 Gynocardin, 94
 Hellicorein, 449
 Heileboretin, 128, 440
 Hellicorein, 78 148 440
 Hesperidin 383
 Hydrochonan, 526
 Ibu, 683
 Indican 106, 419
 Ipomoein, 130, 701
 Irguin, 589
 Isalpin, 702
 Kolanin, 627
 Lappin, 755
 Leptandrin, 733, 726
 Lanacin, 735
 Loganin, 687
 Lolin, 183
 Lotusin, 54, 534
 Luminin, 118, 531, 548
 Menthin, 62
 Nerin, 138, 148
 Oleandrin, 348
 Ononid, 531
 Ononin, 531
 Oxyococin, 666
 Phaeo-lanatin 54
 Phillyrin, 683
 Pisierin, 328
 Populin, 397
 Prophetin, 749
 Purshamin, 623
 Quercetin, 560
 Quercitrin, 506, 510, 584,
 666
 Rhamnetin, 124 620
 Rhamnin, 124, 620
 Rhamno-cathartin, 124
 Rhiananthin, 738
 Robinin, 119 560
 Salicin, 397 505
 Sclatin, 148, 377
 Sedutin, 377
 Sinabin, 489
 Sinigrin, 489
 Solania, 60 83 32 33
 144, 148, 149 7 8, 72,
 722, 724
 Tampelin 702
 Tephrosin, 533 558
 Thevetin, 148, 692
 Turpethin 702
 Tucin, 149 607
 Veratramarin 383
 Villainin, 508
 Volin 632 633
 Waterin 534
 Xanthoxanthin, 37 769
Gluco-tropaeolum 375
 Gluc, 529
 GLUMIFLORAE, 336 360
Glyca Benghar, 803
Glyceria, 90, 826
 aquatica 83
 canadensis, 692
 hyemalis Ergot on 276
Glycine hispida 520 f 1-
 831
 Sosa 520
Glycyrrhiza 73
Glycyrrhizin, 531
Glycyrrhiza 534
 glabra, 527 f 348
 glabra v. *glandulifera* 537
Glycyrrhizin, 527, 531
Gmelin J. P. 3 888
Gnaphalium vrilanense, 814
 GNATACEAE, 825
 GNETALES, 326
Gnetum grandens, 825
 terrestris, 825
Gnidia carina 642
Gnoscopin, 481
 Goat's Death 64
 Goat's Rue, 118 358
 Godet, C. 862, 865
 Godfrey, B. 883
 Godfrey, George, 885, 890
 Goullee, R. S. 885, 908
 Goepfert H. R., 885
 Gnia, G. 885
 Godkin 888
Golden Crydalis, 143
 Glow 756, 779
 Sea 448, f 668, 688
Goldenrod 138, 222 785,
 767 774 776, 865
 Fragrant, 774
 Rid, 775, 776
 Rust, 222, 775
Goldsmith, W. W. 363 384
Gompholobium, 534, 831
 urcinatum 533
Goniobalanus macrophylla,
 806
Gomolobus laevis, 809
 obliquus, 809
Gonorrhea, 170
 Good King Henry 427 f
 427
 Good, Peter 885
 Goodale, G. L., 633
Goodenia grandiflora 821
 GOODENIACEAE, 825
Goodia, 533
 lobifolia, 533 831
 medicaginea, 533

- Gooseberry, Cape, 713
 Cultivated, 500
 Missouri, 500
 Septoria on, 384
 Goosefoot, 427
 Gooseplant, 417
 Family, 424-430
 Gordon, P. R., 871, 885
 Gordonia, 629
 Gossio, B., 885
 Gossypium, 624, 626
 arboreum, 623
 herbaceum, 623, 625
 herbaceum, 74, 124, 622
 f. 425 623 626, 837
 hirsutum, 625
 Gottheil, 885
 Gotti, 263
 Gouania, 2, 52, 89, 621, 847
 amara, 621
 Gouma, 641, f. 641
 Gourd, Dipper, 75
 Family, 135, 749-751
 Missouri, 751
 Nest-egg, 750
 Sponge, 750
 Towel, 750
 Core, 885
 Gracilaria lemaneiformis, 194
 Graham, H., 885
 GRASS NAK, 33, 89, 90, 102
 103, 356-367, 692, 835
 Crinoid, 767
 Grand Traversa Disease, 341
 Grandidia, 833
 Grape, Clinton, 620
 Concord, 620
 European, 620
 Family, 124
 Fox
 Northern, 620
 Southern, 620
 Pruitt, 587
 Janerville, 620
 Mustang, 124, 620
 Oregon, 112, 472, f. 492
 Powdery Mildew on, 269
 Small, 620
 Wild Blue, 620
 Worden, 620
 Graphis scripta, f. 308
 Grass, 868
 Beard, 344-348
 Bent, 223
 Bine, 338, f. 339
 Ergot on, f. 27, 276
 Powdery Mildew on, f. 27, 272
 Rye-joint, Ergot on, f. 27
 Bottle, Ergot on, f. 27
 Bristly Foxtail, f. 320
 Brume, 338
 Chum, 406
 Cord. Fresh-water, f. 69
 Curly Small, 345
 Cuscuta, 345
 Eel Fresh-water, 332
 Family, 89, 90, 336-337
 Foxtail, 207, 349, 865
 Fowl Meadow. Powdery
 Mildew on, 272
 Holy, 339
 Hungarian, 102, 350
 Johnson, f. 344, 345, 346,
 865
 Knot, 421
 Lemon, 344
 Meadow-out Tall, f. 34
 Mexican, 342
 Needle, 66, f. 70, 102, 330,
 f. 355, 356-357
 Western, f. 66, 102
 Orchard, Powdery Mildew
 on, 272
 Rust on, 229
 Spot-disease of, 282, f.
 283
 Pigeon, 98
 Potentilla, 102, 355-356, f.
 356, 865
 Smut on, f. 216
 Poverty
 Long-awned, 352-353, f.
 353
 Short-awned, f. 353
 Quack, 103, 164, 165
 Black-spot Disease of, f.
 280, 280
 Ergot on, 276
 Quaking, 90
 Rame, 406
 Rattlesnake, 692
 Red Top, 338, 845
 Ergot on, 277
 Rice-cut, f. 338, 339
 Rust, f. 223, 224-227, f.
 223
 Rye, 361-364
 Common, 161
 Italian, 161
 Salt, 90
 Sleepy, 102, 355, 357, 358
 845
 Smut, 98, 212, 214-219
 Snake, 492
 Sprowing Smut, f. 214
 Squirrel-tail, 66, 103, 360,
 364-367, f. 368
 Injuries from, 66, 360
 Powdery Mildew on, 272
 Staggere, 21
 Sweet, 865
 Tickle Rust on, f. 217
 Triple-awned, 352
 Tulip, 868
 Vanilla, f. 41, 552
 Vernal, 354
 Sweet, 352
 Wheat, 364
 Western. Frost cut, 276
 Wild, Ergot on, 76
 Wire, 374
 Yellow-eyed, 372
 Gratiola officinalis, 1, 134,
 734, 852
 peruviana, 852
 sp. 134
 Gratiolin, 734
 Gravel-root, 772
 Graves, C. B., 881, 885
 Grawitz, Paul, 260, 295, 298,
 302, 303, 885
 Gray Dr. Ann, 333, 602, 885,
 886
 Greasewood, 68, 107, 429, 430
 Great Angelica, 660
 Lobelia, 753
 Ragweed, f. 136, 137, 762,
 766
 St. John's-wort, 123, 629
 630-631, f. 615
 Willow-herb, 444
 Great-flowered Larkspur, 468
 Magnolia, 473, f. 474
 Great Britain. Poisonous
 plants of, 867
 Green, J. R., 250, 486, 886
 Green, Wesley, 886
 Green Algae, 160
 Ash, 683
 Brier, 104
 Hellebore, f. 45, 75, 109,
 449
 Mold, f. 257, 357, 261
 Pigweed, 107
 Tea, 628
 Greene, E. R., 746
 Greenish, H. C., 886
 Greenleaf, R. W., 886
 Greiner, Karl, 886
 Gresham, M., 54, 65, 84, 88,
 89, 90, 96, 101, 110,
 114, 115, 124, 128, 130,
 135, 141, 315, 436, 447,
 498, 500, 303, 535, 567,
 590, 621, 698, 750, 803,
 804, 886
 Gressler, P. G. L., 886
 Grevillea miconioides, 844
 Grewia, 2, 52, 89, 621
 asiatica, 855
 bracteata, 855
 Malaccana, 855
 orientalis, 855
 pilosa, 855
 pycnantha, 855
 Grey, Geo., 886
 Griffiths, David, 633
 Grobe, 886
 Gromwell, 620
 Grigorenko, G., 886
 Grindelia, 89, 757, 774
 lanceolata, 138
 robusta, 774, 818
 squarrosa, 138, 755, 774
 Tenuifolia, 815
 Grindelin, 755, 774
 Grönlund, C., 886
 Gross, Herman, 886
 Ground Cherry, 715, f. 7-9
 Ivy, 131, 710, f. 711
 Plum, f. 961
 Grounds, 793-797
 Salt, 138
 Grouse poisoned by Laurel,
 669
 "Grub-in-the-head," 138, 776
 Gruby, J., 295, 301, 302, 886
 Grünfeld, 279
 Guaco, 417
 Guaiacum officinale, f. 878, 882
 Guaiacum, 146
 Guapachum, 80
 Guay, 526
 Guayana, 606
 Guayana, J., 886
 Guatteria venenifera, 806
 Guava, 523, 639
 Guayana, J., 886
 Guenther, J., 887
 Guerin, P., 363, 364, 887
 Guiana. Poisonous plants of,
 867
 Guignard, L., 119, 574, 747,
 887
 Guinea Pepper, 725
 Guislotia abyssinica, 755
 Gum, Canoe, 405
 Elastic, 588
 Galbanum, 648
 Kino, 523, 640
 Sandarac, 328
 Sour, 664
 Tragacanth, 520
 Tapelo, 664
 Gumbo, 623
 Gumweed, 755, 774
 Gun Cotton, 623
 Gustaf, 52
 Augusta, 814
 brasiliensis, 814
 Guita, 681
 Guita-percha, 680, 681
 Tree, f. 680
 Gottenberg, G., 96, 435, 887
 Guttifer, 827
 Guttifer, 370
 Guttifer, 874

- HEMIBASIDII**, 209, 210-220
Hemidesmum, 90
indense, 695, 809
Hemileia vastatrix, 222
Hemiphaea microdonta, 138
Hemlock, 327, f. 327, 651-659
865
Bulb-bearing, 126
Pitch, 327
Toison f. 60, 51, 73, 23
648, 651
Purple-stemmed, f. 655
Water, 126, 648, 651-659,
865
Bulbous, f. 658, 659
European, 617
Oregon, 656-658, f. 657
Hemolysis, 15, 36, 242
Hemorrhagic Septicemia, 177
Hemp, 74, 106, 410-411, 531
Common, f. 420
Indian, 72, 129, 411, 692,
693, f. 693
Manila, 391
Mauritius, 386
Queenland, 624
Hempel C., 890
Hemphalea, 2, 4, 718, 726-727
Black, 33, 726-727, f. 727
Henderson, 278
Hensel, Alice, 85, 732, 890
Henna, 638
Hennings, P., 229
Henry Almé, 863, 890
Henry J., 890
Henry T. A., 54, 84, 348,
450, 534, 881, 890
Henry W. A., 313
Henslow G., 862, 890
Hepatica, 310
Hepaticaria emarginatum,
808
acuminata, 808
venulosa, 808
Heraclaea Poisonous honey
n., 84
Heraclaea, 650, 661, 865
lanceata, 126, 648, 663, 865
spondylium, 865
Hernand, 126, 663
Hernand of Grace, 76
Hernandium Mold, 257
Herba, milianoides, **Breata**
discover of, 2
Hernand, J., 890, 895
Herman L., 890
Hernandria, 53, 89, 436
glabra, 83
Hernandria-amarum, 62
Hernandria-amarum, 828
Hernand, 147
Hernand, 75, 17
consueta, 301
Herr Bertha D., 803
Herring M., 890
Hesperian, 583
Hester Dr., 655
Heteromorphia arborescens,
856
Heteropogon confertus, 354
lanceus, 826
Heteropogon stringosifolia,
837
Herber Fr., 890
Hesperia americana, 500
Herringer H., 262, 295
Hesper, 587
brasilensis, 53, 588
Hesper F., 890
Hesper, 637
Hesperia, 687
Hibbertia glaberrima, 819
longifolia, 89
Hiburnia abelmoschus, 624
canadensis, 623
diversifolia, 837
glabra, 623
esculentum, 623
glabra, 623
Yabdariffa, 623, 624
hiocens, 623
Hickman, R. W., 25, 891
Hickory, 402, 403
Hard-shell, 402
Missouri, 401
Shell-hark, 402
Hieracium, 757
tenaxum, 692, 755
virgatum, 816
Hieracium, 90
amarum, 826
botryata, 826
odorata, 539, f. 341
viridula, 826
High-bush Blackberry, 508, f.
308
Hilger A., 891, 892
Hilman, F. H., 378
Hilmer, R. S., 348
Hinebaugh, T. W., 250, 251,
891
Hippocastanum equestre, 805
rutilum, 805
HIPOCASTANACEAE, 121, 607,
616-618, 877
HIPOCASTANACEAE, 52,
63, 89
Hippocastan, 372
Hippocastan, 865
Manimella, f. 586, 587, 823
Hippocastan rhomboides, 125,
f. 640, 641
Hippocastan vulgaris, 400
Hirta, 723
Hitchcock A. S., 891
Honey Pea, 558
Huchetter, C. P., 891
H. H. H., E. R., 891
H. H. H., F. R., 891
H. H. H., R. F., 891
Hog Pea, 204
Plum, 608
Hogweed, 766
Hogwort, 131, 591, f. 301
Holcus, 826
lanceus, 90
Holcus, R. J., 222
Holcus confertus, 805
lanceus, 608
longifolia, 805
Holland, J. W., 0
Hol v. American, 614
European, 614
Family, 123, 614
Sea, 648
Hollyhock, 623
Holm, T., 6-6, 772, 891
Holmes, F. M., 587, 891
Holy Grass, 339
Homalanthus crepitans, 823
Lechenanthus, 588, 823
lanceus, 53
Homalanthus aromaticus, 808
Homen, E. A., 891
Homeria collina, 390, 827
Homocheidom, 483
Honey Locust, 527
Plant, Simpson, 134
Poisonous, 64, 121, 601,
606, 865
Honeycomb Ringworm, 294
Honey-dew, 275
Honey-suckle Bush, 135
Family, 135, 744-748
Fraser's, 744
French, 525
Rocky Mountain Bush, 135
Succulent's, 744
Transpet, 744
Western, 744
Honeywort, 6-6
Hoodia, 277
Hopper D., 891
Hop Clover, 552
Low, 553
Yellow, 553
Tree, 581
Hops, Common, 106, 406
Oil of, 406
Horand, 14
Hordeum, 340, 366-367
cocciferum, 103
distichum, 366
Guzonianum, 103
subatum, 66, f. 67, 103,
366-367, f. 368
Ergot on, 276
maritimum, 36, 367
nodosum, f. 67
arabicum, 103, 367, f. 368
spon. densum, 366
vulgare, 103, 366
Horchard, 709
Horse Gentian, 135
Nettle, 60, 122, 728, f. 729,
724
Horse-chestnut, 123, 617, f.
618
European, 617
Horseshoe, 113, 435, 486,
490-491, f. 497, 656
Hornetail, 37, f. 41, 101, 312,
322-326, 865
Horn-wood, 724, 755, 777,
778, f. 778
Small, 777
Horsfield, Thomas, 891
Hosokawa Parshana, 525, 831
Houghton, Dr., 411
Hound's Tongue, 131, 705,
706, f. 716
Hoselock, 221
Howard, C. D., 517, 891, 900
Howell, Miss J. K., 231
Howa cornuta, 695
Huart, L., 891
Hubbard R., 891
Huck cherry, 663, 719
Huff, E. H., 771
Hugmeyer, 891
Hughes, D. A., 541, 891
Hughes, Richard, 891
Hugonia Myrtus, 835
Humea, 90
Humulus Lupulus, 105, 406,
857
Oil of, 106, 406
Hungarian Grass, 102
Hungary, Poisonous plants
of, 867
Hunt Reid, 686, 891, 893
Hunt, Annie, Oakes,
612, 862, 891
Hurd crepitans, 587, f. 887
Husmann A., 4, 892
Husmann, Th., 4, 891, 892
Hutchinson D., 892
Huyra, P., 13, 26, 692, 893
Hysanth, 375
Hysanthus originalis, 375,
834
Hyasna Poron, 588
Hyasna, C. W., 380, 415, 541,
580, 663, 677, 688, 747,
892
Hyatt, Virgil, Cape of, 49
Hybanthus peractiantha, 632
Hyde J. N., 13, 9, 295, 298,
299, 302, 892, 900

- Hydnocarpus*, 52
onthetismmum, 810
castanea, 810
heterophylla, 811
nebrina, 54
Kuruk, 811
perdis, 811
venenata, 53, 637, 811
Wrightiana, 811
- Hydnium amarissimum*, 860
gracile, 860
imbricatum, f. 232
- Hydrangea*, 84, 89, 300, 629
arborescens, 115, 500, 851
 Cultivated, 500
 Garden, 500
Hortensis, 84, 115, 300, 629, 851
involucrata, 500, 851
Lindleyana, 500, 851
paniculata, 500
Thunbergii, 500, 851
 Wild, 115, 500
- Hydrastis*, 108, 146, 448
Hydrastis, 447, 448
canadensis, 88, 108, 446, 448, f. 448, 846
- Hydrastium*, 146
Hydrocyanic acid, 73
Hydrochinon, 506
Hydrocotarnin, 481
Hydrocotyle asiatica, 856
japonica, 856
umbellata, 856
vulgaris, 856
- Hydrocyanic acid*, 4, 6, 53, 55, 77, 80, 82, 81, 89, 90, 101, 102, 106, 115, 116, 119, 120, 128, 140, 141, 313, 346, 348, 447, 498, 500, 503, 512, 517, 518, 519, 567, 574, 590, 622, 637, 633, 681, 747, 804, 862, 865
- Distribution in Vegetable Kingdom, 33-55, 89-98
- Effect of 54
 Toxicity of, 7
- Hydrodictyon reticulatum*, 190
- Hydrogen, Peroxid of, 72
 Hydrogen in Alkaloids, 144
 Hydrogen peroxide sulphate, 489
- HYDROPHYLACEAE**, 331, 698, 703-704, 827
- Hydroquinon*, 505, 742
Hygrin, 147, 575
Hydrophilus aspidus, 804
Hydrophorus agnoscens, 53
Hymenocallis, 523, 527, 831
- Hymenocallis*, 805
- HYMENOLICHENES**, 154
Hymenogaster tener, f. 246
- HYMENOGASTRINEAE**, 245-247
- HYMENOGASTRACEAE**, 245
- HYMENOMYCETES**, f. 332, 281
- HYMENOMYCETINEAE**, 233-245
- HYMENOPHYLLACEAE**, 313
- Hynson, H. P., 881, 892
- Hyoscyamus*, 823
Hyoscin, 148, 716, 726, 727, 730
Hyoscin-pseudocyanin, 148
Hyoscyamin, 3, 60, 61, 77, 133, 148, 716, 726, 727, 730, 732, 736
- Hyoscyamus*, 2, 52, 60, 71, 78, 148, 716, 717, 726, 727, 865
alba, 853
serotus, 854
Pauciflorus, 854
muticus, 854
niger, 4, 60, 133, 726-727, f. 727, 854
physaloides, 854
reticulatus, 854
Hyoscyamus procumbens, 841
- HYPERICACEAE**, 124-125, 629-631
- HYPERICINEAE**, 827
- Hypericum*, 629-631
Ascyron, 125, 629, 630-631, f. 631
crispum, 827
humiflorum, 827
Kalmianum, 629
maculatum, 124, 630, 827
 Oil of, 630, 631
perforatum, 125, 629-630, f. 630, 631, 827
prolificum, 629
punctatum, 630
Hypha, f. 17
Hyphosia thibetica, 840
- HYPHOMYCETES**, 249
- Hyphosia indica*, 840
Hyphosia muscosissima, 859
- Hypnotic Poison*, 804
Hypocritic facies, 75
- HYPOCREACEAE**, 100, 273-278, 859
- HYPOCREALES**, 273-278
- Hyslop*, Hedge, 134
- I
Iberis, 486
Ibo, 683
Ibota, 683
Icaca Plum, 505
Ica Plant, 423
Ichthyophora, 53
Cucub, 816
Ictogen, 78, 149, 548
Idaho, Poisonous plants of, 866
Idaho Pea, 120
Ilex, S., 248
Ilex Aquifolium, 123, 614
Caryocarpus, 123, 614, 742, 807
glabra, 807
opaca, 614
peruviana, 614, 742
thibetica, 614
verticillata, 807
Ilex, 614
- ILICINEAE**, 827
- ILICACEAE**, 89, 827
- Illicium*, 89
anatum, 149, 474, 837
horridum, 112, 474, 837
peruviana, 474
- Illinois State Board of Health**, 892
- Ilipa Macleania*, 881, 851
Imbert-Gambeyre, 892
Immunity, 165, 862, 865
Impaction, 9, 68, 102, 342
Impatiens, 618, 619-620
amphorata, 850
aurata, 850
Balsamina, 619
bulbosa, 620, 850
fulva, 123
Noli-tangere, 123, 810
Royalia, 123
Imperatoria Ostruthium, 648
Imperatoria, 125
- Inula*, 692
Indaconitin, 450
India, Poisonous plants of, 867
 Wheat, 419
India-rubber Tree, 405
Indian Aconite, 449, 450
 Almond, 638
 Bean White, 574
 Berry, 1, 149
 Cola, 490
 Cucumber, 104
 Currant, 744, 748
 Heliotrope, 705
 Hemp, 72, 129, 411, 692, 693, f. 693
Ipecacuanha, 693
 Laurel, 478
 Mallow, 124, 626-627
 Mustard, 490
 Opium, 85
 Pear, 635
 Pennywort, 648
 Persimmon, 681
 Pink, 129, 688
 Sarsaparilla, 693
 Shot, 391
 Strawberry, 508, 509
 Teak Tree, 708
 Tobacco, 73, 136, 712
 Turnip, 372, f. 372
- Indiana*, Poisonous plants of, 866
- Indians*, Food plants of 106
Indican, 106, 419
Indica, East, Fish-poison of, 1
 Indigo, 434
 Australian, 834
 Blue, 324
 Chinese, 421
 Large White Wild, 117, 541, f. 367
 Large-fracted Wild, 541
Indigo-caster, 333
Indigofera, 534
And, 324, 831
australis, 834, 831
galeoides, 53, 747, 831
hirsuta, 831
Inebrians, 74, 804
 Inflammation of lateral or gans, 172
 respiratory passages, 170
 serous membrane, 171
Iosatin, 752
Influenza, 176
Inga vera, 525
Ink Tree, 608
Inocybe, 244
achnata, 244
infida, 244
scabra, 860
Inula, Y., 237, 892
Insect Powders, 789
Dalmatian, 789
Persian, 789
 Trap, f. 490
Insecticide, 804
 Hydrocyanic acid aa, 54
 Insectivorous plants, 498
 Intoxicant, 804
Inula, 757
Helenium, 754
Royana, 816
- INULEAE**, 757
- Inula*, 753, 754, 756
Ionidium Ipecacuanha, 858
Iowa, Poisonous plants of, 866
- Ipecac*, 633
 Brazilian, 631
 Powdered, 78

- Roots of, 76, 742
Wild, 599 f. 600
Ipecacuanha, Source of, 87
Indian, 693
Ipooh, 863
Ipomoea, 89, 701
 Batatas, 899, f. 699
 Bona-nox, 699
 congesta, 699
 dissecta, 53
 emetica, 817
 fastigiata, 699, f. 700, 701
 Jalapa, 699, 817
 leptophylla, 701
 pandurata, 130, 817
 Purga, 699, f. 700, 701, 817
 purpurea, 699
 Quamoclit, 699
 sinnata, 817
 tuberosa, 817
Ipomoein, 130, 701
Irogenin, 389
IRIDACEAE, 105, 315, 388-390
IRIDACEAE, 827-828
Iridin, 105
Iris, f. 340, 827
 Carolina, 389
 Dwarf Garden, 389
 Family, 388-390
 floribunda, 389, 827
 foetidissima, 827
 germanica, 389, 827
 hybrida, 827
 missouriensis, 390
 neglecta, 827
 palida, 390
 prismatica, 828
 Pseudoacorus, 828
 pumila, 390
 reticulata, 828
 sibirica, 390
 versigata, 828
 versicolor, 105, f. 388, 389, 390, 828
Irish, H. C., 714, 735
Irisin, 389
Iron, Salt of, 7
Ironwood, 763
Iron-wood, 395, 681
Irritant, 76, 804
Irrig., 523, 892
Iscaria farinosa, 273
Iscus corniculata, 818
 incincta, 846
Ischnandra, 53
Isopilocarpin, 148
Isopyrin, 446
Isopyrum fumarioides, 846
 thalictroides, 446, 846
Isorotlerin, 589
Istoria axillaris, 812
 Brownii, 812
 longiflora, 812
Isotropis juncea, 831
Israel, O., 878, 889, 892, 893, 894, 913
Italian Clover, 118
 Millet, 330
 Pharmacopoeia, 846
 Rye Grass, 361
Italie, L. van, 892
Itch, Barber's, 100, 100
 Fungus causing, f. 300
 Dhobie, 299
 Disease of Horae, 293-294
Ithyophallus impudicus, 79, 840
Ito, 757, 763-764
 axillaris, 763
 zonitiformis, 137, 763-764, f. 764
Ivory, Vegetable, 370
Ivy, *Rosina*, 620
 Branch, f. 671
 Ground, 131, 710, f. 711, 865
 Japan, 620
 Poison, 77, 122, 123, 394, 600, 608, 613, 676, 679, 845
 California, 123, 609
 Three-leaved, 609-610
J
Jabberandi, 148, 182, 845
Jabonin, 148, 582
Jacaranda, 52
 procera, 739, 810
Jack Fruit, 405
 Pen, 521
 Pine, 330
Jack-in-the-Pulpit, 183
Jack-my-Lantern, 99
Jackon, J. R., 892
Jacob, J., 891
Jacobi, 278
Jacoby, C., 892
Jacoby, M., 893
Jacquin, 52
 arbores, 839
 armularis, 839
 obovata, 839
Jahn, E., 893
Jain, 130, 699, f. 700, 702
 aspin, 702
Jamaica Dogwood, 74
 Locust Tree, 523
 Scotch, 523
Jambosa malaccensis, 639
 cucurbit, 639
Jambosa Berries, 639
Jamieson, 89, 115
 americana, 115, 500, 851
Jamestown Weed, 332
 jacarandensis, 336
Janiha, 845
Japaneensis, 146, 450
Japan Cloves, 321
 Ivy, 420
 Medicinal plants of, 866
 Plum, 505, 314
 Poisonous plants of, 867
Japanese Ailanth, 647
 Peppermint, 704
 Perrimmon, 681
 Potato, 710
 Sumac, 604
 Walnut, 401
 Yam, 374
Jasmine, 683
 Cape, 742
 Carolina, Alkaloids of, 147
 Pulse, 129
 Large-flowered, f. 684
 Oil, 86
 Yellow, 688-689
Jasminum floribundum, 840
 grandiflorum, 683, f. 684
 officiale, 848
 spicata, 840
Jatobaria Columba, 473
Jatropha, 52, 598, 596-597, 845
 argentea, 53
 Cereus, 587, 595, 596, 823
 glandulifera, 823
 gossypifolia, 587
 macrocarpa, 823
 Manihot, 75
 multifida, 823
 stunlosa, 77, 122, 396, f. 396
 urens, 596, 823
Java Bean, 574
 Cardamon, 391
Javanica, 743
Jeanmire, 893
Jeffersonia diphylla, 89, 469
Jeliffe, S. M., 893
Jenkins, E. H., 130, 701
Jennings, O. P., 893
Jequichy, 534, 863
 Ophthalmia, 56
 Plant, 11, 52
 Seed, 55, 527
Jerusalem Artichoke, 710, 756
Jerrin, 103, 148, 381
Jessamine, See *Jasmine*
Jessup, H. G., 893
Jew Wandering, 374
Jewel Weed, 619
Jewett, 893
Jimson Weed, 4, 60, 61, 131, f. 123, 729-733, f. 730
 Purple, 133, 729, 732-733
Josselyn, 52
 princeps, 823
Jos, T. R., 319
Jos Pye Weed, 771
John, 302
John, J., 26
Johnson, C., 893
Johnson, F. M., 460
Johnson, L., 96, 114, 890, 4, 9, 421, 431, 455, 460, 484, 491, 560, 625, 668, 712, 736, 744, 767, 772, 773, 774, 797, 893
Johnson, H. J., 167
Johnson, T. Y., 454
Johnson, G. W., f. 346, 345, 346
Johns, 179
Johns, J. J., 319
Johns, L. R., 101, 124, 893
Johns, W., 891, 893
Johns, 386
Johnson, A., 893
Johns, K. W., 893
Johns, 327
Johns, F. H., 893
JUGLANDACEAE, 105, 401, 826
JUGLANDACEAE, 136, 400, 402
Juglandic acid, 401
Juglandin, 105
Juglans californica, 401
 cucurbit, 401, 828
 nigra, 105, f. 401
 regia, 401, 828
 Sibboldiana, 401
Jugube, 621
 Chinree, 621
Juncaceae, 374
Juncus tenuis, f. 375, 374
Juniper, 330-332
 Berries, 101
 Common, 101, 330-331
 Low, 332
 Oil of, 101
 Savin, f. 375, 332
Juniperus, 4, 102, 149, 330-332
 communis, 101, 330, 332
 communis v. alpina, 330
 horizontalis, 102
 Knightii, 332
 occidentalis, 182, 331
 Oxycedrus, 330, 816
 Sabina, 4, 102, 149, f. 331, 332, 816
 scopulorum, 101, 330
 virginiana, 101, 330-331, f. 331, 817
Jurabebis, 715
Jurica Grandifolia, 804
Jute, 621
 Plant, f. 622

Lappakoniha, 450
Lappin, 755
Lappula, 706
 echinata, 707
 horibundula 707
 officinalis, 131
 virginiana, 707
Large Arrow-head, 101
Blue Flag, 105
Cranberry, 665
Peppergrass, 494
Rhododendron, 137
Spotted Spurge 597 f. doz
Tupelo, 664
White Wild Indigo, 117,
 341, f. 347
Large-bracted Wild Indigo,
 341
Large-flowered Jasmine, f.
 684
 Trillium, 104
 Winkerbush 385
Larkspur, 43 f. 47 108-109,
 865, 868
 Blue, 460, 466
 Carolina 108 f. 466
 Field, 108, 466
 Great flowered, 460
 Mountain, Tall, 109
 Poisoning, 44-45, 465-467
 Pinnatus, 461
 Prairie, 460, 461
 Purple, 44, 84, f. 46r, 462-
 463, 468, 467
 Rocket, 460
 sp., 446
 Tall, 108, 462, 463, f. 46r,
 465, 466, 467
 Toxicity of 84, 467
 Western Purple, 109
 Wyoming, 109
Larrea mexicana, 818
Larrea, C. 803, 899
Laserpith, 648
Laserpithum latifolium, 648
Lasia oculata 808
Lasiadenia rupestris, 835
Lasiophan anthyllodes, 855
 ariophanta, 835
Latis, Poison elaborated in,
 81
Lathyrism, 44, 45-46, 80, 534,
 572-573
Lathyrismus, See Lathyrism
Lathyrus, 87, 149, 515, 572,
 573
 amurensis, 831
 aphylla, 831, 865
 C. 43 534, 831
 induratus, 45 534 572,
 831
 hirsutus, 572
 ochroleucus, 572
 odoratus, 831
 ovatus, 572
 palustris, 572
 polymorphus, 572
 purpureus, 83
 calycatus, 45, 534, 831
 sulcatus, 572
 turgidus, 572
 venosus, 525, 572
Latic, 894
Latis venenosa, 854
Laudania, 147 481
Laudanum, 481
Laudanum, 627
Laudanum Poisoning, 6
Laudanum purpurifida, 814
LAURACEAE, 812, 444, 477-478,
 828
Laurel, 127 478, 865
 California, 112, 478

Cherry, 85, 117, 323, 504
Family, 112
Indian, 478
Lady, 642-643, f. 647
Mountain, 64, 84, 665, 669,
 671, f. 670
 Great, f. 666, 667
 Spurge, 125, 64, 643, f.
 547
 Swamp, 472
Laurent, E., 895
Lauric acid, 593
Lauricercus, 505
Lauricercus, 865
Lauricercus, 478
Laurus canariensis, 478
 nobilis, 478, 828
Lavandula Spica, 709
 vera, 34, 709
Lavender, 709
 Oil of, 709
Lavender, C. H. 343, 895
Lavender, 3
Law, James, 259, 277, 296,
 293 895
Lawrence, G. W., 245
Lawrence, H., 324
Lawsonia inermis, 638
Laxative, 804
Lazell, F. J., 895
Le Coint, E. R., 505, 506
L'Eglise, Dr., 665, 896, 906
Le Grand, Dr., 793
Lead Poisoning, 6, 72
Lead-color Puffball, f. 266
Leather J W 895
Leatherwood, 123, 443 f. 642
Leaves, Wilted, 102, 116, 119
Liber, Th., 263, 263
Lecanora, 307
 esculenta, f. 308
 rubra, f. 308
Lecanoraceae, 307
Lecanora minor 627
Lectura, M. 895
Lecythaceae, 638
 Fish poisons among, 2
Lecythus, 638
 allura, 638
 amara, 839
 lanceolata, 839
Ledum, 666
 Catchall, 127
 glutulosum, 127 465 821
 groenlandicum 127 466
 latifolium, 639 821
 palustre, 4, 666, 821
Leek Garden, 383
 Wild, 704, 383
Lecrota lenticularis, f. 398
Lera, F. H., 54
Lefebvre, C. 898, 895
Legrand de Saullé, H., 895
Legue C., 896
Legumes, See Leguminosae
LEGUMINOSAE, 52 61, 89 127
 120 149 395 498 502,
 519-574 828-833, 865
 Arrow poisons among, 1
 Famistic plants of, 520-527
 Fish poisons of 1
 Medicinal plants of, 525
 539
 Mildews on, 269
 Ornamental plants of, 527
 Poisonous plants of, 530-
 535
LEGUMINOSAE, PAPI
LIGNACEAE, 53, 89
Lehmann, Otto, 328, 416, 446,
 849, 593, 606, 650, 666,
 719, 721 754, 862 896
Lehmann, K. R., 896, 900

LEITNERIA, 400
 horrida, 400, f. 401
LEITNERIACEAE, 400
LEITNERIACEAE, 136, 400
Lejolinia mediterranea, f. 194
Lemna, 372
LEMNACEAE, 372
Lemon, 582
 Gum Oil, 344
 Oil of, 383
 Salsa of, 845
 Verbena, 708
Lena exulenta, 521
LENTIBULACEAE, 698, 833
Lentil 521
Lentulus stypticus 860
 suavisolens, 860
Lentulus stypticus, 632
Leontice, 86
 Leontopodium, 830
Leonurus, 710, 711-712
 Cardiaca, 131, 711-712, 828
Lepidogathle Wightiana, 804
Lepidium, 52, 487 493-494
 apetalum, 113, 487, f. 494
 494
 campestre, 493
 Diaba, 818
 latifolium, 818
 oleraceum, 818
 Oenanthifolium, 818
 pachylobum, 8 8
 sativum 31, 113 818
 virginicum, 487 494
Lepidoptera, Fungi upon 273
Lepiota, 243 865
 Morganii, 49, 243, 860
 novius, 240
 protera, 235, 243
 Smooth 240
 Tadpole, 243
Leprosy, 180, 181
Leprosy, 89
Leprosy, 733, 756
Leptanthus luteus, f. 308
Leptanthus luteus, 286
Lespedeza striata, 525
Lentulus, 533, 534
 annularis, 831
Lettuce, 756, 760-762
 B. 761 762
 Cultivated, 756, 762
 European Prickly, 756
 Garden, 760
 Prickly, 137, 760-761, f. 761
 Wild, 761 762
Lewand, 1 52
 glauc, 831
Leuca 277
Lepentidii 168
Leucocoryne, 834
Leucocoryne, 104
 montanum, 104, 834
Leucocoryne, 179
Leucocoryne, 169, 179
Leuca in aestiva, 805
 verum, 805
Leucodii 266
Leucodii, 10, 861
LEUCOSPORA, 235
Leucospora, 667, 672-673
 Catesbaei, 127, f. 672, 673,
 821
 racemosa, 128, 673, f. 673,
 821
 Swamp, 673, f. 673
Leukemia, Infectious, 177
Leuterer, Dr., 896
Levadit, A., 894, 896
Levati Wormseed, Oil of 791
Levi, G., 896
Leviticum officinale 648
Lewandowski, F., 896

- Lewin, L. 637, 896
 Lewis, L. L. 9
Lewisia rediviva, 423, 844
Liatris 90, 737, 773-774
 punctata, 733, 774
 psychostachya, 773
 spicata, 733, 773
 squarrosa, 417
 Lignatard, A. 377, 383, 896
Lilania hypoleuca, 847
LICHENS, 54, 307, 861
 ASCO, 307
 BASIDIO, 307
Lichens, f. 307, 308, 309, 865
 Barbical, 307
 Crustaceous, 307
 Foliaceous, 307, f. 309
 Fruticose, 307
 Reindeer, 307
Lichium 199
Lichenostema Beihou, 850
pyrenifolia 856
LICORICE, 531
 European, 527, f. 528
 Extract of, 527
 Russian, 527
 Wild, 531
 Lichu II V 503
 Light, its effect upon toxicity
 of plants, 83
Lignum-vitæ, 573
 Ligustr. 896
LIGULIFLORAE, 757, 758, 762
Ligusticum canadense, 649
Ligustrum 683-684
 robustum, 683, 684
 vilgare, 120, 684, 840
Lilac, 683
 Powdery mildew on, 267, 269
LILIACEAE, 52, 63, 89, 103, 104, 374, 375, 386, 417, 835-836
 Lilienfeld, Fr., 896
LILIIFLORAE, 155, 374, 390
Lilium superbum, 184
 virginicum, 375
Lily, 375
 Atamano, 105, 387-388, f. 389
 Blackberry, 105
 California, 375
 Day, 375
 Family, 175-185
 Mariposa, 375
 Sacred, 444
 Texas, 690
 Tiger, 375
 Turk's-cap, 104
 Water, 445
 Blue Bowered, 445
 Yellow, f. 445
 Lily-of-the-Valley, 64, 184, 148, 375, 384-385, f. 384
Lima Bean, 75, 87, 119, 520, 574
Lime, 582
 Limettin, 583
Limnium, 824
Limnium, 486
Limnium carolinianum, 128
Limnoria, 89
 aquatica, 852
 cymbalaria, 852
 elatina, 852
 spuria, 852
LYNACEAE, 53, 130, 575, 577, 580-58, 835-836
Linacal 713
Linaleol, 709
Linamarin, 120, 865
Linaria, 734, 735
 vilgata, 735, f. 735, 738, 865
Linaria, 735
Lindeberg, A., 896
Lindley, J. 5, 362, 385, 531, 738, 896
Lindau, Gumara, 381, 302
Lindera Benzoin, 828
Lindera cultrata, 825
Lindsay, W. L., 896
Lingon Berry, 665
Linn, 120
Linn Tree, 621
Linnaea, See Linne, C. v.
Linne, Carl v., 277, 642, 643
Linnaea, C., 302, 897, 908
Linoleic acid, 581
Linonay, 581
Linseed, 580, 621
 Cake, 54
 Linseed Oil, 581
Linum, 580-581
 catarticum, 120, 835
 grandiflorum, 580
 perenne, 580
 rigidum, 120, 581, 835
 terreum, 836
 mitissimum, 53, 120, 580, 581, 836
Linum, 584
Lipierre, 169
Lippia ciliolata, 708
 mexicana, 708
 rodifera, 708
 pseudo-lippia, 629
Lippia, 708
Liquidambar orientalis, 580
 styraciflua, 500, f. 301
Liriodendron, 474
Liriodendron chinense, 837
 californica, 474, 837
Lit. A., 897
Litorea borealis, 860
Litchi, 607
Lithospermum, 704
 arvense, 228
 officinale, 619
Lithospermum, 805
Litum, 307
Little, 897
Little Barley, 367, f. 368
Live Oak, 403
Live-for-ever, 501, 502, f. 502
Liverwort, f. 309, 310, f. 311
Lizard's Tail, 396
Llanos Teguana, 855
Lloyd, C. G., 863, 897
Lloyd, J. U., 96, 484, 894, 897
Lloyd, J. U. and C. J., 452, 454, 456, 460, 465, 468, 471, 897
LOBACEAE, 125, 628, 633-634
Lobelia, 752
Lobelia, 378, 752-753
 Blue, 136
 Bryonia, 752
 cardinalis, 136, 753, 812
 erinus, 753
 Family, 751-753
 fulgens, 812
 great, 753
 inflata, 72, 75, 136, 464, 752, 753, 812
 Kalmii, 136, 812
 Kalm's, 136
 micranthaefolia, 752, 812
 Pale Spiked, 136
 pratensis, 812
 purpurea, 752
 rhynchopetalum, 812
 siphilitica, 136, 753, 812
 spicata, 753, 812
 Spiked, 753
 Type, 812
 Worm, 812
LOBELIACEAE, 748, 751, 753
 "Labellae", 378
 Lobelia, 752
 Lobelia, 136
 Lobelia, 78, 752
 Lobelia, 752
 Lobolly Pine, 330
 Lockjaw, 174
 Lockhart, Dr., 25
 Loose acid, 565
 Loose Disease, 501
 Loose poisoning, 804
 Loose Weed, 6, 37, 39, 97, 533, 544, 562-570, 865
 Distribution of, 565, 570
 Loss from poisoning by, 565
 Purple, 363, 567-570, f. 568
 Stomach, 39, 119, 566, 567, 570, f. 568
 Wooly, 39, 562, 567, 568
Locostum, 37, 40, 80, 865
Locust, 526, 746
 Black, 52, 58, f. 57, 64, f. 66, 118, 527, f. 530, 539-560
 Bark of, 53
 Clammy, 119, 560
 Flowers, 64, f. 66
 Honey, 527
 Poisoning, 56-58
 Tree, 118, 527, 558, 559, 560
 of Jamaica, 523
 of West Indies, 527
Lodman, E. G., 897
Lodgepole Pine, 330
Loranthaceae, 843
Loew, O., 897
Logania Family, 129
LOGANIACEAE, 51, 89, 129, 147, 148, 683, 684-689, 836, 865
Logania, 687
Logwood, 528
Loli n, 103
Lolium arvense, 363
 festuacum, 364
 italicum, 363, 364
 Lamarckii, 826
 im. colum, 363
 multiflorum, 364
 perenne, 364, 826
 remotum, 364
 temulentum, 4, 72, 74, 103, 361-364, 826
 Ergot of, 364
Lolium, 362
Lombroso, C., 290
Lonchocarpus, 1, 52
 densiflorus, 831
 hybrida, 831
 latifolia, 831
 Nicot, 831
 racemosa, 831
 violacea, 52, 533, 831
Long-awned Poverty Grass, 352-353, f. 353
Long Smooth-fruited Poppy, 113
Long-leaved Pine, 329
Lonicer, A., 275
Lonicera, 744
 chrysantha, 812
 flava, 744
 fragrantissima, 744
 involucrata, 135, 744, 812
 japonica, 744, 812
 periclymenum, 744
 sempervirens, 744
 Standishii, 812

- Sullicanthus*, 744
atarica, 744, 812
comantella, 812
Xyostium, 812
Lophopetalum pallidum, 813
Lophophorin, 837
Lophophyton, 13
gaulthor, 14
LOTARITHACEAE, 406, 419, 836
Lotus, 534
Lotus, 681
Lotus, 681
Lotus, 54, 90, 865
Barb, 681
apibacua, 53, 534
umbralis, 534, 831
corniculatus, 534, 831
Egyptian, 445
Amurcolens, 90, 55.
Tree, 681
Lotus, 54, 534
London J. C., 587, 601, 612, 897
Louiseville, 134, 734, 738
Mountain, 134
Swamp, 738
Lovage, 648, 649
Low Hop Clover, 553
Lupinus, 332
Lupine, 550
Arabian, 897
Lucas, E., 897
Lucet, A., 200, 202, 203, 877, 897
Lugo-encephalica, 20
Epizootic, 25
Lycenia, 89
Hoplantia, 53, 128, 851
deliciosa, 851
gryphallia, 851
mammata, 128, 679, 851
multiflora, 85.
obovata, 679
ash, 681, 81
Ludwigia virginica, 440
Lurmen, Ch., 897
Luffa, 89
arkhangula, 89
egyptiaca, 750, 89
virgata, 819
Lumpy Jaw, 133
Lunaria amara, 583
philipponia, 849
Lungwort, 704
Lupat, 530
Lupinus, 531
Lupin Oil, 540
Lupine, 6, 118, 149, 524, 546, 550, 863
Blue, 118
European, 118
White, 547
Hairy, 546
Low, 550
Nehruka, 546, 550
Poisoning in Montana, 548, 550
Silvery, 550
Western, 118
Wild, 546
Lupinidin, 118, 540
Lupidin, 118, 531, 540
Lupinin II, 548
Lupinin III, 548
Lupinosis, 20, 37, 40-43, 80, 448-550, 862, 865
Lupinotoxin, 78, 548
Lupinus, 535, 546-550, 831, 865
albus, 118, 392, 375, 546, 548, 831
angustifolius, 546, 548, 831
argenteus, 118, 546, 550
argophyllus, 118
densiflorus, 831
Arsenax, 149, 548
holosericeus, 548
Leucophyllus, 118, 348, 550, 831
Rinzolius, 548
Isleus, 149, 531, 546, 548, 831
ornatus, 550
perennis, 118, 546
Plattensis, 546, 550
pugulus, 550
sericeus, 550
Lupolic acid, 106
Lupulin, 406
Lutembacher, 897, 882
Luteolin, 479, 736
Luzula, 374
Lychnidium, 439
Lychus, 63, 89, 436, 439, 865
chalcidonicus, 439, 813
diosca, 439, 813
Evening, 439
Pisciculus, 439, 140, 813
indica, 813
Meadow, 440
Scarlet, 439
Lycium, 717, 733
harbamm, 854
chinensis, 716
helioselinum, 133, 716, 733
Lycogala epidendron, 149, 160
LYCOPERIDACEAE, 99, 241, 247, 860
LYCOPERIDINACEAE, 241, 247
Lycopodium Borica, 99, 247, 860
cyathiforme, 247
crispuliforme, 246
gemmatum, 860
gigantum, 245, 860
latifolium, 860
sp., 246
Lycopodium, 89
erectum, 132, 713, 714
pimpinellifolium, 715
LYCOPODIACEAE, 816
LYCOPODIACEAE, 312
Lycopodium, 312
clavatum, 712, 836
Selago, 836
Lycorin, 386
Lycoris radiata, 895
sp., 386
Lycopodium, 757, 762
interio, 762
Lycopodium palmatum, 17
lymphadenia, 184
Lymphangitis, 171, 184, 865
lyngbya, 93, 188
arsenaria, 185
Lyonia, 667, 674-675
maritima, 138, 674-675, 675
Lyonia, A. B., 803, 804, 897
Lysipia atropurpurea, 734
Lysimachia Nummularia, 676, 844
Lysin, 10
LYTHRACEAE, 637, 836-837
Lythrum Salicaria, 638

M
Mahala, 681
McCallum, 25
McCartell, Dr., 25
McCarthy, Gerald, 21, 245, 897
McCord, Dr. E. S., 50
MacDougal, Dr. D. T., 386, 394, 395, 897
McFarland, Dr., 184
McFaydean, J., 857
McGrinnis, J. W., 897
McNease, C. M., 243, 897
McKelway, J. L., 897
MacMillan, Conway, 897
McMullen, Dr., 25
McNeil, Dr. J. H., 279, 546, 898, 902
MacOwen, 534, 552
MacQueen James, 898
McWilliams, J. W., 782
Madame de la Roche, 844
Macaranga, 823
Mace, 445
Macura, 410, 865
gamifera, 406, 414, 415
hincoria, 406
Macrocystis pyrifera, 194
Macradiploia, 286
Macrasporium cleadum, 804
Macrasporium Brassicae, 867, 865
Macrasporium, 125, 819, 865
Madder, 524, 742
Family, 135, 741-744
Madera Vine, 424
Madia glomerata, 138
Chil, 755
calida, 755
Madrone, 665
Madsen, B. A., 898, 899
Madsen Thorwald, 898
Madura, 184
Marrua angustifolia, 812
Massa indica, 839
lauroloba, 839
Magar, 623
Magnan, Dr., 793
Magnolia, L., 898
Magnolia, 111
acuminata, 474
Family, 111, 112, 149, 473, 475
grandiflora, 64, 331, 474, 475, 476, 837
Great-flowered, 473, 474
Large-leaved, 474
macrophylla, 474
obovata, 474
Purple, 474
tripetala, 474
virginiana, 474
Yulan, 474
Magnoliaceae, 89, 90, 117, 444, 446, 473, 475, 837
Magnolia, 474
Magnolia, 52, 63, 89, 607
Brazilian, 64, 607
glabrata, 850
pubescens, 850
Mahogany, 575
Mountain, 89, 503
Nigeria, 587
Mahonia Trailing, 112, 472
Marden, J. H., 4, 17, 96, 190, 533, 534, 588, 650, 76, 724, 729, 734, 753, 803, 804, 898
Marden-hair Fern, 10, 33, 418, 536, 537, 537
Maschiana, 11, 80
Mash, J. M., 898, 918
"Ma" trunk, 744
Maize, See Corn
Maisenic acid, 342
Maun, 411
Makoni, 898
Malabar Cardamon, 391
Malassena Purpur, 298
Madia-gemma of orange and lemon trees, 287
Male Shield Fern, 149, 730, 315
Extract of, 77
Malic acid, 144, 50, 505, 52
Mallotus phillyrifolia, 89, 823

- Mallow Dwarf, 624, f. 624
 False, 626
 Family 564, 622-627
 Indian, 124, 626-627
 Marsh, 623
 Poppy, 624
 Maunten 13
Muhlenbergia nitida, 807
Mulphigia oxyrocca, 837
 MALPIGHIACEAE, 575, 837
Morus moschata, 624
rotundifolia, 624, f. 624
 MALVACEAE, 124, 621, 622-627, 837
 MALVACEAE, 157, 621-627
Malvastrum, 626
coccineum, 564, 626
laevigata, 635, 811
Mimosa americana, 637, 827
 Mammy Apple, 627
 Man-of-the-earth, 699, f. 700, 701
 Manacim, 713
 Mauchinea Tree, f. 586, 587, 865
 Mandarin, 582
Mandragora, 713
purpurea, 834
officinaria, 715, 834
vernalis, 4
Mandragora, 713
 Mandrake, 88, 112, 460-471, f. 470, 713
 Mangrove Sida nt, 7
 Mango, Daughy, 301
 Narcotic, f. 292
Pinellia florae, 106
 Mangola, 423, 866
Mangifera indica, 604, 805
Mangifera glauca, 837
 Mango, 608
 Manguid, 107
 Mangrove, 627, 68, 638
 Mangrove, 638
 Mangrove, 638
 Manihot 32, 586, 587
Globose, 823
officinalis, 53, 303, 587, 823
 Manila Hemp, 391
 Man, J. D., 898
 Manna, 327, 683
 Man in Grass, 90
 Ergot on, f. 27
 Mannes, 721
 Mannetel, 681
 Mannheim, 808
 Mannie, 306
 Maple, Curly 615
 Family 613
 Hard, 615
 Red, 615
 Silver, 613
 Sugar, 615, f. 616
 Maqui Fruit, 623
Moronia arundinacea, 391, 587
 MARANTACEAE, 391
Marasmius oreoides, 53, 860
tenerimus, f. 832
spens, 860
 Marattia, 35
 MARATTIALES 315
 Marasmius 606
 Marchand L., 898
 Marchantia, 310
 polymorpha, f. 309, 311
Marcgravia umbellata, 618
 MARCORAYACEAE, 628
 Marcus, H. D., 898
 Marek, J. 13, 26, 892, 898
 Mare's-tail, 640, f. 777
 Marguerite 736
 Marle, 572
 Marygold, 756
 African, 755-7, 6
- Burr, 779
 Fred, f. 139, 140, 784-785, f. 785
 Flowers, 756
 Marsh, 75, 381, 448-449
 Pot, 756
 Marimbo, 791
 Mariposa Lily, 375
 Marjoram, 710
Asteris tatarica, 127, 817
 Marmalade, 679
 Marubon, 709
Marrubium vulgare, 709
Marsdenia Cundurango, 696, 809
 erecta, 809
 Marsh, C. D., 40, 93, 96, 467, 565, 566, 567, 569, 898
 Marsh C. W., 898
 Marsh Cream, 490
 Elder, 137, 763-764, 767
 Small-flowered, 763
 Matlow, 623
 Matgold, 25, 381, 448-449
 Skullcap, 131
 Vetchling, 572
 Marshall, C. E., 344
 Marshall, H. T., 42, 871, 898
 Marula, 315
quadrifolia, f. 317
 MARULIACEAE, 315
 Marten, 290
 Mascha, J. v., 241, 898
 Masole, P., 487, 898
 Mason, W. P., 8
 Massachusetts, Poisonous plants of 864
 Report of State Board of Health of 95
 Massey, W. F., 898
 Mash, Bitter, 403
 Sweet, 403
 Mastie, 608
 Maté, 614, 747
 Matern, Medica, Literature of 866
 Matia, 396
 Matia de Peru, 396
Marrubium champanilla, 754
 Matrimony Vine, 133, 733
 Matru, 330
 Matruhot, L., 13, 15, 252, 879, 894
Matthiola incana, 486
 lida, 818
 Maurandia, 733
 Maurandia Hemp, 386
 Maurandia A., 899
 May Apple, 469-471, f. 470
 Maybole, Record of injury from polluted water in 9
 Mayday Tree, 504, f. 509
 Mayday, Sea Maidman
 Mayo, N. S., 215, 259, 293, 565, 899
 Maypop, 633
 Mayweed, 140, 756, 787-788
 Maximilian's Sunflower f. 767
 Mazza, M. G. de la, 899
 Mead, Richard, J., 899
 Meadow Anemone, 455
 Lychnis, f. 449
 Rue, 88, 111
 Saffron, 148
 Sweet, 629
 Meadow-out Grass, Tall, f. 340
 Mease, James, 277
 Meat poisoning, 172, 174
 Mechanical injuries, 64, 66, 804, 866
 Meconic acid, 144
- Meconium, 481
 Medea, 2
 Medea 89
 virginiana, 104
 virginica, 834
 Medicago f. 515, 522-533
fecrata, 526-532
hispida, 552
inulina, 553
platycarpa, 526
 in America, 526
sericea, 118, 529, 546, 553
 Med. and Jurisprudence, 866
 Medicinal plants among le-gumes, 527-535
 Distribution of 866
 Medick, 552
 Black, 553
 Mee, George, 899
 Meehan, 679
 Magna 14
 Meibthau, 272
 Meiner, 899
 Meiner, 300
Melaleuca leucodendron, 639, 839
Melampyrum belulina, f. 821
populina, 222
 MELASTOMACEAE, 98-99, 221, 222
Melampyrum arvense, 714, 832
lineare, 114
nitaticum, 134
 Melandryum, 89
Melanophthalmum, 210
 MELIANTHACEAE, 235
 MELANTHACEAE, 879
 Melanthin, 52
Melanthium, 376, 377, 379-380
corbinense, 834
latifolium, 380
parviflorum, 380
virginicum, 103, 379-380, f. 880, 834
 MELASTOMACEAE, 53, 640, 837
 Melia Asadarach, 121, 875, 837
 MELIACEAE, 52, 121, 575, 837
 MELIANTHACEAE, 89, 834
 Melanchoma comosa, 838
 major 838
 Melica, 826
Melicope erythronera, 849
 Melilotus, 90, 523, 533, 551, 552
 alba, 831, 118, f. 551, 552
 indica, 552, 831
 officinalis 118, 532, 83
Melissa officinalis, 7, 0
 Melodinus 52
 monogynus, 807
 Melon, Nutmeg, 750
 Sugar, 750
Melothria acrobiculata, 8, 9
 Meltzer, S. J., 899
 Melvin, A. D., 293, 294, 899, 900
 Memecylon, 53
 Menabes venenata, 809, 866
 Mendis, 575
 Menes, 2
 Meningitis, 169, 171, 172, 866
 Cerebro-spinal, 260
 Epidemic, 169
 Epizootic, 20, 259
 Infectious, 21
 Spinal, 291
 Meningo-encephalitis, 866
 MENISPERMACEAE, 52, 89, 112, 444, 472-473, 838
 Arrow poisons among, 1, 52
 Menispermum, 112, 149, 473

- Menispermum canadense*, 112, 472-473, f. 473, 838
coccineus, 149
Menispermum, 112, 473
Mentha, 78, 710, 713
arvensis v. *piperascens*, 709
arvensis v. *canadensis*, 713
crispa, 713
piperita, 709, 713, f. 713
Psiegium, 709
spicata, 709
Menthol, 709
Menisella, 125, 633-634
 Family, 633-634
gromotifolia, 633
oligasperma, 633, 634
ornata, 633-634
ovata, 125
showy, 633-634, f. 634
Menyanthes, 690-691
trifoliata, 129, 690-691, f. 691, 823
Menyanthin, 129, 69.
Morhous, Oil of 6
Merck, E. 899
Merck's Report, 636
Mercurial, n. 83
Mercurialis, 590
anua, 83, 803 f. 803, 823
perennis, 603, 823
tomentosa, 823
Mercury, Annual, 603 f. 603
 Binifolia of 6
 Three-needled, 603-604
Merleourt, 572
Merriampedia, 184, 188
Mermaid-weed, 640
Merrill, E. D., 52, 899
Mertensia alberta, 704
virginica, 704
Mertensia lachrymans, f. 704, 234
Mesca bean 637
 Burton, See Mesquit bean
Mesca n 637
Messerschmiedium anatum-
um, 824
 Crystallinum 434, 824
Mesquit Tree, 120, 527, 866
Metabolism, Products of 82,
 548
Metaplexis Stantoni, 809
Meta-phosphoric acid, 124
Metchnikoff, E., 249
Methyl xanthin, 146
Methylxanthin, 126, 851
METACHLAMYDEAE, 157
Methaemoglobin, 72
Methoxy, 147
Methylamin, 10, 149
Methylostin, 396
Metricis, 179
Metroxylon Ruppin, 370
Mexican Mulberry, 131
 Prickly Poppy, 113, 483
 Tea, 107, 629
Mexico, Economic plants of,
 864
Meyer, 203
Meyer, A., 899
Meyer H. 888, 899
Mexerein, 642
Mexerion, 149
Mexerium, 642
 Family, 125, 642-643
Michael, Helen A., 375
Michael, L. G. 424, 885, 889,
 894, 895, 898, 899, 907
Michelia Champaca, 837
laurata, 837
mongolica, 474
paludosa, 837
Michener C. B., 259
Micrandus, 587
Microcystis, 185
Microstema punctata, 710
Microsorum caprinus, 148
calarrhala, 170
gonorrhoea, 170
lanceolatus, 169-170
meningitidis, 169
pyogenes, 171
pyogenes v. *albus*, 167
pyogenes v. *aureus*, 167
tetragenus, 170
MICROSPERMAL, 155, 392
Microspira Alm, 267, 369
Microspira comma, 181-182
Microspora, 13, 299, 302
adonisi, 301, 382
canis, 299
equi, 299
Farfar, 298
Higley, 299
Microsporidia, 14
Microthyrus microscopium,
 f. 269
Mignone, 479, 734
Migula, W., 161, 163
Mildew, Downy, 98
 On Bean, 208
 Clover 205, 206
 Grape, 205
 Lettuce, 205
 Millet, 205
 Onion, 205
 Sunflower, 205
 Powdery
 On Cherry, f. 269
 Grape, 269
 Grasses, f. 269, 268
 269-272, f. 271
 Hop, f. 269
 Legumes, 267
 Lilac, 267
Mitell, Walter, 640
Milium, 102
effusum, 826
Milk Plant, 589
Vetch, 562, 567
Milk-veep caused by *Rapa-*
torium, 138
Mitchell, 771, 772
Milkweed, 130, 386, 598, 606,
 697
 Family, 130, 605-607
 Narrow-leaved, 130
 Showy 130, 606-607 f. 607
 Swamp, 130
Milkwort, 385-386
 Family, 384-386
Milner S. T., 546
Milner W. H., 454
Millet, 102, 350-351, 866
 Brown Corn, 350
 Korean Foxtail, f. 349
 German, 102 f. 349, 350
 Italian, 350
 Koda, 349
 Poisoning, 350-351
 Small, f. 349, f. 218
Millier, I., 52, 89
cafra, 832
feruginea, 832
pachycarpa, 832
rostrata, 832
sericea, 832
splendens, 832
Millspaugh, C. F. 96, 134,
 411, 419, 429, 433, 440,
 449, 454, 459, 471, 484,
 489, 509, 552, 568, 586,
 602, 617, 632, 655, 659,
 661, 671, 728, 739, 747,
 752, 762, 790, 793, 899
Milner, Dr. 544
Mimosa, 89
MIMOSEAE, 530
Mimosa lutea, 733
moschata, 733
Mimosa Kanki, 68.
Mimosa, poisons, Testing for,
 3
Mint, 713
 Family, 131, 708-713
 Horse, 709, f. 709
Minneer, J. A., 653
Mio Mio, 755
Miquel, Y. A. W., 214, 247,
 287, 803, 804, 862, 899
Mirbela racemosa, 533
Missouri Currant, 498, f. 500
Gooseberry, 500
Gourd, 751
Hickory, 401
Missouri Bottom Disease, 118,
 546
Mistletoe, 866
 European, 106
 False, 106
 Southern, 415, f. 415
Mitcham Oil, 86
Mitchell A., 124, 621
Mitchell S. R., 899
Mitchella, 89
repens, 742, 848
Mitcher, W., 863, 899
Mitracarpum, 90
Mitragnya speciosa, 848
Mium, 310
horum, f. 311
Moccasin Flower, f. 301
Showy, 105
Mocha Coffee, 744
Mock Orange, 300
Modesto, 89
palmaris, 841
trilobata, 841
venenosus, 841
Moel, J. 899
Moeller, Josef, 900, 917
Moeller, J. E., 900
Mohler, J. R., 26, 28, 251, 263,
 265, 283, 384, 291, 294,
 823, 899, 900
Mohr Ch. 900
Molnia carolina, 826
Mille Ph, 900
Mollugo hirta, 824
Polypodium cirratum
 856
Moltana Cattle-disease, 795
Momordica Balanina, 819
cheranta, 819
cochinchinensis, 819
cymbalaria, 819
Momordica citrifolia, 710
didyma, 629, 709
Articulata, 709, f. 709
purpurea, 709, 710
MONASCACRAE, 247, 248
Monascus, 24
heterosporus, 247
purpureus, 247 f. 248
Monilia candida, 302
MONIMIACEAE, 838
Monkey Flower, 733
Monkey-bread Tree, 621
Monkey-pot Tree, 619
Monkhood Wild, 109, 451
Minnid, 89, 843
MONOCOTYLEDONEAE,
 102, 155
Monospora bicuspidata, 249
Monotropa uniflora, 65, 838,
 842
MONOTROPACEAE, 838
Monotropa delioides, 37,
 pertusa, 808
Montana, Economic plants
 of, 864
Montano floribunda, 816
tomentosa, 816
Montgomery, E. G., 342

Myrica, 575, 637-645

Cape, 637
Spurge, 121, 599
Myrica communis, 640
Ugni, 629

MYXOBACTERYALES, 166-167

MYXOGASTERES, 160

MYXOMYCETES, 167

MYXOTHALLOPHYA, 153

150-160

N

Nabias, De, 901 881

Nandina domestica, 53 810

Napoleone Whitfieldii, 839

Nara, 750

Nararia, 582

Narcissus, 146-181

Arcissus, 104, 386

Jonquilla, 386

poeticus, 104, 386, 805

Poet's, 386

Pseudo-Narcissus, 4, 386

805

Tazetta, Jno, 803

Narcosis, 123

Narcotic, 804

Narcotic poisoning, 73, 78

Narcotics, 73

Narcotin, 3, 112, 146, 147

481 482

Nargamia alata, 817

Naringin, 883

Narrow-leaved Milkweed, 138

Succoweed, 783-784

Northam ostryagum, 835

Nosartium, 120, 171

Family, 120

Garden, 375, 1 377

officinalis, 379

palustris, 490

Native Plant, 614

Nauyn, H, 873

Naucaur, 804

Near, H V, 679, 901

Nebraska Lupine, 546, 550

Neckweed, 735 736

Neomorpho, 287

Neotrichia, 179

Necrosis, 179, 265

Nectandra Rodiaei, 478

Nectar Poisonous, 64

Nectria, 273

Necbe C H, 296

Necle Grass, 66, 102, 355-357

Western, 1, 70, 102, 355, 1

345

Negundo acerolites, 615, 629

Nemser, 188

Nekoe, 52

Nekoid, 52, 53

Nelson, Aven, 138, 332, 367,

376, 901, 918

Nelson, G. B, 96, 97, 109,

378, 462, 795, 901

Nelumbo, American, 108

Nera, 108, 444

nucifera, 444

Nematostylis multifidum, 1 309

"Nenta" Disease, 501 511

"Nenta Lemertii" Disease, 37

Nectria Nidulata, 840

Necossia werneri, 219

Nepenthes gracilis, 819

villora, 1 406

Nepenthes, 497

Nepria, 710

cataria, 710

Nederacen, 131, 710, 1 711

Nephelium, 89, 607

Nephelium, 890

Nishi, 607

Longana, 850

Nephelia, 613

Suppurative, 172

Nephrolepis aculeata, 1 314

Nerianthus, 130

Nerium, 130, 148

Nerium, 4 9, 693-695

odorum, 691, 807

alexander, 4, 65, 75, 129, 130,

691, 1 102, 694-695, 807

Nervin, 10, 78

Nervous system, Poisons af-

fecting, 72

Nesaea crystallata, 837

Nestegg Guard, 750

Neuber, A., 364, 395, 901

Netherlands, Poisonous plants

of the, 867

Nettle, 76, 77, 405, 594

Red, 72, 122, 196, 224-225

Common, 594

Dead, 131, 717 1 717

Family, 489-495

Handling of, 78

Slender, 106

Sum. Scinging, 1 412, 432-

413

Spurge, 122, 1 301, 396, 1

406

Stinging, 106, 413

Western, 106

Wood, 77, 106, 414

Nenmann, L. G., 13, 14, 36,

198, 981

Nenmayer, J., 249

Nerudin, 149, 150

Neruin, 150, 237

New England Amer, 276

New Jersey Tea, 124, 621, 629

Nephrolepis laeta, 739

Nerander, 2

Nerandra, 717 726

phylloides, 131 726, 854

Nichols, H. J., 905, 904

Nicholson, 1 4 13, 4

Nidemia diversifolia, 836

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Nidemia, 728

Spreading, 1 721

Three-flowered, 1 722 722

Yellow-flowered, 692

Nine-bark, 504

Nipa fruticans, 370

Nissolia tenuicarpa, 832

Nitraria indentata, 858

Nitric acid, 78

Nitrobenzene, 72

Nitrogen in alkaloids, 144,

145

Nitrous acid, 73

N. account, 901 916

Nicard, Ed., 184

Nickols, C., 901

Nodularia, 97

spumigera, 91

Noel, Ch., 895, 901

Noel, R., 901

Noel, P., 901

N. 858

Nopalea cochenillifera, 635

Norfolk Pine, 127

Norward, V. A., 293

Northern Plum, Wild, 505

Norward, 787

N. or 144, 146

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

N. 97

Oak, 846

Bark, 403

Bark, 403

Chestnut, 403

Cow, 403

English, 403

- European, 103
Evergreen, 403
Gal's, 403
Live, 403
Pin, 403
Poison, 609, 610, f. 611, 612
Red, 403, f. 404
Scarlet, 403
White, 403
- Oat, Rust 99
Smut, 215-217, f. 216
- Oats, 218, 223, 338-339, 538
Black, 66
Common, 102, 359, f. 360
Cultivated, 359
Rust of, 223
Smut, 98, 215-217, f. 216, 218
Wild, 102, 215, f. 358, 359
- Oblong, 560
- Ochne, 839
- OCNACTAZ, 839
- Ochsaner A. J., 183
- Ochrolechia tartarea, f. 308
- Ochroma Lagopus, 621
- Ochroia, 691
Maurel, 807
- OCROSPORACE, 233
- Ocimum Basilicum, 709
Thymus, 713
viride, 828
- Ocotilla Wax, 637
- Odora, Flaky 93
From decomposition, 93
From poisonous flowers, 64, 154
Oily 93
Pig-nen, 98
- Oedema, 177, 866
of glottis, 78
of lungs, 78
malignant, 175
- Oenanthe crocata, 72, 848, 849, 856, 866
Astrucosa, 4, 856
Lachrymilla, 856
Phellandrium, 648, 649, 857
- Oenantheoxia, 648
- Oenothera, 644
- Oenothera, 718, 914
- Oenothera, 642
- Oenothera, 503
- O'Garra, P. J., 96, 107, 124, 432, 433, 530, 563, 593
- Ohio Buckeye 123, 617, f. 619
- Odontomyces, 306
- Odium, 100, 134, 249, 300, 304
albicans 13, 100, 302, 303
Furfur, 298
granulomatigenes 303
hominis, 303-305
lactis f. 7, 200
lithogenes, 305
minimum, 298
moniliforme, 272
tonicicans, 301
- Oil, 866
Almond, 6, 315, 608
Anise, 648
Anthrax, 789
Atter of Roses, 86
Avocado, 478
Bayberry, 639
Bergamot, 86, 583
Bitter Almond, 6, 315, 303
Rome, 145
Bonset, 770
Buttertree, 688
Camphorin, 478
Carapa, 575
Caraway, 648
Cashew, 608, 864
Cassia, 86, 523
- Castor, 52, 56, 594, 595, 864, 866
- Cedar, 181, 338
- Chamomile, 751
- Chenop, 640
- Chinese Wood, 587
- Cinnamon, 86
- Citron, 583
- Citronella, 345
- Clove, 93, 639
- Cocua, 627
- Corn, 342
- Cresson, 330
- Croton, 6, 72, 78, 149, 588, 589, 866
- Cumin, 648
- Curcas, 78
- Dal, 648
- Eucalyptus, 640
- Eupatorium, 778
- Fennel, 86
- Gaultheria, 663, 866
- Geranium, 586
- Hedeoma, 711
- Hops, 406
- Horseweed, 778
- Jasmine, 86
- Juniper, 181
- Kapok, 624
- Lavender, 86, 709
- Lemon, 583
- Lemon Grass, 344
- Linseed, 581
- Lupin, 548
- Madia, 755
- Marcana, 606
- Marbana, 86
- Mitcham, 86
- Mustard, 7, 77, 114, 489, 373
- Myrica, 640
- Myrrh, 575
- Niger, 755
- Nutmeg, 445
- Olive, 683
- Orange, 86
- Palm, 370
- Palmarosa, 344
- Peanut, 521
- Pennyroyal, 131
- Peppermint, 86, 87, 93, 709, 866
- Pine needles, 774
- Pinboen, 587
- Poley, 78
- Poppy, 479
- Ranunculus, 459
- Rhodium, 690
- Rose, 503, 586
- Rosemary, 86
- Sassafras, 86
- Savin, 149, 866
- Sesame, 698
- Sunflower, 756
- Sweet Birch, 86
- Tanacetum, 789
- Tany, 756, 789-790, 866
- Tolene, 528
- Turpentine, 329, 330, 608
- Verrain, 860
- Wintergreen, 484, 621, 665
- Wormseed, 107, 428
- Lavant, 791
- Wormwood, 793
- Okra, 623
- OLACAE, 53, 839
- Old Man Cactus, 635
- Oleaburgaria arborescens, 816
- Oleandria senegalensis, 849
acubellata, 849
- Olea dioica, 840
- europaea, 683
- glandulifera, 683
- OLACEAE, 683-684, 840
- Oleander, 73, f. 180, 130, 148, 691, f. 604, 694-695, 866
- Oleandrin, 148, 866
- Olearia argophylla, 755
macrodonata, 816
moschata, 816
- Oleaster Family, 125, 640-642
Russian, 641
- Olea, 683
- Oleocera of Male Fern, 419
- Oleum, 654
- Olive, E. W., 222, 248
- Olive, 683
Family, 128, 683, 684
Oil, 608, 683
Wild, 641
- Olive, F. F., 894, 901
- Omphalocarpus, 89
procerum, 63, 681, 851
- Omphalophylla, 179
- ORACEAE, 643-647, 840
- Oreocarpus ediculus, 805
- Orion, 150, 375, 383
Poisoning, 384
Wild, 104
- Oreocera, 318, 320-321
sensibilla, 320
Struthocera, 321
- Ornoid, 531
- Ornith, 531
- Oreocera spinosa, 331
- Ornithodora Arachnoid, 816
- ORMYCTES, 134, 204-209
- Ornith, 13, 294
parvifolia, 17, f. 294
- Ophelia acid, 689
- Ophiodon gummifera, 841
- Ornithodora, 313, 692
- Ophiodon gummifera, 313
- Ophiodon, 827
- Ophithia inia Jocularis, 38
- Ophith, 308
- Opium, 2, 6, 7, 72, 84, 112, 144, 145, 393, 479, 481, 482, 866
Alkaloids of, 145, 146, 481
Consumption of, 39
Ind an, 83
Poisoning by, 6, 10, 77, 78
Poppy, 112
Properties of, 482
Smoking, 83
Statistics of deaths from, 39
Toxicology of, 482
Turkey, 481
- Opioid, 602
- Opium, 165, 177, 178, 182
- Opuntia, 125, 635, 636, 637
Dillenii, 635
Englemanni, f. 496
Ficus indica, 635
Kermanshahia, 637
Lacryl, 635
Perezhia, 635
polycantha, 635
Rafinesquii, 125, 635
Striptacantha, 635
Tuna, 635
vulgata, 635
- OPTUNIALES, 187, 634-637
- Opwyrda, R. J., 902, 906
- Orange, 582, f. 482, 866
Bater, 582
Falc, 866
Mock, 500
Oil, 86
Osage, 106, 406, 414-415
Root, 108
Sour, 582
- Orchard Grass, Powdery Mil
dew of, 272
Rust on, 279
Spot disease of, 282, f. 283
- Orchid Family, 392, 395

- [illegible]

- Phallia duplicatus*, 860
impudicus, 239, 245
Ravenscroft, 860
Phaven, Dr., 782
Pharmacognosy, 866
Pharmacographia, 866
Pharmacopoeia, 866
 American, 866
 British, 866, 874
 French, 866
 German, 866
 Italian, 866, 882
 Swedish, 866
Pharyngitis, 170
Phasium cuspidatum, f. 309
Phaseo-lanatin, 54
Phaseolus, 52, 535, 574, 866
aconitifolius, 832
angustatus, 526
innatus, 53, 54, 75, 87
 119, 520, 574, 747, 832
multiflorus, 119, 520, 574, 832
Mungo v. glaber, 520, 574
semipinnatus, 832
trilobus, 520
vulgaris, 520, 574, 832
Philandre, 649
Phellodendron amurense, 581
Phenol, 330
Phladelphus coronatus, 115, 500, 851
grandiflorus, 115, 500, 851
lemonis, 500, 851
Lewisii, 851
microphyllus, 115, 500, 857
Philippian, *Phanodora planis*
 of the, 867
Phillips, 904
Phyllanthus, 583
Phyllirea maritima, 840
Phyllirin, 683
Pseudodendron bipinnatifidum, 808
Andromedum, 808
Imbr., 808
Sinai, 808
Phlegmon, 171
Phleum pratense, Ergot on,
 276
Phlorizin, 405
Phosbus, Philipp. 4. 803.
 904
Phoenix, 90
dactylifera, 370, 840
Photinia maculata, 838
Phollos caperata, 860
radicans, 860
Phoradendron, 106, 415
latescens, 106, 415, f. 416,
 836
Phormium tenax, 375
Phosphorus, 72
 Poisoning by 6
Phosnik, 53
Phragmidium incrassatum,
 225, f. 225
Rubi, 221
subcorticum, 222
Phragmites, 27
 Ergot on, 276
communis, 279
Phrynus, 72
Phycomyces, 195, 197
nitens, 2
PHYCOMYCETES, 98, 154,
 195-209, 281, 859
Phycophacin, 194
Phyllachora, 279-280
graminis, 280, f. 280
trifolia, 279
Phyllanthus, 52, 59
Canna, 823
Emilia, 823
epiphyllanthus, 823
Gastromela, 823
lanceolata, 823
Nurmi, 823
pusillum, 823
uraria, 823
Phyllocactus, 625
Phycosyamin, 248
Physalis Alkekengi, 854
foetida, 854
virginiana, 854
Physarum, 160
Physcia palmulenta, f. 309
Phycosyamin, 184
Phyne Nut, Cuban, 596
Physocarpus opulifolius, 504
Physostigma, 866
tenaculum, 85, 148, 528,
 531, 852
Physostigma orientale, 854
pratense, 854
Physostigmin, 146, 148, 528,
 531, 866
 Poisoning by, 77, 78, 81
Phytolophus maritimus, 520
Phytobezoar, 866
 Caused by Common Oats,
 102, 359
 Crimson Clover, 68, 118,
 556, 567
Helichrysum, 745
Opuntia, 635-636
Phytolacca, 434-435, 866
abyssinica, 43, 434, 841
actinosa, 841
decandra, 167, 150, 433-435,
 f. 434
dioca, 434, 841
icosandra, 841
littoralis, 434
 Tincture of, 434, 435
PHYTOLACCA, 43, 89, 107,
 425, 434, 435-435, 841,
 842
Phytolacca acid, 435
Phytolacca, 107, 435
Phytolaccoside, 435
Phytophthora infestans, 205,
 208
Phaeoli, 208
Phytotoxin, 11
Picea canadensis, 328
Engelmanni, 327
excelsa, 328, 817
mariana, 328
nichensis, 327
Pichi, 216
Pick, F. 1, 295
Pickeringweed, 374
Piculin, 728
Picrocena excelsa, 853
Picrocena quasiusides, 853
Picro acid, 72
Picris hieracoides, 816
Picropodophyllin, 112, 470
Picrotin, 149
Picrotoxin, 1, 77, 80, 112,
 149, 435, 473, 475, 534,
 607
Picrotoxinin, 149
Pictet, A., 728, 878, 904, 917
Pictou Disease, 140, 795
Pie-plant, 170, 418
Pierre, H. C., 440
Pieris floribunda, 824
formosa, 65
marianna, 821
ovata, 821
ovalifolia, 65, 821
Piffard, H. C., 984
Pigeon Grass Sawal, 98
Pigeon Pea, 521
Pignat, 402
Pigweed, 427, 431-433, f. 433
 Common, f. 426, 427
 Green, 107
 Prostrate, 431, f. 433
 Slender, 432
 Thorny, 107
Pilea pumila, 409
Pilocarpus, 582
Pilocarpus, 77, 78, 80, 146,
 148, 582, 751, 752
Pilocarpus, 849
officinalis, 849
pennatifolius, 148, 582, 849
racemosus, 849
spinosus, 849
Pilocereis, 637
Pilocereis Sargentianus, 637
Pilobolus crystallinus, 195
Pix-aropsis, 338
Pilmerkblatt, 904
Pinella trichostachya, 642
Piment, 639
Pimenta acris, 839
officinalis, 639
Pimpernel, 128, 676
Pimpinella Anisum, 648
Saxifraga, 848, 857
Pimpinellin, 648
Pin Oak, 403
PINACEAE, 101
Pine, 101, 328, 329-330
 Austrian, 329
 Jack, 330
 Laidlaw, 330
 Lodgepole, 330
 Long-leaved, 329
 Norfolk, 327
 Scotch, 352
 Sugar, 330
 Western Yellow, 330
 White, 329, f. 330
Pin-apple, 373
 Family, 373
 W. J., 373
Pinus tabularia, 808
Pinus, 330, 640, 710
d-Pinus, 648
 "Pinex" Resin, 627
Pinguicula vulgaris, 698, 838
Pinguicula, 373
Pinhoen Oil, 587
Piper, 328
Pink Deptford, f. 437
 Family, 435-444
 Hardy, 436
 Indian, 129, 688
 Moss, 423
 Rose, 689
 Sweet, 250
Pink-root, 129, 688, 866
 East Tennessee, 688
Pinkham, 904
Pinnularia undulata, f. 189
Pinus, 101, 329-330
Banksiana, 330
contorta, 330
resinosa, 330
heterophylla, 330
Lambertiana, 329
Laricina, 329, 817
palustris, 329, 330
pinaster, 330
ponderosa, 330
ponderosa v. scopulorum,
 330
Strobus, 329, f. 330
sylvestris, 330, 817
Taeda, 330
Pinwood, 627
Piper, C. V., 526, 904
Piper, 52
angustifolium, 396
Bette, 396
Carpunya, 642

- Digitalis, 5
 Dirca, 5
 Drastic Resins, 5
 Emetic alkaloids, 5
 Hot acid, 5
 Hydrocyanic, 5
 Hypericum, 5
 Inebriant, 74, 804
 Inflammatory, 5
 Iota, 5
 Irritant, 5, 72, 76, 804
 Irritant narcotic, 5
 Iolium, 5
 Lycoperdon, 5
 Melica, 5
 Mineral, 2, 5, 866
 More volatile, 5
 Narcissus, 5
 Narcotic, 5, 72, 804
 Narcotic acid, 5
 Narcotic alkaloid, 5
 Nervous, 5
 Oleander, 5
 Pepper, 5
 Phallus, 5
 Plant, 2, 5
 Pokeweed, 5
 Poppy, 5
 Prunus, 5
 Rhum, 5
 Ruta, 5
 Seignic, 5
 Smartwood, 5
 Squatium, 5
 Stupifying, 5
 Unknown, 5
 Urticaria, 77, 804
 Vegetable, 5, 866
 Volatile narcotic, 5
 Sale of 867
 Pulmonum Lachap. 443
 Plantae novae Legumes,
 510-515
 Treatise upon, 864
 Poke, See Pokeweed
 Pokeweed, 424, 746
 Pokeweed, f. sty. 424, 435
 Family, 433-435, 721
 Root, 435
 Poland, Poisonous plants of,
 867
 Potamogeton, 495, 497
 gracilens, 114, 497
 trachysperma, 114
 Potamogeton boronit, 843
 caeruleum, 843
 Family, 130
 detum, 843
 gracile, 843
 humile, 843
 puriflorum, 843
 reptans, 843
 Poley Oil, 78
 Polemoniaceae, 89, 130, 698,
 843
 Polianthes tuberosa, 66, 186,
 835
 Pollinia, 767
 Pollen as cause of Hay-fever,
 747
 Polk, 184
 Polyalthea argentea, 477
 Polyanthus, 386
 Polycarpaea, 89, 813
 Polycythemia, 169
 Polydemon, 285
 callosa, 28, 100, 282-284
 Polygala, 89, 121, 585-586,
 774
 alba, 843
 amara, 843
 angulata, 843
 aspalathes, 843
 Boykin, 843
 Cyperus, 843
 glandulosa, 843
 serena, 843
 Senega, 63, 121 f. sty. 585-
 586, 617, 843
 verna, 586
 virginiana, 586
 Polygalaceae, 63, 89, 121,
 417, 575, 577, 584-586,
 843
 Polygalic acid, 62, 586
 Polygona, 100-102, 418,
 423, 843
 POLYGONALES, 154, 417,
 423
 Polygonatum, 89
 biflorum, 843
 japonicum, 177
 officinale, 843
 nuttallianum, 843
 POLYGONACEAE, 52
 Polygonic acid, 107
 Polygona, 52, 419, 421-423
 arce, 106, 421, 843
 articulata, 421, 423
 barbatum, 843
 Bistorta, 843
 C. ovalifolium, 421, 843
 flexuosum, 843
 Hydrocotyle, 107, 421, 422,
 423 f. sty. 843
 hydrocotylifolia, 422, 843
 Michxelioides, 421
 orientalis, 421
 pennsylvanicum, 421
 Persicaria, 421, f. 421
 sachalinensis, 421
 hectorum, 421
 POLYPTERIS, 395
 POTTOSACEAE, 100, 313, 315,
 321
 Polypodium, 89, 90, 315-316
 aureum, 316
 bicaudum, 316
 laciniatum, 325
 ligulatum, 325
 peruvianum, 325
 Phymatodes, 325
 scandens, 325
 suspensum, 325
 vulgare, f. 309, 311, 316,
 531, 825
 Polypody, Common, 311, 315,
 316
 POTTOSACEAE, 99, 234-235,
 861
 Polyporus anthelminticus, 861
 hypodas, 861
 igneus, f. 233
 perennis, f. 233
 squamosus, 861
 Sulphur, 234
 sulphureus, 234
 Polyscias, 868
 nodosa, 868
 Polystichum spleniosum, 825
 Polystichum versicolor, 234
 Polystichum rubrum, 273
 Polystichum, 279
 trifolium, 100, 861
 Polytichum, 310
 commune, f. 311
 POMACEAE, 301-306
 Pomegranate, 638
 Pomelia, 89
 punctata, 68
 Pomme Blanche, 538
 Pomme de Prairie, 538
 Pond Scum, f. 100
 Pondweed, 332 f. 177
 Pongamia Picta, 631
 Pontederia cordata, 374
 POTTOSACEAE, 374
 Poor-man's Weather-glass,
 128, 676
 Poplar, 397
 Balsam, 105
 Populus piscicarpa, 806
 Poppy, 90, 480-483
 Alkaloids of, 90, 147, 479-
 481
 California, 479, f. 480
 Common, f. 111, 479
 Corn, 112, 483
 Family, 479-483
 Garden, f. 90, 73, 111, f.
 481, 483
 Long Smooth-fruited, 113
 Mallow, 624
 Oil, 479
 Opium, 112
 Prickly, 113, 483
 Mexican, 112, 483
 Red, 479
 Popalm, 397
 Populus balsamifera, 103, 397,
 849
 canadensis, 397, f. 399
 deltoides, f. 396
 Porcher, f. P., 903
 Porcupine Grass, 102, 355-357,
 f. 399
 Smut, f. 208
 Porriga albertina, 16
 Porta, G., 3
 Porter, V. D., 187
 Porto Rico, Economic plants
 of, 864
 Portulaca grandiflora, 423
 oleracea, 423
 Portulacaceae, 423, 844
 Potamogeton, 90, 332, f. 333,
 334
 natans, f. 333
 Potassic chloride, 72
 cyanide, Poisoning by, 6,
 77
 nitrate, 85
 Potio, 332, 863
 Ait., 374
 Common, 710, 718, f. 718,
 724-725
 Japanese, 710
 Sweet, 698
 Wild, 701
 Potato-rot Fungus, f. 803,
 208
 Potentilla deturica, 88
 fruticosa, 88
 Poterium canadense, 847
 officinale, 847
 Potius cantoniensis, 867
 Pouchet, G. 893, 903
 Poverty Grass, Long-awned,
 352-353, f. 353
 Short-awned, f. 353
 Powell, R. D., 879, 905
 Power, F. H., 4, 34, 110, 217,
 453, 463, 470, 532, 546,
 565, 567, 575, 662, 684,
 903, 906
 Power, B. W., 540
 Prang, L. van, 902, 906
 Prain, Col. 490
 Prairie Larkspur, 460-461
 Theriacalis, 539
 Vitchling, 572
 Wake-robin, 385
 Prail, K., 802
 Prachil, W. W., 906
 Pratia erecta, 812
 Praxiteles, C., 906
 Precipitates, Red, 6
 White, 6
 Preis, 533
 Prenanthes alba, 417, 492,
 755

Pyruus, 145 146 728
Pyrois chazaracha, 842
zampira, 842
sumor, 842
rotundifolia, 842
 Pyrophosphoric acid, 24
 Pyroplasmic acid, 608
Pyronema conizans, Fertilization of, f. 247
Pyrus, 53, 506, 512, 848
americana, 504, 512
angustifolia, 512
Arca, 403
4-ncuparia, 113, 504, 506, 512, 629, 848
barlura, 512
aripitima, 113, 504, 512
cyronaria, 504 512
ylanta, 503 504, 512
tochna, 512, f. 573
japonica, 503, 504
lauria, 848
salva, 16, 504 505 / 512, 512
pinatifida, 503
evularia, 512
ambucifolia, 512
shenale, 512
nichensia, 512
Sorbus, 848
Pyruicaxia, 204
Pythium Deffaryennum, 204
proliferum, 204

Q

Quack Grass, 103, 364-365, f. 365
 Black-spot on, 280, f. 280
 Ergot on, 278
Qua n, E., 906
 Quaking Grass, 90
Quamaea, 378
 Quarter evil, 171
Quassia, 583
amra, 583
Quebrachia Balsamea, 608
Lorentali, 608
Quebrachia, 692
Quebrachinatim, 692
Quebrachia, 147, 692
 Queen of the Meadow, 113
Queen's Delight, 604
 Root, 604
Queenaland Hemp, 824
Quelot, 34
Quercifolia, 360
Quercifolia, 306, 310, 384, 846
Quercus, 103, 402 629, 846, 867
alba, 403 824
chrysolepis, 05
corcina, 403
lira, 403
lucida, 403 824
Muhlenbergii, 403
parvifolia, 403
Paradoxa, 403
Robur, 403
rubra, 103 403, f. 804
Tana, 824
virginiana, 403
 Quillagic acid, See Quillanic acid
 Quillanic acid, 2, 62, 505, 586
Quillaja, 89
 Bark, 72
Saponaria, 53, 505, 849
Quilaja-sapotoxin, 62
Quinacina, 742
Quince, 504
 Seeds, 503
Quacke, H., 293
 Quinic acid, 144

Quinofin, 145, 146, 147
Quinia, 77, 78, 87, 145, 147, 741, 742
Quinoa, 427
Quinquaria indica, 814

R

Rabak Frank 88
 Rabbit Septicemia, 177
 Rabunowitch, Lydia, 249
 Rabuteau, 904
Radicula, 490, 491
Armoracia, 113, 486, 490, 491 818
Rasturbum-aquaticum, 486
palustris v. hispida, 490
 Radish, 486
 Radikofler, L., 52, 803, 804.
 Rafinesque, C. S., 449, 906
 Ragged Robin, 439
 Rage, 198
 Ragweed, 137, 735, 764-767, f. 765
 Pale, f. 764
 Great, f. 137 765-766
 Pullen on cause of Hay Fever 137 767
 Small, f. 766, 766-767
 Tall, f. 766
 Ragwort, 793
 Purple, 736
 Raillot, 15
 Raleigh, Sir W., 714
 Rambach, W. T., 907
 Ramo-grass Cloth, 406
 Ramoelli, William, 771
 RANALIS, 154, 444-479
 Randal Dr., 277
 Randa, 89
aculeata, 849
sumatrensis, 849
 Ransom, 62
 Ransom, W. H., 907
 RANUNCULACEAE, 53, 81 89.
 108-111 417 444, 446, 468, 844-846
Ranunculus, 81, 110-111 447, 448, 452 461 867
abortivus, 111, 458-459 f. 459, 460
acris, 4, 77, 78, 110, 457 458, f. 459, 460, 846
alpestris, 4, 846
aquaticus, 846
aquaticus v. capillatus, 446
arvensis, 468, 846
asiaticus, 846
auricomus, 846
bulbosus, 4, 111 457 459, 846
lappaceus, 110
Ficaria, 446, 846
Pinnule, 468, 846
hybridus, 846
laugimorus, 846
lappaceus, 846
leptosepalus, 188
lingua, 846
 Oil of 459
polyanthemos, 846
repens, 4, 457
scleratus, 4, 77 78, 111 459-458, f. 458
septentrionalis, 110, 457 846
 Thoro, 4, 846
 Kape, 486, 867
 Fungus, 281-284
Raphanus, 867
raphanistrum, 277, 818
sativa, 486, 818

Raphia Fiber, 370
pedunculata, 370
ramifera, 370
 Raspberry, 507-508
 Black, 505 508
 Black-cap, 508
 Wild, f. 507 508
 Garden, 505
 Red, Wild, 505, 507, f. 507
 Rat Poison, 377
 Ratshane, 575
 Raven, 907
 Rattle weed, 867
 Rattlebox, 118, 543-546, f. 543, 564
 Rattlesnake Fern, 692
 Grass, 692
 Master, 692
 Plantain, 692
 Rattlesnake-root, 693, 733
 Rattlesnake-wood, 692 783
 Ratteburg, J. T. C., 4, 5, 803, 863, 874, 907
 Raum, J., 249
 Ravenna, 90
Rauwolfia serpentina, 807
trifoliata, 807
verticillata, 807
Ravenna, 391
 Ravenel, M. P. 23, 30, 263
 Ravenna, 747
 Rav 290
 Rawton, Oliver de, 907
 Read, Alex., 907
 Red Raspberry, 178, 467-468
 Buckeye, 123 617 f. 467
 Cedar, 101 102 330-331 f. 331, 332
 Clover, 279, 525, 553, f. 553
 Currant, 498
 Flax, Garden, 580
 Gum, 500 f. 501
 Hlaw 500, f. 514
 Maple, 615
 Oak, 403, f. 814
 Over 684
 Pepper 87 133 714, 743, 725-726
 Poppy 479
 Raspberry, Wild, 505, 507 f. 507
 Rust, 227
 Saweed, f. 794
 Top, 338
 Ergot on 277
 Vetch, 572
 Red-berried Elder 135
 Red-bud, 527
 Redmond, 771
 Red-root, 104, 385 432
 Redwood, 102
 Reed, H. S., 907
 Rees, 302
 Reese, J. J., 907
 Reinders Lichens, 307
 Reinders, 621
 Remak 13, 18, 293
Remia pedunculata, 742
 Remsen, Dr., 512
 Renner, Dr. 525, 907
 Reun, L. 263
 Reproduction of Bacteria, 161
 Reseda, 846
Inteola, 479, 846
odorata, 479
 Respaciae, 479, 486, 846
 Resin 319 330
 Botany Bay 375
 Chalk, 625
 Piney 627
 Retam, 531

Reynolds, C. F., 775

RHAMNACEAE, 157 620-621.

RHAMNACEAE, 53

RHAMNACEAE, 53, 89, 620-621, 846-847

Fish poisons among, 2, 621

Rhamnetin, 124, 620

Rhamnin, 124, 620

Rhamno-cathartin 124

Rhamnose, 612

Rhamnus Alaternus, 847

californica, 847

cuneata, 847

cathartica, 123, 124, 229, 620, 847

dohrniana, 620

Frangula, 53, 124, 229 630, 847

infectoria, 620

lanceolata, 124, 229

Purshiana, 621 847

stricta, 620

Wrightii, 847

Rhein, 418

Rheinnic acid, 419

Rheum Emodi, 843

hybridum, 843

officinale, 418, 843

palmatum, 843

Rhopanum, 418, 843

Rhopan, 640

Rhamnanthra, 90

communis, 692, 804

Rhnanthra, 738

Rhinanthus major, 852

minor, 852

Rhynchos, 617, 811

Rhizobium leguminosarum, f.

luz, f. 103

Rhizocarpon geographicum, f.

joh

Rhizoid of Botrydium graminum, f. 92

Rhizomorph, 202

Rhizophora Mangle, 818, 847

RHIZOPHYLLACEAE, 638, 847

Rhizophora 198, f. 258, 867

myrsina, 195, 196

Rhode Island Poisonous

plants of, 467

Rhodiol, 506

Rhodium Oil of, 699

Rhododendron, 668

Rhododendron, 54, 64, 222, 666, 667, 669, 867

arborescens, 821

arborescens, 821

Austrian, 867

barbatum, 867

Californica, 127, 667

californicum, 127, 667, 821

corymbosum, 821

catalpifolium, 81, 127, 667, 668, 821

concoloratum, 821

corymbosum, 81, 666, 821

corymbosum, 821

dauricum, 821

ferrugineum, 821

fulgens, 65

gracile, 65

Aristatum, 821

indicum, 821

japonicum, 666

Large 127

lucidatum, 821

maximilianii, 127, f. 466, 667, 668-669, 821

nuditatum, 821

occidentale, 127, f. 668, 668-669, 821

pennsylvanicum, 64, 668, 821

pinnatum, 821

swense, 821

sublancoletum, 821

RHODOMELACEAE, 879

Rhodomenia palustris, 859

RHODOPHYCEAE, 154, 194

RHODOSINOREAE, 235

RHODADALES, 157, 479-497

Rhoeadin, 481

Rhodoglyphus hercynicus, 848

Rhodoglyphus, 418, 419

Rhys, 79, 409, 608-614, 805, 806, 865, 867

causina, 806

chinensis, 806

Corymbosa, 608

Corymbosa, 608, 805

diversifolia, 123, 609, f.

611, 806

Auridana, 806

glabra, 608, 611-614, 805

Gracilis, 806

japonica, 806

littoralis, 806

lucida, 806

Metopium, 806

Michauxii, 806

peruviana, 806

radicans, 806

Rydbergii, 123, 806

scutellata, 806

stricta, 806

succedanea, 608, 806

sylvatica, 806

Toxicodendron, 4, f. 12, 77, 78, 122, 394 608-609, f.

409

venusta, 610, 806

vernifera, 608, 806

verna, 122, 609, f. 806

Wallichii, 806

Rhynchosia minima, 832

Ribes aureum, 53, 508, 851

cereum, 851

gracile, 500

gracilior, 508

macrocarpum, 851

nigrum, 498, 851

prostratum, 851

vulgare, 498

Ribbert, M. H. W., 907

Rice, W. S., 907

Rice Paper, 647

Rice, W. H., 276, 338

Rice-cut, 339

Rich, F. L., 101, 334, 325, 907

Richardia africana, 371

Richardson, 907

Richet, C., 907

Rich. 4, 11, 35, 72, 121, 194 626, 687, 867

Rhododendron africanum, 387

Ricinoleic acid glycerid, 594

Ricinolein, 594

Rucinus, 55, 590 594-595, 862, 864, 867

cinnamomum, 53, f. 25, 121, 594-595, 823

"Rickets," 323

Ricketts, H. T., 306

Ricord-Madonna, J. B., 907

Ridinger, 467

Riecke, V. A., 862, 872, 847, 907

Ringworm, 12, 14, 301, 302

of Body, 301

Dog, 12

Horse, 14

Also see Yinea

Scalp, 301

Ritchie, f. 305

Riviera humilis, 842

Rivolta, S., 263

Rivularia, 93

RHODARIACEAE, 186-187

Robbins, E. L., 424

Robbins, E. L., 899, 907

Roberts, G. H., 260

Robin, Ch., 298, 302, 395, 907

Robinson, 534, 535, 558-561, 867

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Roripha, 339

Wila 127, 673
 Rosenthal, D. A., 908
Rosmarinus officinalis, 709
 Roseal, 506
 Rosen, P., 908
 Rosinski, 159
 Rostrop, E., 290
 Rot Cabbage, 163
 Rosenbiller, von, 908
 Rothe, L., 908
 Rotachky, 728
 Rottlerin, 589
 Roeb, J. P., 908
 Rough Phacelia, 131
 Rouni Larumion, 391
 Round leaved Jugwood, 664
 Green Brier, 104
 Roap, 177
Rouppia Pahlia, 844
verrucosa, 844
Rouppia glabra, 817
 Roux, G., 304, 897, 908
 Row, 199
 Royal Fern, f. 300, 313, 322
 On m. 370
Royaltuna borinquena, 370
Rapier gongylifera, 235
 Rubber, Latex, 588
 Plants, 587, 691, 696
 Tree, f. 416
Rubia tinctorum, 742
 RUBIACEAE, 53, 89, 90, 135,
 144, 74, 744, 848-849
 RUBIACEAE, 138, 740-748
Rubi erin, 381
Rubus, 307, 508
chamaemorus, 848
cuneifolia, 505, 848
idaeus, 505
idaeus v. *aculeatus*,
 505, f. 307
occidentalis, 505, f. 307, 508
odoratus, 504
pariflorus, 505
phoenicifolius, 505
 Prickles and spines of, 507-
 508
villosus, 505, f. 304, 848
Rudbeckia, 757, 779
hirta, 779
laciniata, 139, 756, 779
occidentalis, 779
 Rüdel, C., 908
 Rue, 78, 524
 Common, 581
 Family, 120, 581-583
 Meadow, 111
 Rueh, Dr. C., 565
Ruellia elata, 688, 898
patula, 804
strepens, 804
tuberosa, 804
Rupis, 419, 419-421, 867
obovata, 843
4 elata, 150, 846
Areionella, 105, 420, f. 420,
 846
rhodantha, 49, 420-421
repens, 106, 419, f. 420,
 421, 846
Eckloniana, 846
Myriophyllum, 418, 846
obovata, 846
repens, 421
Patentia, 49
acutata, 49
 Romiein, 421
 Ruge, 145
 Rusby, H. H., 96, 101, 105,
 124, 121, 389, 448, 456,
 471, 484, 491, 160, 584,
 612, 633, 746, 793, 803,
 804, 908
 Ruschhaupt F. R., 908

Ruscus aculeatus, 835
 Rush, 374
 Common, 367 f. 373
 Scouring, 333
Russelia juncea, 734
 Russell, D., 775
 Russell, W., 908
 Russia, Poisonous plants of,
 867
 Russian Larice, 527
 Cresset, 641
 Thistle, 68, 150, 424, 430,
 f. 211, 433
 Turpentine, 330
Russula, 90, 343
Russula, 860
emeticus, 237, 243, 860
foetens, 53, 860
fragilis, 860
rubra, 860
 Rust, 20, 160, 220
 Black-stem, 227
 Clover, 28, 99, 230-231, f.
 231
 Corn, f. 230
 Crowned, f. 231, f. 235, f.
 238, 239
 Goldenrod, 222, 775
 Grass, f. 233, 224-227, f.
 235
 Maine, 230-230
 Ony, 99
 Red, 222
 Reproduction, 233
 Strawberry, 281
 Treating, 230
 Tickle Grass, f. 233
 Wheat, 99, 227 f. 238
 Covered, 227 f. 227, 228
 White, f. 230
Ruta graveolens, 74, 120, 581,
 849
montana, 849
 Rutabaga, 486
 RUTACEAE, 53, 89, 120, 148,
 575, 577, 581, 583, 849
 Rutin, 493
 Rue, 338
 Grass, 361, 364
 Common, 361
 Italian, 361
 Spout, f. 368
 Wild, 27, 90, 149, 546
 Black spot on, 280
 Ergot on, 276
Rhynchospora, 369
 Rynning, 662
 Ryan, A. M., 908
Ryparosa caesia, 53

S

Sabalina officinalis, 377
 Sabal, 381
Sabalina, 379, 381
Sabadina, 379, 381
 SABALACEAE, 849
 Sahaniyeff, N., 872, 908
Sabalina angulata, 689
 Sahing, 78
Sabalina florida, 833
 Sahli, 241
 Sahoutand, 13, 16, 301, 908
 Sabourne, 329
 Saccardo, P. A., 288
Saccharomyces, 100, 249
apiculatus, 250
cerevisiae, 250 f. 251
ellipsoides, 250, f. 251
glutinosus, 250
kefir, 250
mycodevina, f. 249, 250
 SACCCHAROMYCETES, 249-
 253

Sacred Bean, 444
 Lary, 444
 Safflower, 756
 Safford, W. E., 325-326, 908
 Saffron, f. 387, 389, 734, 756
 Caps, 754
 Meadow, 148
 South American, 389
Sagaretia theaeana, 629
 Sage, 710
 Lance-leaved, 710
 South American, 710
 White, 424
 Sage brush, 790, 792, f. 797
 Saghalen knotweed, 421
Sagittaria Engelmanniana,
 335, f. 336
latifolia, 102
 Sage, 370
 Poison, 378
 Wild, 378
 Sagot, M. P., 908
 St. Hualer, 908
 St. Ignatius's Bark, 688
 Bean, 74, 145
 St. John's-bread, f. 528, 527
 St. John's-wort, 124, 386,
 629-631
 Common, 125, 629-630, f.
 630
 Euroasian, 631
 Family, 629-631
 Green, 725, 629, 630-631,
 f. 631
 Kalm's, 629
 Shrubby, 629
 Spotted, 630
Sakalava, 867
 Sake, 281
 Sales, 392
 SALICACEAE, 105, 396-397, 849
 SALICACEAE, 156, 396-399
 Salicin, 397, 505
Salicornia, 150
 Salicylic acid, 505
 Salicoida, 967
 Salix, 629
amvgdalioides, f. 707
 Salmon, f. 10, 277
 Salmon Berry, 505
 Salpingit, 171
 Salubry, 756
 Salvia, 426, 430
Salvia tenuifolia, 424, 430,
 f. 431
temerariifolia, 814
 H. chin., 424, 814
 Salt Bush, 107, 423
 Australian, 424
 Grass, 90
 Groundsel, 138
 Salix, Caustic, 72
 Corrosive, 72
 Saltwort, 430
 Shrubby, 424
 Salvia, 756
amvgdalioides, 828
coccinea, 710
laevigata, 710
officinalis, 710, 828
pratensis, 828
splendens, 710
Salvadora peruviana, 849
 SALVADORACEAE, 849
 Salvina, 313
 Salvia, 756
 Salvia, 710, 756
 Samadera indica, 853
 Sambuca, 747
 Sambucus, 745-747, 867
canadensis, 133, 744, 745-
 747, 812
eburnea, 745, 747, 812
mexicana, 812

- Stroicotrachium graminis*, 282,
f. 283
Scopolamin, 60, 77 78 148,
7.6 733
Scopolina atropoides, 4
Scopula, 60, 716
Scopula cornuta, 254
 japonica, 854
 lurida, 854
 physioides, 854
Scopolin, 716
Scorsonera, 737
Scotch Broom, 118, 408, 551
 Heather, 65, 663
Scotium acuminatum, 860
Scott, C. J., 556
Scott, D. H., 910
Scott, J. L., 773, 910
Scouring Rush, 323
"Scourer," 30
Screenings containing cockle,
108, 439, 440
Screw Pine, 332
Scrivner, F. Lamson, 903
Scrophularia aquatica, 852
 mariandica, 134
 nodosa, 852
SCROPHULARIACEAE, 89, 134,
698, 733-738, 852-853
Scrophularin, 134
SCROPHULARINACEAE, 52
Scum, 93
Scutellaria galericulata, 131
 828
 integrifolia, 828
Sea Holly, 548
 Kale, 486
 Mosses, 91
 Weed, Red, f. 104
Sea Island Cotton, 623
Seaside Heliotrope, 705
Seasons, variation affecting
 toxicity of plants, 81
Sebastiania Palmieri, 823
 Purshiana, 823
Secale luxurians, 375
Secalin, 277, 278
Secalin-toxin, 149
Secalinotoxin, 277
Secotium edule, 730
Seear, Eugene, 634
Secotium erythrocephalum, f.
 246
Securidaca longepedunculata,
 843
Securigera Coronilla, 833
Seurinega, 52
 ramiflora, 823
Sedative, 804
Sedge, f. 71, 369
 Family, 367-368, 369
Sedum, 501, 503
 acre, 114, 301, 502, 503, 818
 purpureum, 501, 502, f. 502
 stanopetalum, f. 302
Seeds, 144, 862
 Aconite, 83
 Adzuki Bean, 520
 Aleurites, 887
 Anagyris, 530
 Anise, 648, 651
 Baptisia, 430
 Bassia, 681
 Beans, 867
 Cassia Bean, 528
 Caraway, 648
 Cassia, 535
 Castanospermum, 527
 Castor Oil, 149
 Cerbera, 692
 Cherry, 503
 Coffee, *Astragalus*, 527
 Coral Tree, 527
 Cockle, Corn, 439-441
 Corn, 108
 Colchicum, 867
 Coroso Palm, 378
 Corypha Palm, 370
 Cotton, 867
 Crab's-eye, 527
 Croton, 149
 Commia, 648
 Cynoglossum, 867
 Datura, 729
 Eutade, 525
 Eriobotrya, 867
 Fennel, 648
 Frangula, 527
 Flax, 120, 581
 Galaga, 535
 Garcia, 867
 Ginseng, 395
 Hura, 97
 Hyoisycamus, 726
 Jatropha, 587
 Inquility, 55, 527, 534
 Kapok, 421
 Larkspur, 867
 Leucimnora, 867
 Liriodendron, 581
 Lupinus, 867
 Lycalis, 867
 Mucuna, 524
 Mush, 624
 Nux vomica, 867
 Ocimum, 710
 Olive, 683
 Omphala, 867
 Oxytropis, 867
 Paprika, 523
 Penniset, 521
 Physostigma, 528, 531
 Poppy, 727
 Protopis, 55
 Psoralea, 535
 Quince, 583
 Rattlebox, 546
 Rape, 867
 Ricinus, 867
 Sabadillo, 867
 Sinapis, 867
SIMARUBACEAE, 867
Sintomere, 464
Strychnos, 683
Tamarind, 529
Trigonella, 581
Ulex, 530
Vetch, 120, 527
Vicia, 867
Water Hemlock, 631
Wormwood, 867
Segura, J. C., 910
Sellen, von, 298
SELAGINELLACEAE, 853
Selby, A. D., 138, 286, 701,
 910
Seligman, C. G., 910
Selmi, P. A., 910
Semecarpus Anacardium,
 608, 806
 heterophylla, 806
Sempervivum montanum, 818
Sempie, R. H., 910
Senebiera pinnatifida, 818
Senecio Snakeroot, 585-586,
 f. 585
Senecio, 755, 757, 758, 795,
 797
 aureus, 140
 Bolcomilae, 796
 Burchella, 795
 Cissaria, 756
 cruciatas, 756
 elegans, 756
 Forsteri, 798
 gnodolensis, 140, 816
 Grysmus, 816
 Jacobaea, 140, 795, 797
 816
 M
 latifolia, 795
 platensis, 140
 solacensis, 140
 vulgaris, 816
 vilneraria, 816
SENECIONEAE, 757
SENECIONIDEAE, 758
Senegia, 149, 586
Senf, E., 393
Senn, Nicholas, 183
Senna, 536
 Leaves, 528
 Substitute for, 334
 Wild, f. 577
Sennacrol, 528
Sennapiolin, 528
Sensative Fern, 320-321
Septentrionalis, 450
Septicemia, 167, 171, 867
 Generalized, 172, 177
 Goose, 182
 Hemorrhagic, 177
 hemorrhagica, 343
 House, 176
 Rabbit, 177
 Sceptacolea, 171
Septoria, 284-286
 Brown, f. 285, 285-286
 graminum, 284-286
 rubra, 284
 tritici, 286
Sequoia sempervirens, 102
Serjania, 52, 89, 606, 607
 acuminata, 850
 curassavica, 606, 850
 erecta, 850
 inbriata, 850
 ichthyostoma, 850
 lithalis, 606, 850
 mercedis, 850
 nodosa, 606, 850
 placatoria, 850
 polyphylla, 850
 trifoliata, 606
Serpentaria, 688
Scrivner, F. W. A., 143
Service Berry, 504
Seaside, 698
Sezanium indicum, 698
Seban, 561, 561
Sebania, 534, 535, 561
 aculeata, 524
 platycarpa, 581, f. 457
 vesicaria, 118, 833
Seaside-flowered Wake-robin
 385
Selaria, 339, 350
 germanica, 350
 glauca, f. 339
 italica, 102, f. 349, 350
 scandens, 826
 verticillata, f. 170
 viridis, f. 307, 350
Seymour, 910
Sexton, 662
Shaddock, 582
Shaggy-man Mushroom, 235
Shallan, 663
Shallot, 375, 383
Shallu, 345
Sharp, G., 911
Shaw, Dr., 560
Sheep-pox, 179
Sheep Sorrel, 106, 150, f. 420
Sheldon, J. L., 290
Shell-bark Hickory, 401
Shellac, 406
Shenstone, J. C., 911
Shepherdia argentea, f. 639,
 641

- момтолом, 854
Melangea, 714, 716, 854
 илгит, 4, 60, f. 62, 74,
 131 132, f. 132, 719-721,
 f. 720, 854
rostratum, 718, 722-723, f.
 723
 эфроксум 854
 лобангит, 854
Stramonifolium 854
Styracium 854
 урчиг, 854
tybactum, 3, 32, f. 721, 722,
 854
tuberosum 133, 714, 718,
 f. 718, 72, 855
villosum, 855
Xanthi, 855
Solidanella, 89
 Суидер'н Herb, 306
Solenostemon 494, 810
Solidago, 757, 774-776
canadensis 138, 774-775
 Rust on 222
Drummondii, 774
missouriensis, 774
missouri, 755-774
rigida, 773-776
serotina, Rust on, 222
speciosa, 774
virginica, 816
 Solomon's Seal 377
 Sommerfeld, T. 911
Synchilus, 717 719-760
arvensis 759-760
hieracium, 759
 Rust on, 222
Sophora, 535, 542-543, 868
alopocorymboides, 833
angustifolia, 530
lutea, 80
japonica, 833
malva 833
serotina, 90, 337, 534,
 542-543, 833
aplica, 90, 117, 542, f.
 543, 564, 833
 Milky 117, 542, f. 544
speciosa, 830
lomentosa, 90, 833
 SYPHOCARPEAE, 530
Euphorbia 530, 543
Synanthus, 804
horrida, 506
 SORDARIACEAE, 280
Yorgilium, 54, 102, 338, f.
 143, 345-348, 503, 747
 868
 Struck, f. 47
virgure, 53, 346, 826
Soroporum, 210
 Sorel, 379-380
 French, 410
 Jamaica, 623
 Mountain 4 9
 Sheep, 106, 150, f. 420
 Woad, 120
 Violet, 580
 Souan, Poisonous plants of,
 866
 Sour Cherry 505, 514
 Dock, 106, f. 420
 Orange, 582
 South American Saffron, 389
 Sage, 710
 Southern Buckthorn, 679
 Mistletoe & S. f. 416
 Sow Thistle, 759-760
 Annual, 759
 Field, 759-760
 Soy Bean, 520, 548
 Spanish Chestnut, 403
 Needle, 67 f. 780, 780-781
 Wormseed, 424
 Vetch, 577
Sparanium, 352, f. 334
Sparte n. 147, 551
Spartium cynosuroides, 1 60,
 198, 339
Spartium junceum, 826
spartium, 147
 SPATHIIFLORAE, 155, 370-
 372
Spathiphyllum canaliculatum, 808
 Spatier 633
 Spearwort 709
 Speedwell, 228, 229, 733
 Purslane, 735-736
Spergula arvensis, 436
 Spermaceae, 90
capitata 849
semirecta, 849
 SPERMATOPHYTES, 155,
 323-802
 Sphaerulic acid, 278, 379
Sphaerelia, 775
 Sphaerulic acid, 28
Sphaerulic, 710
Sphaerulic, 277
Sphaerelia Fragariae, 281
 SPHAERIACEAE, 267, f. 268,
 281
 SPHARACEAE, 280,
 281
Sphaerolobos Castagnei, f.
 No. 270
Sphaerolobos, 286
Sphaerolobos angulatus,
 817
 Spier-bush, 477
 Spider Polium, 7
 Spiderwort, 273
 Blue, 373
 Spier, A. J. T. 911
Spigelia, 498, 868
Anthelmia, 836
glabrata, 836
maritima 76, 129, 688,
 836
reticulata, 836
 Spigelia, 688
 Spiked Lobelia, 753
 Pale 136
 Spokenard, 647
Spilanthes Arnica, 816
oleacea, 816
 Spinach, 150, 424, 427, 435
 Australian, 427
 New Zealand, 423
Spinacia oleracea, 424
 Spinal cord, Poisons acting
 on, 74
 Symptoms of 74
 Treatment, 74
 Spiny Amaranth, 432, f. 422
 Clothier f. 107
Spinosa, 189, 304
Aruncus, 848
bella, 848
canescens, 848
Douglasii, 904
Filipendula, 848
Humboldtii, 848
hypericifolia, 848
japonica, 503, 504, 848
lar rigida, 848
palmaria, 848
salicifolia, 504
Thunbergii, 504
tomentosa, 848
Ulmora, 492
Spiranthes 392
 SPILLACEAE, 165, 182
Spirillum, 150
commis, 161
rubrum, 162
Sporobolus australis, 182
Obermayeri, 182
pallda, 182
Sporogynia, 92, 94, 95, 189, 190-
Splenwort, 318-320
 Common, 318
Spondias dulcis, 608
lutea, 608
mangifera, 806
purpurea, 608
 Sorage, Fresh-water, 93
 Cord, 750
Spora virgata, 53
Sporangium 160
 Spores, 160
Sporodinia, 199
Sporotrichosis, 868
Sporotrichum, 13, 15, 297,
 302
Purpure, 100, 297, f. 297
giganteum, 299
micrococcum f. 298, 298-
 299
longum, 106, f. 300, 300-
 303
 Sordaria invaded by, f.
 296
 Spot disease of currant, 281
 Orchard Grass, 282, f. 283
 Strawberry, 281
 Spotted St. John's-wort, 630
 Sp. go. f. 297, 597-598
 Large, 597, f. 600
 Touch-me-not, 123, 620
 Spout, 911
 Spreading Dogbane, 129, 692
 Nightshade, f. 721
Sprecheria formosissima 366,
 805
 Sprech, C. H. E. 912
 Sprouting Grass Smut f. 219
 Sreuce, 868
 Black 328
 Newey 317, 328
 Thirland, 327
 White, 328
 Spurge 597-603, 868
 Carter, 121 f. 300, 603
 Common, 602
 Cypress, 121, 600 f. 601
 Family M9, 586-604
 Flowering, 121, 122, 598,
 f. 598
 Yellow, f. 602
 Laurel, 123, 642, f. 643
 Myrtle, 121, 599
 Nerf, 122, 596, f. 606
 Spotted, f. 597, 597-598
 Large 597, f. 600
 Spurrey, 436
 Squash Bush, 750
 Crockneck, 750
 Hubbard 751
 Winter 750
 Squaw-weed, 138
 Squill 148, 373, 377
 Squirrel-nail Grass, 66, 103,
 366-367, f. 368
 Powdery Mistletoe, 272
 Insults from, 360
 Squinting Cucumber 749
Stachys arvensis, 131, 828
palustris 828
Schneider 710
Stachytarpheta indica 858
 Stadlinger, 482
 Staff-vec, 614-615
 Family 6 4-615
 Stagger-bush, 126, 674, f. 675,
 675
 Stagers, 100, 366
 Mad, 259
 Stomach, 261
 Staggerwort 140

- Staghorn Sumac. 608
 Stahl, A., 3
 Stalagmitis Mangie 827
 Stalberg, S., 912
 Stalker, M., 9 118, 187 277,
 356, 544, 564 912
 Stange, C. H. 24-28
 Stapelia, 84, 695, 810
 Stapf O. 912
 Staphisagrin, 108, 464, 868
 Staphylea trifolia, 604
 Staphyleaceae, 604
 Staphyloselin, 169
 Lysocaulin, 168
 Staphylococcus pyogenes
 aureus, 168
 Star Anise, 149, 648, 863, 868
 Apple, 679, 681
 Cucumber, 751
 of Bethlehem, 377
 Stercia, Cinnam., 367
 Tacca, 374
 Ymn 374
 Strieck brachymeris, 673
 pectinata, 128 842
 Statutes on Poisoning, Iowa
 code, 6
 Stenocera, 446, 460, 463
 Seed, 464
 Stearic acid, 593
 Stearin, 594
 Stearns, Fred, 468
 Stebler, 96, 335, 439, 449
 360
 Stellaria, 436
 crassifolia, 413
 media, 07 442-444, / 446
 868
 Steloneuma, 676
 imbricatum, 676
 Stemless Loos Weed, 39, 119,
 566, 567 570, / 368
 Stemonitis, 159, 160
 Stephanis, 89
 ornata, 838
 discolor, 838
 Sterculia alata, 855
 Sterculiaceae, 33, 621 853
 Stercoraria ramulosum, /
 707
 Sterigmata, 21
 Sterigmatocystis, 258
 Stevens, A. B., 912, 915
 Stevens, Y. L., 206
 Stewart, P. C., 126, 210, 663
 Stielmeier, 131 706-707
 Large-flowered, 707
 Stick-slight, 786
 Stillaria, 390
 linearis, 823
 epicalyx, 604, 824
 Stillmark 395
 Stimulant, 804
 Stinging Nettle, Small, /
 423 412-413
 Nut, Brazilian, 596
 Hida, 124
 Stinkhorn, 99, 239, 245
 Stinking Clover, 114
 Smut of Wheat, / 219-220
 Willie, 797
 Sipa, 339, 353, 578, 579
 anemone, 66, 354
 capillata, 354, 926
 comata, 66, 70, 102, / 355,
 355-357
 hystrixina, 33
 mabiana, 354, 355, 826
 ussuriensis, 90
 Vesilana, 354
 palmata, 876
 pinnata, 826
 robusta, 90, 102, 357 358,
 826
 rubra, 354
 sparsa, 70, 102, 210, 354,
 355, / 356
 tenuicarpa, / 356
 Western, 355, / 355
 Stipe, 160
 Stickswort, 436
 Striobolus Lyonsi, 536
 Strick, 111
 Strimthia, 179, 272, 290, 642,
 649, 868
 Camerhal, 29
 Mycone, 256, / 259, 260,
 262, 263
 Strickberger, W. W., 680, 912
 Stokes, J. S., 912
 Stone Clover, 553
 Stone, C. E., 613
 Stonerog, 114, 221, 501
 Mossy, 501 502
 Stonewort, 93, 194, 688
 Stora, 506
 Storein, 500
 Strickhill, 67 120, 573-579
 Henslock, / 579
 Stowe, W., 912
 Strabo, 64
 Stratonium, 72, 718
 Poisoning by, 6, 78
 Purple, 729
 Strangles, 370
 Stranburger, E., 82, 893, 901,
 912
 Strathaird, R., 912
 Strawberry, 504-505, 629, 868
 Blite, 474
 Bush, 473
 Chilean, 505
 Cultivated, 115, 609
 European, 115
 Wood, 509
 Garden, 305
 Indian, 505, 509
 Rust, 281
 Tomato, 715
 Tree, 645
 Virginia, Wild, 115, 509
 Streptococcus Septicemia, 171
 Streptococcus melanogenes,
 28, 868
 equi, 170
 erysipalae, 171
 pyogenes, 87 171
 Streptococcus, 171
 Strickland, C. W. 912
 Striga capricornis, 852
 String Bean, 520
 Streptanthus callosus, 698,
 807
 Strong, A. B., 912
 Strophanthus, 692
 Strophanthus, 78, 868
 Emmit, 807
 Aspidus, 148, 692, 867
 Kombe, 692
 Pierre, 807
 Strophantia, 148
 Stropharia aeruginosa, 860
 Strampell, Dr. 522
 Strychnia, Statutes regard-
 ing, 6
 Strychnin, 1, 72, 73, 82 129,
 144, 145, 146, 147, 628,
 684, 688, 728, 729, 736,
 868
 Antidotes against, 520
 Discovery of, 3
 Krate, 688
 Poisoning by, 6, 77
 Sulphate, 688
 Toxicity of, 7
 Treatment for poisoning
 by, 81
 Strychnos, 52, 685, 868
 angustifolia, 836
 brachyloca, 836
 Caribbaea, 148, 836
 cogens, 836
 colubrina, 836
 Cerebraria, / 685, 686,
 836
 pudica, 836
 Iraya, 836
 Iguala, 74, 144, 688
 innocua, 836
 malacensis, 688, 836
 Melanocarpa, 836
 ana-nomura, 74, 85, 147,
 685, 688, 836
 paniculata, 836
 polatorum, 685, 836
 Pseudo-quina, 836
 Samhara, 836
 spinosa, 836
 rosifera, 1 148, 687, 836
 vopurata, 836
 Stryphnodendron Barbati-
 man, 835
 polyphyllum, 833
 Strychnos, 629
 Strubleberry, 719
 Studer, 241
 Stuhl, W. A., 17-19, 20-24,
 28-29, 37, 39-40, 393
 Sturdevant, L. B., 912
 Stylophorum, 868
 dichyllum, 113, 480, 841
 Stylopis, 113, 480
 Styphandra glauca, 831
 Styple, 396
 STYRACACEAE, 679, 682-683
 Styacin, 683
 Styrax Benzoin, 682, 683
 Family, 682
 Styrax fruticosa, 424
 Suture, 411
 Succory, 758-759
 Sudworth, G. R., 332
 Sugar Beet, 107, 424-427, 868
 Cane, 338
 Maple, 615, / 616
 Pine, 529
 Suicide by Poisoning, 3, 868
 Sullivan's Honeysuckle, 744
 Sulpho-cyanate of a. yl, 489
 Sulphur Polyposis, 254
 Sulphur dioxide, 72
 Sulphuric acid, Poisoning by,
 6
 Same, 127, 608-610, 868
 Chinese, 583, / 584
 European, 608
 Japanese, 608
 Poison, 609, / 610
 Smooth, 608, 613-614
 Stag-horn, 608
 Summer Chrysanthemum, 756
 Cypress, 107
 Sundew, 77, 14 497
 Condon, / 90
 Family, 86 114, 497-498
 Sunflower, 756
 Great, 756
 Maximilian's, / 763
 Powdery Mildew of, 269
 Superban, 377
 Supple-jack, 620
 Surgeon General of U. S.
 Army, 892
 Swinam, Poisonous plants of,
 867
 Susotoxin, 150, 171
 Suxum anthelminticum, 821
 Sutton, R. L., 912
 Swain, R. E., 888, 912

- Swainsona**, 534
coroniliifolia, 533, 833
Greviana, 533, 833
Quiveri, 833
pharoides, 833
procumbens, 833
Swamp Canas, 103
Hellebore, 103
Laurel, 672
Leucotoxoe, f. 673 673
Lousewort, 738
M. Jeweed, 130
Swedish Tansper, 102, f. 337, 332
Turnip, 486
Swartia triphylla, 833
Sweet Alyssum, 486
Balm, 397
Basil, 709
Bay, 474
Birch, Oil of, 86
Cinnamon, 371
Cassava, 54, 587
Cherry, 514
Cleary, 648
Clover, 552
White, 1 & f. 44, 552
Yellow, 116, 852
Fern, 399
Flam., f. 308
Gale, f. 308
Family, 397
Hay, 108, 389
Potato, 698
William, 437
Sweet-scented Bedstraw, 744
Shrub, 476
Sweetbriar, 115
Sweetia chinensis, 689
Swalethia humilis, 837
Mahagoni, 374 837
Switzerland, Poisonous plants of 867
Sword Bean, 521
Eucalypta, 500
Syraxia parmaria, 301
Sylvestri, 604
Syme, W. A., 612, 613, 869, 912
Symphoricarpos, 157 743, 747, 748
mollis, 135, 813
occidentalis, 744, 747
orbiculatus, 135 744, 748
racemosa, 135, 744 748, 813
Symphitum officinale, 704
Symplocarpus foetidus, 83, 371, 808
Symplocos Aitonii, 629
laureolata, 6 4
racemosa, 681
Symptoms of Poisoning, 73
Synaktonia, 438
Synalissa ramulosa, f. 307
symphorea, f. 309
SYNANTHAE, 459
Synanthus, 505
Syncephalus cornu, 196
intermedia, 96
Synchytium, 204
decipiens, 204
SYNPHETALAE, 157
Synura, 188
Syphus, 182
Syringa, 89, 115
persica, 683
salicaria, 63, 683
Syringia, 684
Syringopieris, 128
Sisymbrium, 486, 487-488
Althia, 618
altissimum, 1.4, f. 487, 488
officinale, 114, 487, 818
Sophia, 818
torophyllum, 818

T

Tabernaemontana Barbolica, 807
nitrofolia, 807
coronaria, 807
maltacensis, 807
mauritiana, 807
Tabernaemontana, 807
Tacca plantifolia, 374
Starch, 374
Taccacium, 374
Tachia guianensis, 825
Tachia, 53, 633
Tachia serrata, 568
Tachifuge, 884
Tagetes erecta, 736
Takonia, 168
Talamus, 99
macrocarpa, 474, 837
ovata, 837
Talinum polyandrum, 844
Talinia stricta, 850
Tall Blackberry, 845
Buttercup, 10, f. 49
Cape Flower, 729
Crowfoot, 458
Gynaphyl, 436
Larkspur, 108, 462, 463, f. 46, 465, 466, 467
Meadow Owl Grass, f. 500
Ragweed, f. 766
Tallow Tree, 587
Tamaricaceae, 627
Tamarind, 529 f. 587
Tamarindus indica, 529, f. 531
Tamarix gallica, 627
manifera, 627
Tampkin, 702
Tamus remanens, 820
Tanacetum, 756
Tanacetum, 790
Tanacetum, 757 758, 790, 790
baianum, 789
umbelliferum, 816
swigera, 756, 789 790, 793, 816
Tanacetum erucigerum, 810
Tanacetum, 364
Tangerine, 582
Tangetin, 868
Tanghinia venenifera, 148
Tangier Pen, 526
Tannic acid, 144, 403, 583, 608
Tanin, 105, 319, 327 403, 418 608
Tanret, Ch., 277, 278
Tanay, 789-790
Common, 140
Oil, 756, 789-790
Tarweed, 421
Tape-worm, Presence in food sheep, 568
Tephrosia, 252, 253
Tapioca, 75, 587
Tapioca, 2, 621
guianensis, 819
Tar weed, 138, 755
Taraxacum Blumei, 53
Taraxacum, 755
Taraxacum, 755
Taraxacum, 757
officinale, 755, 756, f. 729
Taraxacum, 756
Tartaric acid, 529, 868
TAXACEAE, 101
Taxia, 101, 148, 328
Tarax, 32, 148, 328
barcata, 4, 328, 817, 868
brevisolia, 101, 328, 817
canadensis, 101 328, 817
Taylor, A. S., 4, 363, 913
Taylor, H. H., 905, 913
Taylor, J. L., 130
Taylor, J. Q., 703
Taylor, M. K., 357
Taylor, Thomas, 913
Tea, 628, 868
Black, 628
Chinese, 614
Green, 628
Labrador, 127, 629, 646
Mexican, 107
New Jersey, 621, 629
Oswego, 109, 639
Plant, 628, 629
Substitutes for, 436, 629
Wild, 501
Teak Tree, Indian, 708
Teasel, f. 747
Putera, 741
Tecoma, 32
ceramensis, 810
caminioides, 739
leucocorydon, 810
obtusata, 810
radicans, 135, 739, 810, 862
aperiosa, 810
stans, 810
torophora, 810
Tectaria grandis, 708
Telluriz, 573
Telfairia pedata, 751, 819
Temperature affecting toxicity of poison, 868
Templetonia, 868
rasa, 835
purpurea, 833
rostrata, 833
Tenbouch, J., 913
Tennessee, Poisonous plants of, 866
Tenonite, 210
Rust on, 230
Tephrosia, 558
Tephrosia, 52 535, 558
astagaloides, 833
candida, 833
cimaria, 833
coroniliifolia, 833
densiflora, 833
fruticosa, 833
macrocarpa, 833
nitens, 833
periclyon, 833
purpurea, 834
tomentosa, 833
toricaria, 118 533, 558, 833
turgida, 118, 558, 833
torula, 833
Tephrosia, 533, 558
Tetanus labialis, 833
Terminalia Battersi, 814
Cataphe, 638, 640
Chebulis, 640 814
lagifolia, 640
tomentosa, 814
TERNSTRÖMIACEAE, 52 89, 855
Ternier, T. J., 913
Tetularia, 210
Tetania, 10, 150
Tetanoid spasms a symptom of deliriant poisoning, 73
Tetanolysin, 174
Tetanusparvum, 174
Tetanusparvum, 150
Tetanus, 150, 174, 868
Bacillus, 174
Toxin of, 77, 174

- [illegible]

- Urocytis*, 218
agropyri, 219
occulta, f. 218, 219
Uroglana, 168
Uromyces, 230-231
appendiculatus, 230
Fobae, f. 221
plat., 230
striatus, 230
Trifolii, 99, 230-231, f. 231, 861
Ursone, 665
Urtica, 410, 411-413
dioica, 106, 413, f. 413, 857
gracilis, 106, 413, 857
holosericea, 413, 857
membranacea, 413
pilulifera, 412, 867
spatula, 412
urens, f. 412, 412-413, 857
URTICACEAE, 53, 106, 405, 409, 415, 857
 Fish poisons in, 2
URTICALES, 136, 404-415
Urticaria, 12, 78, 412, 419
Urticarial poisoning, 804
Urticating hairs, *Hugloss*, f. 77
Nettle, f. 77
Sedge, f. 77
Grass, f. 77
UTILAGINACEAE, 98, 210-219, 861
USTILAGINEAE, 217, 364
Ustilago, 210, 868
avenae, 98, 213-217, f. 216, 861
bromivora, 210, 215, f. 218
Cenasti, f. 218
Cynamom, f. 214
esculenta, 210
Flacheri, f. 214
Hordei, 98, f. 213, 217, 861
hypodites, f. 216, 217
mauris, 217
minima, 210
neglecta, 98, 211, f. 218, 861
nuña, 98, 217, 861
panicl-glauci, 217
panicl-glauci mollisco, f. 214, 218
perennans, 216, f. 218
segetum, 217
Sorgi, f. 212
strigiformis, f. 218
Tricli, 98, 217, 861
urticulosa, 98, 219, 861
seae, 98, 210-215, f. 213, 212
Utricularia vulgaris, 898

V

- Vaccaria vulgaris*, 108
Vaccinium, 222, 829, 665
oxiloreum, 665
caespitosum, 665
canadense, 665
corymbosum, 665
v. pallidum, 665
macrocarpon, 665
membranaceum, 665
Myrtillus, 629
Oxycoccus, 665, 666
pennsylvanicum, 665
stamineum, 665
vacillans, 665
Vaginaria, 179
Vahlen, E., 914
Valentin, G. C., 914
Valeriana dioica, 857
officinalis, 857
VALERIANACEAE, 857

- Vallisneria spiralis*, 332
Van de Warker, E., 790
Van der Velde, 168
Van Dremmer, H., 663
Van Es, L., 914, 915
Van Hasselt, 601
Vandellia crustacea, 852
minuta, 852
Vangueria spinosa, 849
Vanilla, 392, 773
Grass, f. 341
guianensis, 392
palmarum, 392
planifolia, 392
plant, 772-773
Pompona, 392
Vanillin, 392, 683
Vanquelin, 734
Vapora, Poisoning by, 77
Variolevis amara, 861
Varnell, Prof., 298
Vasry, Geo., 357, 914
Vater, A., 914
Vateria, 627
Vaucheria, 191
Vaughan, V. C., 150, 364, 573, 888, 897, 914
Vaulabelle, A. de, 914
Vegetable, *Beefsteak*, 234
Butter, 681
Calomet, 371
Poisons, 149
Velleia paradoxa, 825
Velvet-leaf, 124, 627
Venus Fly-trap, 498
Veratrin, 379
Veratramarin, 381
Veratridin, 381
Veratrin, 46, 77, 81, 82, 377, 381
Veratrum, 44, 46
Veratrum, 46, 52, 148, 176, 377, 379, 380-383, 835, 848
album, 4, 73, 148, 381, 835
alkaloids of, 148, 381
Arrow poisons, 2
californicum, 104, 381-382, 835
Isobellianum, 381
viride, f. 45, 75, 85, 103, 148, 381, f. 382, 835
Wood, 103
Verbascum, 53, 734, 868
Blattaria, 734-735, 852
crassifolium, 852
dubium, 852
orientale, 852
phlamoidea, 852
pulverulentum, 852
simplex, 852
sinuatum, 852
thapsoides, 852
Thapsus, 134, 733, 734-735, f. 735, 852
Verbena, 708, 868
Ambetia, 708
Family, 131, 707-708
hirsuta, 708, 858
officinalis, 858
stricta, 708
articulifolia, 708
venosa, 858
VERONACEAE, 131, 707-708, 857-858
Verdier, 874, 882, 914
Vermifuge, 804
Vermont, Economic plants of, 864
Vernonia, 757
anthelminticum, 816
nigrifolia, 816
novboracensis, 222
VERNONIEAE, 757

- Veronica*, 734
Beccabunga, 853
officinalis, 853
pergrina, 735-736
Virginica, 134, 733, 736, 853
Veronal Poisoning, 868
Verachaffelt, E., 914
VERTICILLATAE, 156, 395
Verticillium, 24
graphi, 199
Vervain, Blue, 708
 White, 708
Vesicant, 804
Vericaria gnaphalodes, 818
Vestia lycoides, 855
Vetch, 45, 120, 525, 526
 American, 570
 Black-purple, 526
 Common, 120, 570-572, f. 371
 European, 87
 German, 572
 Hairy, 527, 570
 Milk, 562-567
 Red, 572
 Scarlet, 526
 Spanish, 572
 Wild, 87
 Woolly-pod, 526
Vetch-Bread, 573
Vetchling, 572-573
 Marsh, 572
 Prairie, 572
Viburnum Lantana, 813
macrophyllum, 813
Opulus, 813
Vicat, P. R., 914
Vicia, 570-572
americana, 570
atrovirens, 526-531
dasycarpa, 527
Ervilia, 813
Faba, 522, 523, 572
fulgens, 526
hirsuta, 527
sativa, 53, 120, 269, 525, 570, f. 377, 572
villosa, 525, 570
Vicin, 120, 572
Victoria regia, 445
Vigna Catjang, 520, f. 532
Villamilia peruviana, 434
Villarsia Moorei, 839
Villiers, A., 914
Villosa, 508
Vinca major, 691, 807
minor, 691, 807
puzilla, 807
Vincetoxicum, 695
nigrum, 810
officinalis, 696, 810
Vinegar, Acetic acid in, 163
Tarragon in, 756
Viola, 631-632, 858
blanda, 631
canadensis, 631
cucullata, 125
Natellii, 631
odorata, 125, 631, 858
pedata, 631, 858
pedatifida, 631
pubescens, 631, f. 632
sepicola, 858
striata, 631
tricolor, 631, f. 632, 632-633
VIOLACEAE, 125, 417, 628, 631-633, 858
Violaguericrin, 632
Violet, Bird-foot, 631
 Blue, Common, 125
 Canada, 631
 Dog's-tooth, 104

European, 83
Fragrant, 631
Pansy, 632
Sweet, 125, 632
White-flowered, 631
Wood Sorrel, 580
Yellow, 631, f. 632
Violin, 632, 633
Viper's Bugloss, 131
Virebaw, R., 231, 263
Virginia Creeper, 124, 150, 620
Peppergrass, 72
Snakeroot, 417
Strawberry, Wild, 115, 509
Virgin's Bower, 109, 446, 456, f. 457
Pisum album, 106, 836
seedlings, 595
Pisum viridiflorum, 629
VITACEAE, 124, 620, 858
Vitis Heterophylla, 708
Nagundo, 708
pteropoda, 858
vitalis, 858
Vitis elongata, 858
Minahau, 858
Sapunaria, 858
syssylifolia, 858
Vitrol, Oil of, 143
Voandzeia subterranea, 521
Vogel, 3
Volken, G., 915
VOLVOCAEAE, 98, 191-193
Volturna bombycina, 243-244
volucrea, 860
Volvox, 93, 98, 191, 193
globular, 193
minor, 193
Vullemis, P., 290, 299
Vuillet, Jean, 895

W

Wahoo, 123, 148, 614, 615, f. 615
Wager, H., 306
Wainwright, J. W., 915
Wake-robin, f. 371, 385
Erect, 104
Large-flowered, 385
Prairie, 385
Scalloped-flowered, 385
White-flowered, Large, 385
Waldron, Dr. C. B., 56, 560, 561
Waldron, L. R., 915
Wall-flower, 486
Waleff, B., 695, 915
Wallace's Farmer, 915
Wallroth, K. F. W., 13
Walnut, Black, f. 400, 401
California, 401
English, 401
Japanese, 401
Walsch, J. I., 295
Walsh, L. H., 915
Walshia amorphella, 569
Walsura, 52, 89
pisidic, 837
Wandering Jew, 374
Warburton, C. W., 274, 346, 915
Warder, H., 903, 915
Warren, L. E., 805-806, 915
Washington, Flora of, 865
Washingtonia, 370
Palm, 370
Water, Algae in, 91-95
Avena, 505
Cherry-laurel, 503
Chestnut, 640
Chinquapin, 444
Cress, 496

Crowfoot, 446
Dock, Great, 420
Dropwort, 649
Hemlock, 126, 648, 651-659
Lily, 444-445
Blue-flowered, 445
Yellow, f. 445
Moulds, f. 208
Parasit, Cut-leaved, 76
Creeping, f. 650
Pepper, 422-423
Wild, 422-423, f. 423
Plantain, 102, 335, f. 335
Family, 335
Pollution of, 91
Smartweed, 421-422
Water as source of disease, 8-10
Water-bloom, 93, 186, 868
Waterleaf Family, 131, 703, 704
Watermelon, 750, f. 748
Water-weed, 332, 445
Watkins, H. C., 909
Watkins, W. H., 915
Watson, E. B., 403
Watson, Miss C. M., 564
Watson, Sorens, 342, 564, 915
Wattles, Australian, 523
Way, 915
Weatherby, C. A., 915
Weather-glass, Poor-man's, 128
Shepherd's, 128
Webb, Paul C., 915
Webber, H. J., 287, 582, 625
Wechsberg, 168
Weeds, Poisonous, 868
Weems, J. B., 903
Weevers, 82
Weichardt, 482, 767
Wellcome Laboratory, 4
Wellman, F. C., 915
Welwitschia, 326
Wenzell, 277
Weschke, Carl, 915
West India Arrow-root, 77, 391
Cedar, 575
Gherkin, 750
Locust Tree, 527
Westbrook, F. E., 915
Western, Aconite, 450, 452, 453, f. 452
Cleome, 114
Cone-flower, 779
Dogwood, 664
Honeysuckle, 744
Larkspur, Purple, 109
Lupine, 118
Needle Grass, f. 66, 102, 355, f. 355
Nettle, 106
Snakeroot, 773-774
Yellow Pine, 330
Westgate, J. M., 915
Weitsstein, R. von, 915
Wheat, 225-338
Bunt, f. 270, 219-220
Cow, 134, 734
Egyptian, 345
Grass, 274, 364
Slender, 364
Western, 364, 365
India, 419
Rust, 99, 227, f. 228, 228
Smut, 217
Stinking, 98, 219, 220, f. 219
Whelpley, H. M., 915
Whipple, G. C., 93, 189, 915
White, E. A., 915
White, G. R., 916
White, J. C., 96, 104, 112,

134, 215, 317, 455, 456, 464, 471, 485, 490, 503, 583, 587, 600, 601, 608, 609-612, 634, 679, 739, 794, 803, 804, 916
White, R. B., 417
White, Amanita, 34
Ash, 683
Aster, Small, 776
Banberry, 108
Birch, 464
Cedar, 327, 328
Clover, 118, 525, 553, 557, f. 557
Sweet, 118, f. 357
Gentian, 690
Hellebore, 73, 163, 381
Indigo, Wild, 117, 541, f. 347
Jasmine, Large Flowered, f. 684
Lady's Slipper, Small, 105, 393
Lupine, European, 547
Mulberry, 406
Mustard, 77, 486, 488, f. 489, 490
Oak, 403
Pepper, 87
Pine, 329, f. 329
Rust, f. 304
Sage, 424
Snakeroot, 771
Thorn, 513
Vervain, 708
White-flowered Wake-robin, Large, 385
White-wood, 138, 756, 770-779, f. 774
Widal, P., 263, 916
Wigand, Albert, 916
Wikstroemia Chamaedaphne, 855
viridiflora, 855
Wilcox, E. M., 100
Wilcox, R. V., 5, 44, 84, 96, 97, 109, 378, 379, 383, 462, 466, 548, 568, 626, 656, 762, 793, 795, 916
Wittek, R. W., 903, 916
Wilson, T. E., 916
Wild Apple, Balsam, 751
Apple, Crab, 512
Barely, 66, 67, 103, 338, 368, f. 368
Carrot, 663-664
Cherry, Black, 64, 117, 503, 505, f. 315, 515-516
Red, 116, f. 304, 803, 516-519, f. 316
Western, 515
Cinnamon, 627
Coffee, 745
Comfrey, 131, 706
Cucumber, 751
Ginger, f. 415, 416
Grape, Blue, 670
Heliotrope, 131
Hydrangea, 115, 500
Indigo, Large-bracted, 541
Large White, 117, 541, f. 541
Ipecac, 599, f. 600
Leek, 104, 383
Lettuce, 761, 762
Licorice, 531
Lupine, 546
Monkshood, 109, 453
Oats, 102, 215, f. 358, 359
Olive, 641
Onion, 104
Pea, 525
Pineapple, 373

Plum, northern, 116, f. 506
 Potato, 701
 Rice, 276, 338
 Rosemary, 127, 673
 Rye, 27, 90, 149, 276, 280, 346
 Sago, 378
 Senna, f. 337
 Tea, 501
 Tobacco, 729
 Veitch, 87
 Water Parsnip, 49, 650, 656, 661-663, f. 661
 Yam, 374, 688
 Wildeman, K. de, 916
 Wiley, H. W., 916
 Williams, Dr., 10
 Williams, 917
 Williams, A. J., 916
 Williams, S. W., 916
 Williams, T. A., 348, 357, 916
 Williams, W. L., 259, 916
 Williams, W. T., 916
 Willis, G. S. V., 917
 Willoughby, W. C., 917
 Willow, f. 308, 629
 Family, 396-397
 Peach-leaved, f. 397
 Willow-herb, 629
 Great, 644
 Wilmer, B., 917
 Wilson, Dr., 372
 Wilson, Dr., 372
 Wilson, F. W., 23, 94, 130, 694, 917
 Wilson, James, 878, 917
 Wilson, J. C., 917
 Wilson, P., 917
 Will, Corn, 163
 Wind-flower, 109, f. 456, 455-456
 Wine Palm, 376
 Wineberry, 303
 Winegate, J. B., 917
 Winkler, L., 917
 Winnebago Lake, 93
 Winslow, Kenelm, 83, 96, 109, 231, 278, 381, 450, 471, 482, 518, 575, 593, 628, 687, 689, 726-727, 728, 731, 736, 742, 917
 Wintergreen, 621, 665
 Oil of, 404, 621, 665
 Winter's-bark, 474
 Winton, A. L., 917
 Winton Disease, 795
 Wire Grass, 374
Wistaria, 64, 527, 534, 629
chinensis, 527, 534, 629, 833
perfoliata, 527
 Wisteria, 534
 Witch Hazel, 586
 "Witches Broom" on Birch, 253
 on Cherry, 253
 Withania somnifera, 855
 Withania, R. A., 872, 917
 Wittstein, C. C., 917
 Woehler, 585
 Wokas, 444
 Wolfberry, 744
 Wolfenstein, A., 767
 Wolfenstein, R., 904, 917
 Wolfner, Dr., 199
 Wolfgang, 482
 Wolf's-bane, Trailing, 109
 Wood, 276, 503
 Wood, Alphonso, 917
 Wood, H. C., 917
 Wood, Anemone, 77, 109
 Nettle, 77, 106, 413, 414
 Sorrel, 120, 580
 Family, 120

Strawberry, European, 509
 Woods, Dr. A. P., 348
 Woodville, Wm., 917
 Woody, Aster, 138, 774
 Woolly Loco Weed, 39, 562, 567, 568
 Thistle, f. 308, 801
 Woolly-pod Vetch, 526
 Wooden Grape, 620
 Wyoming Larkspur, 189
 Wormseed, 187, 428, f. 426
 Oil of, 187, 428
 Levant, Oil of, 791
 Spanish, 424
 Wormwood, 74, 754, 755, 756, 790, 791-794, f. 791
 Biennial, 140, f. 792
 Common, 140, 793
 Wright, 168
 Wright, J. S., 917
 Wright, R., 917
 Wright's Datura, 133, 712-713
 Wyman, Dr. Walter, 343
 Wynham, R., 881

X

Xanthin, 147
 Bases, 146
 Derivatives, 146
 Di-methyl, 146
 Methyl, 146
Xanthisma texana, 138, 816
Xanthium, 757, 767-770
canadense, 67, f. 68, 137, 762-770, f. 309, 816
spinosum, 137, 767, f. 766, 816
strumarium, 137, 816
Xanthopuccin, 108, 468
Xanthoxstrumarin, 137, 769
Xanthorrhoea, 358
hastilis, 375
 Xenophon, 64
Xeranthemum annuum, 53
Xerophylla Douglasii, 833
Ximenia americana, 53
Xyba dolabriformis, 833
Xylometum pyriforme, 844
Xylopi odoratissima, 806
polycarpa, 806
salicifolia, 806
Xylorhiza Parryi, 776
 XYRIDACEAE, 372, 858
Xyris communis, 858

Y

Yagna, 370
 Yam Bean, 521
 Chinese, 699
 Family, 374
 Japanese, 374
 Starch, 374
 Wild, 374, 688
 Yarrow, 149, 756, 785-787, f. 786, f. 787
 Yampao, 123
 Yeast, f. 249, 250-251
 Chinese, 198
 Common, 250-251, f. 250, f. 251
 Parasitic, 249
 Pink, 250
 Yellow Birch, 404, f. 405
 Cleome, 496
 Clover, Hop, 552
 Sweet, 118, 552
 Flax, Large-flowered, 120, 581
 Flowering Spurge, f. 601
 Jasmine, 688-689
 Lady's Slipper, Small, 105, f. 393

Pine, Western, 330
 Puccoon, 446, 468
 Violet, 631, f. 631
 Water Lily, f. 445
 Yellow-eyed Grass, 372
 Yellow-flowered Alfalfa, 526
 Bitter-weed, 539
 Nightshade, 692
 Yellow-leaf disease of barley, 281, f. 381
 Verba, 614
 Maté, 614, 742
 Santa, 703
 Yew, 4, 72, 101, 146, 326, 328, 668
 American, 101, 328
 European, 101, 328
 Yopon, 742
 York-road poison, 511
 Yucca, 89, 375
angustifolia, 375
filamentosa, 375, 835
glauca, 63, 835
imperialis, 375
 Yuld, 380
 Yulan Tree, 474

Z

Zaluzha, 729
 Zamang, 527
 Zamia, 326
 media, 819
Zanthoxylum, 89
alatum, 120, 849
americanum, 581
caribaeum, 849
Clava-Herculis, 581
Narongillo, 849
scandens, 120, 849
Zea amyloacea, 341
amyloaccharata, 341
canina, 342
erecta, 341
indurata, 341
indurata, 341
Maya, 102, f. 317, 826
 History of, 340-342
saccharata, 341
tunicata, 341
 Zedoury, 392
 Zehl, L., 917
Zenobia speciosa, 821
Zephyranthes, 386-388
atamazeo, 105, 387-388, f. 388, 803
 Ziegler, Ernst, 304
 Zinc, Sulphate of, 6
Zingiber officinale, 77, f. 390, 391, 838
 ZINGIBERACEAE, 391, 838
Zinnia, 756
elegans, 756
Zinnia, Ergot on, 276
latifolia, 210
Zizyphus, 2, 89, 621
Joazeiro, 847
Jujuba, 621
Lotus, 621, 847
sativa, 621, 847
 Zoethout, W., 9, 918
Zoidiogramma, 154
 Zooglossa, 161
 Zoon, J., 917
 Zopf, W., 918
 Zwaluwenberg, A. V., 909, 918
Zygadenus, 377-379
angustifolium, 835
chloranthus, 378-379
elegans, 6, 103, 835
paniculatus, 378
 smooth, 378-379

INDEX

977

венерическая, 6, 103, f. 376,
377-378, 379, 435
Зудящая, f. 92, 189, 190, f.
191
ЗУОНИМАТАСКИ, 190

ZYGOMYCETES, 154, 195,
204
ZYGOHYLLACEAE, 89, 573,
858
Zygothidium coccineum, 858
Fubago, 858

iodocarpum, 858
Zygorhynchus, 193
Zygoxypores, 13
Zylio, ■
Zymase, 250